REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188
Public reporting burgen for this collection of im gathering and maintaining the data needed, an collection of information, including suggestions Davis Highway, Suite 1204, Arlington, VA 22202	formation is estimated to average 1 hour p d completing and reviewing the collection of for reducing this burden, to Washington h 4302 and to the Office of Management al	er response, including the time for r of information. Send comments requested and the second	eviewing instructions, searching existing data s rrding this burden estimate or any other aspect information Operations and Reports, 1215 Je ject (0704-0188), Washington, DC 20503.
1. AGENCY USE ONLY (Leave blar	k) 2. REPORT DATE	3. REPORT TYPE AN	D DATES COVERED
	May 15, 1996	Final Report	01 Jan 91 - 31 Dec 3
4. TITLE AND SUBTITLE Numerical Modelling of Layer	the Continental Shel	f Bottom Boundary	
6. AUTHOR(S) Hsiao-ming Hus and Robe	ert C. Beardsley		
7. PERFORMING ORGANIZATION N	AME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER
Woods Hole Oceanograph Department of Physical 360 Woods Hole Road Woods Hole, MA 02543-19	Oceanography		
9. SPONSORING/MONITORING AG Department of the Navy Office of Naval Researc Resident Representative 496 Summer St., Room 10	ch e	ES)	10. SPONSORING / MONITORING AGENCY REPORT NUMBER
12a. DISTRIBUTION / AVAILABILITY Approved for publi distribution is unlin	c release;		12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 word	de)		
See attached final rep		40	
See attached final rep	9960703 0		15. NUMBER OF PAG 16. PRICE CODE

, N Final Report for ONR Grant N00014-91-J-1473, Drs. Hsiao-ming Hsu and Robert C. Beardsley, Principal Investigators

The primary objective of this project was to examine the nature of wind-driven upwelling and downwelling along a 2-dim shelf with sloping bottom topography. The approach was to use the Semi-spectral Primitive Equation Model (SPEM) developed by D. Haidvogel and coworkers as the basic circulation model, and conduct a series of idealized model runs to investigate the influence of bottom slope, surface windstress and heat flux, and mixing on the surface and bottom boundary layers which form during the initial stages of up/downwelling. The advanced turbulent closure schemes of Mellor and Yamada were added to our version of SPEM and the model studies were conducted. These studies showed that an asymmetry develops in the bottom boundary layer thickness depending of if the flow is up or downwelling, and that if the coast is shallow enough, that the surface and bottom boundary layers can merge near the coast. Two AGU abstracts were presented and several talks were given on this work. A paper co-authored with A. Herman on the technical aspects of the vertical model resolution was also published.

- Hermann, A.J. and H.-M. Hsu, 1989. A semispectral ocean circulation model with mixed layer physics; Part 2: the value of stretched coordinates. *EOS, Transactions, American Geophysical Union*, fall meeting.
- Hsu, H-m., Robert-C. Beardsley, and J. F. Price, 1992. Including surface and bottom boundary layer physics in a primitive-equation model to study coastal circulations. *EOS*, *Transactions, American Geophysical Union*, **72**(52), 59 (abstract).

Hermann, A.J. and H.-M. Hsu, 1993. A vertical coordinate mapping technique for semispectral primitive equation models of oceanic circulation. J. Atmos. Ocean Tech., 10(3), 381-396.

Final Report for ONR Grant N00014-91-J-1473, Drs. Hsiao-ming Hsu and Robert C. Beardsley, Frincipal Investigators

The primary objective of this project was to examine the nature of wind-driven upwelling and downwelling along a 2-dim shelf with sloping bottom topography. The approach was to use the Semi-spectral Primitive Equation Model (SPEM) developed by D. Haidvogel and coworkers as the basic circulation model, and conduct a series of idealized model runs to investigate the influence of bottom slope, surface windstress and heat flux, and mixing on the surface and bottom boundary layers which form during the initial stages of up/downwelling. The advanced turbulent closure schemes of Mellor and Yamada were added to our version of SPEM and the model studies were conducted. These studies showed that an asymmetry develops in the bottom boundary layer thickness depending of if the flow is up or downwelling, and that if the coast is shallow enough, that the surface and bottom boundary layers can merge near the coast. Two AGU abstracts were presented and several talks were given on this work. A paper co-authored with A. Herman on the technical aspects of the vertical model resolution was also published.

- Hermann, A.J. and H.-M. Hsu, 1989. A semispectral ocean circulation model with mixed layer physics; Part 2: the value of stretched coordinates. *EOS, Transactions, American Geophysical Union*, fall meeting.
- Hsu, H-m., Robert~C. Beardsley, and J. F. Price, 1992. Including surface and bottom boundary layer physics in a primitive-equation model to study coastal circulations. *EOS, Transactions, American Geophysical Union*, **72**(52), 59 (abstract).
- Hermann, A.J. and H.-M. Hsu, 1993. A vertical coordinate mapping technique for semispectral primitive equation models of oceanic circulation. J. Atmos. Ocean Tech., 10(3), 381-396.

DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF COLOR PAGES WHICH DO NOT REPRODUCE LEGIBLY ON BLACK AND WHITE MICROFICHE.