

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE AUG 15, 1996	3. REPORT TYPE AND DATES COVERED Quarterly - 1 APR-30 JUNE 96	
4. TITLE AND SUBTITLE THE EFFECTS OF MAGNETIC STORM PHASES ON F-LAYER IRREGULARITIES FROM AURORAL TO EQUATORIAL LATITUDES			5. FUNDING NUMBERS GRANT NO. N00014-89-J-1754	
6. AUTHOR(S) Jules Aarons and Michael Mendillo Co-Principal Investigators				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Boston University Center for Space Physics 725 Commonwealth Avenue Boston, MA 02215			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) ONR Arlington, VA			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release Distribution is Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p style="margin-left: 40px;">Analysis of equatorial observations of GPS from stations on the magnetic equator and in the anomaly region allow us to begin to study the morphology of height of plumes and the occurrence pattern of thin layers of irregularities during years of low solar flux. Very high altitude plumes have been observed during magnetic storms in the data sets analyzed for 1993-1996. At high latitudes a storm expands the irregularity oval moving the irregularity regions producing scintillation and phase fluctuations to lower latitudes and to polar latitudes.</p>				
14. SUBJECT TERMS Ionosphere, Radio Wave Propagation Scintillation			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

ENC (1)

OFFICE OF NAVAL RESEARCH

QUARTERLY REPORT

for

1 April 1996- 30 June 1996

GRANT No.: N00014-89-J-1754

**THE EFFECTS OF MAGNETIC STORM PHASES ON
F-LAYER IRREGULARITIES
FROM AURORAL TO EQUATORIAL LATITUDES**

Jules Aarons and Michael Mendillo, Co-Principal Investigators

**Boston University
Center for Space Physics
Boston, MA 02215**

Reproduction in whole, or in part, is permitted for any purpose of the United States Govern-
ment

19960926 067

A. RESEARCH PAPERS

In this quarter, this grant's studies were divided between high latitude and equatorial region studies. Two papers on these subjects were submitted for publication to the Proceedings of the Ionospheric Effects Symposium of 1996

High Latitude GPS Observations and Receiver Constraints by J. Clynch, Naval PostGraduate School and J. Aarons, Boston University

GPS Phase Fluctuations in the Equatorial Region J. Aarons, M. Mendillo, and R. Yantosca

B. GPS PHASE FLUCTUATIONS AT AURORAL LATITUDES DURING MAGNETIC STORMS

J. Aarons, M. Mendillo, R. Yantosca Center for Space Physics Boston University Boston, MA 02215

A draft of studies involving predominantly auroral latitudes has been completed. The data indicate the domination of the irregularity oval in the timing of irregularity appearance. Even during magnetic storms the entry and exit of the station into the auroral oval dominates although the magnetic storm essentially modulates the pattern.

ABSTRACT

Using simultaneous observations of the GPS satellites from several high latitude stations it was possible to determine the behavior pattern for the development of phase fluctuations in the auroral and sub-auroral irregularity regions. Starting with characteristics of the irregularity development in magnetically quiet days, it was noted that the start and stop times of phase fluctuations correlated with the entry and exit into the irregularity oval. The duration of phase fluctuations during quiet days was a function of how close the station was to the oval. Maximum occurrence took place near magnetic midnight. The irregularity oval while matching the auroral oval in much of its behavior receded to very high latitudes during extended periods of magnetic quiet. The irregularity oval can be developed in terms of intensity whereas the auroral oval optically is only of occurrence.

During magnetic storms the irregularity oval expands equatorwards and polewards and phase fluctuations increase in intensity. While the geographic position of the station relevant to the oval is important during storms, the dynamics of each storm modifies the simple behavior shown during quiet times.

It was not possible to determine the altitude of the irregularities for the group of magnetic storms analyzed in this paper. Magnetic activity increases the number flux of both soft and hard electrons with the former precipitating at F layer altitudes and the latter at E layer altitudes. The phase fluctuation data are of total irregularity intensity. Recent studies have stressed E layer dominance. However the long studies of spread F using ionosondes and the flux increase of soft electrons as well as hard electrons during magnetic storms makes it difficult to evaluate the contributions of the F layer. Earlier studies have indicated that both E layer aurora and F layer irregularities frequently appear simultaneously during both quiet and disturbed periods. The

thickness of the F layer irregularity region may cause that region to dominate in contributions to irregularities when E and F layer critical frequencies are close (which has been shown in some studies of auroral activity).

C. FUTURE STUDY; INTERRELATIONSHIP OF HIGH LATITUDE AND EQUATORIAL IRREGULARITY DEVELOPMENT

Data sets similar to those cited will be used to study the effect of high latitude magnetic variations and high latitude phase fluctuations on the developments of plumes at the equator. At the equator, it is the generation of neutral winds that is presently thought to be the dominating storms effects on the generation or inhibition of equatorial plume structure for equatorial latitudes in all likelihood requires the use of several magnetic parameters and will include localized effects.

If magnetic data in detail correlates with scintillation, then forecasting will be available if we know the right parameters to put into the model. If this proves to be the case, the study of the relationship of magnetic observations and high latitude development of irregularities would be of great assistance in forecasting scintillations at the equator.