

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 9/27/1996	3. REPORT TYPE AND DATES COVERED Final: 01 JAN 1990 through 30 JUN 1996	
4. TITLE AND SUBTITLE The Effect of Langmuir Circulation on the Surface Boundary Layer		5. FUNDING NUMBERS N00014-90-J-1495	
6. AUTHOR(S) Robert A. Weller and Albert J. Plueddemann			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Woods Hole Oceanographic Institution 360 Woods Hole Road Woods Hole, MA 02543-1541		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Department of the Navy Office of Naval Research Resident Representative 496 Summer St., Room 103 Boston, MA 02210-2109		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Final Report attached			
12a. DISTRIBUTION / AVAILABILITY STATEMENT approved for public release distribution is unlimited			
19961008 131			
13. ABSTRACT (Maximum 200 words) The objectives of the effort were to improve our understanding of the dynamics of Langmuir circulation, to examine its role in controlling the vertical fluxes of heat and momentum which determine the mean vertical structure of the upper ocean, and to make progress toward incorporating the effect of Langmuir circulation in models predicting how the ocean responds to atmospheric forcing.			
DTIC QUALITY INSPECTED 6			
14. SUBJECT TERMS dynamics of Langmuir circulation, vertical fluxes of heat and momentum, vertical structure of the upper ocean, Langmuir circulation in models, atmospheric forcing		15. NUMBER OF PAGES 6	
17. SECURITY CLASSIFICATION OF REPORT unclassified		16. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT unclassified	20. LIMITATION OF ABSTRACT UL	

R. Weller
September 27, 1996

Final Report: N00014-90-J-1495

Robert A. Weller and Albert J. Plueddemann
The Effect of Langmuir Circulation on the Surface Boundary Layer

Department of Physical Oceanography
Woods Hole Oceanographic Institution
Woods Hole, MA 02543

Internet: rweller@whoi.edu, aplueddemann@whoi.edu
phone: (508) 457-2000 ext. 2508 (RAW), ext 2789 (AJP)
fax: (508) 457-2181

The objectives of the effort were to improve our understanding of the dynamics of Langmuir circulation, to examine its role in controlling the vertical fluxes of heat and momentum which determine the mean vertical structure of the upper ocean, and to make progress toward incorporating the effect of Langmuir circulation in models predicting how the ocean responds to atmospheric forcing.

In 1990 we participated in the cooperative Surface Wave Processes Program (SWAPP). High vertical and temporal resolution measurements of velocity and temperature were made from the Research Platform FLIP. Surface wave and meteorological measurements were made at the same time. Collaborators (Smith, Pinkel, Farmer, Crawford) made acoustic Doppler current measurements, imaged bubble clouds, and collected profiles of microstructure near the surface. These data provided the information about the surface forcing (air-sea fluxes and surface waves), the Langmuir circulation, and the mixed layer structure needed to address the objectives. Following the field work we have organized meetings of the SWAPP participants to facilitate collaborative analyses, have exchanged data, and carried out analysis and publication.

The work has resulted in:

- the development of an objective measure of Langmuir circulation strength from the SWAPP observations.
- demonstration that Langmuir circulation effects the mean vertical structure of velocity in the surface boundary layer.
- demonstration that the observed time series of Langmuir circulation strength covaries with the Craik-Leibovich instability parameter.
- unique documentation of the sub-inertial variability of the vertical structure of the oceanic boundary layer.

A time series of the strength of the Langmuir circulation was produced for the duration of SWAPP. The observed strength of the circulation was found to be closely related to the magnitude of the instability parameter derived from the Craik-Leibovich theory. This result supports wave-current interaction theories of Langmuir circulation by showing that both wind stress and Stokes drift must be included in order to parameterize the variability of circulation strength. Thus, it was possible for the first time to relate variations in the forcing (in terms of the mean vertical shear and the surface wave Stokes drift) and in the vertical structure of the upper ocean to the strength of Langmuir circulation observed in the open ocean.

The mean vertical structure of the surface boundary layer was found to be not entirely consistent with present models and to have more shear at sub-inertial frequencies than anticipated from the Price-Weller-Pinkel (PWP) mixed layer model. At the inertial frequency, the vertical structure is closer to slab-like, with shear concentrated at the mixed layer base. The low-frequency shear was surface-intensified, coherent with the wind, and strikingly similar to that seen in numerical simulations of Langmuir circulation. Despite the complexity of the shear profile, the vertically integrated velocity agreed well with the

predicted Ekman transport. Because the surface-forced flow was dominated by near-inertial oscillations with shear concentrated at the mixed layer base, shear-driven mixing and mixed layer deepening was not apparently influenced by the low frequency shear. More often than not, PWP did well at predicting mixed layer depth and the vertical structure of temperature.

However, the presence of Langmuir cells did noticeably impact the evolution of the vertical structure during periods (tens of hours) immediately following strong wind events, when local winds were low but waves persisted. During these periods the PWP model predicted restratification in response to the combination of low wind and solar heating. However, the observations show that Langmuir cells persisted after the end of the wind forcing, redistributed the heat from the penetrating shortwave radiation, and kept the surface layer mixed. In addition to these periods following wind events and more generally, it was found that the variance of the high-frequency shear within the mixed layer was related to Langmuir circulation strength (Gnanadesikan et al., submitted). Thus, even when a deep mixed layer was correctly predicted by the PWP model during strong wind and waves, the slab-like structure of the predicted horizontal velocity field was in sharp contrast with the observed strong high frequency variability in near-surface shear.

The implication of these results is that although the bulk properties of the surface boundary layer can be reproduced with some success from existing models, the vertical distribution of heat and momentum within the boundary layer cannot be. The principal physical mechanism which remains unaccounted for in models is Langmuir circulation (Leibovich, 1983). Langmuir circulation arises from an instability of mean-flow shear in the presence of a surface wave Stokes drift. The results from SWAPP show that predictions of SST and near-surface vertical structure can be improved by incorporating into the models wave forcing and a proxy for mixing by Langmuir circulation.

These findings and the measurements of Langmuir circulation made from FLIP in MILDEX and more recently in SWAPP have had significant impact, causing renewed interest in modeling the impact of large eddies within the mixed layer on ocean physics, biology, and chemistry and in explicit consideration of surface wave forcing of the upper ocean. The basic research effort led the more applied modeling effort and to participation in shallow water acoustic reverberation experiments funded by the ONR ocean acoustics program.

Publications resulting from support under this grant:

Dickey, T. A., A. Plueddemann, and R. Weller, 1995. Current and water property measurements in the coastal ocean. Chapter 15 in The Sea, Vol. 10, The Global Coastal Ocean, K. H. Brink and A. R. Robinson, editors, Part A, Processes and Methods, Wiley Interscience, New York, in review.

Gnanadesikan, A., Comment on "Report on statistics and physical oceanography" Statistical Science, 9:208-212, 1994.

Gnanadesikan, A., Dynamics of Langmuir Circulation in oceanic surface layers, Ph.D. Dissertation, MIT/WHOI Joint Program in Physical Oceanography, WHOI Tech. Rep. 94-23, 354 pp, 1994.

Gnanadesikan, A., Modeling the diurnal cycle of carbon monoxide: Sensitivity to physics, chemistry, biology and optics, J. Geophys. Res., 101:12177-12191, 1996.

Gnanadesikan, A., Mixing driven by vertically variable forcing: An application to the case of Langmuir circulation, J. Fluid Mech., 322,81-107, 1996.

- Gnanadesikan, A., A. J. Plueddemann, and R.A. Weller, Observational evidence that Langmuir circulations are driven by wave-current interaction: Scaling the time-varying near-surface Eulerian shear, subm. J. Mar. Res.
- Gnanadesikan, A., and R.A. Weller, Structure and variability of the Ekman spiral in the presence of surface gravity waves, J. Phys. Oceanogr. 25:3148-3171, 1995.
- O'Brien, M. M., A. J. Plueddemann, and R. A. Weller, 1991. The response of ocean mixed layer depth to physical forcing: Modeled vs. observed. Biol. Bull., 181(2), 360-361.
- Park, M. M., R. C. Singer, A. J. Plueddemann, and R. A. Weller, 1991. High-speed, real-time data acquisition for Vector Measuring Current Meters. IEEE J. Ocean Eng., OE-16, 360-367.
- Park, Y.G., J.A. Whitehead, and A. Gnanadesikan, Turbulent mixing in stratified fluids: Layer formation and energetics, J. Fluid. Mech., 279:279-311, 1994.
- Plueddemann, A.J., J.A. Smith, D.M. Farmer, R.A. Weller, W.R. Crawford, R. Pinkel, S. Vagle and A. Gnanadesikan, 1996. Structure and variability of Langmuir circulation during the Surface Waves Processes Program. J. Geophys. Res., 101(C2), 3525-3543.
- Reider, K. F., J. A. Smith, and R. A. Weller, 1994. Observed directional characteristics of the wind, wind stress, and surface waves on the open ocean, Journal of Geophysical Research, 99(C11), 22589-22596.
- Reider, Karl. F., Jerome. A. Smith, and Robert A. Weller, 1995. Some observations of co-linear wind stress and wave breaking. Journal of Physical Oceanography, 25.
- Weller, R. A., M. A. Donelan, M. G. Briscoe, and N. E. Huang, 1991. Riding the crest: A tale of two wave experiments. Bulletin of the American Meteorological Society, 72(2), 163-183.
- Weller, R.A. and A.J. Plueddemann, 1996. Observations of the mean vertical structure of the oceanic surface boundary layer. J. Geophys. Res., 101(C4), 8789-8806.