


**THE SCREECH OWL:
ITS LIFE HISTORY AND
POPULATION ECOLOGY
IN NORTHERN OHIO**



NUMBER 71

**UNITED STATES
DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE**

19970319 043

DTIC QUALITY INSPECTED 1

NORTH AMERICAN FAUNA

This publication series includes monographs and other reports of scientific investigations relating to birds, mammals, reptiles, and amphibians, for professional readers. It is a continuation by the Fish and Wildlife Service of the series begun in 1889 by the Division of Ornithology and Mammalogy (Department of Agriculture) and continued by succeeding bureaus—Biological Survey and the Bureau of Sport Fisheries and Wildlife. The Service distributes these reports to official agencies, to libraries, and to researchers in fields related to the Service's work; additional copies may usually be purchased from the Division of Public Documents, U.S. Government Printing Office.

Reports in NORTH AMERICAN FAUNA since 1950 are as follows (all sale stock is exhausted):

60. Raccoons of North and Middle America, by Edward A. Goldman. 1950. 153 pp.
61. Fauna of the Aleutian Islands and Alaska Peninsula, by Olaus J. Murie; Invertebrates and Fishes Collected in the Aleutians, 1936-38, by Victor B. Scheffer. 1959. 406 pp.
62. Birds of Maryland and the District of Columbia, by Robert E. Stewart and Chandler S. Robbins. 1958. 401 pp.
63. The Trumpeter Swan; Its history, habits, and population in the United States, by Winston E. Banko. 1960. 214 pp.
64. Pelage and Surface Topography of the Northern Fur Seal, by Victor B. Scheffer. 1961. 206 pp.
65. Seven New White-winged Doves from Mexico, Central America, and South-western United States, by George B. Saunders. 1968. 30 pp.
66. Mammals of Maryland, by John L. Paradiso. 1969. 193 pp.
67. Natural History of the King Rail, by Brooke Meanley. 1969. 108 pp.
68. The Sea Otter in the Eastern Pacific Ocean, by Karl W. Kenyon. 1970. 352 pp.
69. Natural History of the Swainson's Warbler. 1971. 90 pp.
70. The Distribution and Occurrence of the Birds of Jackson County, Oregon, and Surrounding Areas, by M. Ralph Browning, 1975. 69 pp.

Library of Congress Cataloging in Publication Data

VanCamp, Laurel F.
The screech owl.

(North American fauna; no. 71)

Bibliography: p.

Supt. of Docs. no.: I 49.30:71

1. Screech owls. 2. Birds—Ohio. 3. Bird populations. I. Henny, Charles J., joint author. II. Title. III. Series.

QL155.A4 no. 71 [QL696.S83]

596'.0097s [598.9'7] 75-619387

**THE SCREECH OWL:
ITS LIFE HISTORY AND POPULATION
ECOLOGY
IN NORTHERN OHIO**

By

Laurel F. VanCamp, *Wildlife Naturalist*
Ohio Department of Natural Resources
Division of Wildlife

Charles J. Henny, *Research Biologist*
Migratory Bird and Habitat Research Laboratory
U.S. Fish and Wildlife Service



NUMBER 71

**UNITED STATES
DEPARTMENT OF THE INTERIOR
FISH AND WILDLIFE SERVICE**



North American Fauna, Number 71

**Published by
FISH AND WILDLIFE SERVICE
1975**

Contents

	Page
INTRODUCTION	1
The Study Population	3
The Study Area	4
Accumulation of Data	6
TAXONOMY	8
FOOD HABITS	9
During the Nesting Season	10
In Autumn and Winter	13
Fish in the Diet	15
Seasonal Variation	16
MIGRATION AND DISPERSAL	18
Dispersal Related to Age and Season	19
Dispersal Distance	20
Directional Pattern of Dispersal	22
Records of Individual Birds	24
BREEDING BIOLOGY	25
Breeding Season	25
Clutch Size	26
Nesting Success and Fledging Rates	27
Causes of Nest Loss	32
POPULATION DYNAMICS	33
Sex Ratio in Population	33
Mortality Rates	33
Causes of Postfledging Mortality	35
Age at Sexual Maturity	37
PESTICIDES AND POLLUTION	38
Eggshell Thinning	39
Residue Levels in Eggs	40
POLYMORPHISM	41
Annual Variation in Color Phase Ratios	43
Sex, Age, and Color Phase	47
Genetic Hypotheses	49
Brood Size and Color Phase	54
STATUS OF POPULATION	55
Index to Annual Abundance	55
Life Equation Approach	58
Summary of Status	59

(Cont'd)

CONTENTS (Cont'd)

SUMMARY	60
ACKNOWLEDGMENTS	62
LITERATURE CITED	63

Illustrations

FIGURE

Frontispiece, The cryptic coloration of the gray phase of the screech owl.....	v
1. A nest box is checked on the study area	1
2. The four-county study area in northern Ohio.....	2
3. An improved nest box which was used on the study area since the early 1960's	4
4. A typical open field adjacent to a wooden creek bottom in the study area	6
5. The directional preference of screech owl dispersal in the northeastern United States.....	23
6. A banded screech owl being released at its nest box by the senior author	24
7. Five downy screech owls	29
8. Changes in the color phase ratios in the northern Ohio study area, 1944-73	43

NOTE: For additional copies, write C. J. Henny, Building 16, Federal Center, Denver, Colorado 80225.



Frontispiece—The cryptic coloration of the gray phase of the screech owl
(photo courtesy of G. Ronald Austing).

INTRODUCTION

The screech owl (*Otus asio*) is native to North America and breeds throughout the United States and in portions of Canada and Mexico. It is a small owl, 20 cm (8 in) in length from the tip of the bill to the tip of the tail, with a wing span of 56 cm (22 in); it has yellow eyes and prominent ear tufts (see Frontispiece). Although the species is common throughout much of North America, it has not been studied intensively, particularly over a long period. The published literature is concerned mostly with food habits, color phase, taxonomy, and miscellaneous observations. Breeding biology and population dynamics have received little attention.



Fig. 1.—A nest box is checked on the study area
(photo courtesy of George Laycock).




Fig. 2.—The four-county study area in northern Ohio.

This report presents the basic life history and population information about screech owls in northern Ohio over a 30-yr period. The owls studied were nesting in boxes (Fig. 1) established for wood ducks (*Aix sponsa*) along rivers, creeks, and marshes in a four-county area (Ottawa, Sandusky, Wood, and Lucas Counties) near Lake Erie (Fig. 2). No special trapping techniques were required as the screech owls readily used these nesting boxes and could be easily captured while in them. More than 3,000 owls were captured and banded; 500 were recaptured after the initial banding, some 10 or 15 times. This process provided a large quantity of basic information for this report.

The banding data provided useful information on dispersal, mortality, and age at sexual maturity; in several instances, however, the data were supplemented by banding data of other workers in the area north and east of Missouri. More than 500 nesting attempts by the screech owls were recorded, and 53 different species of birds were listed as food items in the nest boxes during the nesting season. Supplemental information on the

seasonal food habits of the screech owl was obtained from food habit cards filed at the Patuxent Wildlife Research Center, Laurel, Maryland.

The senior author initiated the study in 1944 and collected the field data through 1973. The junior author analyzed the data, wrote the report and coordinated the pesticide phase of the study with the Patuxent Wildlife Research Center in 1973. We have also brought together the relevant literature on the screech owl with emphasis on population ecology. This subject may be particularly timely since McLane and Hall (1972) showed that a low dietary dosage of DDE (a chlorinated hydrocarbon pesticide) causes eggshell thinning under laboratory conditions. Eggshell thinning has been associated with declines of many raptorial and fish-eating birds in both western Europe and North America (Ratcliffe 1967; Hickey and Anderson 1968; Lockie et al. 1969; Hickey 1969). Furthermore, the screech owl data have been subjected to a modeling process (Henny et al. 1970) to determine population parameters for the species in northern Ohio.

The Study Population

The screech owl population reported herein is one that nested in wood duck boxes in northern Ohio between 1944 and 1973. The fact that the birds were using artificial nesting structures may cause a bias in a portion of the findings, particularly in terms of population trends (indices) and annual recruitment rates. For example, trends in population numbers should not be based on the percentage of the nest boxes occupied each year because the nest boxes were always placed in suitable habitat. A population reduction from the loss of habitat would be undetected by such an approach. Our population index most likely represents the annual population nesting in suitable habitat. We believe, however, that most of the other findings presented in this paper are representative of the wild screech owl population in the study area, and, as such, were not biased by the type of nest site used.

Screech owls began using the nest boxes in northern Ohio much earlier than wood ducks. In fact, we have found wood ducks nesting in boxes after young screech owls have fledged. It is our subjective assessment that screech owls keep starlings (*Sturnus vulgaris*), which are much more competitive than the owls, out of the nest boxes within the owls' feeding areas. Upon occasion, starlings will bury wood duck eggs under their nesting material and cause the hen to desert. We therefore contend that wood duck nesting success is enhanced by the presence of screech owl nesting. Adequate numbers of nest boxes should be provided for both species.




Fig. 3.—An improved nest box which was used on the study area since the early 1960's. One hundred of this type of box were available annually during the last decade.

Beginning early in the 1960's, most of the nest boxes were made with cylinders shaped like rockets (after the design of Bellrose 1955; Fig. 3). The owls used the boxes for nesting, roosting, and feeding.

The Study Area

Nearly all of the study area lies within the boundaries of what was called the "Black Swamp" during the early annals of North American history. The Black Swamp was drained late in the nineteenth century and is so transformed that an observer would have difficulty in estimating its former extent. Mayfield (1962) discussed changes in the natural history of the area since the coming of the white man, but indicated that agriculture had modified this region long before a competent naturalist had studied the plants and animals.

From the beginning of settlement in the Black Swamp (circa 1820), the entire agricultural system revolved around corn. Most of the conversion to agricultural land in northwestern Ohio took place after 1850 and continued until the turn of the century (Kaatz 1953). Within a few decades, the Swamp was transformed into one of the most productive agricultural regions in Ohio. By 1930, the Black Swamp was the most completely cropped area in the State (Sitterley and Falconer 1933). Today, like 60 or 70 yr ago, the Black Swamp lands are dominantly rural. Although there are urban and industrial areas around Toledo and other smaller cities, the rural scene prevails elsewhere.

The Black Swamp, like most dense, unbroken forests, was not rich in wildlife (Kaatz 1953). Information on bird populations is scarce; however, it is almost certain that many of the species common at the time of settlement have been extirpated or greatly reduced in numbers. Conversely, some of the most common birds today were rare or nonexistent in the area when the first settlers arrived. Some species that are abundant today prefer nesting sites provided by the structures of modern man and are scarcely to be found in natural sites. Mayfield (1962) reviews the changed status of a number of bird species. It is unclear how or to what extent habitat changes have influenced the status of screech owls. In reference to central Ohio, Trautman (1940:275-276) reported, "Statements by hunters and farmers who were well acquainted with the screech owl indicate that before 1900 this species was decidedly uncommon in the area and that with the removal of most of the forests and the decrease in numbers, or extirpation, of barred and horned owls, the screech owl became increasingly numerous. These statements agree with my studies." Furthermore, Bent (1938:246) states, "The screech owl also has been known to nest frequently, even regularly, in cavities in trees close to houses in towns and cities, thus showing more confidence in human beings than most other owls show."

An illustration of an open field adjacent to a wooded creek bottom where several screech owls nested annually is shown in Fig. 4. The loss of forested land in the four-county study area—12% between 1958 and 1967 (Table 1)—is continuing. Farmers are using more and more of their land each year for agriculture. The wooded creek bottom and woodlots are being cleared for mechanized agriculture. Only 7.6% of the study area was classified as forested land in 1967. This contrasts with about 90% of the area being forested in 1853 (Diller 1962). In 1967, the four-county area consisted of 85.2% cropland, 1.9% pasture, 7.6% forested, and 5.3% other land (Anon. 1971). The most important agricultural crops are corn and soybeans.



Fig. 4.—A typical open field adjacent to a wooded creek bottom in the study area.

Table 1. *The forest area in the four-county study area in northern Ohio, 1958 vs. 1967 (from Anonymous 1971).*

County	Forest area (acres)	
	1958	1967
Lucas	— ^a	23,873 (19.5) ^b
Ottawa	7,485	8,398 (5.9)
Sandusky	21,000	15,932 (6.6)
Wood	19,217	17,616 (4.9)
Total^a	47,702	41,946

^aThe 1958 Lucas County figure was calculated with an incorrect standard (H. G. Smith, Soil Conservation Service, Columbus, Ohio, personal communication); therefore, the totals are for the three-county area with comparable data.

^bIn parentheses percent of county inventoried which was forested.

Accumulation of Data

A measure of the annual and seasonal effort to conduct this study is of importance. Unfortunately, the person-days spent in the field were not recorded. The number of nest boxes checked during the nesting season provides an index of annual effort during the 30-yr study. This measure is admittedly crude because the number of visits to each nest box was not recorded. For example, some of the boxes could not be visited more than once during several of the years (for additional details see results). Forty-five boxes were checked in 1944, the first year of the study. The average number of nest boxes checked for each 5-yr period during the study was as

follows: 479, 727+, 985+, 730, 650, 678. The exact number of nest boxes checked from 1952 to 1955 is unknown; however, a high percentage of the boxes were checked only once at the time young were about to fledge. From 1956 to 1973, the annual effort was consistent. More than 4,249 boxes were checked during the 30-yr period.

Breeding Season

Pairs appeared in the nest boxes by the first week of February. During the initial years of the study, some nest desertions resulted from visiting the boxes in February and March. Therefore, in later years the nest boxes were not checked before mid-April. Most screech owls were incubating by mid-April, and, unless the incubating birds were handled, the nests were not deserted. Nest boxes were checked at this time and the males banded if they were in the nest box. Females were not handled until the young hatched. The nest boxes were generally checked three times between mid-April and early June. The females were banded during the first check in which young were in the nests. The young were not banded until the color phase could be determined.

Postfledging Period

Nest boxes were little used after the young fledged, but use began to increase in October when the deciduous trees lost their leaves. We visited nest boxes only rarely in late summer or early fall.

Winter Period

November, December, and January was a period of much nest box use by individual birds. The wood duck boxes were checked periodically during these months and a large number of screech owls was banded. Beginning in the early 1960's all boxes were visited during the annual maintenance check in January; not all boxes were visited during the winter in the earlier years.

In addition to roosting, the screech owls used the boxes for feeding stations during winter. The trait of carrying prey to a cavity before feeding may have evolved: (1) to lower screech owl vulnerability to predation by larger owls, (2) to prevent other animals from eating the prey not consumed immediately by the owls, (3) as a food storage technique to carry the owls over periods of inclement weather or periods of low food availability, or (4) from the habit of the male feeding the female (Burton 1973). This food-carrying trait accounts for the large number of prey items recorded in the nest boxes. As mentioned above, the owls did not use the nest boxes in late summer or early fall when the leaves of deciduous trees are present. We believe the dense foliage protected the screech

owl from predation by larger owls during this period. Thus, the need for using a cavity as a feeding station was minimized in the late summer and early fall.

TAXONOMY

The screech owl is confined to North America and has a continuous breeding distribution from southeastern Alaska, southern British Columbia, southern Manitoba, southern Ontario, southern Quebec, and Maine south to the Cape district of Baja California, Jalisco, Hidalgo, southern Tamaulipas, and Florida (American Ornithologists' Union 1957). The American Ornithologists' Union also recognizes the whiskered owl (*O. trichopsis*) in the mountain areas from southeastern Arizona to Honduras and the flammulated owl (*O. flammeolus*) in southern British Columbia south through the mountains west of the Great Plains to the highlands of Mexico and Guatemala. Additionally, 18 subspecies of the nonmigratory screech owl are recognized.

Numerous other classifications were in effect before 1957 and agreement among taxonomists is still not complete. As with most species of birds, subspecies of the screech owl have been described on the basis of variation in the length of the wing, bill, tail, and tarsus; and in the color pattern of the plumage. In addition to geographical variation, the species is polymorphic in the eastern part of its range and has two distinct color forms: one with the plumage mainly bright rufous and the other with plumage mainly gray. An intermediate color phase also exists, but in low frequency. Elsewhere in the range, only gray birds occur. Owen (1963a) concluded that most of the subspecies had been separated by drawing arbitrary lines through clinal variations. He further indicated that new subspecies were still (in 1959) being described, sometimes based upon as few as five specimens, and thought that the situation should be remedied. Until such time as reproductively isolated sympatric populations are discovered in any part of the range, Owen believed that no subspecies should be recognized and the AOU Check-list entry should include geographical variation.

Marshall (1967), on the other hand, stated that taxonomists have been so engrossed with subspecies that they have ignored a species problem existing in north and middle America. He proposed that seven species and four incipient species of *Otus* occupy north and middle America, where they overlap without interbreeding, in such combinations as their distributions and habitats permit. Marshall's conclusion were based on both morphology and song patterns. The Committee on Classification and Nomenclature of

the American Ornithologists' Union has not commented on the screech owl since the 1957 Check-list was published. For the purposes of the present paper, no trinomial is used and the screech owl population is described on the basis of geography, i.e., the nonmigratory population resident in northern Ohio.

FOOD HABITS

The screech owl is one of the most nocturnal of North American owls. Allen (1924) made a series of observations on the feeding of a brood of young and concluded that the earliest time at which feeding began was 2025, and the latest was 2112; the earliest time at which feeding ceased was 0250, and the latest was 0415. He further reported that both parent birds were engaged in caring for the young. These nighttime hunts yielded a variety of food items including many passerine birds that were apparently captured on their roosting sites.

The food habits of most birds of prey were first studied in detail during the latter part of the 19th and early 20th centuries. Notable publications of this era were by Fisher (1893) and Errington (1932). In summary, they reported that the screech owl diet consisted of mice, shrews, rats, other mammals, small birds, lizards, amphibians, fish, crayfish, insects, spiders, and other invertebrates. Bent (1938) indicated that although birds do not form as large a portion of the food as mammals, the list of species is a long one. More recently, James and Martin (1950) and Stewart (1969) have added several species of birds to the list.

McDowell and Luttringer (1948) estimated that the diet (percentage occurrence) of the screech owl consisted of 22.75% mice, 18% other mammals, 18% songbirds, with the remaining 41.25% reptiles, amphibians, fish, and insects, and other invertebrates. An analysis of 419 pellets collected in western Missouri from 1957 to 1967 showed a preponderance of small rodents in screech owl diets (Korschgen and Stuart 1972). Mice and rats together made up 87% of all foods. Songbirds amounted to 8.4% in occurrence and only 4.1% of the volume. Among birds in portions of the southwestern United States and western Mexico, Ross (1969) lists the common screech owl as being both carnivorous and insectivorous, the whiskered screech owl as being primarily insectivorous but on occasion supplementing its diet through carnivorous predation, and the flammulated screech owl, as being entirely insectivorous. Errington (1932) indicated that as a rule screech owls ate what was most convenient to catch and of a size easy to handle. He further noted that their preferred prey seemed to be mice if such were

available, but in the event of a mouse shortage they readily turned to birds. These studies suggest that the screech owl is an opportunistic predator whose diet includes nearly every class of animal life.

During the Nesting Season

Allen (1924) intensively studied a pair of nesting screech owls in New York for 45 days and found that birds represented an important portion of the diet. Allen provided a list of 77 birds of 18 species brought to the young during the 45-day period. He concluded that because the feathers in the nest undoubtedly represent many more than one bird of each species, the grand total of birds required to feed the three young owls from the time of hatching until left by the old birds were certainly more than 100. Recently, Stewart (1969) also indicated that food items found at the nests of screech owls in Ohio and North Carolina consisted primarily of birds. However, Sherman (1911) reported 40 food items during the nesting season (apparently in Iowa), which included 3 frogs, 1 shrew, 28 mice, and only 8 birds.

During our nesting studies, 477 food items were recorded at wood duck nest boxes used by screech owls. Our observations were made between 26 March and 7 June, and nearly all food items found were intact. Most of the food was stored in the nests during the first 2 wk after the young hatched (late April and early May). Fifty-three species of birds were recorded, making up 64.8% of the items found during the nesting season (Table 2). An attempt was made to calculate biomass of the prey; however, sufficient information was not available regarding the age classes for several of the birds, mammals, and fishes which vary greatly in size. Preliminary calculations suggested that the percentage occurrence and percent biomass in the diet (with the exception of invertebrates which were probably eaten immediately and not adequately represented in the sample) were similar.

Of the five most common species of birds found in the nests, three are transients (Campbell 1968): the yellow-rumped warbler (average arrival 19 April, average departure 20 May), white-throated sparrow (average arrival 14 April, average departure 22 May), and the ruby-crowned kinglet (average arrival 11 April, average departure 17 May); one was a permanent resident (the house sparrow); and one was primarily a summer resident (the starling) nesting in northern Ohio in May. Furthermore, 49 of the 53 species of birds found in the diet of the screech owl must be considered migratory. Therefore, the screech owl definitely takes advantage of the spring migration of small birds to feed its young.

Table 2. *Species stored as food by screech owls during the nesting season in northern Ohio^a.*

Family	Species	n
Birds		
Columbidae		
	Mourning Dove (<i>Zenaida macroura</i>)	2
Cuculidae		
	Yellow-billed Cuckoo (<i>Coccyzus americanus</i>)	1
Strigidae		
	Screech Owl (<i>Otus asio</i>)	1
Picidae		
	Common Flicker (<i>Colaptes auratus</i>)	1
	Downy Woodpecker (<i>Dendrocopos pubescens</i>)	1
Tyrannidae		
	Eastern Kingbird (<i>Tyrannus tyrannus</i>)	1
	Unidentified Flycatcher	1
Hirundinidae		
	Purple Martin (<i>Progne subis</i>)	4
	Barn Swallow (<i>Hirundo rustica</i>)	4
	Tree Swallow (<i>Iridoprocne bicolor</i>)	2
	Bank Swallow (<i>Riparia riparia</i>)	1
	Cliff Swallow (<i>Petrochelidon pyrrhonota</i>)	1
Corvidae		
	Blue Jay (<i>Cyanocitta cristata</i>)	4
Certhiidae		
	Brown Creeper (<i>Certhia familiaris</i>)	1
Troglodytidae		
	Winter Wren (<i>Troglodytes troglodytes</i>)	1
Mimidae		
	Gray Catbird (<i>Dumetella carolinensis</i>)	5
	Brown Thrasher (<i>Toxostoma rufum</i>)	3
Turdidae		
	American Robin (<i>Turdus migratorius</i>)	10
	Hermit Thrush (<i>Catharus guttatus</i>)	8
	Swainson's Thrush (<i>Catharus ustulatus</i>)	7
	Wood Thrush (<i>Hylocichla mustelina</i>)	2
	Veery (<i>Catharus fuscescens</i>)	2
	Unidentified Thrush	3
Sylviidae		
	Ruby-crowned Kinglet (<i>Regulus calendula</i>)	18
	Blue-gray Gnatcatcher (<i>Polioptila caerulea</i>)	4
	Golden-crowned Kinglet (<i>Regulus satrapa</i>)	2
	Unidentified Kinglet	1
Sturnidae		
	Starling (<i>Sturnus vulgaris</i>)	19
Parulidae		
	Yellow-rumped Warbler (<i>Dendroica coronata</i>)	40
	Palm Warbler (<i>Dendroica palmarum</i>)	5
	Common Yellowthroat (<i>Geothlypis trichas</i>)	3

(Cont'd)

Table 2 (Cont'd)

Cape May Warbler (<i>Dendroica tigrina</i>)	2	
Black-and-white Warbler (<i>Mniotilta varia</i>)	1	
Chestnut-sided Warbler (<i>Dendroica pensylvanica</i>)	1	
Tennessee Warbler (<i>Vermivora peregrina</i>)	1	
Yellow Warbler (<i>Dendroica petechia</i>)		
Nashville Warbler (<i>Vermivora ruficapilla</i>)	1	
Canada Warbler (<i>Wilsonia canadensis</i>)	1	
Unidentified Warbler	6	
Ploceidae		
House Sparrow (<i>Passer domesticus</i>)	25	
Icteridae		
Red-winged Blackbird (<i>Agelaius phoeniceus</i>)	5	
Common Grackle (<i>Quiscalus quiscula</i>)	5	
Brown-headed Cowbird (<i>Molothrus ater</i>)	1	
Thraupidae		
Scarlet Tanager (<i>Piranga olivacea</i>)	1	
Fringillidae		
White-throated Sparrow (<i>Zonotrichia albicollis</i>)	37	
Song Sparrow (<i>Melospiza melodia</i>)	14	
Cardinal (<i>Cardinalis cardinalis</i>)	7	
American Goldfinch (<i>Spinus tristis</i>)	4	
Dark-eyed Junco (<i>Junco hyemalis</i>)	4	
Rose-breasted Grosbeak (<i>Pheucticus ludovicianus</i>)	3	
Chipping Sparrow (<i>Spizella passerina</i>)	2	
Field Sparrow (<i>Spizella pusilla</i>)	2	
Savannah Sparrow (<i>Passerculus sandwichensis</i>)	2	
Tree Sparrow (<i>Spizella arborea</i>)	1	
Vesper Sparrow (<i>Poocetes gramineus</i>)	1	
Indigo Bunting (<i>Passerina cyanea</i>)	1	
Rufous-sided Towhee (<i>Pipilo erythrophthalmus</i>)	1	
Unidentified Sparrow	4	
Unidentified Family		
Small Bird	16	
Large Bird	1	
Subtotal (percent occurrence)	309 (64.8%)	
Mammals		
Soricidae		
Unidentified Shrew	4	
Chiroptera (Unknown Family)		
Unidentified Bat	2	
Sciuridae		
Gray Chipmunk (<i>Tamias striatus</i>)	1	
Cricetidae		
Deer Mouse (<i>Peromyscus</i> sp.)	37	
Meadow Vole (<i>Microtus</i> sp.)	52	
Unidentified Mouse	3	
Muridae		
Norway Rat [young] (<i>Rattus norvegicus</i>)	6	
House Mouse (<i>Mus musculus</i>)	39	

(Cont'd)

Table 2 (Cont'd)

Leporidae		
Eastern Cottontail [leg] (<i>Sylvilagus floridanus</i>)	1	
Subtotal (percent occurrence)	145	(30.4%)
Amphibians		
Ranidae		
Leopard Frog (<i>Rana pipiens</i>)	5	
Subtotal (percent occurrence)	5	(1.0%)
Fish		
Centrarchidae		
Green Sunfish (<i>Lepomis cyanellus</i>)	5	
Clupeidae		
Gizzard shad (<i>Dorosoma cepedianum</i>)	8	
Cyprinidae		
Unidentified Minnow	1	
Subtotal (percent occurrence)	14	(2.9%)
Invertebrates		
Crayfish (<i>Cambarus</i> sp.)	2	
Leech (<i>Hirudinea</i>)	2	
Subtotal (percent occurrence)	4	(0.8%)
Grand Total	477	(99.9%)

^a Smaller food items were probably eaten immediately rather than being stored in the nest boxes and were undetected.

A nest box containing four young screech owls at the Winous Point Shooting Club had 40 dead birds stored on 5 May 1972. These included 24 yellow-rumped warblers, 3 common yellowthroats, 1 palm warbler, 2 white-throated sparrows, 1 Nashville warbler, 4 blue-gray gnatchatchers, and 5 unidentified birds—2 of which were sparrows but not house sparrows. When the nest was revisited on 22 May 1972 to band the four young, no food was left in the box. Young flightless starlings were also taken from their nests on several occasions. Three young starlings and one adult were found in a nest box on 14 May 1955. Several trips were apparently made by the screech owl to the starling nest to fill its coffers.

In Autumn and Winter

Data collected by Wilson (1938) at Ann Arbor, Michigan, primarily in the autumn and winter, showed a preponderance of

mammals in the screech owl diet. Only four skulls of birds (0.3% frequency) were found in 1,408 pellets containing 1,549 skulls. Animals which were identified included: meadow mice (87%); deer mice (5%); Cooper lemming mice, *Synaptomys cooperii* (3%); short-tailed shrews, *Blarina brevicauda* (3%); the remainder (2%) included jumping mice, *Zapus hudsonius*; least shrews, *Cryptotis parva*; red squirrels, *Sciurus hudsonicus*; eastern moles, *Scalopus aquaticus*; Norway rats, birds, parts of crayfish, and insects. Wilson concluded by saying that the screech owl's diet showed a definite cycle. During the spring many owls fed upon insects and amphibians. Elsewhere in his paper Wilson (1938:193) notes, "The tendency for owls in this vicinity to ignore birds can possibly be explained by the abundance of mice" Similar results were reported from Michigan in a more recent study by Craighead and Craighead (1956) who reported that meadow mice and deer mice made up 95.3% of the diet in 1942 and 87.2% in 1948. The percentage of small birds (1.2 and 11.4%) showed a substantial increase in the latter winter. According to the Craigheads, this trend toward more small birds in the diet reflected the change in the relation of meadow mice to small bird densities of the latter year.

Fall and winter food items found in the nest boxes during our study totaled 121. The information was collected between 15 October and 23 February, with the majority being recorded in December. We estimated that the screech owl diet during the fall and winter in northern Ohio consisted of approximately 60.3% mammals, 26.4% birds, 5.8% fish, 5.0% frogs, and 2.5% crayfish (Table 3). The nonmigratory house sparrow and cardinal formed over half of the avian food items whereas the migratory bird species were most important during the nesting season.

An observation made in January 1971 was of special note. A screech owl's talons had pierced the skull of a blue jay, but both birds were found dead entangled in a hawthorne tree (*Crataegus* sp.). A defective eye was noted in 13 owls captured during the study. Generally, the defective eyes were sealed shut. The defective eyes may have resulted from owls trying to capture prey in heavily wooded areas; however, the incidence of eye injuries was low (only 13 in more than 3,000 birds handled).

Table 3. *Food items found at nest boxes occupied by screech owls in the fall and winter in northern Ohio^a.*

Birds		Mammals		Fish		Amphibians and crayfish	
House sparrow...	9	Meadow mouse	28	Gizzard shad	5	Leopard frog.	6
Cardinal.....	8	Deer mouse .	20	Bullhead.....	1	Crayfish	3
Mourning dove...	4	House mouse	10	Unidentified fish	1		
Blue jay	3	Norway rat .	9				
Tree sparrow.....	2	Shrew	5				
Starling	1	Unidentified mouse	1				
Common redpoll .	1						
Screech owl	1						
Hermit thrush ...	1						
American robin ..	1						
Common flicker ..	1						
Total items	32	73	7	9
Percent							
(occurrence) ..	26.4	60.3	5.8	7.5

^a A few of the food items may have been taken earlier in the year.

Fish in the Diet

During the 1973 nesting season, a nest box on our study area was found with several dozen gizzard shad. Five green sunfish were also noted in a nest on 5 May 1972 and one minnow on 3 May 1973. Several other nest boxes smelled of fish when they were visited, although no fish were present.

The presence of fish in the diet made us curious about 10 band recovery letters of screech owls that were reported to the Bird Banding Laboratory as drowned. It is doubtful that fishing activities by the owls led to their deaths. Six were found in water tanks (specifically, two in horse troughs and one in a windmill tank), two in rain barrels, one in a backyard swimming pool, and no details were given on the remaining recovery. Although it might be suspected that the drowned birds were all young and inexperienced, only three were less than 2 mo old, whereas all of the others were more than 1 yr old. Kelso (1944) has included water as one of the chief considerations in the choice of territory, along with suitable roosting and nesting cavities, and the absence of larger species of

owls. Kelso indicated that screech owls will drink every day and bathe every 3 or 4 days if water is available. The birds that drowned may have been attempting to bathe or drink, or, more likely, attempting to catch insects or rodents in or near the water. Records of fishing appear in the literature (e.g., Treat 1889), and live fish have been found in tree cavities utilized by screech owls (Frazar 1877).

An incident in our study area on 8 December 1946, provides additional information concerning the relationship of screech owls to water. An oil slick was noticed on the Portage River and three screech owls from the area were captured that day with their legs and underparts covered with oil. The oil on one owl was still present on 14 January 1947 but it had been removed by 27 April 1947. We believe that if the owls were bathing in the water, oil would have been found throughout their plumage. These data lend support to the contention that screech owls take live prey from streams and ponds, and that aquatic food items found in nest boxes were not merely carcasses found along the shoreline. Based on our data, we cannot answer the question, "Do all screech owls take fish occasionally or do a few birds specialize in fish?"

Seasonal Variation

A seasonal pattern in the food habits of the screech owl became apparent during this study, although our sampling method was biased, i.e., most of the invertebrates taken by the screech owls were probably undetected in our study. Actually, the mammals and birds exchanged places of importance between the nesting season and the fall and winter (Table 4). Furthermore, the important bird species in the diet were nearly all migratory (Table 2) and not available during the other portions of the year.

Table 4. *The relative importance of birds to mammals in the seasonal diet of the screech owl (data obtained from nest boxes).*

Time collected	No. mammals	No. birds	Percent birds ^a
Nesting season	145	309	68
Fall and winter	73	32	30

^aThe aggregate of birds plus mammals.

Only rarely have authors discussed the food habits of the screech owl during all seasons of the year, although Baird et al. (187:57) stated, "The food is chiefly small quadrupeds, insects, and occasionally, when they have young, small birds." Fisher (1893) presented tabular data on mammals, birds, and other items found

in the stomachs of birds collected during various months of the year, but did not analyze the data quantitatively for seasonal variation. Fisher (1893:166), however, did state, "It must be said that occasionally it is destructive to small birds, especially during the breeding season, when it has a number of hungry mouths to fill, and also in suburban districts where its favorite food is hard to procure." Fisher's data were collected primarily in the eastern United States. A reanalysis of his bird and mammal data suggested findings similar to ours reported in Table 4.

The 479 food habit cards (approximately one-half of which were used by Fisher [1893]) filed at the Patuxent Wildlife Research Center, Laurel, Maryland, were reanalyzed for seasonal trends in the screech owl diet (Tables 5 and 6). Only data based on stomach contents were considered, and small sample sizes in June, July, August, and September made it necessary to pool 2-mo periods. The relative importance of arthropods in the diet throughout the year was evaluated on the basis of presence or absence in each stomach. This eliminated the problem of tremendous fluctuations in numbers present in individual stomachs and the problem of trying to obtain an exact count of the numbers present. The importance of arthropods (primarily insects) peaked in late summer and declined during the winter (Table 5). It must be emphasized that the arthropods were not detected in our study probably because they were eaten immediately instead of stored. The diet pattern followed availability, again indicating that the screech owl is an opportunistic feeder. The relative importance of birds to mammals, based on the food habit cards (Table 6), was nearly identical to that found in our study (Table 4).

Table 5. *The relative importance of arthropods (primarily insects) in the seasonal diet of the screech owl (data obtained from 479 stomach content cards at Patuxent Wildlife Research Center).*

Month collected ^a	Stomachs checked	No. samples with arthropods	Percent with arthropods
January	56	6	11
February	34	9	26
March	38	9	24
April	28	16	57
May	46	34	74
June-July	40	31	78
August-September .	23	20	87
October	40	24	60
November	98	42	43
December	76	12	16

^a Only samples collected in the Northeastern United States and Ontario.

Table 6. *The relative importance of birds to mammals in the seasonal diet of the screech owl (data obtained from 479 stomach content cards at Patuxent Wildlife Research Center).*

Month collected	Stomachs checked	No. mammals	No. birds	Percent birds ^a
January	56	33	15	31
February	34	15	5	25
March	38	21	6	25
April	28	8	11	58
May	46	3	4	57
June-July	40	3	6	67
August-September	23	5	4	44
October	40	21	8	28
November	98	52	17	25
December	76	45	14	24

^aThe aggregate of birds plus mammals. April-September the percentage of birds was 57, whereas the percentage for October-March was 26. Other vertebrates in the stomachs include 7 toads, 7 frogs, 2 lizards, and 3 fish.

The nesting season of the screech owl seems to be timed to take advantage of the spring migration of small birds. Lack (1968) concluded that breeding everywhere is timed in relation to the availability of food for the young. Eleanora's falcon (*Falco eleanorae*) of the bare rocky islands in the Mediterranean is another species whose nesting season is correlated with the migration of small birds (Vaughan 1961). This falcon, however, has young in September or October and feeds on the small birds migrating south at this season of the year.

The food source for screech owls during the nesting season is possibly unique for owls. The species is not dependent upon a local rodent population, but is primarily dependent upon the spring migration of birds moving into its hunting territory from great distances away. A series of migrant species move through the nesting area on slightly different schedules in May and offers the screech owl a predictable prey source during the complete period when young are in the nest.

MIGRATION AND DISPERSAL

The 1957 Check-list of the American Ornithologists' Union states that the screech owl in the northeastern United States is mainly resident, but wanders in winter south to Alabama and

Georgia. Owen (1963a), probably repeating the Check-list, also indicated that a few birds breeding in the North apparently move south in winter. Evidence of such movement between States, based on banding data, is nonexistent. Most interstate recoveries involved 16- to 32-km (10- to 20-mi) movements of birds banded near the State lines, although two definite exceptions were found. One bird was banded in Minnesota and reported 300 km (185 mi) southeast in Iowa and another was banded on our study area in northern Ohio and recovered 235 km (145 mi) northwest in Michigan. Both recoveries were reported during the first year of life from birds banded as nestlings. The first bird was banded on 25 May 1932 and found dead on 20 December 1932, and the latter was banded on 14 May 1957 and found dead on 20 December 1957. Since the two recoveries were in opposite directions, they suggest a dispersal of young rather than migration. Our findings are in agreement with Fisher's (1893:167) statement, "This owl breeds throughout its range, and does not migrate or even wander far during the winter months."

Dispersal Related to Age and Season

Banding data show that young screech owls begin dispersing from their natal area in late summer and early fall (Table 7), and that by about 1 October, only one-fourth of the surviving young remain in the 10' block (approximately 23,500-hectare area¹) where they were hatched. Some young disperse more than 160 km (100 mi). In contrast, about three-fourths of the adult birds remain sedentary and stay in the 10' block where banded (Table 7). In addition, 87% of the adults remained within 16 km (10 mi) of the banding site and no bird moved more than 64 km (40 mi) (Table 8). Furthermore, three of the four birds that moved beyond 32 km (20 mi) were 6 yr old or older; they may have moved the additional distance to obtain a new mate after their original mate died (adult screech owls have a relatively high adult mortality rate). Generally, there is little movement among adult screech owls. Hence, the major means by which a transfer of individuals occurs among populations of the nonmigratory screech owl is dispersal of young birds. The same findings were reported for the nonmigratory song sparrow by Johnston (1956).

¹An arbitrary geographical division of longitude and latitude used to report bandings of birds and band recoveries.

Table 7. *A comparison of the seasonal movement of young and adult screech owls banded in northern Ohio and in the remainder of the northeastern United States during the nesting season.*

Recovery	NESTLINGS (Banded April-June)				
	Time period of recovery ^b				
location ^a	Apr-June	July-Sept	Oct-Dec	Jan-Mar	Later years
0-6 miles	15 (94%)	20 (61%)	4 (24%)	2 (25%)	8 (27%)
More than 6 miles	1 (6%)	13 (39%)	13 (76%)	6 (75%)	22 (73%)
Total	16	33	17	8	30

Recovery	ADULTS (Banded March-June)				
	Time period of recovery ^b				
location ^a	Apr-June	July-Sept	Oct-Dec	Jan-Mar	Later years
0-6 miles	4 (100%)	4 (80%)	4 (80%)	8 (80%)	23 (74%)
More than 6 miles	0 (0%)	1 (20%)	1 (20%)	2 (20%)	8 (26%)
Total	4	5	5	10	31

^a Recoveries within same 10' block as banded were considered within 6 miles of the banding site. Owls recaptured by banders were not included.

^b Months refer to first year after banding.

Dispersal Distance

The mean distance moved by young birds (calculated after 1 October) was 32 km (20 mi), although a majority of the birds were reported to have moved much shorter distances (Table 8). Several authors have attempted to determine if dispersal distance is random. Most of the tests have involved a comparison of the observed frequency distribution with the Poisson distribution (French et al. 1968). If the dispersal distance is random, then the number of animals in successive zones outward from the center should show a Poisson distribution. In evaluating this distribution the only data readily available on the place of banding and the place of recovery were the respective 10' blocks in which these events occurred. To change areas to locations for which distances can be calculated, we used the geographic centers of the 10' blocks involved. By using this approximation in Table 8, the hypothesis of random dispersal distance for young screech owls was rejected ($P < 0.001$). It appears that two different groups of young birds are

present in the population a group showing little dispersal (less than 32 km [20 mi]), and a group wandering considerable distances (100 to 300 km [60 to 185 mi]). Similar findings were presented for song sparrows by Johnston (1956) and for small mammals by Dice and Howard (1951), Howard (1960), and French et al. (1968). These authors all discussed the possibility that certain individuals within a population may desire to move long distances. Whether there is a genetic basis for his behavior, as proposed by Howard (1960), or whether all individuals are subject to this urge at a certain stage of their lives or season of the year, is uncertain. It should be noted, however, that about 6% of the screech owls moved beyond the expected range. The percentage is similar to the 10% reported for song sparrows in Ohio and California (Johnston 1956). Furthermore, Henny and VanVelzen (1972) indicated that approximately 11% of an osprey (*Pandion haliaetus*) population disperses from their natal area to breed.

Table 8. Comparison of the observed dispersal distance of nestling screech owls with the Poisson distribution (includes recoveries from 1 October). The adult data are included in parentheses but are not statistically tested as little or no dispersal is indicated.

Category (miles)	Observed frequency	Expected (Poisson)	X ²
0-10	25 (48) ^a	4.5	93.39
11-20	21 (3)	11.2	8.58
21-30	3 (2)	14.0	8.64
31-40	2 (2)	11.6	7.94
41-50	0	7.2	13.0 6.23
51-60	0	3.6	
61-70	1	1.5	
71-80	0	0.5	
81-90	0	0.2	
91-100	0	T	
101-110	0	T	
111-120	1	T	
121-130	0	T	
131-140	0	T	
141-150	1	T	
151-160	0	T	
161-170	0	T	
171-180	0	T	
181-190	1	T	
Total	55	54.3	124.78 (P<0.001)

^aNumber in parentheses refers to 55 recoveries of screech owls banded as adults in March through June.

The rigors of wandering during their first autumn and winter probably imposes considerable hardship on the young owls. Many have not yet located suitable roosting areas or other essential requirements, which may make them vulnerable to severe weather and predation from larger owls. However, it should be noted that theoretically, the apparent nonrandom distance of the dispersal has evolved because of its advantage to the species. Current ideas on nonrandom vertebrate dispersal can be conveniently divided into two viewpoints: (1) those that suggest an "inherent" or "genetic" difference between individuals causing them to be short-distance or long-distance dispersers (e.g., Johnston 1956, Howard 1960, Lidicker 1962), and (2) those that suggest the only difference between individuals is their ability (relative dominance) to procure a residence (e.g., Murray 1967).

The first group tends to think of innate dispersal in terms of benefiting the *population* (in spite of the high rate of mortality of long-distance dispersers). The advantages to the population from such behavioral characteristics are said to be: (1) the increase of gene flow to ensure a tendency toward greater panmixia, (2) the increase in ability to spread its range rapidly as favorable habitats are created and to quickly reinvade areas that may have been depopulated by catastrophes, and (3) the increase in ability to regulate population density below carrying capacity. Murray (1967) tends to think in terms of benefit to the *individuals*, i.e., the dominants that established a nearby residence would be at a distinct advantage. Furthermore, Murray (1967) believes the advantages cited by the first group are actually unselected consequences of selection for individuals that aggressively procure breeding sites but move away from dominant individuals. He concluded by saying there is no reason to believe that selection has occurred above the level of the individual. Although the mechanisms involved in dispersal are not fully understood, the many advantages of such a trait are readily apparent.

Directional Pattern of Dispersal

The directional pattern of screech owl dispersal in the northeastern United States was determined by using all birds recovered outside the 10' block where banded. Retrapped birds are not included here as they would be biased to the study area. As shown in Table 8, the majority of the birds were reported within 32 km (20 mi) of the banding site.

The apparent dispersal of more birds east and west is a result of the shape of the 10' blocks at the latitude where the data were collected (Fig. 5). The blocks are rectangular (approximately 21 × 11




Fig. 5.—The directional preference of screech owl dispersal in the northeastern United States.

km [13×7 mi]) and it is a shorter distance (by 53.8%) to an adjacent block if one moves east or west. The data in question are frequencies of recovery in eight recovery areas defined with reference to the 10' block of banding (Fig. 5). Again, we must assume that banding occurred at the center of the block. Because of the shape of the banding block and the recovery areas, these eight frequencies of recoveries would not have equal expected values, even if dispersal were random with respect to the direction of movement. The problem here is to ascertain if the data obtained are consistent with random direction of dispersal.

We computed the distribution of such recovery frequencies conditional on the dispersal distance given a random direction of dispersal.¹ This computation then allowed a test of the null hypothesis that direction of dispersal is random.

Given the length and width of the 10' block of banding, and a set of distances for a corresponding set of frequencies, the expected

¹For additional details, see Memorandum in files at Migratory Bird and Habitat Research Laboratory, "Distribution of band recovery frequencies for the screech owl data." dated 20 May 1974.

frequencies can be computed under the hypothesis of random direction of dispersal. The formulae are not complex, except for requiring some inverse cosines. The theoretical frequencies of recovery by area are shown in parentheses in Fig. 5. The null hypothesis of random directional dispersal is not rejected ($X^2=6.84$, 7 df, $P=0.50$). Therefore, we conclude that the direction of dispersal of the screech owl is random.

Records of Individual Birds

Multi-recaptures on our study area in northern Ohio provided some additional information on movement or lack of movement in adult birds. Generally, after an adult bird established a nesting site, it was found only at that site in future years. For example, an adult female was brooding five young in a box on 12 May 1952. The bird was also found in the same box in 1954 (brooding five young), in 1957 (brooding three young), and 1959 (brooding four young). The bird was found dead in the immediate vicinity of the nest site on 31 March 1960. Records of this type were very common during the study. One of the longest records for an individual owl was a female that was banded in a box on 19 November 1945 and was retrapped eight times at the same nest box before it was killed by a car on 12 February 1958 (approximate age 13 yr) 400 m (0.25 mi) from the nest box. Figure 6 shows a banded screech owl being released at its nest box.



Fig. 6.—A banded screech owl being released at its nest box by the senior author (photo courtesy of George Laycock).

BREEDING BIOLOGY

Breeding Season

Carpenter (1883) discussed the breeding of a pair of screech owls that raised a brood of young in captivity in Massachusetts. He indicated that by 1 February 1883 the actions of the adult birds toward each other began to change and they became more attentive to each other. A clutch of three eggs was found on 15 April and incubation lasted 22 days. Kelso (1944) reported paired owls at Ithaca, New York, by 19 February, but egg-laying did not commence until 15 March. Sherman (1911) reported the first egg of a clutch laid on 27 March 1910 (apparently in Iowa) and that a complete clutch of four eggs was found on 4 April. The first two eggs hatched on 27 April and the remaining two on the next 2 days (26-day incubation period). She reported the birds leaving the nest 30, 31, and 32 days after hatching. Kelso (1950) reported screech owls in New York leaving the nest in 31 to 35 days. Bent (1938) indicated that the incubation period is variously reported as lasting from 21 to 30 days, but that the average is probably around 26 days as determined by Sherman (1911). Craighead and Craighead (1956) studied 28 nests in Superior Township, Michigan, in 1942 and 1948. They reported that the first selection of territories began in late February. The earliest egg-laying date was 18 April and the earliest hatching date was 12 May. The mean date for the broods leaving the nest was 1 June.

During our study, pairs of owls were first seen together in the nest boxes as early as 1 and 3 February; egg-laying began considerably later. The chronology of the nesting season in northern Ohio may be determined from the data in Table 9. Small young (1 to 10 days old) were first seen in the nests during 11-20 April; however, the majority of small young (70%) were found during 21 April-10 May. Therefore, most of the eggs were hatched from 15 April-5 May. Assuming a 26-day incubation period, egg-laying began about 15 March for the majority of the population with a few laying 5 to 10 days earlier. A few small young were seen as late as the first week in June. These young were probably the results of renesting efforts. Thus, most young in the northern Ohio fledge during the last week of May or the first week of June. The young attain a bandable size with their color readily apparent at about 3 wk of age (10 days before fledging).

Table 9. *Percentages of three size categories of young screech owls in 10-day periods during the brooding season in northern Ohio.*

Date	Small young (\approx 1-10 days old)	Medium young (\approx 11-20 days old)	Bandable young (\approx 21-30 days old)
11-20 April	6	0	0
21-30 April	33 ^a	4	5
1-10 May	37	62	17
11-20 May	18	28	43
21-31 May	5	6	30
1-10 June	2	0	4
11-20 June	0	0	1
Number of nests in sample	101	71	440

^a Longitudinal line refers to peak period for young in each size category, which in turn documents the progression of the breeding season.

Clutch Size

Bent (1938) indicated that screech owls lay three to seven eggs, but usually four or five, with the average in favor of five. He further indicated that the extremely large or small sets are rare, and the reports of eight or nine seem doubtful. The average clutch in our study was 4.43 (Table 10).

Among egg sets from 12 museums in the eastern half of the United States, the largest clutches were found in the north-central part of the region and the smallest in the Southeast (Table 10). This suggests a clinal pattern in which clutch size, in general, increased from east to west and from south to north. One unusual clutch of 10 eggs was found during our study. The bird was incubating on 2 May 1967 and three medium-sized young were found in the nest on 18 May. Two birds probably laid the eggs. We banded seven young in another nest on 7 May 1954. Since this was our only record of seven young in a nest and there were no infertile eggs found, it is probably safe to assume that the clutch size was seven. The record of this nest is not included in the table, however, because it was not visited when eggs were in the nest. Little effort was made to obtain information on clutch size during our study because disturbance during the incubation period seemed to cause nest abandonment.

Table 10. *Clutch sizes of screech owls in northern Ohio compared with egg sets in 12 museums (nearly all museum sets were collected before 1900).*

Location	Clutch size								n	Mean
	1	2	3	4	5	6	7	8		
Northeast ^a	0	3	17	36	28	8	0	1	93	4.27
Northern Ohio ^b	0	1	10	37	35	8	0	0	91	4.43
Northcentral ^c	0	0	2	7	6	3	0	0	18	4.56
Midwest ^d	0	1	6	5	3	0	0	0	15	3.67
Arkansas-Oklahoma- Texas	0	0	7	9	0	0	0	0	16	3.56
Georgia-South Carolina-Tennessee ...	0	2	5	3	0	0	0	0	10	3.10
Florida	1	13	32	7	4	0	0	0	57	3.00

^aMassachusetts, Rhode Island, New York, Pennsylvania, New Jersey, Maryland, and Washington, D.C.

^bDoes not include one clutch of 10 eggs.

^cOhio, Indiana, Illinois, and Wisconsin.

^dMissouri, Iowa, Kansas, and Nebraska.

Nesting Success and Fledging Rates

Craighead and Craighead (1956) provide the only estimate of the number of young fledged per nesting attempt; however, their estimate was based on several assumptions because the information from most nests was incomplete. They concluded that between 2.6 and 3.0 young were fledged per pair of adults in Michigan and Wyoming.

Much of our information from nests was also incomplete. Many of the nests were not visited early, i.e., during the incubation period, and some unsuccessful nests were not detected. The number of active nests in 1944-73 (Table 11) partially reflects the authors' effort rather than abundance. From these 440 successful nestings, 1,673 young were believed to have been fledged (3.80 young per successful nesting). The annual variation in numbers fledged per successful nesting was small (coefficient of variation [$\times 100$] of 8.2).

Table 11. *Success of screech owl nests in northern Ohio, 1944-73.*

Year	No. active nests	No. nests successful ^a	Percent of nests successful ^b	No. young fledged	Young fledged per successful nest
1944	4	4	100.0	17	4.25
1945 ^c	5	4	80.0	16	4.00
1946 ^c	9	6	66.7	22	3.67
1947 ^c	19	14	73.7	56	4.00
1948 ^c	20	11	55.0	37	3.36
1949 ^c	19	15	78.9	61	4.07
1950	26	24	92.3	87	3.63
1951	11	10	90.9	39	3.90
1952	25	23	92.0	85	3.70
1953	34	31	91.2	116	3.74
1954	49	44	89.8	197	4.48
1955	42	42	100.0	161	3.83
1956	22	20	90.9	81	4.05
1957	16	14	87.5	54	3.86
1958	17	14	82.4	50	3.57
1959	15	12	80.0	49	4.08
1960 ^c	17	11	64.7	40	3.64
1961 ^c	12	9	75.0	32	3.56
1962	20	18	90.0	70	3.89
1963 ^c	16	11	68.8	40	3.64
1964	9	9	100.0	34	3.78
1965	8	7	87.5	29	4.14
1966	9	9	100.0	30	3.33
1967 ^c	18	16	88.9	54	3.38
1968	13	13	100.0	49	3.77
1969 ^c	12	10	83.3	35	3.50
1970	13	12	92.3	45	3.75
1971 ^c	13	11	84.6	34	3.09
1972	18	16	88.9	53	3.31
1973 ^d	—	—	—	—	—
Total	511	440	86.1	1,673	3.80

^aFledged at least one young.^bThis percentage is biased high because many nests were visited only once when the young attained bandable size.^cYears when 75 percent of nests were visited on two or more occasions.^dData excluded because one egg from each clutch was collected for pesticide study.

The banding date disclosed variation in the number of young reported per successful nesting (Table 12). The earliest nesting attempts produced slightly more young than later attempts. This may be partially due to the older, more experienced birds nesting first, whereas the late nests may include more first-year breeders. Also, a few renests with small clutch sizes may have been included in the last category. A brood of five young is shown in Fig. 7.




Fig. 7.—Five downy screech owls (photo courtesy of G. Ronald Austing).

Table 12. Number of young screech owls per successful nest during five banding periods (from early to late nesters) in northern Ohio.

Banding date ^a	Successful nests	Number banded	Mean per nest
22 April — 9 May	88	367	4.17
10 May — 15 May	88	336	3.82
16 May — 19 May	88	335	3.81
20 May — 23 May	88	334	3.80
24 May — 20 June	88	301	3.42
Total	440	1,673	3.80

^aThe banding dates were adjusted so that an equal number of successful nests were reported in each banding period.

Because the data presented in Table 11 include a number of nests that were visited only once, i.e., at the time of banding, a rationale was needed to obtain an estimate of the number of young fledged per breeding pair. The data are obviously an overestimate of the percentage of nesting attempts that were successful. Two methods were used to estimate the number of young fledged per breeding pair: (1) use of only nests with the most complete set of observations, i.e., nests visited initially during incubation, and (2) the Mayfield (1961) Exposure-day Method (and the data which met his criteria).

Nests Visited Initially During Incubation

This method was restrictive and reduced the number of usable nesting records from the total of 511 to 165, which were the most complete ones (Table 13). Because egg-laying in northern Ohio begins about 15 March for most of the population, the nests visited between 18 March and 10 April best represent the fledging rate per nesting attempt (Table 13). The fledging rate estimates for 11-30 April were believed to be inflated because of undetected losses before the first visit to the nest. Thus, we believe the best estimate by this rationale is 2.55 young fledged per pair.

Table 13. *Screech owl nesting success and fledging rates as related to the date the nest was first visited. Only nests with eggs present at first visit were included.*

Date first visited ^a	No. of nests	No. of young fledged	Young fledged per nesting attempt
18 March — 10 April	20	51	2.55
11 April — 20 April	64	178	2.78
21 April — 30 April	81	238	2.94

^a All years combined.

Mayfield Exposure-day Method

Other field workers have often been faced with interpreting nesting success with data from nests that were not observed from the start of incubation. Further, the sample may include some nests for which success of nesting was unknown. If such partially complete records are included (or excluded) from his calculations, there is a danger of distorting the conclusions. Mayfield (1961) presented a way of dealing with this problem by reducing the data to units of exposure, which reflects not only the number of nests but also the length of time each was under observation. A convenient measure of exposure is the nest-day (equivalent to one nest for 1

day). With this method, all single or multiple observations of individual nests can be incorporated into the sample, even though some of these do not go back to the very beginning or do not carry through to the end. Mayfield (1961) points out that after mortality and survival rates are expressed per nest-day, the probability of survival of eggs and young may be calculated for all or any part of this nesting period. If the survival rate per nest-day is S , the probability of survival of a nest for d days is S^d . Mayfield further pointed out that the survival rate during incubation is different from the rate during the nestling period, so these two stages must be treated separately. Also, egg survival must be considered separately from nest survival because of the loss of individual eggs, particularly at hatching time. The probabilities of survival in different stages of nesting may be combined through the mathematical principle that the probability of a succession of events is the product of their separate probabilities.

We found that only 13 young from successful nestings (nests from which at least 1 young is fledged) were lost in 1,090 exposure days between 1944 and 1973 (loss per exposure-day of 0.012). Furthermore, the loss of young per exposure-day was quite consistent throughout the brood rearing period in successful nestings (i.e., 22 April–9 May—0.010, 10–20 May—0.012, and 21 May–20 June—0.015). Based on a 30-day period for the young in the nest, an estimated 0.36 young was lost per successful nesting from the period of hatching to fledging. Thus, we worked backwards from the number of young fledged per successful nesting (3.80 from Table 11) to determine the number of young hatched per successful nest. This approach was a necessity because of our weak data for early in the incubation period. Early visits had the potential to cause nest abandonment and thus, it was not possible to estimate egg loss on an exposure-day basis. Egg loss in successful nests was estimated by difference. With an estimated 3.80 young fledged per successful nesting, we estimate that approximately 4.16 young hatched per successful nesting from the 4.43 eggs laid (Table 10). Thus, by difference, an estimated 94% of the eggs in successful nests hatched. The loss of young after hatching from broods that were not completely destroyed appeared to be low.

As mentioned earlier, the percentage of successful nesting was overestimated in Table 11, particularly during years when less than 75% of the nests were visited two or more times (Table 11, footnote c). It is most important to obtain an estimate of the percentage of nestings that were successful. Mayfield's (1961) method is again believed applicable. Since few nests were visited more than twice, a problem exists regarding the timing of nest loss.

Therefore, we must of necessity assume that the loss rate of complete nests was constant through time. This assumption may or may not be true; therefore, our recruitment estimates based on this rationale must be regarded as tentative. Our pooled data indicate that during 5,189 exposure days, 34 nests were lost or believed lost (young in 2 nests may have fledged before being revisited; however, we assumed that one nest was successful). An estimated 0.006552 nest was lost per nest-day. With an incubation period of 26 days and a nestling period of 30 days, each nest was susceptible to loss for 56 days. By applying Mayfield's formula, (0.993448^{56}) we estimated that 69.2% of the screech owl nesting attempts in northern Ohio were successful during the study. With 3.80 young fledged per successful nest, an estimated 2.63 young were fledged per nesting attempt based on the Mayfield Exposure-day Approach. This estimate is in close agreement with that of 2.55 fledged young based on a much smaller sample of nests observed throughout the breeding season. In summarizing our data on recruitment, we believe that the best estimate of the number of young fledged in northern Ohio during the 30-yr study was about 2.5 to 2.6 young per nesting attempt.

Causes of Nest Loss

Sixty-six unsuccessful nests were found during our study. The actual cause of nest failure was determined for a few of the nests listed in Table 14. Most of the nests (41) were either deserted or destroyed. Raccoons (*Procyon lotor*) were implicated in the loss of 15 nests, and the eggs failed to hatch in 7 nests. Craighead and Craighead (1956) noted that fox squirrels (*Sciurus niger*) destroyed the eggs of two pairs of screech owls in Michigan and appropriated the nest hollows. They also pointed out that screech owl nests and ranges are commonly within the nesting range of the great horned owl (*Bubo virginianus*) and that young suffer their heaviest mortality soon after leaving the protection of the nest hollow.

Table 14. *Causes for screech owl nests being unsuccessful in northern Ohio, 1944-72.*

Cause of loss	Number of nests
Deserted or destroyed (reason unknown)	41
Raccoon predation	15
Eggs failed to hatch	7
Young dead in nest	2
People (children) took young	1
Total	66

POPULATION DYNAMICS

Sex Ratio in Population

Behavioral differences between sexes are to be expected in most species. Bellrose et al. (1961) described biases involved in sampling waterfowl populations to obtain sex ratios. For waterfowl and many other groups of birds, a bias may result from (1) a differential migration of the sexes (time of year sample collected), (2) the type of area where collection was made (characteristics of the habitat), or (3) a differential vulnerability of the sexes to the method of collection (e.g., shot, trapped, etc.).

The information on screech owls concerning sex ratios may also be biased, although the nonmigratory characteristic of the species tends to minimize several potential biases. The sex data were obtained from the stomach content records filed at the Patuxent Wildlife Research Center. Not all records contained information on sex. To eliminate one potential bias, the information from March through June (the nesting season when the female was probably on the nest) was eliminated from the sample. Thus, for July to February, of 234 screech owls collected in the northeastern United States, 113 (48.3%) were males and 121 (51.7%) were females. The sample did not deviate significantly from the expected 50:50 sex ratio (chi-square = 0.13, 1 df).

Data obtained in our study area could not be analyzed for sex ratio information because of the potential bias associated with the sampling scheme (nearly all birds captured at nest boxes were paired).

Mortality Rates

Mortality rates for the screech owl have not been estimated previously. In this study, life tables were made from band recovery data in northeastern United States and Ontario; a large percentage of the data came from our study area. Recoveries were not included if information was insufficient to tell if the bird was dead or alive and released. Also, all retrapped birds, sight records, and skeletal remains were omitted. Nestlings were generally banded about 10 days before fledging and, therefore, recoveries during the first 10 days after banding were not included. An initial date (the date after which recoveries would be used in the life tables) of the first 15 May after banding was used for the adults, as opposed to 1 January, because we believe that the nonmigratory screech owl does not attain an adult mortality schedule by 1 January. Henny (1972) indicated that most migratory passerines achieve a schedule of

adult mortality by the first 1 January after hatching; however, the more sedentary species (e.g., cardinal, black-capped chickadee [*Parus atricapillus*] blue jay) in the northern portion of the United States do not attain their schedule of adult mortality rates until later. This phenomenon was first reported for the sedentary California quail (*Lophortyx californicus*) by Emlen (1940).

Nestlings and adults banded through 1964 and recoveries processed through August 1974 were included in the analysis. It is believed that the life span of the banded cohort was nearly completed by 1974 as only two recoveries greater than 9 yr of age have been reported during the last 50 yr. However, the possibility still exists that the mortality rate estimates from the composite dynamic life table (Hickey 1952) may be biased slightly on the high side.

Mortality rate estimates shown in Table 15 suggest that approximately 32.8 to 39.0% of the adult population dies annually. The range in the estimates of adult mortality rate is undoubtedly due to small sample sizes, particularly for the cohort banded as nestlings. Probably the best estimate of adult mortality rate would be based on the pooled recoveries from bandings of both nestlings (second year and later) and adults. The combined data yielded an annual adult mortality rate estimate of 33.9%. The first-year mortality rate for young birds is much higher, about 69.5%.

Table 15. Estimates of mortality rates for screech owls banded as nestlings and adults in the northeastern United States and Ontario during 1915-64.

Age interval years	Adults ^a			Nestlings		
	Number of recoveries	Alive at beginning	Mortality rate	Number of recoveries	Alive at beginning	Mortality rate
0-1	55	134	0.328 ^b	73	105	0.695 ^c
1-2	25	79		13	32	0.390 ^b
2-3	13	54		7	19	
3-4	9	41		4	12	
4-5	7	32		2	8	
5-6	8	25		2	6	
6-7	6	17		3	4	
7-8	3	11		1	1	
8-9	5	8		0	0	
9-10	1	3		0	0	
10-11	1	2		0	0	
11-12	0	1		0	0	
12-13	1	1		0	0	

^a An initial date of the first 15 May after banding was used for all birds not banded as nestlings. On this date, the birds were arbitrarily classified as adults.

^b Annual adult mortality rate estimate.

^c First-year mortality rate estimate.

Causes of Postfledging Mortality

Band recoveries provide some information on causes of mortality, although dead owls found by humans are not a random sample of the mortality occurring in a population. For example, predation by other raptors would most likely be undetected because the carcasses would disappear before being found. Actually, most bands reported to the Bird Banding Laboratory are from birds that died as a result of man's activities (his vehicles, pets, etc.). Furthermore, the majority of the bands are reported as found on dead birds with no cause of death mentioned.

Screech owls hit by motor vehicles and found along roads appear frequently in the sample reported to the Bird Banding Laboratory (Table 16). Sutton (1927) reported 113 requests for permits to retain specimens of the screech owls which had been found dead or in weakened or wounded condition in Pennsylvania. The causes of death included 6 which may have died of starvation or illness, 2 which had flown into window panes, 7 which were caught by steel traps, 1 which had been killed by the felling of a tree, 13 which had been shot, 2 which were alive but in poor condition, and 82 which evidently were killed by flying into automobiles. Schorger (1954) recorded 235 road-killed screech owls during a period of 18 yr between Madison, Wisconsin, and Freeport, Illinois; Stupka (1953) examined 41 screech owls along the highways in or near Great Smoky Mountains National Park. Both Sutton (1927) and Stupka (1953) believed that in most instances these birds were struck and killed while feeding upon prey, primarily insects, along highways.

The recovery data submitted to the Bird Banding Laboratory, as anticipated, suggested that predation on the screech owl by other raptors was of minor importance. Craighead and Craighead (1956), however, indicated that the screech owl was fairly vulnerable to the great horned owl because both often hunt the same habitat.

The importance of the various mortality factors affecting a screech owl population cannot now or perhaps ever be determined. However, it is important to recognize that man, his machinery, his vehicles, his pets, and his buildings and other structures do take their toll; at the same time, it should be understood that many of man's alterations of the habitat have benefited the screech owl. Numerous authors have stated that screech owls seem to take kindly to man's occupation of the land and breed and thrive in the vicinity of his dwellings, in spite of the injudicious warfare which is sometimes waged against them.

Table 16. *Circumstances that led to the deaths of banded screech owls reported to the Bird Banding Laboratory.*

	Month of recovery												Total
	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	
Found dead	15	21	27	21	22	25	16	17	12	16	23	15	230
(no cause reported)													
Killed by motor vehicle	3	7	9	3	3	5	7	4	5	4	3	15	68
Misc. or no information	4	7	10	4	8	5	7	3	2	3	7	2	62
Shot	6	3	4	1	3	8	4	2	3	4	4	5	47
Found in building	2	3	5	1	1	1	2	2	4	5	—	—	26
Drowned	—	—	1	2	2	—	2	3	—	—	—	—	10
Killed by dog or cat	2	1	—	1	1	—	—	—	1	1	—	—	7
Flew into object	2	1	—	—	1	—	1	1	—	—	1	1	8
Trapped or snared													
(Not used for bird banding)	3	1	1	1	—	—	—	—	—	—	1	1	8
Sick bird	1	—	1	—	—	2	1	—	—	2	—	1	8
Killed by train	1	—	2	—	—	—	—	—	—	1	—	—	4
Killed by hawk or owl	1	—	—	—	—	—	—	—	1	—	—	1	3
Total	40	44	60	34	41	46	40	32	28	36	39	41	481

Age at Sexual Maturity

The age at sexual maturity is one of the most important and difficult parameters to determine for a wild population of birds, and published information on the subject for the screech owl was not available. Screech owls were recorded breeding as 1-yr-olds on 10 occasions (7 females and 3 males) during our study, and 3 other 1-yr-olds were found in nest boxes in April but with no sign of eggs or a mate. These birds were all banded as nestlings and found the following spring. Of particular interest was a 1-yr-old male that mated with its mother in 1971 and produced one young. Nestings where a 1-yr-old was involved fledged 3.55 young per successful attempt (slightly lower than the 3.80 average for the population, Table 11). Ten is a small number to have of definite records of 1-yr-olds breeding considering the amount of time spent in the field during the 30-yr study; however, it must be remembered that the young disperse during the late summer and early fall (Table 7). Thus, the chance of recapturing a banded 1-yr-old screech owl in another nest box is remote.

As well as establishing the capability of 1-yr-olds to breed, we need to know the percentage that do breed at this age. Of 13 1-yr-olds found during the nesting season, 10 (77%) were nesting. This estimate may be biased because other nonbreeding 1-yr-olds may not associate with nest boxes and therefore be undetected. Another approach is to compare the number of breeding 2-yr-olds found nesting during the study with the number of 1-yr-olds found nesting. Then, given the average annual adult mortality rate of approximately 33.9% (from previous section), and given that eight 2-yr-olds were found nesting, we estimated that there should be a total of 12.1 breeding 1-yr-olds located if all 1-yr-olds and 2-yr-olds attempted to nest. Of course, only 10 were found, which provides an estimate of 83% (10/12.1) of the 1-yr-olds attempting to nest annually.

Conclusions regarding the percentage of the 1-yr-old screech owls that attempt to nest annually must be regarded as tentative at this time. Furthermore, the percentage may fluctuate from year to year. Our two estimates (based on very small sample sizes) suggest that an average of possibly 77 to 83% of the 1-yr-olds attempt to nest annually, but the percentage may be lower.

In comparison, Henny (1972) summarized information from several published and unpublished sources on age at sexual maturity of great horned owls. He concluded that most individuals do not breed until they are 2-yr-olds, although an average of about one-fourth of the population nests as 1-yr-olds, with the percentage

varying from year to year depending upon local conditions. Similar but more dramatic findings have been reported for other owls that are primarily dependent upon one food source. Barn owls (*Tyto alba*) were reported to nest at the end of their first year of life (Schneider 1937, Stewart 1952); however, in general, barn owls seem to breed irregularly. Wallace (1963:209) stated, "The barn owl may breed almost continuously during peak years in the *Microtus* (meadow mouse) cycle, but slow down or skip a breeding season when its staple prey is scarce." Similar observations were recorded for the tawny owl (*Strix aluco*) in England. Lack (1966) reported that 59% of the population was breeding pairs, but the percentage that nested in specific years ranged from 0 to 90%. Increases in clutch size also were reported for some species of owls in years when rodents were abundant. Lack (1968) reported that the short-eared owl (*Asio flammeus*) may lay nine eggs during a vole plague, but only about half that many during other years.

The number of young fledged per successful screech owl pair was very consistent from year to year (Table 11). This is probably the result of the species being an opportunistic feeder. We do not believe that large numbers of breeding-age screech owls fail to produce in years when the local rodent cycle is low, although rodent cycles may have some influence on 1-yr-olds. Unlike the barn owl, tawny owl, and short-eared owl, the screech owl is not totally dependent upon local rodents (or one genus of mouse) for its food supply. In fact, the young are fed a diet which consists largely of small birds of many species which migrate through the area (Table 2). Thus, the varied food supply for the young is not produced locally; it is highly dependable and actually migrates to the owl's hunting areas annually from numerous places, some being considerable distances away. The opportunistic feeding habits of the screech owl apparently tend to maintain a very uniform reproduction annually and, to our knowledge, nonbreeding is limited to a small percentage of the 1-yr-olds. We must caution, however, that more research is needed on sexual maturity.

PESTICIDES AND POLLUTION

Eggshell thinning has been found to occur widely and is well documented, especially in the United States and England. In the United States, natural populations of at least 22 species representing seven orders have been affected. In seven of eight species where shell thinning exceeded 20%, there was an associated population decline (Anderson and Hickey 1972). The thinning

began to occur in the United States and England in the mid-forties at the time DDT and other organochlorine insecticides were first available for widespread use (Ratcliffe 1967, Hickey and Anderson 1968). The possible relationship of shell thinning to the use of DDT led to a series of studies to test the effects of DDT and its metabolites on shell thinning and reproduction of birds.

Initial laboratory studies were conducted on mallards (*Anas platyrhynchos*), black ducks (*Anas rubripes*), and American kestrels (*Falco sparverius*) and showed a direct causal relationship between DDE and eggshell thinning (Heath et al. 1969; Longcore et al. 1971; Wiemeyer and Porter 1970). The next species to be studied in a similar manner was the screech owl (McLane and Hall 1972). The owls were given a dietary dosage of 2.8 ppm of DDE (equivalent to 10 ppm dry weight). Birds fed untreated food in 1970 and DDE dosage in 1971 laid eggs with shells that were 12% thinner than in 1970. McLane and Hall (1972) concluded that screech owls seem to show greater shell thinning than American kestrels which showed 10% thinning on a similar diet (Wiemeyer and Porter 1970).

Eggshell Thinning

The work by McLane and Hall (1972) indicated that screech owls were susceptible to the eggshell thinning phenomenon. In cooperation with the Patuxent Wildlife Research Center, one egg was collected from each of the 21 clutches on our study area in 1973. The eggs were measured for eggshell thickness and analyzed for chlorinated hydrocarbon residues. The remainder of the clutch was observed for hatchability and fledging rates. Klaas and Swineford (In preparation, Patuxent Wildlife Research Center) compared shell thickness in these 1973 eggs with shell thickness in eggs collected before 1947 in Ohio (Table 17). No significant decrease in eggshell thickness in the 1973 population was detected.

Eggshells collected from the field in 1973 were about 8% thicker than the controls in the laboratory experiment (Table 17). This is of interest because many of the owls in the laboratory experiment were obtained from our study area in northern Ohio in 1967. The reason for the thinner control eggs in the laboratory study is unknown, but it may be due to a number of factors (e.g., age of birds, diet, etc.).

Table 17. *Shell thickness of screech owl eggs.*

Location	No. of females	No. of eggshells measured	Shell thickness (mm)		
			Mean	(S.E.)	Range
Laboratory experiment ^a					
Controls	14	66	.218		.180-.253
Experimental (DDE)	6	28	.189 ^b		.170-.205
Field study ^c					
Pennsylvania pre-1946 ^d	37	37	.241	.003	.197-.277
Ohio pre-1946 ^d	12	12	.244	.004	.230-.287
Ohio 1973 (random, early) ^d	19	19	.234 ^f	.004	.197-.260
Ohio 1973 (addled) ^e	8	16	.243 ^f	.007	.157-.270

^a Patuxent Wildlife Research Center (McLane and Hall, 1972).

^b Significantly different from shell thickness of controls ($P < 0.05$).

^c Klaas and Swineford (In preparation, Patuxent Wildlife Research Center).

^d One egg selected at random from each clutch in early stages of incubation, but after clutch was completed.

^e Unhatched eggs removed from nest after incubation was complete. Includes 11 eggs from three clutches in which no eggs hatched.

^f No statistical differences were detected in the means for the four groups of eggs. t-test ($P > 0.1$).

Residue Levels in Eggs

The 19 fresh eggs collected early in the nesting season and an additional 16 addled eggs found in the same nests at the end of the 1973 nesting season were analyzed for chlorinated hydrocarbons (Klaas and Swineford, in preparation, Patuxent Wildlife Research Center). DDE and polychlorinated biphenyls (PCB's) were found in all 35 samples (Table 18), although the levels were generally low, as expected in view of the lack of eggshell thinning in the population (Table 17). Eight eggs in the sample (all from separate clutches) contained dieldrin (range 0.10-0.24 ppm).

Table 18. *Residue levels of DDE and PCB's in 35 screech owl eggs collected in northern Ohio, 1973 (from Klaas and Swineford in preparation, Patuxent Wildlife Research Center).*

Category	Arithmetic mean ^a (ppm)	Range (ppm)
DDE	1.29	0.33-2.8
PCB	1.32	0.26-3.4

^a Wet weight.

Numerous authors have stated that bird-eating and fish-eating raptors usually build up higher residues and have greater shell thinning than do rodent-eaters (review by Stickel 1973). Why did the screech owls, whose diets contained large quantities of migrant birds, not display these characteristics? We can only speculate that the lack of shell thinning and high egg residues in the Ohio population was because the eggs were laid before the predominance of avian prey became available to the owls. The owls just prior to and during the egg-laying period were feeding primarily on rodents. Thus, the seasonal pattern in the food habits of the screech owl may be offering the species some protection against the accumulation of high pesticide loads.

Eggs in three clutches contained the organochlorine pesticide mirex (three eggs in one clutch, mean 0.10 ppm, one egg in another clutch, 0.12 ppm, and one in another clutch 0.16 ppm). Mirex has stimulated considerable controversy during the past several years because of its actual and proposed widespread aerial application throughout the southeastern United States to control the imported fire ant (*Solenopsis saevissima richteri*). Recent laboratory studies by Heath and Spann (1973), however, showed no perceptible reproductive effects on bobwhite quail (*Colinus virginianus*) or mallards. Since mirex spray activities have been confined to the southeastern United States, we believe the screech owls in northern Ohio obtained their mirex from migratory passerine birds that wintered in the Southeast. The diet of screech owls during certain periods of the year consists mainly of migratory passerines (Table 2).

POLYMORPHISM

The existence of bimodal or polymodal variation within a species is referred to as polymorphism, in which intermediate forms occur at low frequency or are even absent (Owen 1963b). Polymorphism may be defined as the occurrence together in the same habitat of two or more distinct genetic forms of a species of animal or plant in such proportions that the rarest of them cannot be maintained by recurrent mutation (Ford 1940). Differences between the sexes, between young and older individuals, and among seasons are excluded from this definition. Hence, an extremely variable species may not necessarily be polymorphic. Huxley (1955) provided an outline for the analysis of polymorphism which included among other items: (1) the ecogeographical mapping of morph-frequency and determination of ratio-clines (if present, this should be repeated at regular intervals), (2) experimental and statistical

evaluation of viability and other intrinsic or extrinsic selective advantages of the morphs concerned, and (3) genetical analysis of the underlying balance and stability mechanisms. In evaluating polymorphism in the screech owl, we attempt to follow the outline presented by Huxley.

The existence of two distinct color forms of the screech owl has been known since about 1874 when Ridgway (Baird et al. 1874) realized that rufous and gray birds were of the same species and that the forms were independent of age, sex, or season. Earlier, rufous and gray birds had been variously interpreted as belonging to different sexes or to different age classes (review by Hasbrouck 1893). Ridgway later published in many natural history and scientific journals an appeal for information on the frequency of rufous and gray birds, nesting information, and any other data that might help to solve the problem. The information was passed over to E. M. Hasbrouck who published an account of his findings (Hasbrouck 1893). A number of inaccuracies in Hasbrouck's paper were pointed out at the time in a critical review (Allen 1893). It was 70 yr until another full survey of the geography of color forms in the screech owl was attempted (Owen 1963b), although several authors discussed small geographic areas (Martin 1950, Schorger 1954, Hrubant 1955).

Owen (1963b:189) summarized his description of geographical trends in the occurrence of color forms of the screech owl as follows:

"Throughout most of North America east of about 104° there are two forms of the screech owl: one with the plumage mainly bright rufous and the other with the plumage mainly gray. Birds of intermediate coloration also exist, but in most areas they are rare. The relative frequency of rufous birds varies geographically in the form of a cline from north to south; about a quarter or less of the northern population is rufous, while in the south (the Gulf coast and Florida excepted) up to three quarters of the population may be rufous. . . . The geography of polymorphism in the screech owl appears to be an unusual pattern of variation within an animal species."

Owen (1963b) indicated that the presence of polymorphism in the screech owl probably represents balanced adaptation of the forms to varying environmental conditions. After noting that the adaptive significance of the polymorphic forms was not understood, Owen indicated that a cline in the relative frequency of the forms as opposed to random or irregular distribution supports the view that polymorphism in the species is maintained by selection operating along environmental gradients. Fisher (1930)

indicated that it is extremely unlikely that two or more very different phenotypes would be equally adapted to the environment in which they live; a balance of selective forces must be involved, for if not, one form would rapidly replace the other and there would be no polymorphism.

Annual Variation in Color Phase Ratios

Hasbrouck (1893) and Owen (1963b) studied the general patterns in the geographical occurrence of the two color phases of the screech owl. However, to explore the factors responsible for the patterns observed, we believed it would be fruitful to study yearly fluctuations in a given population, preferably one near the edge of its range. Although the screech owl's range extends into southern Ontario (Godfrey 1966), our study area in northern Ohio was sufficiently close to the northern edge of the range that yearly fluctuations were probable. During our 30-yr study, more than 4,000 observations of screech owls allowed us to assess fluctuations in the composition of the two morphs within the population.




Fig. 8.—Changes in the color phase ratios of screech owls in the northern Ohio study area, 1944-73 (mean with 95% CI).

The population during the first 8 yr of the study (1944-51) consisted of about 25% red-phase birds. During December 1951, the percentage of reds decreased significantly (Table 19, Fig. 8). No significant annual change in the color phase ratio occurred during the remainder of the 30-yr period, although the red phase seemed to increase slightly during the early and mid-1960's. Owen's (1963b) suggestion that environmental gradients were probably responsible for the clinal pattern observed in North America led us to review weather data. The Toledo weather station (Toledo Express Airport) is within the study area and monthly summaries for the 30 winters (November-April) were obtained for that station from the National Weather Records Center (Asheville, N.C.). The deepest snowfall during the 30-yr study occurred in December 1951 as a snowcover of 10.16 cm (4 in) or more lasted for 17 days, averaging 21.34 cm (8.4 in) (25.40 to 33.02 cm [10-13 in] on the ground for 6 days). The snow was accompanied by below-normal temperatures; the lows during the first 15 days of the snowcover averaged -15C and for 4 days the temperature was below -18C.

Table 19. *A summary of the color phase percentages of screech owls by 2-year periods in the Ohio study area. (Includes retrapped birds banded in previous years.)*

Years	Percent red	Percent gray	Number observations	Standard deviation	Percent red 95% C.I.
1944-45	26.8	73.2	149	3.63	19.7 to 33.9
1946-47	25.5	74.5	267	2.67	20.3 to 30.7
1948-49	26.2	73.8	267	2.69	20.9 to 31.5
1950-51	23.3	76.7	275	2.55	18.3 to 28.3
1952-53	14.7	85.3	537	1.53	11.7 to 17.7
1954-55	9.8	90.2	695	1.13	7.6 to 12.0
1956-57	13.0	87.0	300	1.94	9.2 to 16.8
1958-59	9.3	90.7	323	1.62	6.1 to 12.5
1960-61	16.5	83.5	176	2.80	11.0 to 22.0
1962-63	13.6	86.4	206	2.39	8.9 to 18.3
1964-65	16.7	83.3	120	3.40	10.0 to 23.4
1966-67	19.2	80.8	156	3.15	13.0 to 25.4
1968-69	12.3	87.7	179	2.46	7.5 to 17.1
1970-71	9.8	90.2	174	2.25	5.4 to 14.2
1972-73	11.9	88.1	218	2.19	7.6 to 16.2

The combination of heaviest snowfall during the study and extremely cold weather seemed to be correlated with the only significant change in color phase ratios during the 3 decades. Thus, we hypothesize that heavy snowfall in combination with low temperatures plays a direct or indirect role in the relative abundance of the two morphs in the northern portion of the range. The gray phase, which is more abundant in the northern latitudes where winters are more extreme, seemed to survive the critical period much better than the red phase.

Retrap records of adult screech owls banded on the study area before the extreme winter provided information about relative survival of the two color phases during December 1951. Adult birds banded during 1947-50 were used in the analysis, and retrap records during 1952 or later were considered evidence of possible differential survival of the two morphs during the critical period of December 1951 (Table 20). This admittedly small sample suggests that 44% ($10.8 - 6.0 / 10.8 = 44\%$) more red-phased birds than grays were lost during the interval which corresponds quite well with the drop in the ratio of reds from 1950-51 to 1952-53 (change from 23.3 to 14.7, a drop of 37% [$23.3 - 14.7 / 23.3 = 37\%$]).

Table 20. *A comparison of adult screech owls banded in 1947-50 and retrapped in 1952 or latter^a.*

Category	Red	Gray
Number banded	50	157
Number retrapped	3	17
Percentage retrapped	6.0	10.8

^aOnly banded birds retrapped at least once after banding were included (they may have been retrapped initially at any time).

A comparison between the two phases was then made for adult birds banded between 1953 and 1970 to determine if the gray phase birds were always retrapped at a higher rate than the red phase. The percentage of birds retrapped 2 or more yr after banding was nearly identical for both color phases during the 18-yr period (Table 21). This lends further support for our differential survival hypothesis for December 1951. Differential survival, of course, would lead to a change in the ratio of the two morphs. No significant change in the ratio of the two morphs was observed between 1952-53 and 1972-73 when each morph was retrapped in nearly identical percentages.

Table 21. *A comparison of adult screech owls banded in 1953-70 and retrapped 2 or more years after banding^a.*

Category	Red	Gray
Number banded	120	831
Number retrapped	9	65
Percentage retrapped	7.5	7.8

^aOnly banded birds retrapped at least once after banding were included (they may have been retrapped initially at any time).

It is tempting to speculate about the population composition over the last 40 or 50 yr in light of snowfall patterns. Based on records for the study area during the last 85 yr, snowfall was less than normal during 1920-50 and more than normal during the last 20 yr. The less than normal snowfall during the 25 yr before the study began may have allowed the red phase to increase in abundance; then, the severe weather in December 1951 directly or indirectly seems to have caused a decrease in the red-phased segment of the population. The more than average snowfall during the next 20 yr may be responsible for maintaining the population composition at the lower ratio of red to gray phases, although a hint was present that the red phase attempted to increase in abundance at times.

The ruffed grouse (*Bonasa umbellus*), another polymorphic species in North America, has a somewhat similar clinal pattern in color phase ratios (red and gray) within its range and some knowledge is available for this species. Gordon W. Gullion has worked for many years with survival rates of grouse, including survival related to color phase. Gullion and Marshall (1968:157) stated, "As a generalization, the prevalence of red-phase grouse in the population increases as the climate grades from the Arctic south. This suggests that the difference in mortality may be related to the severity of the winter; or more specifically, that there may be a color-linked susceptibility to cold or other climatic conditions, which selects against the red-phase grouse." Gullion (1966) indicated that during winters which lacked sufficient snow for burrowing, mortality among red-phase grouse was greater than among gray birds, but with better snow conditions the red-phase birds outlived the gray.

Gullion and Marshall (1968:157) indicated, "Earlier we believed that this differential survival represented periodically increased vulnerability to predation among red-phase birds (primarily as a result of insufficient burrowing snow), and we still cannot discount this idea entirely. . . . We have suggested that the red-phase

grouse may be more conspicuous to raptors hunting overhead than are the gray-phase birds." Thus, in summary, the increased mortality of the red-phase grouse may be due to a color-linked susceptibility to severe weather or a predation problem associated with color-linked vulnerability. The same conclusions seem to be applicable to the screech owl; in years of heavy snowcover, the red phase screech owls appear to be more prone to mortality because they have not developed the burrowing behavior of the ruffed grouse. The most important natural predator of the screech owl would probably be the great horned owl (Craighead and Craighead 1956), but we are doubtful that predators over a very short term could be responsible for the major change in color phase ratios in 1951. Lustick (1969) studied the effects of artificial radiation on bird energetics and concluded that radiant energy reduces oxygen consumption in birds and that the reduction is correlated with feather color, being greater in dark birds. Mosher and Henny (in press) recently conducted metabolic experiments with screech owls to attempt to ascertain the mechanism responsible for the change in color phase ratios in the population in 1951 by exposing birds of the two color phases to various temperature regimes. These data demonstrated a significant difference in oxygen uptake between color phases at -10 and -5°C. This supports the hypothesis that red-phase screech owls are restricted in their northern distribution by color related metabolic differences from the gray-phase birds. It also suggests that the extremely cold temperatures for a prolonged period of time in December 1951 were responsible for excessive mortality of red-phase birds.

Sex, Age, and Color Phase

Baird et al. (1874:51) first concluded that plumage color was independent of sex and age when they stated, "These two very different plumages are entirely independent of age, sex, or season, and that they are purely individual there can be no doubt, since in one nest there may often be found both red and gray young ones, while their parents may be either both red or both gray, the male red and the female gray, or *vice versa*." To our knowledge a large sample of birds from a given area has not been available to determine if the color is independent of sex. Hasbrouck (1893) presented data showing the relation of color to sex based on 646 birds from throughout the breeding range, but since it is doubtful that each sex was sampled equally throughout the species range, valid comparisons could not be made. Martin (1950) compared 75 study skins from Ontario and found no significant relationship

between sexes and color phases. Similarly, we detected no significant relationship between sex and color phase in our sample of 760 nesting screech owls (Table 22). Fewer males were in the sample because they were often away from the nest at the time it was visited.

Table 22. *The color phase composition of adult nesting screech owls, 1944-73 (includes pairs in which the male was unknown).*

	Gray	Red	Percent red ^a
Males	220	40	15.4
Females	403	97	19.4

^aNo significant difference in color phase composition of nesting males and females (chi-square 1.87, 1 df NS). The intermediates were not included here.

The red and gray birds were found in almost identical ratios in both breeding populations and the young birds produced (Table 23). The intermediates were also consistent in both age categories (2.6 and 2.3%).

Table 23. *Classification of screech owl color morphs from northern Ohio. The data include only pairs in which the color phase of both parents was known. Percentages are in parentheses.*

Category	Red	Intermediate	Gray	Total
1944-73				
Breeding population	81 (16.1)	13 (2.6)	408 (81.3)	502 (100.0)
Offspring	126 (15.3)	19 (2.3)	681 (82.4)	826 (100.0)
Totals				
Including intermediates	207 (15.6)	32 (2.4)	1,089 (82.0)	1,328 (100.0)
Excluding intermediates	207 (16.0)	—	1,089 (84.0)	1,296 (100.0)
1952-73				
Breeding population	62 (15.4)	12 (3.0)	328 (81.6)	402 (100.0)
Offspring	91 (14.1)	17 (2.6)	536 (83.2)	644 (99.9)
Totals				
Including intermediates	153 (14.6)	29 (2.8)	864 (82.6)	1,046 (100.0)
Excluding intermediates	153 (15.0)	—	864 (85.0)	1,017 (100.0)

Genetic Hypotheses

Hrubant (1955) analyzed screech owl data by the gene frequency method to test whether the matings occurred at random, and whether the offspring were produced in agreement with any of several genetic hypotheses. The genetic hypotheses tested were: (1) one pair of genes with dominance, (2) one pair of genes without dominance, (3) multiple alleles, and (4) sex associated inheritance. He concluded that lack of dominance may be dismissed as an acceptable mode of inheritance since neither of the ways in which this mode could be applied could account for the distribution of colors of the young produced and that none of the sex-associated modes are acceptable since the two attributes, color and sex, are independent of each other. This left Hrubant (1955) with but two modes of inheritance as possible explanations of the observed data. In testing the assumption of simple dominance, it was noted that red by red produce only red. Some grays were expected, but the lack of grays may be attributed to the small sample analyzed (only 10 offspring). Under the hypothesis of multiple alleles, the observed value of the intermediate class from red by gray matings deviated from the hypothetical value by a significant amount; however, small samples (only 10 offspring from red by red and 55 offspring from red by gray) made Hrubant's findings somewhat inconclusive.

One Pair of Genes with Dominance

Using the same approach as Hrubant, but with considerably larger sample sizes, we again tested the hypothesis of simple dominance. The method requires the assumption that the population is in equilibrium with respect to mating type, phenotype, and genotype. The significant change in color phase ratios between 1951 and 1952 suggested that the data collected during the first 8 yr of the study (1944-51) should not be pooled with the latter data. Therefore, genetic hypotheses were tested with information collected during the last 22 yr of the study (1952-73). To test this hypothesis the owls may be classified merely as red or gray, and the difference assumed to be caused by one pair of genes with the gene for red exhibiting dominance over the gene for gray. The intermediate owls may be regarded as misclassified gradations of the red and gray. Removal of the intermediates from the sample total in Table 23 leaves 864 gray and 153 red owls. Since an overwhelming number of the matings were gray by gray (Table 24), and since these matings produced only gray offspring, the gray phenotype may be assumed to be due to the recessive gene, and red due to its dominant allele.

Table 24. *Mating types of progeny of screech owls in northern Ohio with respect to color, 1952-73.*

Mating type adult	Number of successful matings	Progeny		
		Red	Inter.	Gray
1. Red × red	8	23	0	5
2. Red × gray	46	68	1	63
3. Red × Inter.	0	0	0	0
4. Inter. × Inter.	0	0	0	0
5. Inter. × gray	12	0	16	29
6. Gray × gray	135	0	0	439
Totals	201	91	17	536

Let R stand for the dominant gene for red pigment and r stand for its recessive allele for gray pigment. Let p be the frequency of r in the sample, such that $p + q = 1$. Values of q and p may be obtained from the sample so that they are the maximum likelihood estimates of θ and $1-\theta$, the true frequencies of r and R in the population.

Following the discussion by Hrubant (1955), the estimates may be based upon the parents only, the offspring only, or upon the entire sample of 1,017 owls, omitting the intermediates. The entire sample of 1,017 appears to be the best choice since there is no marked evidences of heterogeneity within it and since the parents and offspring contain about the same frequencies of red and gray owls. Of 1,017 owls in the entire sample (omitting intermediates), 864 were gray. The frequency of r thus $q = \sqrt{864/1,017} = 0.9217$ and of R is $p = 0.0783$.

The theoretical mating type frequency, determined by the expansion of the binomial $((1-\theta) + \theta)^4$ using $q = 0.9217$ as an estimate of θ , and the observed mating frequencies are compared in Table 25. The data in this table show that the various kinds of matings occurred with frequencies in agreement with those expected if mating is at random. The computed chi-square is smaller (3.30) than the critical (5.99) at the 5% level of significance of 2 degrees of freedom.

Table 25. *Testing specific mating types of screech owls under hypothesis of one pair of genes with dominance.*

Mating type	Theoretical frequency	Expected proportions	Expected frequency	Observed frequency
1. Red × red RR × RR	$(1-\theta)^4$	0.00004	0.02264	4.3
2. Red × red RR × Rr	$4\theta(1-\theta)^3$	0.00177		
3. Red × red Rr × Rr	$4\theta^2(1-\theta)^2$	0.02083		
4. Red × gray RR × rr	$2\theta^2(1-\theta)^2$	0.01042	0.25566	48.3
5. Red × gray Rr × rr	$4\theta^3(1-\theta)$	0.24524		
6. Gray × gray rr × rr	θ^4	0.72170	0.72170	136.4
		1.00000	189.0	189
$X^2 = 3.30, 2 \text{ d.f.}$				

Table 26 contains the test of the observed distribution of young produced by individual mating types compared to their expected distribution. The expected numbers of offspring are found from the relative frequencies of the matings given in Table 25. The data show that in all three mating combinations the observed numbers of offspring occur with frequencies in agreement with those calculated if the difference between red and gray is due to only one pair of genes with red dominant to gray. In fact, the observed and expected frequencies for red by gray and gray by gray are in complete agreement, whereas the red by red pairings are in nearly complete agreement (chi-square 0.22). The critical level for rejection of the hypothesis is 3.84 at the 5% level of significance for 1 degree of freedom. The observed data are nearly a perfect fit.

Table 26. *Distribution of young screech owls produced by mating phenotypes under hypothesis of one pair of genes with dominance.*

Mating	Observed number offspring		Expected number offspring		X ²
	Red	Gray	Red	Gray	
1-3 Red × red	23	5	21.6	6.4	0.22
4-5 Red × gray	68	63	68.2	62.8	—
6 Gray × gray	0	439	0	439.0	—

Multiple Alleles

The other possible mode of inheritance (Hrubant 1955) is a multiple allelic system. Assume the red, intermediate, and gray phenotypes are inherited in this manner with a graded order of dominance of Y (red) dominant to y' (intermediate) dominant to y (gray). Table 27 illustrates the observed and expected sample frequencies and theoretical population frequencies used in deriving the values for the statistics $r = 0.0760$, $s = 0.0152$, and $t = 0.9088$, as estimates of the true frequencies of genes.

Table 27. *Sample frequencies and estimates of population gene frequencies of screech owls under hypothesis of a system of multiple alleles.*

Genotype	Theoretical frequency	Observed frequency	Expected frequency
YY	p^2		0.0058
Yy' Red	$2p\sigma$	0.146	0.0023
Yy	$2p\gamma$		0.1381
y'y'	σ^2		0.0002
Inter.		0.028	
y'y	$2\sigma\gamma$		0.0276
yy Gray	γ^2	0.826	0.8259
$t = \sqrt{864/1046} = 0.9088$, frequency of y(γ) $s = \sqrt{864+29/1046} - 0.9088 = 0.0152$, frequency of y'(σ) $r = 1 - (t+s) = 0.0760$, frequency of Y(p)			

The theoretical mating type frequency determined by the expansion of the trinomial $(p + \sigma + \gamma)^4$ and the observed mating type frequencies are compared in Table 28. The data in this table show that the various kinds of matings occurred with frequencies in agreement with those expected if mating is at random. The computed chi-square (5.99) is smaller than the critical (11.07) at the 5% level of significance for 5 degrees of freedom; therefore, we concluded that the matings were at random.

Table 28. *Testing specific mating types of screech owls under hypothesis of a system of multiple alleles.*

Mating type by phenotypes	Theoretical frequency	Expected proportions	Expected frequency	Observed frequency
1. Red × red	$p^4 + 4p^3\sigma + 4p^3\gamma + 4p^2\sigma^2 + 8p^2\sigma\gamma + 4p^2\gamma^2$	0.0213	4.3	8
2. Red × gray	$2p^2\gamma^2 + 4p\sigma\gamma^2 + 4p\gamma^3$	0.2415	48.5	46
3. Red × inter.	$2p^2\sigma^2 + 4p\sigma^3 + 12p\sigma^2\gamma + 4p^2\sigma\gamma + 8p\sigma\gamma^2$	0.0081	1.6	0
4. Inter. × inter.	$\sigma^4 + 4\sigma^3\gamma + 4\sigma^2\gamma^2$	0.0008	0.2	0
5. Inter. × gray	$2\sigma^2\gamma^2 + 4\sigma\gamma^3$	0.0460	9.2	12
6. Gray × gray	γ^4	0.6821	137.1	135
		0.9998	200.9	201
			$X^2 = 5.99, 5 \text{ d.f.}$	

The observed distribution of young produced by the specific phenotypic mating types as compared to their expected distribution is shown in Table 29. The data show agreement in all cases between the frequencies of the observed numbers of offspring and those calculated if the difference between red, intermediate, and gray were due to a multiple allelic genetic system. This system involves a graded order of dominance of red over intermediate over gray.

Table 29. *Distribution of young screech owls produced by mating phenotypes under hypothesis of a system of multiple alleles.*

Mating	Observed numbers offspring			Expected numbers offspring			X ²
	Red	Inter.	Gray	Red	Inter.	Gray	
1. Red × red	23	0	5	21.3	0.1	6.6	0.63
2. Red × gray	68	1	63	68.6	1.1	62.3	0.02
3. Red × inter.	—	—	—	—	—	—	—
4. Inter. × inter.	—	—	—	—	—	—	—
5. Inter. × gray.	0	16	29	0	22.3	22.7	3.53
6. Gray × gray	0	0	439	0	0	439.0	—

Unfortunately then, neither of the two hypotheses of gene action (one pair of genes with dominance and multiple alleles) that appeared satisfactory to Hrubant (1955) can be eliminated on the basis of breeding data presented here. In fact, the evidence for the two hypotheses was strengthened by the new data as the expected values were in nearly perfect agreement with the observed breeding data. The fact that red by red produced approximately 25% grays was a most important finding because Hrubant noted that red by red produced only reds in his small sample. Therefore, the following hypotheses remain for the gene action: (1) If the intermediate phenotype is assumed to be due to genetic modifiers, the difference between red and gray is due to one pair of alleles, the gene for red exhibiting dominance over its allele for gray; (2) if all three colors are considered, the colors are inherited on the basis of a series of three alleles with a graded order of dominance of red → intermediate → gray.

Brood Size and Color Phase

Data were insufficient to provide a fledging rate estimate (per nesting attempt) for each color phase; however, a pooled estimate of the number of young fledged per successful nest was obtained. The color phase of both parents was documented for 211 successful nesting attempts during 1944–72 (Table 30). The data for 1973 were not included because one egg was taken from each nest for pesticide analysis. No significant difference in the number of young fledged for the various breeding combinations could be detected.

Table 30. *A summary of the estimated number of young screech owls fledged per successful nest for the various combinations of color phase pairs, 1944-72.*

Category	Young fledged						Number nests	Young fledged per successful nest	95% C.I.
	1	2	3	4	5	6			
Gray × gray	18	7	35	49	30	8	147	3.61	3.39-3.83
Gray × red	1	5	8	15	13	1	43	3.86	3.52-4.20
Red × red	1	0	1	4	3	0	9	3.89	3.06-4.72
Inter. × gray	1	0	1	5	3	1	11	4.09	3.32-4.86
Inter. × red	0	0	0	1	0	0	1	4.00	—
Total	21	12	45	74	49	10	211	3.70	

STATUS OF POPULATION

The status of any wildlife population may be evaluated in three general ways: (1) a complete count or direct enumeration on an annual basis, (2) an index to abundance at some time period during the life cycle, or (3) a life-equation approach which provides an indirect evaluation of the population's condition. These approaches are not new and fall within the general outline presented by Leopold (1933:139): "(1) *Census* (Measuring the Stock on Hand), and (2) *Measuring the Productivity* of the stock and comparing it with a standard." Furthermore, it is always useful to compare results obtained from several approaches. The complete count or direct enumeration of screech owls is impossible due to their habitat preferences. Therefore, we will rely upon an index to annual abundance and the life-equation approach to evaluate the status of the screech owl population in northern Ohio.

Index to Annual Abundance

The occupancy rate of wood duck boxes by screech owls during the nesting season may provide some insight into the relative annual abundance of the species on the study area. However, since the nest boxes were always placed in suitable habitat and loss of habitat has occurred on the study area (Table 1), the occupancy rate for nest boxes would provide only an index to abundance in suitable habitat. Thus, the possibility of an undetected decline due to loss of habitat exists. To further complicate the matter, we have been forced to use an index to annual abundance based on successful nests. This modification was necessary because during

some years the boxes were visited early in the nesting season (a chance to observe nests that were later abandoned and unsuccessful), whereas in other years nests were visited only at banding time (successful nests only recorded). Therefore, because of the fluctuations in annual effort we believe that a more realistic annual comparison of abundance would be obtained from successful nests only.

Although indices based on wood duck box usage have many potential biases, two factors present during this study tend to minimize these biases: (1) the same observer checked the boxes annually, and (2) approximately the same number of boxes were available each year.

The information collected during the 30-yr period suggested no long-term trend in abundance in suitable habitat, as measured by numbers of successful nests (Table 31). The 4 yr (1952-55) when students assisted in the study could not be evaluated because the number of nests checked was unknown. Excluding those 4 yr, the 10-yr indices to abundance were as follows: 1944-51, 10.1; 1956-63, 8.4; and 1964-73, 9.0. Successfully nesting screech owls were found annually in approximately 8.9% of the nest boxes during the 30-yr study. Assuming that 69.2% of the screech owl nesting attempts were successful (see section on BREEDING BIOLOGY), another 4.0% of the nest boxes were used annually, but without success, making a total of about 12.9% of the boxes utilized annually during the study.

Table 31. *An index to the abundance of screech owls in suitable habitat in northern Ohio, 1944-73.*

Year ^a	No. of boxes checked	No. of successful nests	Percent with successful nests
1944	45	4	9
1945	78	4	5
1946	95	6	6
1947	128	14	11
1948	133	11	8
1949	139	15	11
1950	140	24	17
1951	145	10	7
Total 1944-1951	903	88	10.1 ^b
1956	210	20	10
1957	208	14	7
1958	190	14	7
1959	180	12	7
1960	155	11	7
1961	140	9	6
1962	130	18	14
1963	125	11	9
Total 1956-63	1,338	109	8.4 ^b
1964	125	9	7
1965	135	7	5
1966	130	9	7
1967	130	16	12
1968	130	13	10
1969	130	10	8
1970	130	12	9
1971	130	11	8
1972	135	16	12
1973	153	19	12
Total 1964-73	1,328	122	9.0 ^b
Total 1944-73	3,569	319	8.9 ^b

^aNumbers of boxes checked in 1952-55 are unknown.^bEach year given equal weight.

Life Equation Approach

A mathematical model showing the relations between population parameters that yield stable populations was developed by Henny et al. (1970). Information needed for the model includes (1) mortality rate schedules (obtained from recoveries of banded birds), (2) recruitment rates, and (3) age at sexual maturity. Estimates of these parameters have been presented in this paper for the screech owl.

We know that screech owls are capable of breeding at the end of their first year of life (as 1-yr-olds), although the percentage in this age class that breeds is not known with certainty. However, we still believe that the exercise in constructing a life-equation model is worthwhile, because it will point out future research needs and act as a check on the internal consistency of the input data. Adult mortality rate estimates for the screech owl range from 32.8 to 39.0% (Table 15), with a pooled estimate of 33.9%. The mortality rate estimate for the first year of life was 69.5% (Table 15). The observed recruitment rate per breeding pair was estimated at between 2.55 (Table 13) and 2.63 based on the exposure-day method.

Initially, let us assume that all screech owls breed as 1-yr-olds, although we have some evidence that a portion of the 1-yr-olds do not nest. The estimated number of young that must be fledged per breeding-age pair to maintain a stable population may be estimated by the following formula (from Henny 1972):

$$\bar{m} = \frac{1 - s}{s_0 (1 - s + s_1)}$$

where

\bar{m} = the average number of female fledglings produced per breeding-age female ($2\bar{m}$ = the total number of young produced per breeding female assuming an equal sex ratio of fledglings)

s = third year and later survival rate

s_0 = first year survival rate

s_1 = second year survival rate

By using the pooled estimate for adult mortality and assuming that all 1-yr-olds attempt to nest, it is estimated that only 2.22 young are required per breeding pair to maintain a stable population. This is slightly less than the observed recruitment (2.55 to 2.63), and, although the mortality rates may not be precise, the modeling approach suggests that not all 1-yr-old screech owls need to attempt

to nest annually in order to maintain a stable population. To pursue this matter a little further, let us assume for the moment that the recruitment rates and the mortality rates are correct and, through the modeling process, let us calculate the percentage of 1-yr-olds (solve for p_1) that must attempt to nest annually to balance the life equation (see Equation 1 in Henny 1972:6). Given the above constraints and assuming that 2-yr-olds and older attempt to nest each year, the population could remain stable if at least an average of 60% of the 1-yr-olds nested annually. This estimate is in fairly close agreement with the 77 to 83% estimate based on a small sample of field data which may be biased upward.

Summary of Status

In summary, we believe that the screech owl population in northern Ohio has fluctuated in abundance during the last 30 yr, with no long-term trend being apparent in suitable habitat (Table 31). In addition, the recruitment rate observed during the long-term study seemed to be adequate to maintain the population numbers. We have indirect evidence that all 1-yr-olds are not required to nest (from modeling approach) and field evidence suggesting that a sufficient percentage of 1-yr-olds are nesting. Furthermore, we have no evidence of eggshell thinning in the population (Table 17), although laboratory tests have shown that the species is susceptible to the eggshell thinning phenomenon. Residue levels of DDT and its metabolites along with PCB's were low (Table 18). It should be cautioned, however, that even though screech owls seem to be adaptable to people and their structures, continued habitat loss may gradually reduce the population numbers in northern Ohio as the woodlots and creek bottoms become cleared and farmed more intensively. The forested lands in the four-county study decreased 12% between 1958 and 1967.

SUMMARY

1. The life history and population ecology of the screech owl was studied in northern Ohio between 1944 and 1973. The owls nested in boxes established for wood ducks. The birds were banded and periodically recaptured. Food habits, productivity information, and color phase of the parent birds and offspring were recorded.
2. The screech owl is an opportunistic feeder. The diet changes with the seasons of the year, i.e., during the nesting season migrant birds replace mammals in importance, and during the late summer insects become important.
3. There is no evidence from banding data to suggest that screech owls in the northeastern United States migrate.
4. Young screech owls begin dispersing from their natal areas in late summer or early fall, with only about one-fourth of the young birds remaining within 10 km (6 mi) of the banding site. On the contrary, adult birds remain close to the area where they previously nested.
5. The hypothesis of random dispersal distance in young screech owls was rejected. There appear to be two groups in the population: (1) a group showing little dispersal, and (2) a group wandering considerable distances.
6. The directional pattern of dispersal was random.
7. Pairs of screech owls were noted in the nest boxes in early February, but egg laying did not peak until about 15 March. Hatching takes place in mid-April to early May and most of the young leave the nest the last week of May or the first week in June.
8. The mean clutch size was 4.43 with a mean of 3.80 young fledged per *successful* nest. An estimated 69.2% of the nesting attempts were successful. Including unsuccessful nesting attempts, an estimated 2.55 to 2.63 young were fledged per *breeding pair*.
9. Annual variation in the number of young fledged per successful nest was small. The lack of variability was probably due to the predictable food source of passerine birds that migrated through the area each spring when young were in the nest. This contrasts with the highly variable productivity reported for owls dependent upon local cyclic rodent populations.

10. Raccoon predation was implicated in a number of unsuccessful nesting attempts, but the overall effect of the raccoon on the screech owl population was not fully assessed.
11. The sex ratio in the population **did not** differ significantly from a 50:50 ratio.
12. The first year mortality rate of the screech owl was estimated at 69.5%; adult mortality was estimated at 33.9%.
13. Male and female screech owls banded as nestlings were known to nest successfully at the end of their first year of life (as 1-yr-olds). Two estimates of the average proportion of 1-yr-olds nesting ranged from 77 to 83%, but the percentage may be lower.
14. No significant eggshell thinning was detected in the population in 1973, and residue levels of DDE and PCB's were low.
15. In northern Ohio, the screech owl population consists mainly of a gray- and red-phased birds, but about 2 to 3% of the birds are intermediate in color.
16. A significant change in the ratio of red- to gray-phased birds occurred in December 1951. This change coincided with the lowest temperatures during the 30-yr study and the heaviest snowfall. The red-phased birds decreased from 23.3% of the population to 14.7%. Additional evidence from retrap information suggests that the grays survived much better than the reds during the stressful period. The ratio of reds in the population failed to increase to pre-1951 levels during the next 20 yr.
17. No significant relationship existed between sex and color phase in our sample of 760 nesting screech owls.
18. The following hypotheses of gene action remain: (1) If the intermediate phenotype is assumed to be due to genetic modifiers, the difference between red and gray is due to one pair of alleles, the gene for red exhibiting dominance over its allele for gray; or (2) if all three colors are considered, the colors are inherited on the basis of a series of three alleles with a graded order of dominance of red \rightarrow intermediate \rightarrow gray.
19. No significant difference in brood size and color phase of the parents could be detected.
20. An index to annual abundance suggests that the screech owl population fluctuated in northern Ohio during the last 30 yr, but with no long-term trend apparent in available habitat. The life-equation approach also suggests that, over the long term, productivity probably balanced mortality.

ACKNOWLEDGMENTS

The assistance of Michael Smith, Larry Shafer, Lee Garling, Denny Genzman, and Gene Walston in the field aspects of the study during the mid-1950's is appreciated. Informative discussions with Milton B. Trautman and Ronald L. Stuckey of the Ohio State University, and Theodore A. Bookhout of the Ohio Cooperative Wildlife Research Unit, led to the location of some interesting references concerning changes in the flora and fauna of the region. Kenneth P. Burnham of the Migratory Bird and Habitat Research Laboratory assisted in the analysis of directional dispersal, and Erwin E. Klaas of the Patuxent Wildlife Research Center conducted the pesticide analysis. William H. Stickel of the Patuxent Wildlife Research Center brought to our attention the vast food habit files that are stored at the Center. Richard D. Porter of the Denver Wildlife Research Center kindly reviewed the manuscript and Richard Wilmot reviewed the section concerning polymorphism. G. Ronald Austing and George Laycock provided several photographs to illustrate the publication. Our sincere thanks is expressed to all who assisted.

LITERATURE CITED

- Allen, A. A. 1924. A contribution to the life history and economic status of the screech owl (*Otus asio*). *Auk* 41(1):1-16.
- Allen, J. A. 1893. Hasbrouck on "Evolution and dichromatism in the Genus *Megascops*." *Auk* 10(4):347-353.
- American Ornithologists' Union 1957. Check-list of North American birds. Fifth Ed. Lord Baltimore Press. 691 pp.
- Anderson, D. W. and J. J. Hickey 1972. Eggshell changes in certain North American birds. *Proc. Int. Ornithol. Congr.* 15:514-540.
- Anonymous 1971. Ohio soil and water conservation needs inventory. Ohio Soil and Water Conservation Needs Committee. Columbus, Ohio (sponsored by U.S. Dep. Agric.). pp.14-15.
- Baird, S. F., T. M. Brewer, and R. Ridgway 1874. A history of North American birds. Vol. 3. Little, Brown and Co., Boston, 560 pp.
- Bellrose, F. C. 1955. Housing for wood ducks. *Ill. Nat. Hist. Surv., Circ.* 45 (rev.). 48 pp.
- Bellrose, F. C., T. G. Scott, A. S. Hawkins, and J. B. Low 1961. Sex ratios and age ratios in North American ducks. *Ill. Nat. Hist. Surv. Bull* 27(6):391-474.
- Bent, A. C. 1938. Life histories of North American birds of prey. U.S. Natl. Mus. Bull. 170 (Repr. by Dover Publ. Co., N.Y. 1961). 482 pp.
- Burton, J. A., ed. 1973. Owls of the world: their evolution, structure and ecology. E. P. Dutton Co., New York. 216 pp.
- Campbell, L. 1968. Birds of the Toledo area. *The Blade*, Toledo, Ohio. 330 pp.
- Carpenter, F. H. 1883. Screech Owls breeding in confinement. *Ornithol. and Oologist* 8(12):93-94.
- Craighead, J. J. and F. C. Craighead, Jr. 1956. Hawks, owls, and wildlife. Stackpole Co., Harrisburg, Pa. 443 pp.
- Dice, L. R. and W. E. Howard 1951. Distance of dispersal by prairie deer mice from birthplace to breeding sites. *Contri. Lab. Vertebr. Biol. Univ. Mich.* 50:1-15.
- Diller, O. D. 1962. More land in trees. *The Ohio Farmer*. May 5. p. 29, 49.
- Emlen, J. T., Jr. 1940. Sex and age ratios in survival of the California quail. *J. Wildl. Manage.* 4(1):92-99.
- Errington, P. L. 1932. Food habits of southern Wisconsin raptors, Part I. Owls. *Condor* 34(4):176-186.
- Fisher, A. K. 1893. Hawks and owls of the U.S. U.S. Dep. Agric., Div. Ornithol. Mammal., Bull. 3:163-174.
- Fisher, R. A. 1930. The genetical theory of natural selection. London, Oxford Univ. Press. 291 pp.
- Ford, E. B. 1940. Polymorphism and taxonomy. Pages 493-513 in J. Huxley, ed. *The New Systematics*. Clarendon Press, Oxford. 583 pp.
- Frazar, A. M. 1877. The mottled owl as a fisherman. *Bull. Nuttall Ornithol. Club* 2(2):80.
- French, N. R., T. Y. Tagami, and P. Hayden 1968. Dispersal in a population of desert rodents. *J. Mammal.* 49(2):272-280.
- Godfrey, W. E. 1966. The birds of Canada. *Natl. Mus. Can. Bull. Biol. Ser.* 73, 203. Queen's Printer, Ottawa. 428 pp.

- Gullion, G. W. 1966. The use of drumming behavior in ruffed grouse population studies. *J. Wildl. Manage.* 30(4):717-729.
- Gullion, G. W. and W. H. Marshall 1968. Survival of ruffed grouse in a boreal forest. *Living Bird* 7:117-167.
- Hasbrouck, E. M. 1893. Evolution and dichromatism in the genus *Megascops*. *Am. Nat.* 27(6-7):521-533, 638-649.
- Heath, R. G. and J. W. Spann 1973. Reproduction and related residues in birds fed mirex. *Inter-Am. Conf. Toxicol. Occup. Med.* 8:421-435.
- Heath, R. G., J. W. Spann, and J. F. Kreitzer 1969. Marked DDE impairment of mallard reproduction in controlled studies. *Nature* 224(5214):47-48.
- Henny, C. J. 1972. An analysis of the population dynamics of selected avian species. U.S. Fish Wildl. Serv., Wildl. Res. Rep. 1. 99 pp.
- Henny, C. J. and W. T. VanVelzen 1972. Migration patterns and wintering localities of American ospreys. *J. Wildl. Manage.* 36(4):1133-1141.
- Henny, C. J., W. S. Overton, and H. M. Wight 1970. Determining parameters for populations by using structural models. *J. Wildl. Manage.* 34(4):690-703.
- Hickey, J. J. 1952. Survival studies of banded birds. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Wildl. 15. 177 pp.
- Hickey, J. J., ed. 1969. Peregrine falcon populations their biology and decline. Univ. of Wisconsin Press, Madison. 596 pp.
- Hickey, J. J. and D. W. Anderson 1968. Chlorinated hydrocarbons and eggshell changes in raptorial and fish-eating birds. *Science* 162(3850):271-273.
- Howard, W. E. 1960. Innate and environmental dispersal of individual vertebrates. *Am. Midl. Nat.* 63(1):152-161.
- Hrubant, H. E. 1955. An analysis of the color phases of the eastern screech owl, *Otus asio*, by the gene frequency method. *Am. Nat.* 89(4):223-230.
- Huxley, J. 1955. Morphism in birds. *Acta Inter. Ornithol. Congr.* (1954) 11:309-328.
- James, R. F. and N. D. Martin 1950. A study of screech owls in southern Ontario. *Can. Field Nat.* 64(5):177-180.
- Johnston, R. F. 1956. Population structure in salt marsh song sparrows. Part I. Environment and annual cycle. *Condor* 58(1):24-44.
- Kaatz, M. R. 1953. The settlement of the Black Swamp of northwestern Ohio. *Northwest Ohio Q.* 25:23-36, 134-156, 201-217.
- Kelso, L. H. 1944. Behavior of the eastern screech owl (*Otus asio naevius*). *Biol. Leafl.* 23. Cornell Univ. 7 pp.
- Kelso, L. H. 1950. The post juvenile molt of the northeastern screech owl. *Biol. Leafl.* 50. Cornell Univ. 3 pp.
- Korschgen, L. J. and H. B. Stuart 1972. Twenty years of avian predator-small mammal relationships in Missouri. *J. Wildl. Manage.* 36(2):269-282.
- Lack, D. 1966. Population studies of birds. Clarendon Press. Oxford. 341 pp.
- Lack, D. 1968. Ecological adaptations for breeding in birds. Methuen & Co., Ltd., London. 409 pp.
- Leopold, A. 1933. Game management. Charles Scribner's Sons. New York. 481 pp.
- Lidicker, W. Z., Jr. 1962. Emigration as a possible mechanism permitting the regulation of population density below carrying capacity. *Am. Nat.* 96(1):29-33.
- Lockie, J. D., D. A. Ratcliffe, and R. Balharry 1969. Breeding success and organochlorine residues in golden eagles in west Scotland. *J. Appl. Ecol.* 6(3):381-389.
- Longcore, J. R., F. B. Samson, and T. W. Whittendale, Jr. 1971. DDE thins eggshells and lowers reproductive success of captive black ducks. *Bull. Environ. Contam. Toxicol.* 6(6):485-490.

- Lustick, S. 1969. Bird energetics: effects of artificial radiation. *Science* 163(3865):387-390.
- Marshall, J. T., Jr. 1967. Parallel variation in north and middle American screech owls. *West. Found. Vertebr. Zool., Monogr.* 1. 72 pp.
- Martin, N. D. 1950. Colour phase investigations on the screech owl in Ontario. *Can. Field Nat.* 64(6):208-211.
- Mayfield, H. 1961. Nesting success calculated from exposure. *Wilson Bull.* 73(3):255-261.
- Mayfield, H. 1962. Changes in the natural history of the Toledo region since the coming of the white man. *Jack-Pine Warbler* 40(2):36-52.
- McDowell, R. D. and L. A. Luttringer, Jr. 1948. Pennsylvania birds of prey. *Pa. Game Comm., Harrisburg.* 32 pp.
- McLane, M.A.R. and L. C. Hall 1972. DDE thins screech owl eggshells. *Bull. Environ. Contam. Toxicol.* 8(2):65-68.
- Mosher, J. A. and C. J. Henny. (In Press) Thermal adaptiveness of plumage color in screech owls. *Auk* 94.
- Murray, B. G., Jr. 1967. Dispersal in vertebrates. *Ecology* 48(6):975-978.
- Owen, D. F. 1963a. Variation in North American screech owls and the subspecies concept. *Syst. Zool.* 12(1):8-14.
- Owen, D. F. 1963b. Polymorphism in the screech owl in eastern North America. *Wilson Bull.* 75(2):183-190.
- Ratcliffe, D. A. 1967. Decrease in eggshell weight in certain birds of prey. *Nature* 215(5097):208-210.
- Ross, A. 1969. Ecological aspects of the food habits of insectivorous screech owls. *Proc. West. Found. Vertebr. Zool.* 1(6):301-344.
- Schneider, W. 1937. Beringungsergebnisse an der mitteleuropäischen Schleiereule (*Tyto alba guttata* Brehm). *Vogelzug* 8:159-170.
- Schorger, A. H. 1954. Color phases of the screech owl between Madison, Wisconsin, and Freeport, Illinois. *Auk* 71(2):205.
- Sherman, A. R. 1911. Nest life of the screech owl. *Auk* 28(2):155-168.
- Sitterley, J. and J. Falconer 1933. Type of farming areas in Ohio. *Ohio State Univ. and Ohio Agric. Exp. Stn. Bull.* 56 (mimeo).
- Stewart, P. A. 1952. Dispersal, breeding behavior, and longevity of banded barn owls in North America. *Auk* 69(3):227-245.
- Stewart, P. A. 1969. Prey in two screech owl nests. *Auk* 86(1):141.
- Stickel, L. F. 1973. Pesticide residues in birds and mammals. Pages 254-312 in C.A. Edwards, ed. *Environmental pollution by pesticides*. Plenum Press, London and New York. x + 542 pp.
- Stupka, A. 1953. Some notes relating to the mortality of screech owls in Great Smoky National Park. *Migrant* 24(1):3-5.
- Sutton, G. M. 1927. Mortality among screech owls of Pennsylvania. *Auk* 44(4):563-564.
- Trautman, M. B. 1940. The birds of Buckeye Lake, Ohio. *Misc. Publ. Mus. Zool. Univ. Mich.* 44. 466 pp.
- Treat, W. E. 1889. A fishing screech owl. *Auk* 6(2):189-190.
- Vaughan, R. 1961. *Falco eleonora*. *Ibis* 103a:114-128.
- Wallace, G. J. 1963. An introduction to ornithology. Second Ed. MacMillan, New York. 491 pp.
- Wiemeyer, S. N. and R. D. Porter 1970. DDE thins eggshells of captive American kestrels. *Nature* 227(5259):737-738.
- Wilson, K. A. 1938. Owl studies at Ann Arbor, Michigan. *Auk* 55(2):187-197.