|  | OCUMENIAIIUN FA  | UE I  | OMB No. 0704-0188   |
|--|--|---|---|
| the resorting surgen for this conscision of int  |  | sectorized the tame lot the   | wing instructions, searching earling data search.   |
| hering and maintaining the data needed, and<br>ection of information, including suggestions<br>ection of information, including suggestions  | for reducing this burgen, to Washington Head<br>4382, and to the Office of Management and B  | unget, Paserwark Reduction Projec   | (#704-6186), Weshington, OC 20103.  |
| AGENCY USE ONLY (Leave blan  | 2. REPORT DATE 2/21/97   | I JA KEFURL I I FL. MAN   | DATES COVERED<br>/90 - 10/31/96   |
|  |  |   | FUNDING NUMBERS   |
| TITLE AND SUBTITLE   | ,  |   | G   |
| Mesoscale Studies of   | Explosive Cyclogenesis   |   | N00014-91-J-1068 P00007   |
| AUTHOR(S)  |  |   |   |
| Roger Wakimoto   |  |   |   |
| PERFORMING ORGANIZATION N  | AME(S) AND ADDRESS(ES)   |   | 8. PERFORMING ORGANIZATION<br>REPORT NUMBER   |
| University of Califor  | rnia, Los Angelos  |   |   |
| Department of Atmosp   | heric Sciences   |   | A90-1751B-06  |
| 405 Hilgard Avenue   |  |   |   |
| Los Angeles, CA 9009   |  |   | 10. SPONSORING / MONITORING   |
| . SPONSORING/MONITORING AG   |  |   | AGENCY REPORT NUMBER  |
| Department of the Na   | vy<br>arch   |   |   |
| Office of Naval Rese<br>800 North Quincy St.   | al Uli   | ·   |   |
| Arlington, VA 22217  |  |   | 5AA   |
| 1. SUPPLEMENTARY NOTES   | 10   | 970428  | JIN   |
| I. SUPPLEMENIANT RUIES   | 19   | 19/0420 1   |   |
|  | , <b>A</b>   |   |   |
|  | CTATEMENT  | 1   | 126. DISTRIBUTION CODE  |
| 2a. DISTRIBUTION / AVAILABILITY  |  |   |   |
| ADDDOVED FOR PURITC  | RELEASE; DISTRIBUTION  | IS UNLIMITED.   |   |
| WLIVOARD FOR TOPHTO  |  |   |   |
| ALLKOVED LOK LOBIC   |  |   |   |
| ALLVARD LOW LODIEC   |  |   |   |
|  | cs)  |   |   |
| 13. ABSTRACT (Maximum 200 wor  |  | vses of the convec  | tive and frontal  |
| 13. ABSTRACT (Maximum 200 wor<br>During the past four<br>attracture within rai   | years, extensive analy   | s over the Atlanti  | c Ucean has been  |
| 13. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within raj  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have   | s over the Atlanti<br>undergone a scale   | c Ocean has been<br>contraction were  |
| 13. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense   | years, extensive analy<br>oidly deepening cyclone<br>warm fronts that have<br>rst time. In addition.   | s over the Atlanti<br>undergone a scale<br>mesoscale vortice  | c Ocean has been<br>contraction were<br>s, approximately  |
| <b>3. ABSTRACT</b> (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fi:<br>25-40 km wide, were  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along   | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts   | c Ocean has been<br>contraction were<br>s, approximately<br>. When these  |
| <b>13. ABSTRACT</b> (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into a   | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| <b>13. ABSTRACT</b> (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into a   | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| <b>13. ABSTRACT</b> (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into a   | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a  | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| B. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| B. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| B. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| B. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and  |
| B. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and<br>en published in   |
| B. ABSTRACT (Maximum 200 wor<br>buring the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc<br>the literature.   | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and<br>en published in   |
| B. ABSTRACT (Maximum 200 wor<br>During the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc  | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and<br>en published in   |
| B. ABSTRACT (Maximum 200 wor<br>buring the past four<br>structure within ray<br>documented. Intense<br>observed for the fir<br>25-40 km wide, were<br>vortices come into<br>deepening of the cer<br>thermodynamic struc<br>the literature.   | years, extensive analy<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ptral low pressure appe                             | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and<br>en published in   |
| <ul> <li>ABSTRACT (Maximum 200 wor<br/>During the past four<br/>structure within ray<br/>documented. Intense<br/>observed for the fit<br/>25-40 km wide, were<br/>vortices come into<br/>deepening of the cent<br/>thermodynamic struct<br/>the literature.</li> <li>SUBJECT TERMS</li> <li>SECURITY CLASSIFICATION</li> </ul> | years, extensive analyoidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ntral low pressure appe<br>ture of these fronts an      | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The                       | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and<br>en published in<br>15. NUMBER OF PAGES<br>3<br>16. PRICE CODE                                     |
| <ul> <li>ABSTRACT (Maximum 200 wor<br/>During the past four<br/>structure within ray<br/>documented. Intense<br/>observed for the fit<br/>25-40 km wide, were<br/>vortices come into<br/>deepening of the cent<br/>thermodynamic struct<br/>the literature.</li> <li>14. SUBJECT TERMS</li> </ul>                              | r years, extensive anal<br>pidly deepening cyclone<br>warm fronts that have<br>rst time. In addition,<br>shown to develop along<br>a proper phasing with a<br>ntral low pressure appe<br>ture of these fronts an | s over the Atlanti<br>undergone a scale<br>mesoscale vortice<br>these warm fronts<br>baroclinic wave a<br>ars to occur. The<br>d vortices have be | c Ocean has been<br>contraction were<br>s, approximately<br>. When these<br>loft, rapid<br>kinematic and<br>en published in<br>15. NUMBER OF PAGES<br>3<br>16. PRICE CODE<br>CATION 20. LIMITATION OF ABSTRAC |

## UNIVERSITY OF CALIFORNIA, LOS ANGELES

| EGO SAN FRANCECO SANTA BARBARA SANTA CRUZ | RIVERSID | LOS ANGELES | RVINE  | •     | · DAVIS | ERKELSY |
|---|----------|-------------|--------|-------|---------|---------|
|   |          | ······      |        |       |         |         |
|   | + =      | ÷ .         | 144 A. |       |         | -       |
| DEPARTMENT OF ATMOSPHERIC SCIENCES        |          |             |        | ~ · · |         |         |
| 405 HILGARD AVENUE                        |          |             |        |       |         |         |
| LOS ANGELES, CALIFORNIA 90095-156         |          |             |        |       |         |         |
| (310) 825-1751                            |          |             |        |       |         |         |
| INTERNET: ROGER @ ATMOS.UCLA.EDU          |          |             |        |       |         |         |
| FAX: (310) 206-5219                       |          |             |        |       |         |         |

February 21, 1997

Dr. Robert F. Abbey, Jr. Program Manager for Marine Meteorology Dept. of the Navy Office of Naval Research 800 N. Quincy St. Arlington, VA 22217

Dear Bob,

Enclosed is a final report for the grant from the Office of Naval Research that recently ended (#N00014-91-J-1068). An overview of the research objectives and a listing of the publications over the last 4 years are included. Thank you for your support over the years.

Sincerely,

Rog. M. With

Roger M. Wakimoto Professor of Meteorology

Enclosure

i

•• ••

Mesoscale Studies of Explosive Cyclogenesis and the state of the state

of a constraint data constraint  $\partial\omega \partial y \hat{x}$  is least of the task .

In recent years the midlatitude, over ocean winter storm than can explosively develop has gained a great deal of interest owing to its unpredictability and resultant devastating effect of commercial shipping and naval operations. In an attempt to understand the mechanisms that cause this large deepening rate in a relatively short period of time (10 mb/6 hr), ERICA (Experiment on Rapidly Intensifying Cyclones over the Atlantic) was organized for the winter 1988/1989 season.

The most important research data platforms deployed during ERICA were the NOAA P-3 aircraft equipped with Doppler radar. It was recognized early in the program that airborne dual-Doppler analyses would be important to identify the location of the convection within these cyclones and to provide detailed kinematic motion in select portions of the storm. The former objective was to identify the areas of latent heat release which is critical to understand how the cyclone is maintained. The latter was important to complete the three-dimensional wind structure in more detail than could be provided by in-situ measurements recorded by the P-3 aircraft.

Over the past four years, extensive analyses of the IOP #4 and 5 storms have been performed. Perhaps, the most important result of these studies is the documentation of mesoscale vortices, approximately 25-40 km wide, developing along intense warm fronts that have undergone a scale contraction. When these vortices come into a proper phasing with a baroclinic wave aloft, rapid deepening of the central low pressure appears to occur. This details of this process have not been documented before in the literature.

One of the limitations of Doppler radar analysis is that it only reveals the kinematic structure of the phenomenon under investigation. Recent techniques have

1

been developed that allow for a thermodynamic retrieval (from the dual-Doppler wind the Rest inclusion with the numerical simulations of extratropical cyclones.

. .

The following papers have been published that address the above objectives and acknowledge funding from the Office of Naval Research -

2

## Results from based on Support from the Office of Naval During the Past 4 Years

- Wakimoto, R.M., 1993: Mesoscale analysis of convective phenomena. International Workshop on Mesoscale Meteor. and TAMEX Program Review, Taipei, 84-93. INVITED TALK.
- Wakimoto, R.M., and J.K. Lew, 1993: Observations of a Florida waterspout during CaPE. Weather and Forecasting, 8, 412-423.
- 3. Wakimoto, R.M., and N.T. Atkins, 1993: Airborne Doppler analyses of an intense extratropical cyclone during ERICA., *Preprints*, 26th Conf. on Radar Meteor., Norman, Amer. Meteor. Soc., 282-284.
- 4. Wakimoto, R.M., and P.G. Black, 1993: Damage survey of Hurricane Andrew and its relationship to the radar-detected eyewall. Preprints, 20th Conf. on Hurricanes and Tropical Meteor., San Antonio, Amer. Meteor. Soc., 54-57.
- 5. Blier, W., and R.M. Wakimoto, 1994: Syncptic and mesoscale analysis of the early evolution of the ERICA IOP 5 explosive oceanic cyclone. Proceedings, The Life Cycles of Extratropical Cyclones, Vol. III, Bergen, Norway, 25-30.
- 6. Paldor, N., C.-H. Liu, M. Ghil, and R.M. Wakimoto, 1994: A new frontal instability: Theory and ERICA observations. J. Atmos. Sci., 22, 3227-3237.
- 7. Wakimoto, R.M., and P.G. Black, 1994: Damage survey of Hurricane Andrew and its relationship to the eyewall. Bull. Amer. Meteor. Soc., 75, 189-200.
- Wakimoto, R.M., N.T. Atkins, and C.-H. Liu, 1994: Airborne Doppler analysis of the mesoscale circulation and frontal structure of an explosive oceanic cyclone during ERICA IOP 5. Proceeding, The Life Cycles of Extratropical Cyclones, Vol. III, Bergen, Norway, 236-241.
- 9. Blier, W., and R.M. Wakimoto, 1995: Observations of the early evolution of an explosive oceanic cyclone during ERICA IOP 5. Part I: Synoptic overview and meso-a-scale structure. Mon. Wea. Rev., 123, 1288-1310.
- 10. Wakimoto, R.M., N.T. Atkins, and C. Liu, 1995: Observations of the early evolution of an explosive oceanic cyclone during ERICA IOP 5. Part II: Airborne Doppler analysis of the mesoscale circulation and frontal structure. Mon. Wea. Rev., 123, 1311-1327.

11. Liu, C.-H., R.M. Wakimoto, N.T. Atkins, and F. Roux, 1995: Dynamic and thermodynamic retrievals of airborne dual-Doppler radar analyses for the ERICA IOP 4 and 5 storms. Preprints, 27th Conference on Radar Meteorology, Vail, Amer. Meteor. Soc., 808-810.

3 - -----

4

- 12. Liu, C.-H., 1996: Analysis of mesoscale circulations within explosivelydeepening marine cyclones - ERICA Observations. Ph.D. Dissertation, University of California, Los Angeles, 149 pp.
- 13. Liu, C.-H., R.M. Wakimoto, and F. Roux, 1997: Observations of mesoscale circulations within extratropical cyclones over the North Atlantic Ocean during ERICA. Mon. Wea. Rev., 125, 341-364.