

AD_____

GRANT NUMBER DAMD17-94-J-4496

TITLE: The Identification and Cloning of the Wnt-1 Receptor

PRINCIPAL INVESTIGATOR: Xi He, Ph.D.

CONTRACTING ORGANIZATION: National Institutes of Health
National Cancer Institute
Bethesda, Maryland 20892

REPORT DATE: September 1997

TYPE OF REPORT: Final

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

DTIC QUALITY INSPECTED 4

19980416 146

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE September 1997	3. REPORT TYPE AND DATES COVERED Final (30 Sep 94 - 30 Sep 97)	
4. TITLE AND SUBTITLE The Identification and Cloning of the Wnt-1 Receptor			5. FUNDING NUMBERS DAMD17-94-J-4496	
6. AUTHOR(S) Xi He, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) National Institutes of Health National Cancer Institute Bethesda, Maryland 20892			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, MD 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200) <p>Ectopic activation of members of the wnt family, such as wnt-1, causes the formation of mammary tumors in mice, providing a model for breast cancer. However, Wnt receptors, essential components mediating Wnt oncogenic functions, had not been identified previously. In <i>Drosophila</i>, a member of the Frizzled (Fz) family of seven transmembrane receptors, the DFizzled 2 (DFz2), was shown to function as a receptor for the Wingless, the ortholog of Wnt-1. This implies that the large family of Fz proteins are likely receptors for Wnt molecules. However, the scarcity of soluble Wnt proteins complicates the study of ligand-receptor relationships and their specificity. We developed an approach in <i>Xenopus</i> embryos to pursue such study.</p> <p>In <i>Xenopus</i> embryos, the Wingless/Wnt-1 subclass of Wnt molecules induces axis duplication whereas the Wnt-5A subclass does not. This difference could be explained by distinct signal transduction pathways or by a lack of Wnt-5A receptor(s) during axis formation. We found that Wnt-5A induces axis duplication in the presence of hFz5, a member of the Frizzled family of seven transmembrane receptors. Wnt-5A/hFz5 signaling is antagonized by glycogen synthase kinase-3 and by the N-terminal ectodomain of hFz5. These results identify hFz5 as a receptor for Wnt-5A. In addition, we found that a secreted Frizzled related protein, FRP, is an antagonist for Wnt. This study illustrated a general approach for studying Wnt-Fz interactions.</p>				
14. SUBJECT TERMS Humans, Anatomical Samples, Wnt-1, Mammary Oncogene, Receptor Tyrosine Kinase, Breast Cancer			15. NUMBER OF PAGES 30	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

_____ Where copyrighted material is quoted, permission has been obtained to use such material.

XH Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

XH Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

XH In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

_____ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

XH In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

XH In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

XH In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Y. He 11-18-97
PI - Signature Date

TABLE OF CONTENTS

Front cover	1
Form SF 298	2
Foreword	3
Table of contents	4
Introduction	5
Results and Discussion	7
Conclusions	13
Figure legends	14
Methods	16
References	18
Acknowledgments	21
Bibliography	22
Figures	23

INTRODUCTION

Wnt genes encode a large family of secreted signaling molecules essential for development and tumorigenesis (1, 2). Of particular interests to human breast cancer research, ectopic activation of certain wnt genes, such as wnt-1, causes mammary carcinogenesis in mice (3). How Wnt-1 induces mammary tumors is not understood. However, given that Wnt molecules are secreted growth factors, it is likely that a receptor-mediated signaling transduction pathway is involved. The aim of this project was to identify Wnt receptors.

In *Drosophila*, wingless (*wg*) gene, the ortholog of murine wnt-1, is required for many stages of embryogenesis (4, 5). Interestingly, upon gene transfer *wg* cDNA can transform murine mammary epithelial cells as does wnt-1 cDNA, demonstrating a striking functional conservation between *wg* and wnt-1 (6). Recently, a receptor for Wg was identified to be a member of the *Drosophila* Frizzled (Fz, ref. 7) family of seven transmembrane receptors. DFz2, when introduced by transfection, confers recipient cells the ability to bind and to respond to Wg (8). The identification of Dfz2 as a Wg receptor implies that the fz family of genes, which include nine mammalian members identified so far (9, 10), are likely to encode receptors for Wnt molecules. However, the scarcity of soluble Wnt proteins (11) complicates the study of Wnt-Fz interactions and their specificity. We have developed an approach to address this issue using *Xenopus* embryos without soluble Wnt proteins.

In *Xenopus*, several Wnt molecules, such as Wg, Wnt-1, the *Xenopus* wnt-3A and wnt-8 (Xwnt-8), have been shown to mimic an early dorsalizing signal that induces the gastrula

organizer and in doing so are able to induce embryonic axis duplication (12-17), thus offering a unique opportunity to study Wnt function and Wnt signal transduction pathway. However, several other Wnt proteins, such as the *Xenopus* Wnt-5A, were found unable to induce axis duplication, and have been classified as a distinct subclass that activates a different pathway(s) (17-19).

In this report we show that Wnt-5A can induce axis duplication in the presence of a member of the mammalian Fz family, hFz5, but not other Fz proteins tested. Wnt-5A signaling via hFz5 is antagonized by GSK-3, and by the hFz5 N-terminal ectodomain that presumably binds and titrates Wnt-5A. These results identify hFz5 as a receptor for Wnt-5A. Importantly, because Wnt-5A is expressed during the development of the mammary gland (20), the revelation of Wnt-5A-hFz5 ligand-receptor relationship will enhance the understanding of normal mammary development and physiology. Moreover, this study establishes axis induction in *Xenopus* as a general approach to investigate Wnt-Fz relationships which play pivotal roles in mammary neoplasia.

In addition, we also identified a novel secreted protein, FRP (for Frizzled related protein), as a functional antagonist to Wnt proteins. FRP shares significant homology with the amino-terminal region of Fz proteins but lacks any transmembrane domain. FRP can completely block axis duplication induced by Wg, Wnt-1 and Xwnt-8 in *Xenopus* embryos. Given the role of Wnt-1 as an oncogene in mammary cancer, the identification of FRP as an inhibitor of Wnt-1 may have implications on genetic intervention of tumor formation.

RESULTS AND DISCUSSION

hFz5 mediates Wnt-5A axis induction

Mouse Wnt-1, *Xenopus wnt-3A* (Xwnt-3A), Xwnt-8, and *Drosophila Wg* induce dorsal axis duplication when low levels, usually 1-10 pg, of their corresponding RNAs are injected into the ventral side of early *Xenopus* embryos (12-15, and data not shown). In contrast, Xwnt-5A RNA fails to do so even after ventral injection at higher doses (75pg to 1ng per embryo); instead, dorsal injection of Xwnt-5A RNA generates head and tail defects that may result from perturbation of cell movements during gastrulation (18). Xwnt-4 and Xwnt-11 behave similarly to Xwnt-5A (19). The Xwnt-8 dorsalizing function is observed before the mid-blastula transition (MBT) when zygotic transcription begins, whereas the Xwnt-5A effect occurs after MBT (13-15, 17). The difference between the effects of Xwnt-8 and Xwnt-5A may reflect the activation of distinct signaling pathways or the lack of a functional Xwnt-5A receptor(s) during axis formation.

To examine whether a particular Fz protein can function as an Xwnt-5A receptor, synthetic RNAs corresponding to Dfz2 (ref. 8) and six mammalian fz cDNAs - mfz3, 4, 6, 7, 8 (from mouse) and hFz5 (from human, ref. 10) - were pooled into two groups and co-injected with 10 pg of Xwnt-5A RNA into the ventral side of 4-cell stage embryos. Injection of Xwnt-5A alone, either fz group alone, or Xwnt-5A together with fz group 2 (mfz3, 4, 6, and 7) produced no phenotypic effects (Fig. 1A). However, co-injection of Xwnt-5A with fz group 1 RNAs (Dfz2, hFz5 and mfz8) induced extensive dorsal axis duplication; in many cases, duplication was complete as determined by the presence of

anterior structures such as the eyes and the cement gland (Fig. 1A and 3A). When the three fz RNAs in group 1 were individually tested, Xwnt-5A plus hFz5 generated similar degrees of axis duplication, whereas Xwnt-5A plus Dfz2 or mfz8 did not (Fig. 1B). Thus, hFz5 alone among the Fz proteins tested is responsible for mediating axis induction by Xwnt-5A. The mature Wnt-5A proteins (after cleavage of the signal peptides) are 100% identical between mouse and human, and 95% identical between mouse and *Xenopus* (18, 21, 22). Given this high degree of sequence identity, it is not surprising that murine wnt-5A RNA also induced axis duplication when co-injected with hFz5 RNA, albeit less efficiently (Fig. 1C). The lower efficiency might be due to effects of untranslated regions in the murine wnt-5A construct on RNA stability and/or translation efficiency.

Axis duplication by Xwnt-5A plus fz group 1 or hFz5, as described above, was observed in fourteen of twenty embryo batches tested during the course of this study. In these cases, 90-100% of the injected embryos showed axis duplication, of which 17-82% included eyes. In the remaining six of twenty embryo batches, the same co-injection induced no or few axis duplications (less than 30%, none complete). The reason for this poor response in some batches is unknown. Possible explanations include variations in the stability of injected RNAs and/or translated proteins, the efficiency of Wnt-5A secretion, the assembly and/or localization of hFz5 protein, or the availability of unknown co-receptor molecules.

It should be noted that while hFz5 is more closely related to DFz2 and mFz8 than to any other known Fz proteins (8, 10), neither DFz2 nor mFz8 cooperated with Xwnt-5A in axis induction. However, DFz2 appeared to be functional in *Xenopus* embryos since the same concentration of Dfz2 RNA as used for co-injection with Xwnt-5A substantially enhanced axis induction by suboptimal amounts of wg RNA (data not shown). The failure

of the other Fz proteins to mediate Xwnt-5A function could be due to an inability to bind Xwnt-5A or an inability to signal in this context.

A dose response curve illustrated that in the presence of 0.4 ng hfz5 RNA per embryo, 1 pg Xwnt-5A RNA induced partial secondary axes whereas 20 pg of Xwnt-5A RNA sometimes hyperdorsalized embryos (Fig. 2). Although the relative protein levels have not been determined, the dose of Xwnt-5A RNA required for axis induction was in a similar range to effective doses of Xwnt-3A, Xwnt-8, wnt-1 and wg RNAs. Together these data suggest a high degree of specificity of the interaction between Xwnt-5A and hFz5.

Histological examination of embryos with duplicated axes revealed that Xwnt-5A plus hFz5 RNAs induced a full array of dorsal tissues, including notochord, neural tube and somites (Fig.3B). There is one notable difference between axes induced by Xwnt-8 and those induced by Xwnt-5A plus hFz5: while the ectopic axes induced by Xwnt-8 are often indistinguishable from the endogenous ones, the axes induced by Xwnt-5A and hFz5 are shorter in most cases, even when eyes and the cement gland are present (Fig. 3A). This might reflect the previously described ability of Xwnt-5A to inhibit cell movements during gastrulation (18).

Xwnt-8 and other Wnt proteins in its group induce dorsal development via the formation of an ectopic Spemann organizer (13-15). We tested whether Xwnt-5A plus hFz5 act similarly by examining the expression of the organizer-specific gene *gooseoid* (*gsc*, ref. 23), using wholemount in situ hybridization. Embryos injected with Xwnt-5A or hfz5 RNA alone expressed *gsc* only dorsally, as did uninjected controls; in contrast,

embryos co-injected with Xwnt-5A and hFz5 RNAs exhibited two opposing domains of *gsc* expression, indicating the formation of an ectopic organizer (Figure 3C).

Drosophila Wg function is mediated by inhibition of the *zeste-white 3* (*shaggy*) gene product, the homolog of vertebrate glycogen synthase kinase-3 (GSK-3, ref. 24). Likewise, the dorsalizing function of Xwnt-8 RNA in *Xenopus* is mimicked by dominant-negative mutant forms of GSK-3 and antagonized by overexpression of wild type GSK-3 (25-27). We therefore asked whether Xwnt-5A signaling via hFz5 is also transduced by inhibition of GSK-3. Co-injection of wild type GSK-3 β RNA blocked dorsal axis duplication by Xwnt-5A plus hFz5 RNAs (Fig. 4A), suggesting that Xwnt-5A signaling through hFz5 requires the inhibition of GSK-3.

All Fz proteins contain a N-terminal extracellular domain composed of a conserved cysteine-rich domain (CRD) and a variable linker region before the first putative transmembrane helix (10). The N-terminal ectodomain of DFz2, when anchored to the membrane, promoted binding of Wg to the cell surface, and deletion of CRD from mFz4 abolished Wg binding (8). Thus the N-terminal extracellular domain appears involved in ligand binding. If the N-terminal ectodomain of hFz5 is involved in binding Xwnt-5A, its overexpression as a secreted molecule might prevent Xwnt-5A from binding to and signaling through hFz5. Indeed, co-injection of RNA coding for the hFz5 N-terminus (hFz5N), without provision of a membrane anchor, antagonized axis induction by Xwnt-5A plus hFz5 in a dose-dependent manner (Fig. 5). Dorsal injection of hFz5N RNA alone did not affect endogenous dorsal axis formation (although gastrulation defects were seen in some embryos); the injected embryos appeared to have an intact neural groove at the neurula stage, and the anteriormost structures (eyes and the cement gland) were present at

the tadpole stage (not shown). These results argue for a specific effect of hFz5N on blocking axis duplication by Xwnt-5A plus hFz5 .

FRP: an antagonist of Wnt signaling

While we were studying Wnt-5A-hFz5 interactions, Dr. Jeff Rubin at NCI purified a secreted polypeptide from conditioned media of a human cell line. cDNA cloning revealed that this peptide shares significant homology with the cysteine-rich domain (CRD) of the amino-terminal region of Fz proteins but lacks any transmembrane domains (Fig.6). This protein, therefore, was named Frizzled related protein (FRP). We and Dr. Rubin initiated a collaboration to study the function of this novel secreted protein.

From the study on Wg-DFz2 (8) and our study on Wnt-5A-hFz5 as described above, it appears that the amino-terminal region of Fz proteins are involved in binding to Wnt ligands. Since FRP is homologous to this region, we reasoned that it may bind to and modulate the signaling activity of Wnt proteins. We envisioned two alternatives: FRP might antagonize Wnt function by binding the protein and blocking access to its cell surface receptor such as Fz proteins; or FRP might enhance Wnt activity by facilitating the presentation of ligand to the Fz receptors, analogous to the action of soluble receptors for certain cytokines (28).

To test these possibilities, we examined the effect of FRP on Wnt-dependent axis duplication. As shown in Figure 7, injection of Wnt-1, Wg or Xwnt-8 mRNA induced partial or complete axis duplication in at least 75% of the embryos. However, when similar quantities of FRP and Wnt RNA were co-injected, the incidence and extent of axis duplication were significantly reduced. Higher concentration of FRP RNA resulted in

almost complete inhibition of Wnt signaling (Fig. 7). It should be noted that FRP inhibited axis duplication by sub-optimal amount of Wnt-1 which induces mostly partial axis duplication. This suggests that even under the sub-optimal condition of Wnt signaling, FRP does not enhance the signaling thus does not function as a soluble “co-receptor” but rather behaves as an antagonist. Surprisingly, FRP was much less effective in antagonizing Xwnt-3A (Fig. 7), suggesting a degree of specificity regarding interactions of FRP with different members of the Wnt family.

CONCLUSIONS

Our data indicate that in the presence of hFz5, Wnt-5A can transduce a signal similar to that of Xwnt-8 or Wnt-1, mediated by a pathway involving down-regulation of GSK-3. The simplest interpretation of these results is that hFz5 functions as a receptor for Wnt-5A.

We further demonstrated that FRP, a molecule related to the Fz extracellular domain, functions as an antagonist for Wnt signal transduction. Given that certain wnt genes, such as wnt-1, are oncogenic, this raises the possibility that FRP may function as a anti-oncogene or tumor suppresser gene.

At least sixteen *wnt* and eight *fz* , and four FRP genes (29) have been identified in mammals thus far. The ligand-receptor relationships among Wnt and Fz and FRP proteins are likely to be complex, as seen in other large families of signaling molecules and their receptors. Biochemical study of the Wnt-Fz interaction remains challenging because of the difficulty in obtaining soluble Wnt proteins. By reconstituting a Wnt-Fz signaling system in *Xenopus* embryos and blocking this signaling system via a dominant-negative form of Fz ectodomain or FRP, the experiments described here provide a general assay to address the relationships among Wnt, Fz and FRP proteins.

FIGURE LEGENDS

Figure 1. Dorsal axis induction by Xwnt-5A mediated via hFz5. n represents the total number of embryos scored from two to six independent experiments; each bar represents the average percentage of axis duplication; the solid portion within each bar represents the average percentage of extensive axis duplication, which is defined by the presence of the cement gland and at least one eye in the duplicated axis. Unless otherwise specified, 10 pg of wnt RNA and/or 400 pg of each fz RNA were injected per embryo. **(A)** Ventral injection of Xwnt-5A RNA together with RNA for fz group 1 (Dfz2, hfz5 and mzf8) induces axis duplication. Xwnt-5A RNA alone, fz group 1 or group 2 (mfz3, 4, 6, and 7) alone, and Xwnt-5A RNA plus fz group 2 did not. **(B)** Xwnt-5A RNA induces axis duplication in the presence of hfz5 RNA, but not with Dfz2 or mzf8 RNA. **(C)** Murine wnt-5A RNA plus hFz5 also induce axis duplication.

Figure 2. Dose response curve of Xwnt-5A and hfz5 RNAs for axis duplication. The bar legend is as in Figure 1.

Figure 3. Induction of anteriormost structures, dorsal axial tissues, and the Spemann organizer by Xwnt-5A plus hfz5. **(A)** Xwnt-5A plus hfz5 induce axis duplication. An example of complete axis duplication with eyes in both axes is shown (stage 41). **(B)** Xwnt-5A plus hfz5 induce a complete set of dorsal tissues. Cross section in the trunk region of an embryo at stage 41 reveals the presence of a neural tube, notochord and

somites in both axes. (C) Xwnt-5A plus hzf5 induce ectopic *gooseoid* (*gsc*) expression in embryos at stage 10.5, as visualized by whole mount *in situ* hybridization.

Figure 4. GSK-3 β antagonizes axis induction by Xwnt-5A plus hzf5. RNA for human GSK-3 β was injected at 1ng per embryo. (A). Examples of control (uninjected) and injected embryos at stage 19. (B) Note the complete suppression of axis duplication in the presence of GSK-3. The Bar legend is as in Fig. 1.

Figure 5. The N-terminal ectodomain of hFz5 (hFz5N) as a secreted protein suppresses axis duplication by Xwnt-5A plus hzf5 in a dose dependent manner. Dorsal injection of same doses of hzf5N RNA did not affect the endogenous dorsal development (data not shown). At higher doses (0.8-2ng RNA per embryo) some embryos with gastrulation defects were observed.

Figure 6. Comparison of the cysteine-rich domain (CRD) of FRP and other members of the Fz family. Solid black shading highlights identities present in FRP and any other Fz family member. The consensus sequence indicates residues present in at least 8 of the 16 Fz or Fz-related proteins. **, the ten invariant cysteine residues; *, other invariant residues.

Figure 7. The hypothetical structures of Fz and FRP proteins. CRD: cysteine-rich domain.

Figure 8. FRP inhibits axis duplication induced by Wnt-1, Wg or Xwnt-8. Note that FRP is much less effective in inhibiting Xwnt-3A. The Bar legend is as in Fig. 1. The amount of mRNA injected per embryo is shown below the bars.

METHODS

Plasmids

All fz cDNAs were subcloned in the pRK5 vector with a optimal Kozak consensus sequence for translation at the initiator ATG (CCACCATG, preceded by different restriction sites for subcloning), and with different lengths of 3' untranslated regions. Xwnt-5A and human GSK-3 were in pSP64 vector as described (18, 26). Murine wnt-5 cDNA (ref. 21) was cloned in pCS2+ vector as the EcoRI and XbaI fragment, which contains about 60 bp 5' and 360 bp 3' untranslated regions. hfz5N was generated by introduction of a stop codon just before the first putative transmembrane helix (changing the preceding amino acids 237 and 238 from phenylalanine and tryptophan to threonine and arginine). The corresponding DNA fragment was subcloned in pCS2+. The FRP NaeI-SalI cDNA fragment, which includes the full coding sequence, was subcloned into the StuI and XhoI sites of pCS2+.

RNA injection

All RNAs for injection were synthesized as capped transcripts *in vitro* with SP6 RNA polymerase (Ambion Megascript). Unless otherwise specified, RNAs were injected into the two cells near the equatorial midline region at the 4-cell stage.

Embryo handling

Embryo preparation, staging, fixation and sectioning, and wholemount in situ hybridization were performed as described (26).

Scoring axis duplication

Percentages of axis duplication were obtained by dividing the total number of embryos with a duplicated axis by the total number of embryos, scored from two to six independent experiments. Extensive axis duplication is defined by the presence of the cement gland and at least one eye in the duplicated axis.

REFERENCE

1. Nusse R, Varmus HE: Wnt genes. *Cell* 69: 1073-87 (1992).
2. Parr BA, McMahon AP: Wnt genes and vertebrate development. *Curr Opin Genet Dev* 4(4):523-8 (1994).
3. Tsukamoto AS, Grosschedl R, Guzman RC, Parslow T, Varmus HE: Expression of the int-1 gene in transgenic mice is associated with mammary gland hyperplasia and adenocarcinomas in male and female mice. *Cell* 55: 619-25 (1988).
4. Klingensmith J, Nusse R: Signaling by wingless in *Drosophila*. *Dev Biol* 166: 396-414 (1994).
5. Siegfried E, Perrimon N: *Drosophila* wingless: a paradigm for the function and mechanism of Wnt signaling. *Bioessays* 16: 395-404 (1994).
6. Ramakrishna NR, Brown AMC: Wingless, the *Drosophila* homolog of the proto-oncogene Wnt-1, can transform mouse mammary epithelial cells. *Development (suppl)*: 95-103 (1993).
7. Vinson CR, Conover S, Adler PN: A *Drosophila* tissue polarity locus encodes a protein containing seven potential transmembrane domains. *Nature* 338: 263-4 (1989).
8. Bhanot P, Brink M, Samos CH, Hsieh J-C, Wang Y, Macke JP, Andrew D, Nathans J, Nusse R: A new member of the frizzled family from *Drosophila* functions as a Wingless receptor. *Nature* 382: 225-230 (1996).
9. Chan SDH, Karpf DB, Fowlkes ME, Hooks M, Bradley MS, Voung V, Bambino T, Liu, MY, Arnaud CD, Strewler GJ, Nissenson RA: Two homologs of the *Drosophila* polarity gene frizzled (fz) are widely expressed in mammalian tissues. *J Biol Chem* 267: 25202-207 (1992).
10. Wang Y, Macke JP, Abella BS, Andreasson K, Worley P, Gilbert DJ, Copeland NG, Jenkins NA, Nathans J: A large family of putative transmembrane receptors homologous to the product of the *Drosophila* tissue polarity gene frizzled. *J Biol Chem* 271: 4468-76 (1996).
11. Smolich BD, McMahon JA, McMahon AP, Papkoff J: Wnt family proteins are secreted and associated with the cell surface. *Mol Biol Cell* 4: 1267-75 (1993).
12. McMahon A, Moon RT: Ectopic expression of the proto-oncogene int-1 in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* 58: 1075-84 (1989).

13. Smith WC, Harland RM: Injected Xwnt-8 RNA acts early in *Xenopus* embryos to promote formation of a vegetal dorsalizing center. *Cell* 67: 753-65 (1991).
14. Sokol S, Christian JL, Moon RT, Melton DA: Injected wnt RNA induces a complete body axis in *Xenopus* embryos. *Cell* 67: 741-52 (1991).
15. Chakrabarti A, Matthews G, Colman A, Dale L: Secretory and inductive properties of *Drosophila* wingless protein in *Xenopus* oocytes and embryos. *Development* 115: 355-369 (1992).
16. Wolda SL, Moody CJ, Moon RT: Overlapping expression of Xwnt-3A and Xwnt-1 in neural tissue of *Xenopus laevis* embryos. *Devel Biol* 155: 46-57 (1993).
17. Moon RT, Christian JL, Campbell RM, McGrew LL, DeMarais AA, Torres M, Lai CJ, Olson DJ, Kelly GM: Dissecting Wnt signalling pathways and Wnt-sensitive developmental processes through transient misexpression analyses in embryos of *Xenopus laevis*. *Development (Suppl)*: 85-94 (1993)
18. Moon RT, Campbell RM, Christian JL, McGrew LL, Shih J, Fraser S: Xwnt-5A: a maternal Wnt that affects morphogenetic movements after overexpression in embryos of *Xenopus laevis*. *Development* 119: 97-111 (1993).
19. Du SJ, Purcell SM, Christian JL, McGrew LL, Moon RT: Identification of distinct classes and functional domains of Wnts through expression of wild-type and chimeric proteins in *Xenopus* embryos. *Mol Cell Biol* 15: 2625-34 (1995).
20. Gavin B, McMahon AP: Differential regulation of the Wnt gene family during pregnancy and lactation suggests a role in postnatal development of the mammary gland. *Mol Cell Biol* 12: 2418-23 (1992).
21. Gavin B, McMahon JA, McMahon AP: Expression of multiple novel Wnt-1/int-1 related genes during fetal and adult mouse development. *Genes Dev* 4: 2319-32 (1990).
22. Clark CC, Cohen I, Eichstetter I, Cannizzaro LA, McPherson JD, Wasmuth JJ, Iozzo RV: Molecular cloning of the human proto-oncogene Wnt-5A and mapping of the gene (WNT5A) to chromosome 3p14-p21. *Genomics* 18: 249-60 (1993)
23. Cho KWY, Blumberg B, Steinbeisser H, DeRobertis, EM: Molecular nature of Spemann's organizer: the role of the homeobox gene goosecoid. *Cell* 67: 1111-1120 (1991).
24. Siegfried E, Chou TB, Perrimon N: wingless signaling acts through zeste-white 3, the *Drosophila* homolog of glycogen synthase kinase-3, to regulate engrailed and establish cell fate. *Cell* 71: 1167-79 (1992).
25. Pierce SB, Kimelman D: Regulation of Spemann organizer formation by the intracellular kinase Xgsk-3. *Development* 121: 755-65 (1995).
26. He X, Saint-Jeannet JP, Woodgett JR, Varmus HE, Dawid IB: Glycogen synthase kinase-3 and dorsoventral patterning in *Xenopus* embryos. *Nature* 374: 617-622 (1995).

27. Dominguez I, Itoh K, Sokol SY: Role of glycogen synthase kinase 3 beta as a negative regulator of dorsoventral axis formation in *Xenopus* embryos. *Proc Natl Acad Sci USA* 92: 8498-502 (1995).
28. Kishimoto T, Tada H, Akira S: Cytokine signaling. *Cell* 76: 253-62 (1994).
29. Rattner A, Hsieh J-C, Smallwood PM, Gilbert DJ, Copeland NG, Jenkins NA, Nathans J: A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc Natl Acad Sci USA* 94: 2859-63 (1997).

ACKNOWLEDGMENTS

This work was carried out from September 1995 to December 1996 in Dr. Harold Varmus' laboratory at the National Cancer Institute (NCI), in collaboration with Dr. Igor Dawid's laboratory at the National Institute of Child Health and Human Development, Dr. Jeff Rubin's laboratory at NCI, and with Dr. Jeremy Nathan's laboratory at the Johns Hopkins University School of Medicine. We are grateful to Randy Moon for Xwnt-5A plasmid and Bruce Huang for technical help, and Bart Williams, Jay Debnath, Pamela Schwartzberg, and Mike Rebagliati for discussion.

BIBLIOGRAPHY

Publications

He, X., Saint-Jeannet, J-P., Wang, Y., Nathans, J., Dawid, I.B., and Varmus, H. E. 1997. A member of the Frizzled protein family mediating axis induction by Wnt-5A. Science 275: 1652-4.

Finch, P.W., **He, X.**, Kelley, M.J., Uren, A., Schaudies, P., Popescu, N.C., Rudikoff, S., Aaronson, S.A., Varmus, H.E., and Rubin, J. S. (1997). Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action. Proc. Natl. Acad. Sci. USA 94: 6770-5.


Saint-Jeannet, J-P., **He, X.**, Varmus, H. E., and Dawid, I. B. 1997. Dorsoventral patterning of the neuraxis: induction of neural crest by Wnt-1 and Wnt-3A. Proc. Natl. Acad. Sci. USA, in press.

Meeting abstract


He, X., Saint-Jeannet, J-P., Wang, Y., Nathans, J., Dawid, I.B., and Varmus, H. E. Identification of a member of the Frizzled protein family mediating axis induction by Wnt-5A. Presented at: Era of Hope, the Department of Defense Breast Cancer Research Program Meeting, Washington, DC, November, 1997.

Figure 1 (He et al.)

A



B



C




Figure 2 (He et al.)




Figure 3 (He et al.)





Figure 4 (He et al.)

A



B




Figure 5 (He et al.)




Figure 6

hFRP	57	CVDLEPADLRL	CHNVGYKKMV	MPNLLGHEHM	AEVKQQAASSW	VELLNKNCHA
hFZ	39	COPLS..IPL	CTDIAYNOTI	MPNLLGHTNQ	EDAGLEVHOF	YPLVKVQCSP
hFZ5	33	COEIT..VPM	CRGIYGNLTH	MENQFNHDIQ	DEAGLEVHOF	WELVEIQCSS
mFZ3	28	CEFIT..LRM	QODLRYNTTF	MPNLLNHVDQ	QTAALAMEPF	HEMVLNDCSR
mFZ4	45	CDPIR..IAM	CONLYNVTK	MPNLVGHDLQ	TDDELQLTTF	TELIQYGCSS
mFZ6	24	CEFIT..VPR	CMKMTYNTTF	FPNLMGHDYDQ	GIAAVEMGHF	LHLANLECSF
mFZ7	49	COPLS..IPL	CTDIAYNOTI	MPNLLGHTNQ	EDAGLEVHOF	YPLVKVQCSP
mFZ8	35	COEIT..VPL	CRGIYNYTY	MENQFNHDIQ	DEAGLEVHOF	WELVEIQCSS
rFZ1	111	COPLS..IPL	CTDIAYNOTI	MPNLLGHTNQ	EDAGLEVHOF	YPLVKVQCSPA
rFZ2	44	COPLS..IPL	CTDIAYNOTI	MPNLLGHTNQ	EDAGLEVHOF	YPLVKVQCSP
dFZ	53	CEFIT..ISI	CKNIPYNTTI	MENLIGHTKQ	DEAGLEVHOF	APLVKIGCSD
dFZ2	64	CEEIT..IPM	CRGIYNTLS	FPNEMNHETQ	DEAGLEVHOF	WELVEIKCSP
cFZ	26	CKQVD..HEM	CNDLRYNLT	FPNLDVDESW	KDASESILTY	KPLLSVVCSE
mCOL	370	CLPLEPTLIT	CSRLGIHFW	LENHLGHTDS	VEVYATVQAM	GRFLHTNCHP
hFRZB	35	CEPVR..IPL	CKSLPWNMTK	MENHLGHTSQ	ANAILAIEQF	EGLLGTHCSP
mSDF5	40	CKPTEANLCL	CEGIEYQNM	MPNLLGHEHM	KEVLEQAGAM	IPLVMKQCHP

* * * * *
 CONSENSUS C-PI-. .IPL C--I-YN-T- MPNLLGH--Q --AGLEVHOF -PLV---CSP

hFRP	GTOVFLCSLF	APVC..LD.	RPTYPGRWLC	BAVRDSCEEV	MOFFEGFYWPE
hFZ	ELRFFLCSMY	APVC.TVLE.	QALPPCRSLC	ERARQGCAL	MNKEGEQWPE
hFZ5	DLRFFLCTMY	TEICLPDYH.	KELPFCRSVC	ERAKAGCSL	MRQYGEFANPE
mFZ3	DLRFELCALY	AEICME..YG	RVTLECRRLC	QRAYSECSKL	MEMEGVPWPE
mFZ4	QLQFFLCSVY	VEMCTEKINI	PIGFCGMC	LSVKRRCEEV	LREFFGANPD
mFZ6	NIEMFLCAE	IEICTE..QI	HVVLECRKLC	EKIVSDCKKL	MDTFGLRWPE
mFZ7	ELRFFLCSMY	APVC.TVLD.	QALPPCRSLC	ERARQGCAL	MNKEGEQWPE
mFZ8	DLKFFLCSMY	TEICLEDYK.	KELPFCRSVC	ERAKAGCAPL	MRQYGEFANPD
rFZ1	ELKFFLCSMY	APVC.TVLE.	QALPPCRSLC	ERA.QGCEAL	MNKEGEQWPD
rFZ2	ELRFFLCSMY	APVC.TVLE.	QALPPCRSLC	ERARQGCAL	MNKEGEQWPE
dFZ	DLQFFLCSLY	VEVC.TILE.	RPTPFCRSIC	ESAR.VCEKL	MKTYNENWPE
dFZ2	DLKFFLCSMY	TEICLEDYH.	KELPFCRSVC	ERARSGCAPL	MOQYSFEWPE
cFZ	QLKFFLCSVY	FEMCNEKLAN	PIGFCRFLC	LSVQEKCLEV	LESEGEKWED
mCOL	FLAWFCLLL	APSCGPG.PP	PELPECRQFC	EALEDEC...	...WNYLAGD
hFRZB	DLLEFLCAMY	AEICTIDFOH	ETLNECKSVC	ERARQGCPEI	LIKYRHSWPE
mSDF5	DTKKFLCSLF	AEVCLDDLD.	ETLQECHSLC	MOVKDRCAEV	MSAFEGPWED

* * * * *
 CONSENSUS -L-FFLCSMY AP-C---L-. -PIPPCRSLC ERA--GCEPL M--FGF-WPE

hFRP	MLK..CDKEP	.EG...DVC	166
hFZ	RIR..CEHFP	RHG..AEQIC	150
hFZ5	RMS..CDRLP	VLGRDAEVLG	147
mFZ3	DME..CSRFP	D.....C	133
mFZ4	TEN..CSKEP	PQN.DHNHMC	158
mFZ6	ELE..CNRLP	H.....C	129
mFZ7	RIR..CENFP	VHG..AGEIC	160
mFZ8	RMR..CDRLP	EQG.NPDTLC	148
rFZ1	TIK..CEKEP	VHG..AGELC	221
rFZ2	RIR..CEHFP	RHG..AEQIC	155
dFZ	NLE..CSKEP	VHG..GEDLC	163
dFZ2	RMA..CEHLP	LHG.DPDNLC	177
cFZ	VIR..CDKEP	LEN.NREKMC	139
mCOL	RIPVV.CASLP	SQE...DGYC	479
hFRZB	NLA..CEELP	VYDR...CVC	147
mSDF5	MTE..CDRFP	QDN...DLC	152

* * * * *
 CONSENSUS -L-. .C--FP --G.C

Figure 7




Figure 8

