GRANT NUMBER DAMD17-94-J-4339

TITLE: A Genetic Approach to Identifying Signal Transduction Mechanisms Initiated by Receptors for TGF-B-Related Factors

PRINCIPAL INVESTIGATOR: F. Michael Hoffman, Ph.D.

CONTRACTING ORGANIZATION: University of Wisconsin Madison, Wisconsin 53706-1490

101 Märs och distrettig skylliger och die model til polytike. Dierteraret i so Section and Critical Contraction and Contraction and Contraction and Contraction and Contraction and Contraction REPORT DATE: October 1997

TYPE OF REPORT: Annual

PREPARED FOR: Commander U.S. Army Medical Research and Materiel Command

Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

1

19980416 124

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for re /iewing instructions, searching a gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments rega ding this burden estimate or an collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate to Information, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington				
1. AGENCY USE ONLY (Leave blank,	2. REPORT DATE October 1997	3. REPORT TYPE AND Annual (1 Oct	-	
4. TITLE AND SUBTITLE A Genetic Approach to Identify Signal Transduction Mechanisms Initiated by Receptors for TGF-Beta-Related Factors			5. FUNDING NUMBERS DAMD17-94-J-4339	
6. AUTHOR(S) F. Michael Hoffman, Ph	I.D.		i	
na u chai dhe Miller Miller (1991)			an de banave de l'alternet al la sua a construction a la construction de la debande de la construction de la co En la construction de la construction	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Wisconsin Madison, Wisconsin 53706-1490			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, MD 21702-5012			(の話) なたり・タット スティーション	
an a	a na mangana na mangana na mangana na mangana na kana na mangana na mangana na mangana na mangana na mangana na	en e		
	 A second sec second second sec	n an	n An an	
12a. DISTRIBUTION / AVAILABILITY	STATEMENT	· · ·	126. DISTRIBUTION CODE	
Approved for public re	lease; distribution u	nlimited		
n <mark>an antara antar 19 meter de la constante de la c 19 meter de la constante de la c</mark>		میں بار میں بین کر ہے۔ میں میں میں میں ایک اور ایک ایک اور ایک ایک اور ایک	n an	
to TGF-beta-related fac sensitized using a mutat caused by the sensitized mapped to six different in the signaling pathway pathway. DNA sequence mutations in the Mad ge identify the gene affected TGF-beta-related ligand beta-related ligand in se screen and the phenotyp	w mutations affecting the sitors was carried out in Dro ion in the Type I receptor. I genetic background were genees. Three of these gen y, confirming the ability of cing was used to establish the ene and in the type II recepted by three mutations recoved a called 60A. Phenotypic a veral developmental proce- bic analysis reveal that 60A illa decapentaplegic protein	sophila. The genetic New mutations affere recovered. The new hes were previously hes were previously hes the screen to recover the molecular lesion tor gene. Positional ered in the screen. The nalysis confirmed the sses in Drosophila. Hes may function as part	background was cting the phenotypes mutations were known to participate mutations in the in 5 of the new cloning was used to 'he gene encodes a e role of this TGF- Recovery in the	
ng ana ang ang ang ang ang ang ang ang a	na na ana ao		n na na manainte na sua na manainte na sua na sua sua sua sua sua sua sua sua sua su	
14. SUBJECT TERMS Transfor	ming Growth Factor, S	Signal Transduct:	Lon, 15. NUMBER OF PAGES	
Genetic Approaches, Gr Receptors, Genetic Str			16. PRICE CODE	
17. SECURITY CLASSIFICATION 1 OF REPORT	8. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIN OF ABSTRACT	ICATION 20. LIMITATION OF ABSTRAC	
	Unclassified 2	Unclassified	Unlimited	
NSN 7540-01-280-5500			Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. Z39-18 298-102	

I.

1

FOREWORD

a second of the

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

A second control of spirit lines.
 A second control of spirit lines.

n an ann an chuirte ann an chuireann. An ann ann an Airte Airte An Airte Airte Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

MAIN conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

M Hin conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

mH In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules. energia de la composición de la compos Composición de la comp

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

TABLE OF CONTENTS

	Page
Front Cover	1
Form 298	2
Foreword	3
Table of Contents	4
Introduction	5
Body	5-6
Conclusions	6

INTRODUCTION:

Cancer cells have multiple genetic defects that alter the cells' normal growth regulation by disrupting or activating the regulatory pathways that normally maintain an appropriate cell number in a specific tissue. Identification of the genes and proteins that participate in these regulatory pathways has provided and continues to provide insights into the molecular basis of cancer. Identification of the proteins, and the regulatory circuitry in which they participate, provides reagents for cancer diagnosis, prognostic opportunities, and targets for the discovery and development of new therapeutic agents with which to treat cancers.

Over the past 10 years, it has been established that many of the critical regulatory genes altered in human cancer cells have been highly conserved during evolution such that homologous genes exist in other organisms, including the genetically accessible organisms yeast, the nematode *C. elegans* and the fruit fly *Drosophila melanogaster*. This has provided an opportunity to use genetic strategies in these model organisms to identify new genes that function on the regulatory pathways. Dr. Hoffmann's laboratory has used this strategy to identify new components on the TGF-beta-related signal transduction cascade in Drosophila. The overall goal of these studies is to identify new gene products that are essential to the function of this pathway and to understand the molecular mechanisms by which the gene products interact. The working hypothesis, which has been confirmed now by Dr. Hoffmann's laboratory and several other laboratories, is that new genes on regulatory pathways, identified by genetic strategies in model organisms, can be used as molecular probes to identify new human genes that participate in the regulatory pathways altered in cancer cells.

RESULTS:

In the initial years of the project we identified a sensitized genetic background in Drosophila for the TGF-beta-related pathway and carried out a mutagenesis screen of 10,000 genomes for modifier mutations. Genetic characterization of the new mutations that were recovered indicated that some may be new alleles of genes previously identified on the pathway. These included five new alleles of Mad, two new alleles of punt and 1 new allele of medea. During the past year we have characterized the Mad and punt alleles by DNA sequencing. The results provided molecular confirmation of the genetic interaction and a molecular identification of the mutation as follows:

Mutant designation	Gene	Amino acid change
D14	Mad	Glutamine 168 to stop codon
D16	Mad	Tyrosine 255 to Asparagine
D24	Mad	Aspartic acid 437 to Asparagine
D13	Punt	Cysteine 67 to tyrosine
D18	Punt	Glutamic acid 378 to valine

The Mad mutation at 437 occurs in the "hot-spot" of the Mad protein that is conserved in human Smad proteins and is the region of most Smad mutations reported in human colon cancer and human pancreatic cancer. The punt mutation at 67 occurs in one of the cysteines that is conserved in human type receptor kinases.

In addition to the molecular characterization of the alleles reported above, a major effort over the past year has been the positional cloning of one of the new modifier genes.

Three alleles were recovered in a single complementation group that mapped to the distal end of the right arm of the third chromosome. Genetic crosses to stocks containing chromosome deletions were used to narrow the location of the gene to the polytene chromosome bands in subdivision 60A. A P-element transgene in that region was mobilized with transposase and a screen for P-element alleles of into the modifier gene was carried out successfully. DNA adjacent to the new P-element insertion site was cloned and cosmid clones of the region isolated. A candidate gene in the region was identified as 60A, a Drosophila TGF-beta superfamily member most similar in amino acid sequence to the human protein bone morphogenetic protein 7. To determine whether the mutations occurred in the 60A gene itself (versus in an adjacent gene), 60A DNA was amplified from the mutant flies by RT-PCR of mRNA or PCR of genomic DNA. Mutations were identified in 60A for all three sequences.

The mutant phenotype of the 60A mutations was characterized. Absence of 60A was lethal to the animal and led to absence of fat body, absence of the first constriction in the midgut and reduced gastric caeca length (structures at the anterior end of the midgut). These phenotypes are similar to those reported for mutations in the Mad gene, a protein critical for the signal transduction activated by TGF-beta-related factors.

The most interesting finding was that mutations in 60A combined with a weak mutation in the type I receptor thick veins, led to mutant phenotypes much more severe than either mutant gene alone and that the more severe mutant phenotypes were very similar to those produced by mutations in the Drosophila TGF-beta-related factor decapentaplegic (dpp). We interpret these finding to mean that dpp and 60A form a heterodimeric ligand that is important for the biological effects of both genes. The heterodimer may be largely functionally redundant to dpp homodimers. A manuscript on this work has just been submitted for publication.

Finally, two other modifier mutations recovered in the original screen are being mapped in order to pursue their positional cloning.

CONCLUSIONS:

We have successfully characterized a number of new mutations from the modifier screen to the molecular level. This includes the identification of the molecular defect present in new alleles of Mad and punt. We have identified mutant alleles of the 60A gene and discovered the importance of 60A/dpp heterodimers for Drosophila development. TGF-beta family members are known to have the capacity to form heterodimers, which may have very different functions than the homodimers, e.g., activins and inhibins, but there are few examples of these occurring *in vivo*. Our results should stimulate new interest in the existence and function of heterodimeric forms of TGF-beta-related ligands. Our progress is consistent with the goals, time table and Work Plan initially proposed.