

**INTERNET DOCUMENT INFORMATION FORM**

**A . Report Title: Summary of Notifiable Diseases, United States  
1994**

**B. DATE Report Downloaded From the Internet: 17 Jun 98**

**C. Report's Point of Contact: (Name, Organization, Address,  
Office Symbol, & Ph #: U.S. Department of Health and Human  
Services**

**D. Currently Applicable Classification Level: Unclassified**

**E. Distribution Statement A: Approved for Public Release**

**F. The foregoing information was compiled and provided by:  
DTIC-OCA, Initials: \_\_PM\_\_ Preparation Date: 17 Jun 98**

The foregoing information should exactly correspond to the Title, Report Number, and the Date on the accompanying report document. If there are mismatches, or other questions, contact the above OCA Representative for resolution.

<b>DISTRIBUTION STATEMENT A</b>
<b>Approved for public release; Distribution Unlimited</b>

19980618 129



Published October 6, 1995, for  
1994 / Vol. 43 / No. 53

# MNWR

MORBIDITY AND MORTALITY WEEKLY REPORT

---

## Summary of Notifiable Diseases, United States

1994

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

Public Health Service  
Centers for Disease Control  
and Prevention (CDC)  
Atlanta, Georgia 30333

DTIC QUALITY INSPECTED



The statistical summary of notifiable diseases in the United States is published to accompany each volume of the *Morbidity and Mortality Weekly Report* by the Centers for Disease Control and Prevention (CDC), Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. Summary of notifiable diseases, United States, 1994. MMWR 1994;43(53): [inclusive page numbers].

Centers for Disease Control and Prevention ..... David Satcher, M.D., Ph.D.  
*Director*

The material in this report was collected and forwarded to CDC by the 57 reporting areas and was received and prepared by:

Epidemiology Program Office ..... Stephen B. Thacker, M.D., M.Sc.  
*Director*

Richard A. Goodman, M.D., M.P.H.  
*Editor, MMWR Series*

Division of Surveillance  
and Epidemiology ..... Donna F. Stroup, Ph.D., M.Sc.  
(Through April 1995) *Director*  
Scott F. Wetterhall, M.D., M.P.H.  
(May 1995) *Acting Director*

Scientific Information and Communications Program  
Public Health Publications Branch ..... Suzanne M. Hewitt, M.P.A.  
*Chief*

Rachel J. Wilson  
*Project Editor*

Information Resources Management Branch ..... Elizabeth E. Rubery  
*Chief*  
Morie M. Higgins  
*Visual Information Specialist*

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or the U.S. Department of Health and Human Services.

Copies can be purchased from Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402-9325. Telephone: (202) 783-3238.

The following CDC staff members contributed to this report:

Denise T. Koo, M.D., M.P.H.  
Andrew G. Dean, M.D., M.P.H.

Ruth W. Slade  
Carol M. Knowles  
Deborah A. Adams  
Wanda K. Fortune  
Patsy A. Hall  
Robert F. Fagan  
Barbara Panter-Connah  
Harry R. Holden  
Gerald F. Jones  
Clarence Lee Maddox

*Division of Surveillance and Epidemiology  
Epidemiology Program Office*

Dana J. Milk  
*Scientific Information and Communications Program  
Public Health Publications Branch  
Epidemiology Program Office*

**Consultant**  
Willie J. Anderson  
*Office of the Vice President for Health Affairs  
Emory University*

## **Table of Contents**

<b>Foreword .....</b>	<b>ii</b>
<b>Background .....</b>	<b>iii</b>
<b>Data Sources .....</b>	<b>v</b>
<b>1994 Highlights for Selected Diseases .....</b>	<b>vii</b>
<b>Bibliography .....</b>	<b>x</b>

### **Part 1**

<b>Summaries of Notifiable Diseases in the United States, 1994</b>	
Reported Cases, by Month, 1994 .....	3
Reported Cases, by Geographic Division and Area, 1994 .....	4
Reported Cases, by Age Group, 1994 .....	10
Reported Cases, by Race, 1994 .....	11
Reported Cases, by Ethnicity, 1994 .....	12

### **Part 2**

<b>Graphs and Maps for Selected Notifiable Diseases in the United States.....</b>	<b>13</b>
---	-----------

### **Part 3**

<b>Historical Summary Tables Covering the Period 1945–1994</b>	
--	--

<b>Notifiable Diseases —</b>	
Summary of Reported Cases, per 100,000 Population,	
United States, 1985–1994 .....	69
Summary of Reported Cases, United States, 1985–1994 .....	70
Summary of Reported Cases, United States, 1975–1984 .....	72
Summary of Reported Cases, United States, 1965–1974 .....	74
Summary of Reported Cases, United States, 1955–1964 .....	76
Summary of Reported Cases, United States, 1945–1954 .....	78
Deaths From Specified Notifiable Diseases, United States, 1983–1992.....	80

<b>State and Territorial Epidemiologists and Laboratory Directors .....</b>	<b>inside back cover</b>
---	--------------------------

## **Foreword**

### ***MMWR Summary of Notifiable Diseases, United States, 1994***

This publication contains summary tables of the official statistics for the reported occurrence of nationally notifiable diseases in the United States for the year 1994. This information is collected and compiled from reports to the National Notifiable Diseases Surveillance System (NNDSS). Because the dates of onset and dates of diagnosis for notifiable diseases are often unknown, these surveillance data are presented by the week that they were reported to public health officials. These data are then finalized and published in the *MMWR Summary of Notifiable Diseases, United States*, for use by state and local health departments; schools of medicine and public health; communications media; local, state, and federal agencies; and other agencies or persons interested in following the trends of reportable conditions in the United States. Publication of the annual summary also ensures documentation of diseases that are considered national priorities for notification and of the annual number of cases of such diseases.

Part 1 contains information on morbidity for each of the conditions considered nationally notifiable during 1994. In all tables, leprosy is listed as Hansen disease and typhus fever (tick-borne) as Rocky Mountain spotted fever (RMSF). The tables show the number of cases of notifiable diseases reported to CDC for 1994, as well as the distribution of cases by month and geographic location, and by patient's age, race, and ethnicity. The data are final totals as of July 7, 1995, unless otherwise noted.

Part 2 contains graphs and maps depicting summary data for many of the notifiable conditions described in tabular form in Part I.

Part 3 includes tables showing the number of cases of notifiable diseases reported to CDC and to the National Office of Vital Statistics since 1945. It also includes a table on deaths associated with specified notifiable diseases reported to the National Center for Health Statistics, CDC, for the period 1983–1992.

## **Background**

As of January 1, 1994, 49 infectious diseases were designated as notifiable at the national level. *A notifiable disease is one for which regular, frequent, and timely information on individual cases is considered necessary for the prevention and control of the disease.* This section briefly summarizes the history of national notifiable disease reporting in the United States.

In 1878, Congress authorized the U.S. Marine Hospital Service (the precursor to the Public Health Service, PHS) to collect morbidity reports on cholera, smallpox, plague, and yellow fever from U.S. consuls overseas; this information was to be used for instituting quarantine measures to prevent the introduction and spread of these diseases into the United States. In 1879, a specific Congressional appropriation was made for the collection and publication of reports of these notifiable diseases. The authority for weekly reporting and publication was expanded by Congress in 1893 to include data from states and municipal authorities. To increase the uniformity of the data, Congress enacted a law in 1902 directing the Surgeon General to provide forms for the collection and compilation of data and for the publication of reports at the national level. In 1912, state and territorial health authorities—in conjunction with PHS—recommended immediate telegraphic reporting of five infectious diseases and monthly reporting by letter of 10 additional diseases. The first annual summary of *The Notifiable Diseases* in 1912 included reports of 10 diseases from 19 states, the District of Columbia, and Hawaii. By 1928, all states, the District of Columbia, Hawaii, and Puerto Rico were participating in national reporting of nearly 30 specified conditions. At their annual meeting in 1950, the State and Territorial Health Officers authorized a conference of state and territorial epidemiologists whose purpose was to determine which diseases should be reported to PHS. CDC assumed responsibility for the collection and publication of data on nationally notifiable diseases in 1961.

The list of nationally notifiable diseases is revised periodically. For example, diseases may be added to the list as new pathogens emerge or deleted as their incidence declines. Public health officials at state health departments and CDC continue to collaborate in determining which diseases should be nationally notifiable; the Council of State and Territorial Epidemiologists (CSTE), with CDC input, makes recommendations annually for additions and deletions to the list of nationally notifiable diseases. However, reporting of nationally notifiable diseases to CDC by the states is voluntary. Reporting is currently mandated (by state legislation or regulation) only at the state level. The list of diseases that are considered notifiable, therefore, varies slightly by state. All states generally report the internationally quarantinable diseases (i.e., cholera, plague, and yellow fever) in compliance with the World Health Organization's International Health Regulations. During 1994, 49 infectious diseases were considered notifiable at the national level and were reported to CDC; 41 were reported on a weekly basis, and eight were reported monthly.

CSTE and CDC held a national surveillance conference November 30-December 2, 1994 to review the state of national infectious disease surveillance. Infectious diseases that have been approved for addition to national surveillance during 1995 are *Chlamydia trachomatis* (genital infections), coccidioidomycosis (for regional surveillance), cryptosporidiosis, hantavirus infection, (post-diarrheal) hemolytic uremic syndrome,

pediatric infection with the human immunodeficiency virus, invasive group A streptococcal infections, streptococcal toxic-shock syndrome, and drug-resistant *Streptococcus pneumoniae*. These conditions currently are not reportable in all states, and the mechanism for reporting may not involve clinicians or consist of reports of individual cases (i.e., traditional notification methods). Reports of the number of cases of these conditions will not appear in the current year tables before 1996.

At the 1994 conference the following diseases were also proposed as deletions from the list of infectious diseases under national surveillance: amebiasis, aseptic meningitis, primary encephalitis (except for arboviral encephalitis), postinfectious encephalitis, granuloma inguinale, unspecified hepatitis, leptospirosis, lymphogranuloma venereum, rheumatic fever, and tularemia. These changes were confirmed by a vote of the full membership of CSTE in early 1995. The number of reported cases of these diseases will not appear in the current year tables after 1994.

The list of 49 infectious diseases that were designated as notifiable at the national level during 1994 appears below:

AIDS	Measles
Amebiasis*	Meningococcal disease
Anthrax	Mumps
Aseptic meningitis*	Pertussis
Botulism**	Plague
Brucellosis	Poliomyelitis
Chancroid**	Psittacosis
Cholera	Rabies, animal
Diphtheria	Rabies, human
Encephalitis, primary*	Rheumatic fever*
Encephalitis, postinfectious*	Rocky Mountain spotted fever (RMSF)
<i>Escherichia coli</i> O157:H7**	Rubella
Gonorrhea	Rubella, congenital syndrome
Granuloma inguinale*	Salmonellosis**
<i>Haemophilus influenzae</i> , invasive	Shigellosis**
Hansen disease (leprosy)	Syphilis
Hepatitis A	Syphilis, congenital
Hepatitis B	Tetanus
Hepatitis, non-A, non-B	Toxic-shock syndrome
Hepatitis C, unspecified*	Trichinosis
Legionellosis	Tuberculosis
Leptospirosis*	Tularemia*
Lyme disease	Typhoid fever
Lymphogranuloma venereum*	Yellow fever**
Malaria	

\*Deleted from the nationally notifiable disease list in 1995

\*\*Not published in the weekly tables

## Data Sources

Provisional data on the reported occurrence of notifiable diseases are published weekly in *MMWR*. After each reporting year, staff in state health departments finalize reports of cases for the year with local or county health departments, as well as reconcile the data with reports previously sent to CDC throughout the year; these data are compiled in final form in this summary. Notifiable disease reports (published in the annual *MMWR Summary of Notifiable Diseases* only after the approval of each state epidemiologist) are the authoritative and archival counts of cases. *MMWR Surveillance Summaries* or other surveillance reports produced by CDC programs, which are useful for detailed epidemiologic analyses, may not agree exactly with numbers reported in the annual *Summary of Notifiable Diseases* due to differences in the timing of reports, the source of the data, and the use of different case definitions.

Data in this summary are derived primarily from reports transmitted to the Division of Surveillance and Epidemiology, Epidemiology Program Office, CDC, by the 50 state, two city, and five territorial health departments through the National Electronic Telecommunications System for Surveillance (NETSS). Final data for other diseases are from surveillance program records of the CDC programs listed below; requests for further information regarding these data should be directed to the source specified.

### **National Center for Health Statistics**

Office of Vital and Health Statistics Systems (deaths from selected notifiable diseases)

### **National Center for Infectious Diseases**

Division of Bacterial and Mycotic Diseases (toxic-shock syndrome and laboratory data on botulism, *Escherichia coli* O157:H7, *Salmonella*, and *Shigella*)

Division of HIV/AIDS (human immunodeficiency virus/acquired immunodeficiency syndrome)

Division of Vector-Borne Infectious Diseases (laboratory data on arboviral encephalitis)

Division of Viral and Rickettsial Diseases (animal rabies)

### **National Center for Prevention Services**

Division of Sexually Transmitted Diseases and HIV Prevention (gonorrhea, syphilis, chancroid, granuloma inguinale, and lymphogranuloma venereum)

Division of Tuberculosis Elimination (tuberculosis)

### **National Immunization Program**

Epidemiology and Surveillance Division (poliomyelitis)

Disease totals for the United States, unless otherwise stated, do not include data for American Samoa, Guam, Puerto Rico, the Virgin Islands, and the Commonwealth of the Northern Mariana Islands (CNMI).

Data on notifiable diseases before 1960 (before CDC assumed responsibility for collection and publication of these data) are obtained from publications of the National Office of Vital Statistics.

Population estimates for states are based on the July 1, 1994, post-censal estimates made by the U.S. Department of Commerce, Bureau of the Census, Population Division, Population Estimates Branch, Press Release CB94-204.

Population estimates for territories are from the 1990 census, U.S. Department of Commerce, Bureau of the Census, Press Releases CB91-142, 242, 243, 263, and 276.

Rates in the *1994 Summary of Notifiable Diseases* were based on data for the U.S. total resident population. However, population data from states in which diseases were not notifiable or disease data were not available were excluded from rate calculations.

### **Interpreting Data**

The data reported in this summary are useful for analyzing disease trends and determining relative disease burdens. However, these data must be interpreted in light of reporting practices. Some diseases that cause severe clinical illness (e.g., plague or rabies), if diagnosed by a clinician, are most likely reported accurately. However, persons who have diseases that are clinically mild and infrequently associated with serious consequences (e.g., salmonellosis) may not even seek medical care from a health-care provider; even if these diseases are diagnosed, they are less likely to be reported. The degree of completeness of reporting also is influenced by the diagnostic facilities available; the control measures in effect; the public awareness of a specific disease; and the interests, resources, and priorities of state and local officials responsible for disease control and public health surveillance. Finally, factors such as changes in the case definitions for public health surveillance, the introduction of new diagnostic tests, or the discovery of new disease entities may cause changes in disease reporting that are independent of the true incidence of disease.

Public health surveillance data are published for selected racial and ethnic population groups because these variables may be risk markers for certain notifiable diseases. Risk markers can identify potential risk factors for investigation in future studies. Data on race and ethnicity also can be useful for identifying groups to target for prevention efforts. Year 2000 objectives for racial and ethnic groups (standardized to the 1940 U.S. population) have been established for several of the notifiable diseases.

#### **EXPLANATION OF SYMBOLS USED IN TABLES, GRAPHS, AND MAPS**

Data not available .....	NA
No reported cases.....	-
Report of disease not required in that jurisdiction (not notifiable) .....	NN

## **HIGHLIGHTS FOR SELECTED INFECTIOUS DISEASES NOT NATIONALLY NOTIFIABLE DURING 1994**

### **Chlamydia**

*Chlamydia trachomatis* infections are common among adolescents and young adults. An estimated 10% of sexually active adolescent females are infected with chlamydia. In 1994, 47 states reported 448,984 chlamydia infections. From 1984 through 1994, reported cases increased from 3.2 cases per 100,000 to 188.4. This trend may reflect increasing recognition and interest among health-care providers and public health officials.

### **Coccidioidomycosis**

An outbreak of coccidioidomycosis occurred in Ventura County, California, following the 1994 Northridge earthquake. From January 24 through March 15, 1994, 203 infected persons were identified, compared with 52 cases that were reported through passive surveillance in the county in 1993. The National Center for Infectious Diseases (NCID/CDC) Emerging Infections Program (EIP), in collaboration with the State of California Department of Health Services, is conducting active surveillance for coccidioidomycosis in 10 California counties.

### **Cryptosporidiosis**

In 1994, waterborne outbreaks of cryptosporidiosis were investigated in Las Vegas, Nevada, Walla Walla, Washington, and Lake Nummy, New Jersey, focusing national attention on the potential for waterborne transmission of *Cryptosporidium*. In September 1994, a national workshop on waterborne cryptosporidiosis was held at CDC, which resulted in guidelines and recommendations for prevention of cryptosporidiosis in severely immunosuppressed persons, appropriate public health responses to the problem, and epidemiologic and laboratory-based surveillance and research.

### **Hantavirus**

Hantavirus Pulmonary Syndrome (HPS) is a recently recognized hantaviral illness caused by Sin Nombre virus and the newly identified Black Creek Canal and Bayou viruses. The identified rodent reservoirs for these viruses — *Peromyscus maniculatus* and *leucopus* (deer and white-footed mice) for Sin Nombre virus and its variants and *Sigmodon hispidus* (cotton rat) for Black Creek Canal virus — extend across the continental United States. As of July 20, 1995, national surveillance for HPS has identified 113 confirmed case-patients in 23 states (case fatality rate: 52%); 31 of these cases occurred in 1994.

### **Invasive group A streptococcal infections**

Prospective and retrospective active surveillance data for invasive group A streptococcal (GAS) infections were analyzed and several risk groups identified, including: persons who have human immunodeficiency virus (HIV) infections and acquired immunodeficiency syndrome (AIDS), injecting-drug users, persons who have cancer, diabetes mellitus, heart disease or chronic lung disease, alcohol abusers, and children who have varicella. Although different GAS strains have been identified from individual case patients, M-type 1 strains predominated.

### **Drug-resistant *Streptococcus pneumoniae***

In the United States, the prevalence of drug-resistant *Streptococcus pneumoniae* (DRSP) has increased since 1987 from 3.6% to 14.5%, according to limited voluntary reporting by 12 sentinel hospitals. Limited 1994 surveillance data from these hospitals indicate the proportion of invasive disease (bacteremia and meningitis) caused by penicillin-resistant pneumococci ranges from 3% to 30% and shows widespread geographic variation. Information regarding community-specific DRSP prevalence is needed to assist clinicians in choosing optimal empiric therapy. To enhance efficient and timely reporting, CDC is currently piloting an electronic laboratory-based surveillance system.

### **Vancomycin-resistant enterococci**

In 1994, the percentage of nosocomial enterococci reported as resistant to vancomycin increased from 11.5% in 1993 to 13.6% among Intensive Care Unit (ICU) isolates and from 4.9% to 9.1% among noncritical care unit isolates. The increase was more dramatic among isolates from noncritical care units, suggesting that vancomycin-resistant enterococci are spreading from their focus in ICUs.

### **Pneumonia of unknown etiology**

From 1979 to 1994, the overall crude death rate for pneumonia and influenza increased 59%, from 20.0 to 31.8 deaths per 100,000. Through 1992 (the most recent year for which complete data are available), pneumonia of unspecified etiology (ICD-9 code 486) accounted for most of the overall increase—the age-adjusted death rates in this diagnostic category increased 74%. Since the 1970s, several previously unrecognized infectious agents have been identified as causes of lower respiratory infections, including *Legionella pneumophila*, *Chlamydia pneumoniae*, and Sin Nombre virus. Recent prospective studies of community-acquired pneumonia indicate that an etiology cannot be identified in up to 50% of cases.

### **Transfusion-associated infectious diseases**

An Institute of Medicine committee recently released the report, "HIV and the Blood Supply: An Analysis of Crisis Decisionmaking," calling for the establishment of a surveillance system at CDC to detect, monitor, and warn of adverse effects in the recipients of blood and blood products. CDC is reviewing existing surveillance systems to highlight and address areas that need improvement. Diseases that are being examined to evaluate the level of risk associated with transfusion include HIV/AIDS, Chagas disease, babesiosis, Creutzfeld-Jacob disease, the hepatitis viruses, malaria, and transfusion-associated sepsis.

## **INTERNATIONAL NOTES**

### **Dengue**

Although dengue fever is not endemic in the United States, its incidence is increasing in most tropical areas throughout the world. In 1994, CDC processed serum samples from 91 residents of 27 states and the District of Columbia who had travelled to countries where dengue is endemic. Among these 91 persons, 37 (40.7%) cases of dengue were diagnosed serologically or virologically.

### **Plague**

During September and October 1994, outbreaks of bubonic and pneumonic plague were reported from sites east and north of Bombay, India, respectively. A lack of reliable epidemiologic information contributed to the ensuing international health emergency. Evidence revealed that plague did not occur in international travelers or spread beyond the original foci.

## Bibliography

### General

- Benenson AS. Control of communicable diseases in man. 15th ed. Washington, DC: American Public Health Association, 1990.
- CDC. Manual of procedures for the reporting of nationally notifiable diseases to CDC. Atlanta: US Department of Health and Human Services, Public Health Service, CDC, 1995.
- CDC. Case definitions for public health surveillance. MMWR 1990;39(No. RR-13).
- CDC. Mandatory reporting of infectious diseases by clinicians, and mandatory reporting of occupational diseases by clinicians. MMWR 1990;39(No. RR-9).
- CDC. National electronic telecommunications system for surveillance—United States, 1990–1991. MMWR 1991;40:502.
- CDC. Update: graphic method for presentation of notifiable disease data—United States, 1991. MMWR 1991;40:124–5.
- CDC. Use of race and ethnicity in public health surveillance. MMWR 1993;42(No. RR-10).
- CDC. Update: changes in notifiable disease surveillance data—United States, 1992–1993. MMWR 42(42):824–6.
- CDC. Sexually transmitted disease surveillance, 1993. Atlanta: US Department of Health and Human Services, Public Health Service, 1994.
- Stroup DF, Wharton M, Kafadar K, Dean AG. An evaluation of a method for detecting aberrations in public health surveillance data. Am J Epidemiol 1993;137(3):373–80.
- Teutsch SM, Churchill RE, eds. Principles and practice of public health surveillance. New York: Oxford University Press, 1994.
- Thacker SB, Choi K, Brachman PS. The surveillance of infectious diseases. JAMA 1983; 249:1181–5.

### AIDS

- CDC. AIDS among racial/ethnic minorities—United States, 1993. MMWR 1994;43(35):644–7, 653–5.
- CDC. Heterosexually acquired AIDS—United States, 1993. MMWR 1994;43(9):155–60.
- CDC. HIV/AIDS surveillance report. Atlanta: US Department of Health and Human Services, 1994;6(2):1–39.
- CDC. Recommendations of the U.S. Public Health Service Task Force on the use of zidovudine to reduce perinatal transmission of human immunodeficiency virus. MMWR 1994;43(No. RR-11):1–20.
- CDC. Update: acquired immunodeficiency syndrome—United States, 1994. MMWR 1995;44(4): 64–7.
- CDC. Update: trends in AIDS diagnosis and reporting under the expanded surveillance definition for adolescents and adults—United States, 1993. MMWR 1994;43(45):826–31.
- CDC. Update: AIDS among women—United States, 1994. MMWR 1995;44(5):81–4.

### Amebiasis

- Katzenstein DA. Drug treatment of amebiasis. In: Peterson PK, Verhoef J, eds. Antimicrobial agents annual 2. New York: Elsevier Science Publishers, 1987:247–52.
- McAuley JB, Herwaldt BL, Stokes SL, et al. Diloxanide furoate for treating asymptomatic *Entamoeba histolytica* cyst passers: 14 years' experience in the United States. Clin Infect Dis 1992;15:464–8.
- Ravdin JI, ed. Amebiasis: human infection by *Entamoeba histolytica*. New York: John Wiley and Sons, Inc., 1988.
- Ravdin JI. Amebiasis. Clin Infect Dis 1995;20:1453–66.

### Anthrax

- Brachman PS. Anthrax. In: Hoeprich PD, Jordan MC, Roland AR, eds. Infectious diseases. 5th ed. Philadelphia: JB Lippincott Co., 1994:1003–8.
- Meselson M, Guillemin J, Hugh-Jones M, et al. The Sverdlovsk anthrax outbreak of 1979. Science 1994 Nov;266(5188):1202–8.

### **Arboviral Infections**

#### **( California serogroup viruses, Eastern equine encephalitis, St. Louis encephalitis, and Western equine encephalitis )**

Monath TP, ed. The arboviruses: epidemiology and ecology. Boca Raton, FL: CRC Press, 1983.  
Tsai TF. Arboviral infections in the United States. Infect Dis Clin North Am 1991;5:73-102.  
Tsai TF. Arboviruses and related zoonotic viruses. In: Oski FJ, ed. Principles and practice of pediatrics. 2nd ed. Philadelphia: JB Lippincott Co., 1994:1266-88.

### **Aseptic Meningitis**

CDC. Aseptic meningitis—New York State and United States, weeks 1-36, 1991. MMWR 1991; 40(45):773-5.  
Moore M. Enteroviral disease in the United States, 1970-1979. J Infect Dis 1982;146:103-8.  
Morens DM, Zweighaft RM, Bryan JA. Nonpolio enterovirus disease in the United States, 1971-1975. Int J Epidemiol 1979;8:49-54.  
Wilfert CM, Lehrman SN, Katz SL. Enteroviruses and meningitis. Pediatr Infect Dis J 1983;2: 333-41.

### **Botulism**

St. Louis ME, Peck SHS, Bowering D, et al. Botulism from chopped garlic: delayed recognition of a major outbreak. Ann Intern Med 1988;108:363-8.  
Webber JT, Hatheway CL, St. Louis ME. Botulism. In: Hoeprich PD, Jordan MC, Ronald AR. Infectious diseases: a treatise of infectious processes. 5th ed. Philadelphia: JB Lippincott Co., 1994:1185-94.  
Woodruff BA, Griffin PM, McCroskey LM, et al. Clinical and laboratory comparison of botulism from toxin types A, B, and E in the United States 1975-1988. J Infect Dis 1992;166:1281-6.

### **Brucellosis**

CDC. Brucellosis outbreak at a pork processing plant —North Carolina, 1992. MMWR 1994; 43(7):113-6.  
Chomel BB, DeBess EE, Mangiamele DM, et al. Changing trends in the epidemiology of human brucellosis in California from 1973 to 1992: a shift toward foodborne transmission. J Infect Dis 1994 Nov;170(5):1216-23.  
Kaufmann AF, Fox MD, Boyce JM, et al. Airborne spread of brucellosis. Ann NY Acad Sci 1980;353:105-14.  
Kaufman AF, Wenger JD. Brucellosis. In: Last JM, Wallace RB, eds. Public health and preventive medicine. Norwalk, CT: Appleton and Lange, 1992:263-4.  
Staskiewicz J, Lewis CM, Colville J, Zervos M, Band J. Outbreak of *Brucella melitensis* among microbiology laboratory workers in a community hospital. J Clin Microbiol 1991;29:287-90.

### **Chancroid**

CDC. Chancroid in the United States, 1981-1990: evidence for underreporting of cases. MMWR 1992;41(No. SS-3):57-61.  
Schmid GP. Chancroid. In: Evans AS, Brachman PS, eds. Bacterial infections of humans: epidemiology and control. 2nd ed. New York: Plenum Medical Book Company, 1991;173-8.  
Schmid GP. Chancroid and *Granuloma inguinale*. In: Kelley WB, ed. Textbook of internal medicine. 2nd ed. Philadelphia: JB Lippincott Co., 1991:1412-4.

### **Cholera**

Blake PA. Epidemiology of cholera in the Americas. Gastroenterol Clin North Am 1993;22:639-60.  
Swerdlow DL, Ries AA. *Vibrio cholerae* non-O1—the eighth pandemic? Lancet 1993;342:382-3.  
Wachsmuth IK, Blake PA, Olsvik O, eds. *Vibrio cholerae* and cholera: molecular to global perspectives. Washington, DC: American Society for Microbiology, 1994.  
World Health Organization. Guidelines for cholera control. Geneva: World Health Organization, 1993.

### **Congenital Syphilis**

CDC. Surveillance for geographic and secular trends in congenital syphilis—United States, 1983-1991. MMWR 1993;42(No. SS-6):59-71.  
CDC. Evaluation of congenital syphilis surveillance system—New Jersey, 1993. MMWR 1995; 44:225-7.

- Zenker PN. Syphilis. In: Schdlower M, Shafer M-A, eds. Adolescent medicine: state of the art reviews. Philadelphia: Hanley and Belfus, Inc., 1990;1:511-29.
- Zenker PN, Berman SM. Congenital syphilis: trends and recommendations for evaluation and management. *Pediatr Infect Dis J* 1991;10(7):516-22.

#### Diphtheria

- CDC. Diphtheria acquired by U.S. citizens in the Russian Federation and Ukraine—1994. *MMWR* 1995;44:237,243-4.
- CDC. Diphtheria epidemic—New Independent States of the Former Soviet Union, 1990-1994. *MMWR* 1995;44:177-81.
- Chen RT, Broome CV, Weinstein RA, Weaver R, Tsai TF. Diphtheria in the United States, 1971-81. *Am J Public Health* 1985;75:1393-7.
- Farizo KM, Strebel PM, Chen RT, Kimbler A, Cleary TJ, Cochi SL. Fatal respiratory disease due to *Corynebacterium diphtheriae*: case report and review of guidelines for management, investigation, and control. *Clin Infect Dis* 1993;16:59-68.

#### *Escherichia coli* O157:H7

- Bell BP, Goldoft M, Griffin M, et al. A multistate outbreak of *Escherichia coli* O157:H7—associated bloody diarrhea and hemolytic uremic syndrome from hamburgers. The Washington experience. *JAMA* 1994;272:1349-53.
- Besser RE, Lett SM, Webber JT, et al. An outbreak of diarrhea and hemolytic uremic syndrome from *Escherichia coli* O157:H7 in fresh-pressed apple cider. *JAMA* 1993;269:2217-20.
- Griffin PM. *Escherichia coli* O157:H7 and other enterohemorrhagic *Escherichia coli*. In: Blaser MJ, Smith PD, Ravdin JI, Greenberg HB, Guerrant RL, eds. *Infections of the gastrointestinal tract*. New York: Raven Press, 1995:739-61.
- Griffin PM, Tauxe RV. The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. Coli*, and the associated hemolytic uremic syndrome. *Epidemiol Rev* 1991;13:60-98.

#### Gonorrhea

- CDC. Special focus: surveillance for sexually transmitted diseases. *MMWR* 1993;42(No. SS-3): 1-11, 29-39.
- CDC. Decreased susceptibility of *Neisseria gonorrhoeae* to fluoroquinolones—Ohio and Hawaii, 1992-1994. *MMWR* 1994;43:325-7.
- CDC. Increasing incidence of gonorrhea—Minnesota, 1994. *MMWR* 1995;44:282-6.

#### Granuloma inguinale

- Schmid GP. Chancroid and Granuloma inguinale. In: Kelley WB, ed. *Textbook of internal medicine*. 2nd ed. Philadelphia: JB Lippincott Co., 1989:1555-6.

#### *Haemophilus influenzae*, invasive

- CDC. Recommendations for use of Haemophilus b conjugate vaccines and a combined diphtheria, tetanus, pertussis, and Haemophilus b vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 1993;42(No. RR-13).
- Adams WG, Deaver KA, Cochi SL, et al. Decline of childhood *Haemophilus influenzae* type b (Hib) disease in the Hib vaccine era. *JAMA* 1993;269:221-6.
- CDC. Progress toward elimination of *Haemophilus influenzae* type b disease among infants and children—United States, 1987-1993. *MMWR* 1994;43(8):144-8.

#### Hansen disease (Leprosy)

- Mastro TD, Redd SC, Breiman RF. Imported leprosy in the United States, 1978 through 1988; an epidemic without secondary transmission. *Am J Public Health* 1992 Aug;82(8):1127-30.
- Noordeen SK. Epidemiology and control of leprosy—a review of progress over the last 30 years. *Trans R Soc Trap Med Hyg* 1993 Sep-Oct;87(5):515-7.
- Smith PG. Recent trends in the epidemiology of tuberculosis and leprosy. *Trop Geogr Med* 1991 Jul;43(3):S22-9.

#### Hepatitis

- Alter MJ, Mares A, Hadler SC, Maynard JE. The effect of under reporting on the apparent incidence and epidemiology of acute viral hepatitis. *Am J Epidemiol* 1987;125:133-9.
- CDC. Hepatitis surveillance report no. 55. Atlanta: US Department of Health and Human Services, Public Health Service, 1994.

### **Hepatitis A**

Shapiro CN, Coleman PJ, McQuillan GM, et al. Epidemiology of hepatitis A: seroepidemiology and risk groups in the U.S.A. *Vaccine* 1992;10(suppl 1):S59-62.

### **Hepatitis B**

Margolis HS, Alter MJ, Hadler SC. Hepatitis B: evolving epidemiology and implications for control. *Semin Liver Dis* 1991;11:84-92.

### **Hepatitis, C/Non-A, non-B**

Alter MJ, Hadler SC, Judson FN, et al. Risk factors for acute non-A, non-B hepatitis in the United States and association with hepatitis C virus infection. *JAMA* 1990;264:2231-5.

Alter MJ, Margolis HS, Krawczynski K, et al. The natural history of community-acquired hepatitis C in the United States. *N Engl J Med* 1992;327:1899-905.

### **Legionellosis**

CDC. Legionnaires' disease associated with cooling towers—Massachusetts, Michigan, and Rhode Island, 1993. *MMWR* 1994;43(27):491-9.

Edelstein PH. Legionnaires' disease. *Clin Infect Dis* 1993;16:741-9.

Hoge CW, Breiman RF. Advances in the epidemiology and control of *Legionella* infections. *Epidemiol Rev* 1991;13:329-40.

Marston BJ, Lipman HB, Breiman RF. Surveillance for Legionnaires' disease: risk factors for morbidity and mortality. *Arch Intern Med* 1994;154:2417-22.

World Health Organization. Epidemiology, prevention and control of legionellosis: memorandum from a WHO meeting. *Bull World Health Organ* 1990;68:155-64.

### **Leptospirosis**

Anderson DC, Folland DS, Fox MD, Patton CM, Kaufmann AF. Leptospirosis: a common-source outbreak due to leptospires of the Grippotyphosa serogroup. *Am J Epidemiol* 1978;107:538-44.

Faine S. Guidelines for the control of leptospirosis. Geneva, Switzerland: World Health Organization, 1982.

Takafuji ET, Kirkpatrick JW, Miller RN, et al. An efficacy trial of doxycycline chemoprophylaxis against leptospirosis. *N Engl J Med* 1984;310:497-500.

### **Lyme disease**

CDC. Lyme disease surveillance—United States, 1993. *MMWR* 1994;43(31):564-5,571-2.

Kalish R. Lyme disease. *Rheum Dis Clin North Am* 1993;19:399-426.

Steere AC. Lyme disease. *N Engl J Med* 1989;321:586-96.

### **Lymphogranuloma venereum**

Perine PL, Osoba AO. Lymphogranuloma venereum. In: Holmes KK, Mardh PA, Sparling PF, et al. eds. Sexually transmitted disease. New York: McGraw-Hill, 1990:195-204.

### **Malaria**

CDC. Health information for international travel 1994. Atlanta: CDC, 1994: US Department of Health and Human Services, Public Health Service, CDC. DHHS publication no. (CDC) 94-8280.

CDC. Local transmission of *Plasmodium vivax* malaria — Houston, Texas, 1994. *MMWR* 1994;44(15):295.

Lobel HO, Miani M, Eng T, Bernard KW, Hightower AW, Campbell CC. Long-term malaria prophylaxis with weekly mefloquine. *Lancet* 1993;341:848-51.

Zucker JR, Campbell CC. Malaria: principles of prevention and treatment. *Infect Dis Clin North Am* 1993;7:547-67.

### **Measles**

CDC. Measles prevention: recommendations of the Immunization Practices Advisory Committee. *MMWR* 1989;38(No. SS-9):1-18.

CDC. Measles—United States, first 26 weeks, 1993. *MMWR* 1993;42:813-6.

CDC. Absence of reported measles—United States, November 1993. *MMWR* 1993;42:925-6.

### **Meningococcal disease**

- CDC. Meningococcal vaccines. MMWR 1985;34(18):255-9.  
CDC. Laboratory-based surveillance for meningococcal disease in selected areas—United States, 1989–1991. MMWR 1993;42(No. SS-2):21–30.  
Jackson LA, Schuchat A, Reeves MW, Wenger JD. Serogroup C meningococcal outbreaks in the United States: an emerging threat. JAMA 1995;273(5):383–9.  
Schwartz B, Moore PS, Broome CV. Global epidemiology of meningococcal disease. Clin Microbiol Rev 1989;2(suppl):S118–24.

### **Mumps**

- Briss PA, Fehrs LJ, Parker RA, et al. Sustained transmission of mumps in a highly vaccinated population: assessment of primary vaccine failure and waning vaccine-induced immunity. J Infect Dis 1994;169:77–82.  
CDC. Mumps prevention. MMWR 1989;38(22):388–92,397–400.  
CDC. Mumps—United States, 1985–1988. MMWR 1989;38(7):101–5.  
CDC. Mumps Surveillance — United States, 1988–1993. MMWR 1995;44(No. SS-3):1–14.  
Hersch BS, Fine PEM, Kent WK, et al. Mumps outbreak in a highly vaccinated population. J Pediatr 1991;119:187–93.

### **Pertussis**

- CDC. Pertussis surveillance—United States, 1989–1991. MMWR 1992;41(No. SS-8):11–19.  
Cherry JD, Brunell PA, Golden GS, et al. Report of the Task Force on Pertussis and Pertussis Immunization, 1988. Pediatrics 1988;81(suppl):S939–84.  
Davis SF, Sutter RW, Strebel PM, et al. Concurrent outbreaks of pertussis and *Mycoplasma pneumoniae* infection: clinical and epidemiological characteristics of illnesses manifested by cough. Clin Infect Dis 1995;20(3):621–8.

### **Plague**

- Craven RB, Barnes AM. Plague and tularemia. Infect Dis Clin North Am 1991;5:165–75.  
Poland JD. Plague. In: Hoeprich PD, ed. Infectious Diseases. 2nd ed. New York: Harper and Row, 1989:151;1296–306.

### **Poliomyelitis**

- CDC. Lack of evidence for wild poliovirus circulation—United States, 1993. MMWR 1993;43:957–9.  
Prevots DR, Sutter RW, Strebel PM, Weibel RE, Cochi SL. Completeness of reporting for paralytic poliomyelitis, United States, 1980 through 1991. Archives of Pediatrics and Adolescent Medicine 1994;148:479–85.  
Strebel PM, Sutter RW, Cochi SL, et al. Epidemiology of poliomyelitis in the United States: one decade after the last reported case of indigenous wild virus-associated disease. Clin Infect Dis 1992;14:568–79.

### **Psittacosis**

- Anonymous. Compendium of chlamydiosis (psittacosis) control, 1994. National Association of State Public Health Veterinarians, Inc. JAVMA 1993;203:1673–80.  
Barnes RC. Laboratory diagnosis of human chlamydial infections. Clin Microbiol Rev 1989; 2:119–36.  
Kuritsky JN, Schmid GP, Potter ME, Anderson DC, Kaufmann AF. Psittacosis: a diagnostic challenge. J Occup Med 1984;26:731–3.  
Wong KH, Skelton SK, Daugherty H. Utility of complement fixation and microimmunofluorescence assays for detecting serologic responses in patients with clinically diagnosed psittacosis. J Clin Microbiol 1994;32:2417–21.

### **Rabies**

- CDC. Rabies prevention—United States, 1991: recommendations of the Immunization Practices Advisory Committee (ACIP). MMWR 1991;40(No. RR-3).  
Baer G. Natural history of rabies. 2nd ed. Boca Raton, FL: CRC press, 1991.  
CDC. Compendium of animal rabies control, 1995. MMWR 1995;44(No. RR-2).  
Krebs JW, Strine TW, Smith JS, Rupprecht CE, Childs JE. Rabies surveillance in the United States during 1993. JAVMA 1993;205:1695–1709.

### Rheumatic fever

- Dajani AS. Prevention of rheumatic fever: a statement for health professionals by the Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease of the Council on Cardiovascular Disease in the Young, the American Heart Association. *Pediatr Infect Dis J* 1989;8:263–6.
- Kaplan EL, Johnson DR, Cleary PP. Group A streptococcal serotypes isolated from patients and sibling contacts during the resurgence of rheumatic fever in the United States in the mid-1980's. *J Infect Dis* 1989;159:101–3.
- Veasy GL, Wiedmeier SE, Orsmond GS, et al. Resurgence of acute rheumatic fever in the intermountain area in the United States. *N Engl J Med* 1987;316:421–8.

### Rocky Mountain spotted fever (RMSF)

- Burgdorfer W, Anacker FL, eds. *Rickettsiae and rickettsial diseases*. New York: Academic Press, 1981.
- Dalton MJ, Clarke MJ, Holman RC, et al. National surveillance for Rocky Mountain spotted fever, 1981–1992, epidemiologic summary and evaluation of risk factors for fatal outcome. *Am J Trop Med Hyg* 1995;52(5):405–13.
- McDade JE, Fishbein DB. Rickettsiaceae: the rickettsiae. In: *Laboratory diagnosis of infectious diseases: principles and practice*. Vol II. Viral, rickettsial, and chlamydial diseases. New York: Springer-Verlag, 1988:864–89.

### Rubella

- CDC. Rubella prevention: recommendations of the Immunization Practices Advisory Committee (ACIP). *MMWR* 1990;39(No. RR-15).
- CDC. Increase in rubella and congenital rubella syndrome—United States, 1988–1990. *MMWR* 1991;40(6):93–9.
- CDC. Outbreaks of rubella among the Amish—United States, 1991. *MMWR* 1991;40(16):264.
- CDC. Congenital rubella syndrome among the Amish—Pennsylvania, 1991–1992. *MMWR* 1992; 41(26):468–9,475–6.
- CDC. Rubella and congenital rubella syndrome—United States, January 1, 1991–May 7, 1994. *MMWR* 1994;43(21):391,397–401.
- Lindgren ML, Fehrs LJ, Hadler SC, Hinman AR. Update: rubella and congenital rubella syndrome, 1980–1990. *Epidemiol Rev* 1991;13:341–8.

### Salmonellosis

- Lee LA, Puhr ND, Maloney EK, Bean NH, Tauxe RV. Increase in antimicrobial-resistant *Salmonella* infections in the United States, 1989–1990. *J Infect Dis* 1994;170:128–34.
- Mishu B, Griffin PM, Tauxe RV, Cameron DN, Hutcheson RH, Schaffner W. *Salmonella enteritidis* gastroenteritis transmitted by intact chicken eggs. *Ann Intern Med* 1991;115:190–4.
- Mishu B, Koehler J, Lee LA, et al. Outbreaks of *Salmonella enteritidis* infections in the United States, 1985–1991. *J Infect Dis* 1994;169:547–52.
- St. Louis ME, Morse DL, Potter ME, et al. The emergence of grade A eggs as a major source of *Salmonella enteritidis* infections: new implications for the control of salmonellosis. *JAMA* 1988;259:2103–7.
- Tauxe RV. *Salmonella*: a postmodern pathogen. *Journal of Food Protection* 1991;54:563–8.

### Shigellosis

- Lee LA, Shapiro CN, Hargrett-Bean N, Tauxe RV. Hyperendemic shigellosis in the United States: a review of surveillance data for 1967–1988. *J Infect Dis* 1991;164:894–900.
- Parsonnet J, Greene KD, Gerber AR, et al. *Shigella dysenteriae* type 1 infections in U.S. travellers to Mexico. *Lancet* 1989;543–5.
- Ries AA, Wells JG, Olivola D, et al. Epidemic *Shigella dysenteriae* type 1 in Burundi: panresistance and implications for prevention. *J Infect Dis* 1994;169:1035–41.

### Syphilis

- Aral SO, Wasserheit JN, Green SB, Judson FN, Sparling FP. The NIAID/NIH working group on integrated behavioral research for prevention and control of STDs. Part III: Issues in evaluating behavioral interventions. *Sex Transm Dis* 1990;17:208–10.
- CDC. Primary and secondary syphilis—United States, 1981–1990. *MMWR* 1991;40(19):314–5, 321–3.
- CDC. Regional and temporal trends in the surveillance of syphilis, United States, 1986–1990. *MMWR* 1991;40(No. SS-3):29–33.

CDC. Special focus: surveillance for sexually transmitted diseases. MMWR 1993;42(No. SS-3):13-9.  
Nakashima AK, Rolfs RT, Flock ML, Kilmarx P, Greenspan GR. Epidemiology of syphilis in the United States, 1941-1993. Sex Transm Dis 1995 (in press).

#### Tetanus

CDC. Surveillance of tetanus—United States, 1989-1990. MMWR 1992;41(No. SS-8):1-9.  
Gergen PJ, McQuillian GM, Kiely M, et al. A population-based survey of immunity to tetanus in the United States. N Engl J Med 1995;332:761-6.  
Sutter RW, Cochi SL, Brink EW, Sirotnik BI. Assessment of vital statistics and surveillance data for monitoring tetanus mortality, United States, 1979-1984. Am J Epidemiol 1990;131:132-42.

#### Toxic-shock syndrome

CDC. Reduced incidence of menstrual toxic shock syndrome—United States, 1980-1990. MMWR 1990;39(25):421-3.  
Gaventa S, Reingold AL, Hightower AW, et al. Active surveillance for toxic shock syndrome in the United States, 1986. Rev Infect Dis 1989;(suppl):S28-34.  
Schuchat A, Broome CV. Toxic shock syndrome and tampons. Epidemiol Rev 1991;13:99-112.

#### Trichinosis

Bailey TM, Schantz PM. Trends in the incidence and transmission patterns of human trichinosis in the United States, 1982-1986. Rev Infect Dis 1990;12:5-11.  
Campbell WC, ed. *Trichinella and trichinosis*. New York: Plenum Press, 1983.  
CDC. Trichinosis surveillance—United States, 1987-1990. MMWR 1991;40(No. SS-3):35-42.  
McAuley JB, Michelson MK, Hightower AW, Engeran S, Wintermeyer LA, Schantz PM. A trichinosis outbreak among Southeast Asian refugees. Am J Epidemiol 1992;135:1404-10.

#### Tuberculosis

American Thoracic Society/CDC. Treatment of tuberculosis and tuberculosis infection in adults and children. Am J Respir Crit Care Med 1994;149:1359-74.  
CDC. Recommendations for counting reported tuberculosis cases. Atlanta: US Department of Health and Human Services, Public Health Service, 1977.  
CDC. Tuberculosis morbidity—United States, 1994. MMWR 1995;44:(20)387-95.

#### Tularemia

Evans ME, Gregory DW, Schaffner W, McGee ZA. Tularemia: a 30 year experience with 88 cases. Medicine 1985;64:251-69.  
Schmid GP, Kornblatt AN, Connors CA, et al. Clinically mild tularemia associated with tick-borne *Francisella tularensis*. J Infect Dis 1983;148:63-7.  
Taylor JP, Istre GR, McChesny TC, Satalowich FT, Parker RE, McFarland LM. Epidemiologic characteristics of human tularemia in the southwest-central states, 1981-1987. Am J Epidemiol 1991;133:1032-8.

#### Typhoid fever

CDC. Typhoid immunization: recommendations of the Advisory Committee on Immunization Practices. MMWR 1994;43(No. RR-14).  
Ryan CA, Hargrett-Bean NT, Blake PA. *Salmonella typhi* infections in the United States, 1975-1984: increasing role of foreign travel. Rev Infect Dis 1989;11(1):1-8.  
Woodruff BA, Pavia AT, Blake PA. A new look at typhoid vaccination: information for the practicing physician. JAMA 1991;265(6):756-9.

#### Varicella

CDC. Varicella outbreak in a women's prison—Kentucky. MMWR 1989;38(37):635-6,641-2.  
Gershon AA, LaRussa P, Hardy I, Steinberg S, Silverstein S. Varicella vaccine: the American experience. J Infect Dis 1992;166(suppl 1):S63-8.  
Lieu TA, Cochi SL, Black SB, et al. Cost-effectiveness of a routine varicella vaccination program for U.S. children. JAMA 1994;271:375-81.  
Varicella Vaccine. Medical Letter 1995;37(950):55-7.

# **PART 1:**

**Summaries of  
Notifiable Diseases  
in the United States**

## NOTIFIABLE DISEASES — Summary of reported cases, by month, United States, 1994

NAME	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Unk.
AIDS*	78,279	7,058	4,446	8,774	5,954	6,136	7,100	6,305	7,814	7,563	5,800	6,056	5,273	-
Amebiasis	2,983	158	167	196	315	215	258	226	197	280	202	488	-	-
Anthrax	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Aseptic meningitis	8,932	351	322	423	516	497	690	1,043	1,067	1,030	1,086	7,180	1,189	-
Botulism, total	143	13	8	12	7	19	11	12	8	15	9	9	10	-
Brucellosis	119	4	3	6	9	8	5	18	11	10	1	9	35	-
Chancroid†	773	.....	198	.....	204	.....	1	1	3	1	4	8	8	-
Cholera	39	-	2	3	3	.....	.....	.....	.....	.....	.....	.....	.....	-
Diphtheria	2	-	2	2	-	-	-	-	-	-	-	-	-	-
Encephalitis, primary infections	717	42	56	43	47	36	43	67	66	86	80	52	99	-
Post-infectious	143	7	9	12	17	8	13	14	12	10	12	11	18	-
<i>Escherichia coli</i> O157:H7	1,420	14	28	23	51	75	88	227	208	150	195	139	222	-
Gonorrhea†	418,068	.....	92,183	.....	99,347	.....	111,415	.....	115,123	.....	115,123	.....	1	-
Granuloma inguinale†	3	84	98	91	107	145	72	111	55	69	107	69	166	-
<i>Haemophilus influenzae</i> , invasive	1,174	136	11	7	17	5	15	18	8	6	15	12	15	-
Hansen disease (leprosy)	26,796	1,235	1,788	1,669	2,250	1,863	1,835	2,597	2,042	2,191	2,820	2,128	4,378	-
Hepatitis A	12,517	708	936	959	1,159	906	834	1,093	891	883	1,054	975	2,119	-
Hepatitis B	4,470	309	276	388	375	296	361	352	297	387	329	318	782	-
Hepatitis, Cytom-A non-B	444	21	30	22	48	34	38	42	33	38	39	32	67	-
Hepatitis, unspecified	1,615	85	112	105	141	100	122	161	175	145	148	120	201	-
Legionellosis	38	6	1	1	5	-	4	4	2	6	6	3	4	-
Leptospirosis	13,043	386	260	377	513	450	831	2,622	1,841	1,528	1,549	1,158	1,528	-
Lyme disease	235	.....	74	.....	54	.....	54	.....	52	.....	55	.....	55	-
Lymphogranuloma venereum†	1,229	48	91	99	85	56	71	101	126	104	122	70	256	-
Malaria	1,963	6	38	67	241	298	97	71	15	23	19	11	77	-
Measles (rubella)	2,886	245	278	274	290	285	198	212	140	124	199	197	444	-
Meningococcal disease	1,537	71	123	108	153	134	114	126	163	70	141	90	244	-
Mumps	4,617	257	344	260	274	281	274	362	402	306	603	312	942	-
Pertussis (whooping cough)	17	1	-	-	-	2	6	6	1	-	-	1	-	-
Plague	38	3	5	4	-	-	-	-	-	-	-	-	-	-
Poliomyelitis, paralytic§	8,147	474	500	867	759	667	692	628	845	764	759	645	547	-
Poxtacosis	6	1	-	-	-	-	1	-	-	-	2	2	-	-
Rabies, animal	112	6	3	8	13	6	29	7	5	3	10	6	16	-
Rheumatic fever, acute	465	5	6	11	18	20	54	82	84	46	57	28	54	-
Rocky Mountain spotted fever	227	3	31	50	50	12	25	27	4	5	2	2	16	-
Rubella (German measles)	7	-	2	-	-	-	-	-	-	-	-	-	-	-
Rubella, congenital syndrome	43,323	1,560	1,656	1,899	2,799	2,529	3,040	4,776	4,666	4,436	6,057	3,620	6,275	-
Salmonellosis	29,769	1,293	1,406	1,734	2,319	2,070	2,126	2,969	2,597	2,533	3,190	2,541	4,981	-
Shigellosis	81,696	20,981	20,981	21,544	21,544	20,462	20,462	20,462	20,462	20,462	20,462	18,709	18,709	-
Syphilis, total all stages†	20,627	5,308	5,308	5,154	5,154	5,396	5,396	5,396	5,396	5,396	5,396	4,769	4,769	-
Primary and secondary†	2,204	535	535	597	597	525	525	525	525	525	525	547	547	-
Congenital <1 year†	51	1	2	4	4	3	4	3	1	5	8	3	13	-
Tetanus	192	13	21	13	28	16	11	10	19	14	13	11	23	-
Toxic-shock syndrome	32	10	3	5	1,877	2,138	1,949	2,105	2,063	1,964	2,009	2	5	2
Trichinosis	24,361	567	1,302	1	1	7	14	24	10	14	7	1,906	4,621	-
Tuberculosis	96	-	-	-	-	-	-	-	-	-	-	-	18	-
Tularemia	441	17	29	35	51	39	32	44	33	53	43	18	47	-
Typhoid fever	151,219	11,939	14,564	19,132	27,154	22,550	19,723	6,554	2,238	1,984	4,241	7,415	13,715	-

\* AIDS total updated through December 31, 1994.  
 † Cases updated through February 28, 1995.  
 § Two suspected cases of paralytic poliomyelitis were reported in 1994. Confirmation of these cases is pending review by external panel.

**SUMMARY TABLES — 1994**

**NOTIFIABLE DISEASES — Reported cases, by geographic division and area,  
United States, 1994**

Area	Total resident population (in thousands)	AIDS	Amebiasis	Anthrax	Aseptic meningitis	Botulism		Brucellosis	Chancroid
						Foodborne	Infant		
United States	260,341	78,279*	2,983	-	8,932	50	85	119	773†
New England	13,270	2,836	80	-	340	-	1	-	1
Maine	1,240	117	10	-	33	-	-	-	-
N.H.	1,137	92	2	-	47	-	-	-	-
Vt.	580	38	3	-	36	-	-	-	-
Mass.	6,041	1,401	61	-	103	-	-	-	1
R.I.	997	276	4	-	121	-	-	-	-
Conn.	3,275	912	NN	-	NN	-	1	-	-
Mid. Atlantic	38,125	22,465	876	-	957	2	16	2	365
N.Y. (excl. NYC)	10,905	2,220	120	-	466	1	-	-	8
N.Y.C.	7,264	12,724	701	-	150	-	1	-	357
N.J.	7,904	4,993	25	-	NN	-	2	-	-
Pa.	12,052	2,528	30	-	341	1	13	2	-
E.N. Central	43,184	6,324	187	-	1,652	2	9	8	48
Ohio	11,102	1,184	22	-	399	1	-	2	8
Ind.	5,752	622	21	-	222	-	2	-	-
III.	11,752	3,104	45	-	472	1	2	5	38
Mich.	9,496	1,035	42	-	538	-	1	1	-
Wis.	5,082	379	57	-	21	-	-	-	2
W.N. Central	18,210	1,638	131	-	476	-	2	1	8
Minn.	4,567	422	39	-	43	-	-	-	-
Iowa	2,829	130	21	-	121	-	-	1	1
Mo.	5,278	713	38	-	175	-	-	-	2
N. Dak.	638	20	8	-	14	-	-	-	-
S. Dak.	721	19	4	-	3	-	-	-	-
Nebr.	1,623	89	6	-	41	-	-	-	-
Kans.	2,554	245	15	-	79	-	2	-	5
S. Atlantic	46,398	18,857	203	-	2,000	-	4	17	30
Del.	706	271	3	-	41	-	1	-	-
Md.	5,006	2,722	14	-	246	-	1	1	-
D.C.	570	1,399	2	-	53	-	-	-	-
Va.	6,552	1,162	39	-	337	-	2	2	-
W. Va.	1,822	96	6	-	39	-	-	-	-
N.C.	7,070	1,187	19	-	240	-	-	3	10
S.C.	3,664	1,158	NN	-	31	-	-	-	-
Ga.	7,055	2,245	58	-	80	-	-	3	-
Fla.	13,953	8,617	62	-	933	-	-	8	20
E.S. Central	15,890	2,099	10	-	582	3	3	2	27
Ky.	3,827	320	6	-	181	-	1	-	-
Tenn.	5,175	764	NN	-	164	2	1	1	3
Ala.	4,219	582	3	-	174	-	-	-	24
Miss.	2,669	433	1	-	63	1	1	1	-
W.S. Central	28,404	7,671	124	-	1,072	25	4	32	260
Ark.	2,453	284	4	-	62	-	1	-	-
La.	4,315	1,239	2	-	40	-	-	2	209
Okla.	3,258	269	8	-	-	1	-	-	-
Tex.	18,378	5,879	110	-	970	24	3	29	51
Mountain	15,215	2,287	150	-	353	-	8	20	3
Mont.	856	30	-	-	8	-	1	-	-
Idaho	1,133	61	6	-	6	-	2	-	-
Wyo.	476	18	-	-	4	-	1	-	-
Colo.	3,656	816	39	-	135	-	1	1	-
N. Mex.	1,654	211	27	-	20	-	-	1	-
Ariz.	4,075	612	61	-	79	-	-	17	3
Utah	1,908	152	6	-	55	-	3	1	-
Nev.	1,457	387	11	-	46	-	-	-	-
Pacific	41,645	13,949	1,222	-	1,500	18	38	37	31
Wash.	5,343	932	78	-	NN	3	2	-	1
Oreg.	3,086	606	109	-	NN	-	3	1	5
Calif.	31,431	12,136	991	-	1,350	4	30	36	25
Alaska	606	59	5	-	19	11	-	-	-
Hawaii	1,179	216	39	-	131	-	3	-	-
Guam	133	1	1	-	13	-	-	-	-
P.R.	3,522	2,359	1	-	72	-	-	-	32
V.I.	102	52	-	-	-	-	-	-	1
C.N.M.I.	43	-	-	-	-	-	-	-	-
American Samoa	47	-	-	-	-	-	-	-	-

\*Total reported to Division of HIV/AIDS Prevention, National Center for Prevention Services, through December 31, 1994. Total includes 153 cases with unknown state of residence.

†Cases updated through Division of Sexually Transmitted Diseases and HIV Prevention, National Center for Prevention Services, as of February 28, 1995.

**NOTIFIABLE DISEASES — Reported cases, by geographic division and area,  
United States, 1994 (continued)**

Area			Encephalitis		<i>Escherichia coli</i> O157:H7	Gonor- rhea	Granuloma inguinale	<i>Haemophilus</i> <i>influenzae</i> , invasive	Hansen disease (leprosy)
	Cholera	Diphtheria	Primary infections	Post- infectious					
United States	39	2	717	143	1,420	418,068*	3*	1,174	136
New England	2	1	18	6	223	8,640	-	45	4
Maine	-	-	5	-	NN	93	-	5	-
N.H.	-	-	-	2	NN	103	-	4	-
Vt.	-	-	3	1	12	40	-	-	-
Mass.	1	1	8	1	134	3,159	-	21	4
R.I.	-	-	2	2	9	478	-	4	-
Conn.	1	-	-	-	68	4,767	-	11	-
Mid. Atlantic	6	-	63	20	160	49,450	-	149	16
N.Y. (excl. NYC)	3	-	38	3	149	11,506	-	70	2
N.Y.C.	1	-	6	5	11	19,491	-	34	13
N.J.	2	-	-	-	-	5,269	-	16	1
Pa.	-	-	19	12	NN	13,184	-	29	-
E.N. Central	5	-	169	22	269	87,065	-	199	9
Ohio	-	-	55	4	109	24,746	-	109	-
Ind.	-	-	11	1	57	9,757	-	26	-
Ill.	2	-	65	5	103	26,571	-	43	6
Mich.	3	-	38	12	-	18,215	-	19	-
Wis.	-	-	-	-	NN	7,776	-	2	3
W.N. Central	1	-	52	12	366	22,834	-	103	-
Minn.	-	-	22	-	147	3,346	-	34	-
Iowa	1	-	1	1	56	1,645	-	9	-
Mo.	-	-	8	6	40	12,557	-	51	-
N. Dak.	-	-	4	-	31	35	-	-	-
S. Dak.	-	-	4	-	18	243	-	2	-
Nebr.	-	-	6	5	74	1,335	-	4	-
Kans.	-	-	7	-	NN	3,673	-	3	-
S. Atlantic	3	-	161	65	23	104,591	2	234	2
Del.	-	-	1	-	NN	2,038	-	1	-
Md.	1	-	28	5	NN	15,137	-	87	-
D.C.	-	-	-	1	-	6,827	-	-	-
Va.	1	-	34	6	NN	13,414	-	22	1
W. Va.	-	-	51	-	NN	805	-	7	-
N.C.	-	-	44	1	6	28,936	2	32	-
S.C.	-	-	-	-	17	13,067	-	3	-
Ga.	-	-	2	-	NN	NA	-	67	-
Fla.	1	-	1	52	-	24,367	-	15	1
E.S. Central	-	-	41	4	6	48,208	-	34	1
Ky.	-	-	16	2	6	5,127	-	4	1
Tenn.	-	-	12	-	NN	15,745	-	10	-
Ala.	-	-	9	1	-	15,881	-	15	-
Miss.	-	-	4	1	-	11,455	-	5	-
W.S. Central	4	-	63	2	98	53,529	-	74	33
Ark.	-	-	1	-	10	6,892	-	5	2
La.	-	-	10	-	-	11,992	-	4	-
Okla.	-	-	-	-	16	4,888	-	45	-
Tex.	4	-	52	2	72	29,757	-	20	31
Mountain	-	-	14	3	100	10,669	-	128	-
Mont.	-	-	-	-	-	85	-	1	-
Idaho	-	-	-	-	-	98	-	5	-
Wyo.	-	-	3	1	NN	82	-	5	-
Colo.	-	-	4	-	76	3,632	-	17	-
N. Mex.	-	-	-	-	-	1,130	-	12	-
Ariz.	-	-	-	1	NN	3,603	-	26	-
Utah	-	-	3	1	NN	303	-	10	-
Nev.	-	-	4	-	24	1,736	-	52	-
Pacific	18	1	136	9	175	33,082	1	208	71
Wash.	-	-	1	-	174	2,893	-	10	7
Oreg.	-	-	-	-	1	978	1	26	-
Calif.	17	1	131	8	NN	27,593	-	165	43
Alaska	-	-	4	-	NN	918	-	3	-
Hawaii	1	-	-	1	NN	700	-	4	21
Guam	1	-	-	1	NN	110	-	-	11
P.R.	-	-	1	3	NN	500	-	3	1
V.I.	-	-	-	-	-	60	-	-	-
C.N.M.I.	-	-	-	-	-	-	-	26	-
American Samoa	-	-	-	-	-	-	-	-	4

\*Cases updated through Division of Sexually Transmitted Diseases and HIV Prevention, National Center for Prevention Services, as of February 28, 1995.

**SUMMARY TABLES — 1994**

**NOTIFIABLE DISEASES — Reported cases, by geographic division and area,  
United States, 1994 (continued)**

Area	Hepatitis A	Hepatitis B	Hepatitis C/non-A, non-B	Hepatitis unsp.	Legionellosis	Leptospirosis	Lyme disease	Lympho-granuloma venereum
<b>United States</b>	<b>26,796</b>	<b>12,517</b>	<b>4,470</b>	<b>444</b>	<b>1,615</b>	<b>38</b>	<b>13,043</b>	<b>235*</b>
<b>New England</b>	<b>296</b>	<b>354</b>	<b>168</b>	<b>15</b>	<b>79</b>	<b>3</b>	<b>2,827</b>	<b>6</b>
Maine	25	11	-	-	5	2	33	-
N.H.	17	28	11	-	-	-	30	-
Vt.	14	12	16	-	1	-	16	-
Mass.	112	200	121	13	55	-	247	6
R.I.	30	8	20	2	18	-	471	-
Conn.	98	95	-	-	NN	1	2,030	-
<b>Mid. Atlantic</b>	<b>2,007</b>	<b>1,761</b>	<b>489</b>	<b>10</b>	<b>264</b>	<b>2</b>	<b>8,171</b>	<b>108</b>
N.Y. (excl. NYC)	543	402	230	6	62	1	5,105	2
N.Y.C.	941	543	4	-	11	-	95	106
N.J.	306	410	211	-	49	1	1,533	-
Pa.	217	406	44	4	142	-	1,438	-
<b>E.N. Central</b>	<b>2,777</b>	<b>1,221</b>	<b>320</b>	<b>16</b>	<b>433</b>	<b>1</b>	<b>530</b>	<b>9</b>
Ohio	1,203	164	24	-	194	-	45	9
Ind.	361	215	9	-	48	-	19	-
III.	615	315	81	9	44	1	24	-
Mich.	352	432	206	7	82	-	33	-
Wis.	246	95	-	-	65	-	409	-
<b>W.N. Central</b>	<b>1,222</b>	<b>714</b>	<b>100</b>	<b>12</b>	<b>106</b>	<b>2</b>	<b>347</b>	<b>2</b>
Minn.	261	82	20	-	4	-	208	-
Iowa	64	27	14	11	34	-	17	-
Mo.	619	538	32	1	41	1	102	2
N. Dak.	6	1	1	-	4	-	-	-
S. Dak.	39	4	-	-	2	-	-	-
Nebr.	122	31	15	-	15	1	3	-
Kans.	111	31	18	-	6	-	17	-
<b>S. Atlantic</b>	<b>1,466</b>	<b>2,240</b>	<b>485</b>	<b>32</b>	<b>413</b>	<b>3</b>	<b>855</b>	<b>65</b>
Del.	22	14	2	-	31	-	106	-
Md.	198	354	21	8	82	1	341	-
D.C.	27	53	2	-	9	-	9	15
Va.	193	142	26	10	17	-	131	-
W. Va.	23	48	47	-	4	-	29	-
N.C.	145	291	59	-	28	1	77	44
S.C.	40	33	10	-	29	-	7	-
Ga.	43	555	220	-	118	-	127	3
Fla.	775	750	98	14	95	1	28	3
<b>E.S. Central</b>	<b>784</b>	<b>1,211</b>	<b>945</b>	<b>2</b>	<b>83</b>	<b>3</b>	<b>43</b>	<b>2</b>
Ky.	221	78	32	-	9	1	24	-
Tenn.	347	1,042	893	1	45	2	13	1
Ala.	139	91	20	1	13	-	6	1
Miss.	77	-	-	-	16	-	-	-
<b>W.S. Central</b>	<b>3,719</b>	<b>1,830</b>	<b>599</b>	<b>94</b>	<b>63</b>	<b>1</b>	<b>174</b>	<b>12</b>
Ark.	253	60	8	3	16	1	15	-
La.	170	203	215	2	20	-	4	12
Okla.	419	141	62	3	12	-	99	-
Tex.	2,877	1,426	314	86	15	-	56	-
<b>Mountain</b>	<b>5,296</b>	<b>694</b>	<b>454</b>	<b>73</b>	<b>97</b>	<b>-</b>	<b>18</b>	<b>7</b>
Mont.	25	21	13	-	16	-	-	-
Idaho	380	77	71	1	2	-	3	-
Wyo.	41	24	177	-	5	-	5	-
Colo.	584	97	79	14	19	-	1	1
N. Mex.	1,100	218	45	11	4	-	5	-
Ariz.	2,159	102	31	27	17	-	-	5
Utah	754	96	18	6	8	-	3	-
Nev.	253	59	20	14	26	-	1	1
<b>Pacific</b>	<b>9,229</b>	<b>2,492</b>	<b>910</b>	<b>190</b>	<b>77</b>	<b>23</b>	<b>78</b>	<b>24</b>
Wash.	1,119	255	294	9	13	-	4	3
Oreg.	1,241	158	46	2	-	1	6	2
Calif.	6,602	2,038	565	176	59	-	68	19
Alaska	209	13	-	-	-	-	-	-
Hawaii	58	28	5	3	5	22	-	-
Guam	23	5	-	9	1	-	-	-
P.R.	86	415	215	3	-	2	-	-
V.I.	3	9	1	-	-	-	-	-
C.N.M.I.	12	1	-	-	-	-	-	-
American Samoa	11	-	-	-	-	-	-	-

\*Cases updated through Division of Sexually Transmitted Diseases and HIV Prevention, National Center for Prevention Services, as of February 28, 1995.

**NOTIFIABLE DISEASES — Reported cases, by geographic division and area,  
United States, 1994 (continued)**

Area	Malaria	Measles		Meningo- coccoal disease	Mumps	Pertussis	Plague	Polio- myelitis, paralytic
		Indigenous	Imported					
United States	1,229	746	217*	2,886	1,537	4,617	17	†
New England	78	15	12	141	26	760	-	-
Maine	6	2	3	23	3	21	-	-
N.H.	3	1	-	8	4	107	-	-
Vt.	3	2	1	5	-	46	-	-
Mass.	38	1	6	68	3	534	-	-
R.I.	10	5	2	-	4	8	-	-
Conn.	18	4	-	37	12	44	-	-
Mid. Atlantic	261	200	27	312	116	695	-	-
N.Y. (excl. NYC)	60	12	16	105	33	254	-	-
N.Y.C.	106	11	4	40	12	224	-	-
N.J.	57	172	3	65	13	15	-	-
Pa.	38	5	4	102	58	202	-	-
E.N. Central	117	60	46	397	267	615	-	-
Ohio	20	15	2	121	77	162	-	-
Ind.	15	-	1	55	7	97	-	-
Ill.	48	18	41	125	110	111	-	-
Mich.	31	24	2	59	59	96	-	-
Wis.	3	3	-	37	14	149	-	-
W.N. Central	48	127	44	174	71	273	-	-
Minn.	16	-	-	23	5	142	-	-
Iowa	5	6	1	21	16	23	-	-
Mo.	14	119	42	78	44	45	-	-
N. Dak.	1	-	-	1	4	5	-	-
S. Dak.	-	-	-	9	-	26	-	-
Nebr.	5	1	1	14	1	14	-	-
Kans.	7	1	-	28	1	18	-	-
S. Atlantic	247	65	9	455	257	431	-	-
Del.	3	-	-	5	-	3	-	-
Md.	83	2	2	35	65	74	-	-
D.C.	15	-	-	7	-	11	-	-
Va.	37	1	2	69	48	37	-	-
W. Va.	-	37	-	14	5	6	-	-
N.C.	12	2	1	57	73	140	-	-
S.C.	5	-	-	40	8	14	-	-
Ga.	43	5	-	82	18	37	-	-
Fla.	49	18	4	146	40	109	-	-
E.S. Central	32	28	-	195	32	129	-	-
Ky.	12	-	-	42	-	60	-	-
Tenn.	10	28	-	40	9	22	-	-
Ala.	9	-	-	77	12	35	-	-
Miss.	1	-	-	36	11	12	-	-
W.S. Central	119	16	7	392	302	246	-	-
Ark.	5	5	-	55	7	33	-	-
La.	12	-	1	47	38	15	-	-
Okla.	9	-	-	53	23	38	-	-
Tex.	93	11	6	237	234	160	-	-
Mountain	41	163	55	178	162	609	15	-
Mont.	-	-	-	6	-	12	-	-
Idaho	2	1	-	17	10	172	-	-
Wyo.	1	-	-	9	3	-	-	-
Colo.	19	24	37	41	4	228	2	-
N. Mex.	3	2	-	17	NN	35	7	-
Ariz.	10	5	4	58	99	122	5	-
Utah	4	131	5	19	28	37	1	-
Nev.	2	-	9	11	18	3	-	-
Pacific	286	72	17	642	304	859	2	-
Wash.	45	3	1	111	23	140	-	-
Oreg.	17	-	2	143	NN	106	-	-
Calif.	207	51	10	374	258	594	2	-
Alaska	2	10	-	5	4	-	-	-
Hawaii	15	8	4	9	19	19	-	-
Guam	-	228	-	2	7	2	-	-
PR.	5	46	-	7	2	3	-	-
V.I.	1	-	-	-	4	-	-	-
C.N.M.I.	1	29	-	-	2	-	-	-
American Samoa	-	-	-	-	3	1	-	-

\*For 1994, includes both 142 cases of out-of-state importations and 75 cases of international importations.

†Two suspected cases of paralytic poliomyelitis were reported in 1994. Confirmation of these cases is pending review by external panel.

**SUMMARY TABLES — 1994**

**NOTIFIABLE DISEASES — Reported cases, by geographic division and area,  
United States, 1994 (continued)**

Area	Psitta-cosis	Rabies		Rheumatic fever, acute	RMSF*	Rubella		Cong. syndrome	Salmonel-llosis	Shigel-llosis
		Animal	Human			Rubella	Cong. syndrome			
United States	38	8,147	6	112	465	227	7	43,323	29,769	
New England	2	2,009	-	2	12	132	2	3,439	508	
Maine	-	10	-	1	-	-	-	191	10	
N.H.	1	221	-	NN	-	-	-	213	20	
Vt.	1	143	-	-	-	-	-	119	4	
Mass.	-	734	-	1	4	126	2	2,009	243	
R.I.	-	153	-	-	-	3	-	248	55	
Conn.	-	748	-	-	8	3	-	659	176	
Mid. Atlantic	6	2,249	-	5	24	8	-	7,066	3,163	
N.Y. (excl. NYC)	1	1,569	-	NN	8	6	-	1,977	1,120	
N.Y.C.	-	16	-	NN	3	1	-	1,889	1,007	
N.J.	1	278	-	5	5	1	-	1,160	522	
Pa.	4	386	-	NN	8	-	-	2,040	514	
E.N. Central	7	69	-	30	40	10	-	5,678	3,648	
Ohio	-	4	-	8	19	-	-	1,337	740	
Ind.	1	14	-	2	8	-	-	581	544	
Ill.	-	21	-	8	11	1	-	1,821	1,494	
Mich.	3	14	-	8	2	9	-	869	432	
Wis.	3	16	-	4	-	-	-	1,070	438	
W.N. Central	4	232	-	12	42	2	-	2,624	2,361	
Minn.	-	22	-	3	1	-	-	759	554	
Iowa	-	90	-	3	-	-	-	404	338	
Mo.	4	27	-	3	22	2	-	642	654	
N. Dak.	-	14	-	NN	-	-	-	68	59	
S. Dak.	-	44	-	2	13	-	-	143	207	
Nebr.	-	-	-	NN	1	-	-	209	426	
Kans.	-	35	-	1	4	-	-	399	123	
S. Atlantic	4	2,083	2	-	224	22	-	9,165	8,352	
Del.	-	74	-	NN	1	-	-	168	38	
Md.	2	520	-	NN	21	-	-	1,178	323	
D.C.	-	4	-	NN	-	-	-	118	70	
Va.	2	428	-	NN	22	-	-	1,135	656	
W. Va.	-	84	1	-	2	-	-	152	15	
N.C.	-	175	-	NN	88	-	-	1,137	1,970	
S.C.	-	179	-	NN	20	-	-	599	505	
Ga.	-	367	-	NN	62	7	-	1,583	1,886	
Fla.	-	252	1	NN	8	15	-	3,095	2,889	
E.S. Central	1	242	2	-	47	-	-	1,777	1,706	
Ky.	-	28	-	NN	10	-	-	380	208	
Tenn.	-	82	1	NN	29	-	-	441	418	
Ala.	1	128	1	NN	2	-	-	507	617	
Miss.	-	4	-	-	6	NN	-	449	463	
W.S. Central	-	741	1	1	63	13	1	3,578	3,259	
Ark.	-	38	-	1	18	-	-	534	193	
La.	-	73	-	NN	1	-	-	591	416	
Okla.	-	39	-	NN	37	4	-	470	240	
Tex.	-	591	1	NN	7	9	1	1,983	2,410	
Mountain	3	154	-	39	13	5	2	2,226	1,953	
Mont.	-	22	-	NN	4	-	-	145	4	
Idaho	-	4	-	1	-	-	-	130	59	
Wyo.	-	24	-	1	2	-	-	60	8	
Colo.	3	18	-	8	4	-	-	709	530	
N. Mex.	-	8	-	1	1	-	1	353	347	
Ariz.	-	56	-	NN	1	-	1	427	680	
Utah	-	13	-	28	-	4	-	202	240	
Nev.	-	9	-	NN	1	1	-	200	85	
Pacific	11	368	1	23	-	35	2	7,770	4,819	
Wash.	4	15	-	-	-	-	-	863	478	
Oreg.	2	13	-	NN	-	4	-	313	165	
Calif.	4	294	1	18	-	27	2	6,235	3,953	
Alaska	1	46	-	5	-	-	-	55	21	
Hawaii	-	-	-	NN	-	4	-	304	202	
Guam	-	-	-	3	-	1	-	76	33	
P.R.	-	77	-	-	-	-	-	737	48	
V.I.	-	-	-	-	-	-	-	2	4	
C.N.M.I.	-	-	-	4	-	-	-	78	60	
American Samoa	-	-	-	-	-	-	-	14	9	

\*Rocky Mountain spotted fever.

**NOTIFIABLE DISEASES — Reported cases, by geographic division and area,  
United States, 1994 (continued)**

Area	Syphilis				Toxic-shock syndrome	Trich-inosis	Tuber-culosis	Tularemia	Typhoid fever	Varicella (chicken-pox)
	Primary & secondary	Cong. (<1 yr.)	All stages	Tetanus						
United States	20,627*	2,204*	81,696*	51	192	32	24,361	96	441	151,219
New England	219	14	1,128	2	5	1	595	1	25	11,676
Maine	4	-	9	-	1	1	35	-	-	711
N.H.	4	-	18	1	-	-	17	-	-	3,408
Vt.	-	-	1	-	2	-	10	-	-	NN
Mass.	90	6	622	-	2	-	329	1	20	5,903
R.I.	16	2	141	-	-	-	56	-	1	1,654
Conn.	105	6	337	1	-	-	148	-	4	NN
Mid. Atlantic	1,446	681	14,302	7	30	5	5,112	2	121	5,978
N.Y. (excl. NYC)	173	59	1,375	3	16	2	641	1	12	NN
N.Y.C.	629	329	8,001	-	-	-	2,995	1	78	5,978
N.J.	240	178	2,188	2	-	2	855	-	25	NN
Pa.	404	115	2,738	2	14	1	621	-	6	NN
E.N. Central	3,162	386	9,492	8	43	3	2,236	6	68	77,332
Ohio	1,187	71	2,740	1	10	-	337	1	7	5,495
Ind.	286	11	844	2	3	-	211	-	4	NN
III.	1,099	258	3,877	1	15	-	1,117	3	42	33,889
Mich.	292	28	1,234	4	15	1	462	1	6	37,948
Wis.	298	18	797	-	-	2	109	1	9	NA
W.N. Central	1,203	82	2,663	4	28	2	610	38	2	18,210
Minn.	56	2	201	1	2	-	140	1	1	NN
Iowa	75	6	235	1	8	1	66	-	-	4,197
Mo.	987	72	1,985	1	7	1	260	24	1	10,147
N. Dak.	-	-	1	-	1	-	10	1	-	48
S. Dak.	2	-	8	-	-	-	28	2	-	619
Nebr.	10	-	46	-	5	-	22	3	-	2
Kans.	73	2	187	1	5	-	84	7	-	3,197
S. Atlantic	5,362	322	18,942	7	12	1	4,448	2	56	8,653
Del.	27	5	138	-	-	-	57	-	-	2
Md.	325	9	1,538	1	-	-	363	1	14	NN
D.C.	170	28	967	-	-	-	121	-	1	16
Va.	796	18	1,919	2	1	-	372	-	9	2,844
W. Va.	8	2	179	-	-	-	80	-	-	5,656
N.C.	1,672	44	4,023	1	1	-	566	-	1	NN
S.C.	799	100	1,945	1	-	-	387	NN	-	135
Ga.	820	42	3,185	-	1	1	740	1	2	NN
Fla.	745	74	5,048	2	9	-	1,762	-	28	NN
E.S. Central	3,997	144	9,992	-	6	-	1,578	3	6	4,701
Ky.	208	13	534	-	2	-	347	2	1	984
Tenn.	1,044	57	2,978	-	2	-	520	-	3	3,717
Ala.	661	18	1,933	-	1	-	433	-	2	NN
Miss.	2,084	56	4,547	-	1	-	278	1	-	NN
W.S. Central	4,124	355	16,275	15	2	-	3,500	27	17	16,159
Ark.	446	29	1,328	-	-	-	264	23	-	NN
La.	1,608	87	5,422	2	-	-	433	1	4	NN
Okla.	157	15	497	1	2	-	261	3	3	NN
Tex.	1,913	224	9,028	12	-	-	2,542	-	10	16,159
Mountain	242	23	1,137	2	12	4	654	9	13	7,286
Mont.	3	-	9	-	-	-	24	3	-	53
Idaho	2	-	10	-	3	-	13	-	-	NN
Wyo.	-	-	3	-	-	2	12	-	-	NN
Colo.	126	4	296	1	6	1	94	1	3	NN
N. Mex.	18	-	178	-	-	-	81	1	1	NN
Ariz.	50	16	419	-	1	-	249	-	4	6,783
Utah	12	-	51	1	2	-	55	2	2	450
Nev.	31	3	171	-	-	1	126	2	3	NN
Pacific	872	197	7,765	6	54	16	5,628	8	133	1,224
Wash.	36	3	281	1	7	-	264	1	12	NN
Oreg.	22	-	100	-	-	-	165	4	5	NN
Calif.	807	194	7,321	5	43	12	4,859	2	111	NN
Alaska	3	-	22	-	-	4	93	1	-	NN
Hawaii	4	-	41	-	4	-	247	-	5	1,224
Guam	2	-	7	-	-	-	21	-	1	952
P.R.	311	20	2,018	2	-	-	274	-	-	9,193
V.I.	7	-	30	-	-	-	10	-	-	551
C.N.M.I.	-	-	-	-	-	-	14	-	1	121
American Samoa	-	-	-	-	-	-	5	-	1	74

\*Cases updated through Division of Sexually Transmitted Diseases and HIV Prevention, National Center for Prevention Services, as of February 28, 1995.

## SUMMARY TABLES — 1994

### NOTIFIABLE DISEASES — Summary of reported cases, by age group, United States, 1994

NAME	Total	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-39	40-49	50-59	60+	Age not stated
AIDS*	78,279	318	418	148	146	324	2,604	10,162	35,466	20,676	5,940	2,077	-
Amebiasis	2,983	19	116	178	144	102	142	208	474	243	123	122	1,112
Anthrax	-	-	-	-	-	-	-	-	-	-	-	-	-
Aseptic meningitis	8,932	2,148	734	762	550	493	686	833	1,367	561	242	322	234
Botulism, total	143	81	2	5	4	3	7	11	6	8	14	10	5
Brucellosis	119	2	-	-	-	-	-	10	15	25	19	7	12
Cholera	39	-	1	-	-	-	-	1	2	4	7	6	11
Diphtheria	2	-	1	-	-	-	-	-	-	-	-	-	-
Encephalitis, primary infections	717	54	71	86	72	42	40	52	76	50	56	114	4
Post-infectious	143	7	17	31	11	8	5	5	13	11	6	25	1
<i>Escherichia coli</i> O157:H7	1,420	28	304	176	132	111	64	60	109	101	103	167	65
Gonorrhea <sup>§</sup>	1,174	-	-	-	8,508	123,079	121,084	60,204	63,985	.....	25,068	.....	9,670
<i>Haemophilus influenzae</i> , invasive	136	210	119	33	18	26	18	23	24	18	18	18	23
Hansen disease (leprosy)	26,796	130	1,911	4,076	2,492	2,036	2,770	2,981	4,904	2,390	1,049	1,279	778
Hepatitis A	12,517	55	73	82	165	807	1,618	1,940	3,615	2,027	840	845	450
Hepatitis B	4,470	19	15	14	21	21	70	172	420	1,668	1,259	331	401
Hepatitis, C (non-A non-B)	4,444	6	23	47	26	30	38	50	109	48	18	41	8
Hepatitis, unspecified	1,615	6	5	4	3	19	28	39	170	259	276	768	38
Legionellosis	38	-	-	1	2	3	2	7	9	8	1	5	-
Leptospirosis	13,043	27	650	1,142	831	632	473	640	2,062	2,207	1,612	2,540	227
Lyme disease	1,229	11	52	65	61	79	149	120	297	196	86	75	38
Malaria	963	72	174	95	128	255	112	43	60	15	2	1	6
Measles (rubella)	2,886	479	537	248	183	332	180	87	157	162	122	333	66
Meningococcal disease	1,537	13	237	473	271	128	59	61	108	52	20	12	103
Numps	4,617	1,640	837	496	548	301	66	76	187	129	46	34	257
Pertussis (whooping cough)	17	-	2	2	1	-	1	1	2	3	3	-	1
Plague	-	-	-	-	-	-	-	-	-	-	-	-	-
Poliomyelitis, paralytic <sup>§</sup>	38	-	1	-	2	-	1	3	10	15	3	3	-
Psittacosis	6	-	-	-	1	-	1	-	-	4	-	-	-
Rabies, human	112	1	4	28	26	10	3	4	8	3	-	4	21
Rheumatic fever, acute	465	3	29	51	24	34	23	25	85	58	61	65	7
Rocky Mountain spotted fever	227	11	17	13	4	11	32	33	54	38	10	4	-
Rubella (German measles)	5,323	5,265	6,425	2,804	1,600	1,541	2,011	2,261	4,406	2,879	1,975	4,511	7,645
Salmonellosis	29,769	645	8,094	5,856	1,585	873	1,472	1,702	2,476	1,059	542	668	4,497
Shigellosis	-	-	-	-	118	2,234	4,067	3,814	6,638	.....	3,692	.....	41
Tetanus	51	-	-	-	1	1	3	3	9	3	5	24	2
Toxic-shock syndrome	192	2	6	7	20	25	19	22	48	25	8	10	-
Trichinosis	32	-	1	2	3	6	6	1	8	2	4	4	-
Tuberculosis	24,361	164	860	399	272	544	1,281	1,808	4,859	4,375	2,920	6,831	48
Tularemia	96	1	7	11	8	4	5	3	3	12	16	22	4
Typhoid fever	441	10	43	51	32	38	59	49	90	28	19	16	6
Varicella (chickenpox)	151,219	227	2,772	4,119	771	256	207	218	219	78	33	37	142,282

\* AIDS total updated through December 31, 1994.

<sup>†</sup> Age data are collected on aggregate forms different from those used for numbers of reported cases. Therefore, total cases reported on this table may differ slightly from other tables. Cases among persons ages <9 years are not shown because some of these may not be due to sexual transmission; however, they are included in the totals.

<sup>§</sup> Two suspected cases of paralytic poliomyelitis were reported in 1994. Confirmation of these cases is pending review by external panel.

## SUMMARY TABLES — 1994

### NOTIFIABLE DISEASES — Summary of reported cases, by race,\* United States, 1994

Name	Total	American Indian or Alaskan Native	Asian or Pacific Islander	Black	White	Other	Race not stated <sup>t</sup>
AIDS <sup>s</sup>	78,279	227	576	31,457	33,180	-	12,839
Amebiasis	2,983	10	144	123	628	14	2,064
Anthrax	-	-	-	-	-	-	-
Aseptic meningitis	8,932	19	155	1,338	5,229	23	2,168
Botulism, total	143	11	4	1	97	1	29
Brucellosis	119	-	-	5	64	3	47
Cholera	39	-	-	2	17	1	19
Diphtheria	2	-	-	-	2	-	-
Encephalitis, primary infections	717	1	11	80	484	3	138
Hepatitis C, primary infections	143	-	2	20	96	-	25
<i>Escherichia coli</i> O157:H7	1,420	3	9	31	856	6	515
Gonorrhea <sup>1</sup>	413,647	1,829	1,416	271,195	43,618	-	95,589
<i>Haemophilus influenzae</i> , invasive	1,174	16	13	174	701	1	269
Hansen disease (leprosy)	136	-	57	3	29	-	47
Hepatitis A	26,796	2,330	386	1,737	15,215	98	6,970
Hepatitis B	12,517	101	837	2,745	5,158	50	3,626
Hepatitis, C (non-A non-B)	4,470	53	35	723	2,400	9	1,250
Hepatitis, unspecified	444	22	14	33	2,412	1	132
Legionellosis	1,615	7	14	138	999	1	456
Leptospirosis	38	-	8	-	23	-	7
Lyme disease	13,043	41	82	205	9,589	7	3,119
Malaria	1,229	3	167	407	327	22	303
Measles (rubella)	963	1	37	80	641	8	204
Meningococcal disease	2,886	34	32	352	1,889	8	591
Nombrilosis	1,337	8	55	115	887	1	471
Pertussis (whooping cough)	4,617	20	57	281	2,454	9	1,796
Plague	17	8	-	-	7	-	2
Poliomyelitis, paralytic*	-	-	-	-	-	-	-
Pituitary	38	1	-	-	19	-	18
Rabies, human	6	-	-	1	4	-	1
Rheumatic fever, acute	112	1	2	1	58	-	50
Rocky Mountain spotted fever	465	14	2	19	324	1	105
Rubella (German measles)	227	-	12	9	131	48	27
Rubella, congenital syndrome	7	-	1	1	5	-	-
Salmonellosis	43,323	264	512	3,713	19,702	49	19,083
Shigellosis	29,769	731	161	5,169	12,159	34	11,515
Syphilis, primary and secondary <sup>1</sup>	20,638	43	73	17,619	1,830	-	1,073
Tetanus	51	-	1	9	32	-	9
Toxic-shock syndrome	192	-	5	4	145	-	38
Trichinosis	32	-	1	1	12	-	17
Tuberculosis	24,361	341	3,868	8,622	11,309	-	221
Tularemia	96	2	1	2	69	-	22
Typhoid fever	441	5	126	44	73	12	181
Variola (chickenpox)	151,219	6	232	1,997	853	-	148,131

\*Some information on race/ethnicity was reported using one variable with the following categories: American Indian or Alaska Native, Asian or Pacific Islander, Black non-Hispanic, White non-Hispanic, and Hispanic. Race not stated includes cases originally reported as Hispanic.

<sup>1</sup>Includes cases originally reported as Hispanic: 12,699 for AIDS; 14,907 for Gonorrhea; and 794 for Syphilis, primary and secondary.

<sup>s</sup>AIDS total reported through December 31, 1994.  
¶Race data are collected on aggregate forms different from those used for numbers of reported cases. Therefore, total cases reported on this table may differ slightly from other tables.  
\*\*Two suspected cases of paralytic poliomyelitis were reported in 1994. Confirmation of these cases is pending review by external panel.

## SUMMARY TABLES — 1994

### NOTIFIABLE DISEASES — Summary of reported cases, by ethnicity,\* United States, 1994

NAME	Total	Hispanic	Non-Hispanic	Ethnicity not stated
AIDS <sup>†</sup>	78,279	12,699	64,637	943
Amebiasis	2,983	386	541	2,056
Anthrax	-	-	-	-
Aseptic meningitis	8,932	1,030	4,369	3,533
Botulism, total	135	26	73	36
Brucellosis	119	79	15	25
Cholera	39	17	10	12
Diphtheria	2	-	2	-
Encephalitis, primary infections	717	42	343	332
Post-infectious	143	12	75	56
<i>Escherichia coli</i> O157:H7	1,420	34	708	678
Gonorrhea <sup>s</sup>	413,647	14,907	314,813	83,927
<i>Haemophilus influenzae</i> , invasive	1,174	66	712	396
Hansen disease (leprosy)	136	30	70	36
Hepatitis A	26,796	5,386	13,940	7,470
Hepatitis B	12,517	1,031	6,580	4,906
Hepatitis, C/non-A non-B	4,470	387	2,259	1,824
Hepatitis, unspecified	444	123	214	107
Legionellosis	1,615	26	823	766
Leptospirosis	38	-	20	18
Lyme disease	13,043	165	6,118	6,760
Malaria	1,229	91	716	422
Measles (rubeola)	963	55	651	257
Meningococcal disease	2,886	298	1,684	904
Mumps	1,537	188	684	665
Pertussis (whooping cough)	4,617	403	2,050	2,164
Plague	17	-	16	1
Poliomyelitis, paralytic <sup>¶</sup>	-	-	-	-
Psittacosis	38	2	12	24
Rabies, human	6	1	2	3
Rheumatic fever, acute	112	5	28	79
Rocky Mountain spotted fever	465	6	244	215
Rubella (German measles)	227	78	118	31
Salmonellosis	43,323	2,564	16,052	24,707
Shigellosis	29,769	2,909	12,149	14,711
Syphilis, primary and secondary <sup>§</sup>	20,638	794	19,449	395
Tetanus	51	1	28	22
Toxic-shock syndrome	192	5	127	60
Trichinosis	32	1	7	24
Tuberculosis	24,361	5,074	19,084	203
Tularemia	96	2	22	72
Typhoid fever	441	114	204	123
Varicella (chickenpox)	151,219	1,697	1,393	148,129

\*Some information on race/ethnicity was reported using one variable with the following categories: American Indian or Alaska Native, Asian or Pacific Islander, Black non-Hispanic, White non-Hispanic, and Hispanic. Ethnicity not stated includes cases originally reported as American Indian or Alaska Native and Asian or Pacific Islander.

† AIDS total reported through December 31, 1994.


§Ethnicity data are collected on aggregate forms different from those used for numbers of reported cases. Therefore, total cases reported on this table may differ slightly from other tables.

¶Two suspected cases of paralytic poliomyelitis were reported in 1994. Confirmation of these cases is pending review by external panel.

# **PART 2:**


**Graphs and Maps  
for Selected  
Notifiable Diseases  
in the  
United States**

## GRAPHS AND MAPS



## GRAPHS AND MAPS

**ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported cases, per 100,000 population, United States and Puerto Rico, 1994\***




\*The denominator for Puerto Rico is based on extrapolations from U.S. Bureau of Census population data from 1990 and 1992 post-censal estimates.

- In 1994, 84% of reported cases of AIDS were among residents of large metropolitan areas (500,000 or more population).

## GRAPHS AND MAPS

ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported pediatric cases,\* United States and Puerto Rico, 1994




\* Children <13 years of age.

- AIDS cases reported among children ages <13 years during 1994 (1,017) were 8% greater than that reported during 1993 (942). Of the 1,017 children affected, 62% were black, 23% were Hispanic, and 50% were female.

## GRAPHS AND MAPS


18 ARBOVIRAL INFECTIONS (of the central nervous system) — reported laboratory-confirmed cases caused by California serogroup viruses, by month of onset, United States, 1985–1994



• California serogroup viruses consistently produce primarily pediatric clinical encephalitis cases in various areas of the eastern United States.

## GRAPHS AND MAPS


**ARBOVIRAL INFECTIONS (of the central nervous system) — reported laboratory-confirmed cases caused by eastern equine encephalitis virus, by month of onset, United States, 1985–1994**



• Human eastern equine encephalitis cases, often associated with high mortality rates (20%) and severe neurologic sequelae, occur in low frequency in states along the Atlantic coast.

**GRAPHS AND MAPS**


**ARBOVIRAL INFECTIONS (of the central nervous system) — reported laboratory-confirmed cases caused by St. Louis encephalitis virus, by month of onset, United States, 1985-1994**



• St. Louis encephalitis, which has historically produced large epidemics, frequently causes intense local outbreaks (e.g., New Orleans 1994).

## GRAPHS AND MAPS


ARBOVIRAL INFECTIONS (of the central nervous system) — reported laboratory-confirmed cases caused by western equine encephalitis virus, by month of onset, United States, 1985–1994



• Human western equine encephalitis cases, for unclear reasons, have occurred only sporadically since the outbreaks in 1987.


## GRAPHS AND MAPS


**BOTULISM (foodborne) — by year, United States, 1975-1994**



\* Data not yet available for 1994.

- Although cases of foodborne botulism remain uncommon, outbreaks (including restaurant-associated ones) continue to occur and, because of the severity of the illness, constitute public health emergencies.




**BRUCELLOSIS — by year, United States, 1965–1994**

Recent data suggest that reported cases of foodborne brucellosis, usually caused by *Brucella melitensis*, are increasing in the United States.

## GRAPHS AND MAPS


**CHOLERA — reported cases, United States and territories, 1994**



• In recent years, most U.S. cholera cases have been associated with foreign travel or food brought into the country from cholera-affected areas. Previously, cholera was more often acquired domestically from shellfish harvested in the Gulf of Mexico.

## GRAPHS AND MAPS

DIPHTHERIA — by year, United States, 1965–1994




NOTE: DTP vaccine licensed 1949.

•An ongoing epidemic of diphtheria is occurring in 14 of the 15 countries of the former Soviet Union. Two cases were reported among U.S. citizens residing in the former Soviet Union. No importation of diphtheria into the United States related to these outbreaks was reported in 1994.

## GRAPHS AND MAPS


### *ESCHERICHIA COLI O157:H7 — reported cases, United States and territories, 1994*



• States in which *E. coli* O157:H7 infection is a notifiable disease increased from nine in 1992 to 33 in 1994. However, because fewer than 60% of clinical laboratories routinely test all stools or albloody stools for *E. coli* O157:H7, many infections are not recognized or reported.

## GRAPHS AND MAPS

### *ESCHERICHIA COLI* O157:H7 — reported isolates,\* United States, 1994



\*Data from the Public Health Laboratory Information System (PHLIS).

•From 1993 to 1994, the number of states reporting *E. coli* O157:H7 isolates to the PHLIS increased more than threefold. Identification of *E. coli* O157:H7 must be confirmed by the state public health laboratory to be reported to PHLIS.

## GRAPHS AND MAPS


GONORRHEA — by sex, United States, 1981-1994



• In 1994, the rate of gonorrhea among men continued to decline; among women it increased from 147.1 per 100,000 in 1993 to 153.7.

## GRAPHS AND MAPS


30 GONORRHEA — by race, United States, 1981-1994



•In 1994, gonorrhea rates increased slightly among all racial and ethnic groups except Hispanics.

## GRAPHS AND MAPS

GONORRHEA — reported cases, per 100,000 population, United States, 1994




NOTE: The Year 2000 Objective is ≤225 per 100,000 population.

- The overall U.S. gonorrhea rate in 1994 was 168.4 per 100,000 and 37 states reported rates below the Healthy People 2000 national objective.

## GRAPHS AND MAPS


### *HAEMOPHILUS INFLUENZAE*, INVASIVE — by age group, United States, 1994



• Invasive *H. influenzae* disease was reported in 329 children ages <5 years. Of 52 cases with known serotype, 31 were reported as serotype B. Lack of information on serotype prevented accurately determining whether most of the cases were vaccine-preventable or represented vaccine failures.

## GRAPHS AND MAPS


HANSEN DISEASE (leprosy) — by year, United States, 1965-1994



• During the past 30 years in the United States, new Hansen disease cases have continued to occur at a fairly constant rate, except for the increase related to foreign immigration between 1978-1988.

## GRAPHS AND MAPS

### HEPATITIS — by year, United States, 1965–1994




\*The first hepatitis B vaccine was licensed June 1982.

†Anti-HCV antibody test available May 1990.

- Non-A, non-B hepatitis reporting was artificially increased after introduction of the anti-hepatitis C virus antibody test in 1990, and has since declined.

## GRAPHS AND MAPS


**HEPATITIS A — reported cases, per 100,000 population, United States and territories, 1994**



• Although the western states have overall higher incidence rates of hepatitis A than the remaining states, they also account for substantial reports. Such cases are often reported in connection with communitywide outbreaks.


## GRAPHS AND MAPS

**HEPATITIS B — reported cases, per 100,000 population, United States and territories, 1994**




• Hepatitis B continues to decline in most states, primarily because of changes in high-risk behaviors among injecting-drug users. High incidence rates are often associated with cities and counties with large urban populations.

## GRAPHS AND MAPS



## GRAPHS AND MAPS


**LYME DISEASE — reported cases, per 100,000 population, United States and territories, 1994**



• The number of reported Lyme disease cases increased 58% from 1993 to 1994, coincident with increases in the density of vector tick populations in several northeastern states where Lyme disease is endemic.


## GRAPHS AND MAPS

MALARIA — by year, United States, 1965–1994



•In 1994, *Plasmodium vivax* infection was diagnosed in three persons in Houston, Texas. These persons did not have recent bloodborne exposure and had not recently travelled to areas where malaria is endemic, strongly suggesting mosquito-borne transmission of malaria in the United States.


**MEASLES (rubeola) — by year, United States, 1960–1994**



• In 1994, 963 cases of measles were reported—the second lowest annual total ever. The largest outbreaks were reported among communities that do not accept vaccination.

## GRAPHS AND MAPS


### MENINGOCOCCAL DISEASE — by year, United States, 1965–1994



- Although overall rates of meningococcal disease have remained relatively constant, serogroup C outbreaks are occurring more frequently in the United States, and vaccine use for outbreak control has increased. In addition, Oregon and part of Washington state reported a substantial increase of serogroup B disease in 1994. The meningococcal vaccine licensed in the United States provides protection against serogroups A, C, Y, and W-135, but not against serogroup B.

## GRAPHS AND MAPS

MUMPS — by year, United States, 1968–1994




NOTE: Mumps vaccine licensed December 1967.

• In 1994, 1,537 mumps cases were reported—the lowest number ever reported in the United States.


## GRAPHS AND MAPS

PERTUSSIS (whooping cough) — by year, United States, 1965–1994



NOTE: DTP vaccine licensed 1949.


- The number of reported pertussis cases declined 30% from 1993–1994—a pattern consistent with the previously observed 3–4 year periodicity in pertussis incidence.

**PERTUSSIS (whooping cough) — by age group, United States, 1994**

• Of 734 pertussis cases among children 7 months to 4 years of age with known vaccination status, 320 (43.6%) had received fewer than three doses of diphtheria-tetanus-pertussis vaccine—the minimum number of doses necessary for clinical protection.

## GRAPHS AND MAPS


PLAUE — among humans, by year, United States, 1965–1994



•Plague continued to be enzootic throughout the 14 contiguous western states, resulting in sporadic human cases caused primarily by the bites of infected rodent fleas, but also by direct contact with infected animals.

## GRAPHS AND MAPS

POLIOMYELITIS (paralytic) — by year, United States, 1965–1994




NOTE: Inactivated vaccine licensed 1955. Oral vaccine licensed 1961.

• Since 1980, all confirmed cases of indigenously acquired paralytic poliomyelitis in the United States have been vaccine-associated.


## GRAPHS AND MAPS

PSITTACOSIS — by year, United States, 1965–1994



• The decline of psittacosis cases reported during recent years may reflect a return to baseline incidence, as cases of *Chlamydia pneumoniae* may have been attributed to *Chlamydia psittaci* infection in the mid-1980s.


**RABIES — wild and domestic animals, by year, United States and Puerto Rico, 1955–1994**




- The number of humans administered rabies post-exposure treatment continues to increase as animal rabies cases increase, especially in the northeastern United States where raccoon rabies continues to spread.

## GRAPHS AND MAPS


ROCKY MOUNTAIN SPOTTED FEVER (RMSF) — by year, United States, 1965–1994



•RMSF is the most common fatal tick-borne disease in the United States, with a case fatality rate of 2%.


**RUBELLA (German measles) — by year, United States, 1966–1994**

• In 1994, 227 cases of rubella were reported in the United States. Of these cases 171 (75%) occurred among persons ages  $\geq 20$  years, and 126 (56%) were reported from one state. Of seven cases of congenital rubella syndrome reported, four were caused by international importation and three were acquired in the United States.



## GRAPHS AND MAPS

SALMONELLA — serotype of isolate by year,\* United States, 1968–1994




\*Data from Public Health Laboratory Information System.

•In 1994, an outbreak of *Salmonella* serotype Enteritidis infection was linked to a nationally distributed ice cream brand. Illnesses were documented in 41 states, and more than 200,000 persons were estimated to have been ill.


## GRAPHS AND MAPS

SHIGELLOSIS — by year, United States, 1965-1994



## GRAPHS AND MAPS

*SHIGELLA* — species of isolate by year,\* United States, 1968–1994




\*Data from Public Health Laboratory Information System.

•Community outbreaks of *Shigella sonnei*, often involving multiple child care centers, continue to be a substantial public health problem.

## GRAPHS AND MAPS


**SYPHILIS (primary and secondary) — reported cases, per 100,000 population, United States, 1994**



- In 1994, the rate of primary and secondary syphilis was 8.1 per 100,000. Rates were below the Healthy People 2000 national objective in 38 states, with 10 states reporting fewer than five cases.

## GRAPHS AND MAPS


### SYPHILIS (primary and secondary) — by sex, United States, 1981–1994



•In 1994, the rate of primary and secondary syphilis among men decreased from 11.3 per 100,000 in 1993 to 8.6, and from 9.5 to 7.6 among women.

## GRAPHS AND MAPS


SYPHILIS (primary and secondary) — by race, United States, 1981-1994



• Since 1990, the rates of primary and secondary syphilis have declined among all racial and ethnic groups. However, the 1994 rate of 59.5 cases per 100,000 for non-Hispanic blacks was 60 times greater than for non-Hispanic whites.

## GRAPHS AND MAPS

CONGENITAL SYPHILIS — in infants <1 year of age, United States, 1963–1994\*




\* The surveillance case definition for congenital syphilis changed in 1989.

• Between 1993 and 1994, the rate of congenital syphilis decreased from 80.7 to 55.6 cases per 100,000 live births.

## GRAPHS AND MAPS


TETANUS — by year, United States, 1965–1994



NOTE: Tetanus toxoid was first available in 1933.

• The 1996 goal for tetanus disease reduction in the United States is zero cases among children ages <15 years; in 1994, one tetanus case was reported in a child 12 years of age.

**TOXIC SHOCK SYNDROME (TSS) — by quarter, United States, 1979–1994**




\* Includes only cases that meet the CDC case definition for *staphylococcal* TSS.

† First available through NETSS beginning 1983.

- The number of reported cases of TSS continued adownward trend. In 1994, 42% (10) of confirmed TSS cases were nonmenstrual.

## GRAPHS AND MAPS


### TRICHINOSIS — by year, United States, 1965–1994



• In 1994, 32 cases were reported, below the mean number reported for the years 1990–1994.

## GRAPHS AND MAPS


TUBERCULOSIS — reported cases, per 100,000 population, United States and territories, 1994




•Fifteen states have tuberculosis rates ≤3.5 cases per 100,000, the interim tuberculosis elimination goal for the year 2000.

## GRAPHS AND MAPS

### TUBERCULOSIS — by year, United States, 1975-1994




•In 1994, 24,361 cases of tuberculosis (9.4 per 100,000) were reported to CDC from the United States, representing a 3.7% decrease from 1993.

**TUBERCULOSIS — by year, among foreign-born persons and persons born in the United States, 1986–1994**

• The number and percent of tuberculosis cases among the foreign-born has increased from 4,925 (21.6%) in 1986 to 7,627 (31.3%) in 1994.

## GRAPHS AND MAPS


TYPHOID FEVER — by year, United States, 1965–1994



•A new single dose parenteral typhoid vaccine was licensed for use in the United States in 1994. The new vaccine is an alternative to the four-dose oral vaccine available.

## GRAPHS AND MAPS

66 VARICELLA (chickenpox) — by month, United States\*, 1986-1994



\*Varicella is reportable in 25 states.

•Approximately 3.7 million cases of varicella occur annually in the United States; of these, an estimated 4%-5% are reported.

# **PART 3:**

## **Historical Summary Tables**

BLANK PAGE

## HISTORICAL TABLES — 1985–1994

**TABLE 1. NOTIFIABLE DISEASES — summary of reported cases, per 100,000 population, United States, 1985–1994**

Disease	1994	1993	1992	1991	1990	1989	1988	1987	1986	1985
AIDS	30.07	40.20	17.83	17.32	16.72	13.58	12.61	8.66	5.36	3.46
Amebiasis	1.20	1.21	1.21	1.23	1.38	1.34	1.20	1.33	1.47	1.92
Anthrax	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Aseptic meningitis	3.71	5.39	5.18	6.26	4.77	4.14	2.94	4.72	4.72	4.50
Botulism, total (including wound and unsp.)	0.06	0.04	0.04	0.05	0.04	0.04	0.03	0.03	0.05	0.05
Foodborne	0.02	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.02
Brucellosis	0.05	0.05	0.04	0.04	0.03	0.04	0.04	0.05	0.04	0.06
Chancroid*	0.30	0.54	0.80	1.40	1.70	1.90	2.04	2.07	1.57	0.87
Cholera	0.02	0.00	0.04	0.01	0.00	0.00	0.00	0.00	0.01	0.00
Diphtheria	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Encephalitis, primary	0.28	0.36	0.30	0.40	0.54	0.40	0.36	0.58	0.54	0.58
Post-infectious	0.06	0.07	0.05	0.03	0.04	0.04	0.05	0.05	0.05	0.07
<i>Escherichia coli</i> O157:H7	0.82	172.40	201.60	249.48	276.60	297.36	287.74	323.14	376.37	384.51
Gonorrhea*	168.40	172.40	201.60	249.48	276.60	297.36	287.74	323.14	376.37	384.51
Granuloma inguinale*	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.01	0.03	0.02
<i>Hemophilus influenzae</i> , invasive	0.45	0.55	0.55	1.10	0.08	0.07	0.10	0.11	0.11	0.15
Hansen disease (leprosy)	10.29	9.40	9.06	9.67	12.64	14.43	11.60	10.39	10.02	10.03
Hepatitis A	4.81	5.18	6.32	7.14	8.48	9.43	10.65	11.17	11.17	11.50
Hepatitis B	1.78	1.86	2.36	1.42	1.03	1.02	1.07	1.23	1.55	1.81
Hepatitis, C/non-A, non-B <sup>§</sup>	0.17	0.24	0.35	0.50	0.67	0.93	1.00	1.27	1.69	2.38
Hepatitis, unspecified	0.63	0.50	0.53	0.53	0.55	0.48	0.44	0.43	0.43	0.37
Legionellosis	0.02	0.02	0.02	0.02	0.03	0.04	0.02	0.02	0.02	0.02
Leprosy	5.01	3.20	0.12	3.80	3.80	3.80	3.80	3.80	3.80	3.80
Lyme disease	0.10	0.10	0.10	0.19	0.10	0.08	0.07	0.13	0.16	0.10
Lymphogranuloma venereum*	0.47	0.55	0.43	0.51	0.52	0.51	0.45	0.39	0.47	0.44
Malaria	0.37	0.12	0.88	3.82	11.17	7.33	1.38	1.50	2.61	1.18
Measles (rubella)	1.11	1.02	0.84	0.84	0.99	1.10	1.21	1.20	1.08	1.04
Meningococcal disease	0.60	0.66	1.03	1.72	2.17	2.34	2.05	5.43	3.37	1.30
Mumps	1	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.03	0.02
Murine typhus fever	1.77	2.55	1.60	1.08	1.84	1.67	1.40	1.16	1.74	1.50
Pertussis (whooping cough)	0.01	0.00	0.01	0.01	0.00	0.00	0.00	0.01	0.00	0.01
Plague	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Poliomyelitis, paralytic	0.02	0.02	0.04	0.04	0.04	0.05	0.05	0.04	0.09	0.05
Psittacosis	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Rabies, human	0.09	0.08	0.06	0.12	0.09	0.13	0.14	0.13	0.12	0.07
Rheumatic fever, acute	0.18	0.18	0.20	0.25	0.26	0.25	0.25	0.25	0.25	0.30
Rocky Mountain spotted fever	0.69	0.67	0.66	0.56	0.45	0.46	0.45	0.45	0.45	0.26
Rubella (German measles)	16.64	16.15	16.04	19.10	19.54	19.26	19.91	20.92	20.73	21.37
Salmonellosis, excluding typhoid fever	11.44	12.48	9.38	9.34	10.89	10.07	12.46	9.80	7.11	7.14
Shigellosis	..	..	..	..	..	..	..	..	..	..
Smallpox	8.10	10.40	13.70	17.26	20.10	18.07	16.43	14.54	11.65	11.45
Syphilis, primary and secondary*	32.00	39.70	45.30	51.69	53.80	44.94	42.37	35.81	28.50	28.50
Total, all stages*	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.03	0.03
Tetanus	0.10	0.08	0.10	0.11	0.13	0.16	0.16	0.15	0.19	0.18
Toxic-shock syndrome	0.01	0.01	0.02	0.02	0.05	0.01	0.02	0.02	0.02	0.03
Trichinosis	9.36	9.82	10.46	10.33	9.46	9.13	9.25	9.44	9.30	9.30
Tuberculosis	0.04	0.05	0.06	0.08	0.06	0.06	0.08	0.09	0.07	0.07
Tularemia	0.17	0.17	0.16	0.20	0.22	0.19	0.18	0.16	0.15	0.17
Typhoid fever	135.76	118.54	176.54	135.82	120.06	121.77	122.43	126.68	122.42	123.23
Vaccinia (chickenpox)	..	..	..	..	..	..	..	..	..	..
Yellow fever	..	..	..	..	..	..	..	..	..	..

NOTE: Rates <0.01 after rounding are shown as 0.00.

\*Post-censal population data from 1993 were used to calculate 1994 rates.

<sup>t</sup> Not previously nationally notifiable.

<sup>§</sup> Anti-HCV antibody test available May 1990.

<sup>\*</sup>No longer nationally notifiable.

† Last indigenous case reported 1911; last imported, 1924.

HISTORICAL TABLES — 1985–1994

TABLE 2. NOTIFIABLE DISEASES — Summary of reported cases, United States, 1985–1994

Disease	1994	1993	1992	1991	1990	1989	1988	1987	1986	1985
AIDS	78,279	103,533	45,472	43,672	41,595	33,722	31,001	21,070	12,932	8,249
Anemias	2,983	2,970	2,942	2,989	3,328	3,217	2,860	3,123	3,532	4,433
Anthrax	—	—	1	—	—	—	—	2	1	—
Aseptic meningitis	8,932	12,848	12,223	14,526	11,852	10,274	7,234	11,487	11,374	10,619
Botulism, total (including wound and unsp.)	143	97	91	114	92	89	84	82	109	122
Foodborne	50	27	21	27	23	23	28	17	23	49
Infant	85	65	66	81	65	60	50	59	79	70
Brucellosis	119	120	105	104	85	95	96	129	106	153
Chancroid	773	1,399	1,886	3,476	4,212	4,692	5,001	4,998	3,756	2,067
Cholera	39	18	103	26	6	—	8	6	23	4
Diphtheria	2	—	4	5	4	3	2	3	—	3
Encephalitis, primary	717	919	774	1,021	1,341	981	882	1,418	1,302	1,376
Post-infectious	143	170	129	82	105	88	121	121	124	161
<i>Escherichia coli</i> O157:H7	1,420	.....	.....	.....	.....	.....	.....	.....	.....	.....
Gonorrhea	418,068	439,673	501,409	620,478	690,169	733,151	719,536	780,905	900,868	911,419
Granuloma inguinale	3	19	6	29	97	7	11	22	61	44
<i>Haemophilus influenzae</i> , invasive	1,174	1,419	1,412	2,764	.....	.....	.....	.....	.....	.....
Hansen disease (leprosy)	136	187	172	154	198	163	184	238	270	361
Hepatitis A	29,796	24,238	23,112	24,378	31,441	35,821	28,507	25,280	23,430 <sup>t</sup>	23,210 <sup>t</sup>
Hepatitis B	12,517	13,361	16,126	18,003	21,102	23,419	23,177	25,916	26,107	26,611 <sup>t</sup>
Hepatitis, C/non-A, non-B <sup>§</sup>	4,470	4,786	6,010	3,582	2,553	2,529	2,619	2,999	3,634 <sup>t</sup>	4,184 <sup>t</sup>
Hepatitis, unspecified	444	627	884	1,260	1,671	2,306	2,470	3,102	3,940 <sup>t</sup>	5,517 <sup>t</sup>
Legionellosis	1,615	1,280	1,339	1,317	1,370	1,190	1,085	1,038	948	830
Leptospirosis	38	51	54	58	77	93	54	43	41	57
Lyme disease	13,043	8,257	9,895	9,465	.....	.....	.....	.....	.....	.....
Lymphogranuloma venereum	235	285	302	471	277	189	185	303	396	226
Malaria	1,229	1,411	1,087	1,278	1,292	1,277	1,099	944	1,123	1,049
Measles (rubeola)	963	312	2,237	9,643	27,786	18,193	3,396	3,655	6,282	2,822
Meningococcal disease	2,886	2,637	2,134	2,130	2,451	2,727	2,964	2,594	2,479	2,982
Mumps	1,537	1,692	2,572	4,264	5,292	5,712	4,866	12,848	7,790	2,982
Murine typhus fever	¶	25	43	50	41	54	49	67	67	37

## HISTORICAL TABLES — 1985–1994

Pertussis (whooping cough)	4,617	6,586	4,083	2,719	4,570	4,157	3,450	2,823	4,195	3,589
Plague	17	10	13	11	2	4	15	12	10	17
Poliomyelitis, paralytic**	-	3	6	9	6	9	9	9	9	7
Pittagosis	38	60	92	94	113	114	98	224	224	119
Rabies, animal	8,147	9,377	8,589	6,910	4,826	4,724	4,651	4,658	5,504	5,565
Rabies, human	6	3	1	3	1	1	1	1	—	1
Rheumatic fever, acute	112	112	75	127	108	144	158	141	147	90
Rocky Mountain spotted fever	465	456	502	628	651	623	609	604	760	714
Rubella (German measles)	227	192	160	1,401	1,125	396	225	306	551	630
Rubella, congenital syndrome	7	5	11	47	11	3	6	5	14	—
Salmonellosis, excluding typhoid fever	43,323	41,641	40,912	48,154	48,603	47,812	48,948	50,916	49,984	65,347
Shigellosis	29,769	32,198	23,931	23,548	27,077	25,010	30,617	23,860	17,138	17,057
Smallpox	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
Syphilis, primary and secondary	20,627	26,498	33,973	42,935	50,223	44,540	40,117	35,147	27,883	27,131
Total, all stages	81,696	101,259	112,581	128,569	134,255	110,797	103,437	86,545	68,215	67,563
Tetanus	51	48	45	57	64	53	53	48	64	83
Toxic-shock syndrome	192	212	244	280	322	400	390	372	412	384
Trichinosis	32	16	41	62	129	30	45	40	39	61
Tuberculosis	24,361	25,313	26,673	26,283	25,701	23,495	22,436	22,517	22,768	22,201
Tularemia	96	132	159	193	152	152	201	214	170	177
Typhoid fever	441	440	414	501	552	460	436	400	362	402
Varicella (chickenpox)	151,219	134,722	158,364	147,076	173,099	185,441	192,857	213,196	183,243	178,162
Yellow fever	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

\* Not previously nationally notifiable.

† Reports from New York City are not available.

§ Anti-HCV antibody test available May 1990.

¶ No longer nationally notifiable.

\*\* Annual case reports from state health departments; numbers may not reflect changes based on retrospective case evaluations or late reports (see MMWR 1986;35:180-2). At time of publication, suspect cases from 1994 had not been confirmed by external review panel.

HISTORICAL TABLES — 1975–1984

TABLE 3. NOTIFIABLE DISEASES — summary of reported cases, United States, 1975–1984

Disease	1984	1983	1982	1981	1980	1979	1978	1977	1976	1975
AIDS	4,445	6,658	7,304	6,632	5,271	4,107	3,937	3,044	2,906	2,775
Amebiasis	5,252	—	—	—	1	—	6	2	—	—
Anthrax	1	—	—	—	—	—	—	—	—	4,475
Aseptic meningitis	8,326	12,696	9,680	9,547	8,028	8,754	6,573	4,789	3,510	20
Botulism, total (including wound and unsp.)	123	133	97	103	89	45	105	129	55	310
Brucellosis	131	200	173	185	183	215	179	232	296	700
Chancroid	665	847	1,392	850	788	840	521	455	628	—
Cholera	1	1	—	19	9	1	12	3	—	—
Diphtheria	1	5	2	5	3	59	76	84	128	307
Encephalitis, primary <sup>†</sup>	1,257	1,761	1,464	1,492	1,362	1,504	1,351	1,414	1,651	4,064
Post-infectious <sup>†</sup>	108	34	36	43	40	84	78	119	175	237
Gonorrhea	878,556	900,435	960,633	990,864	1,004,029	1,004,058	1,013,436	1,002,219	1,001,994	999,937
Granuloma inguinale	30	24	17	66	51	76	72	75	71	60
Hansen disease (leprosy)	290	259	250	256	223	185	168	151	145	162
Hepatitis A	22,040	21,532	23,403	25,802	29,087	30,407	29,500	31,153	33,288	35,855
Hepatitis B	26,115	24,318	22,177	21,152	19,015	15,452	15,016	16,831	14,973	13,121
Hepatitis, non-A, non-B	3,871	3,470	—	—	—	—	—	—	—	—
Hepatitis, unspecified	5,531	7,149	8,564	10,975	11,894	10,534	8,776	8,639	7,488	7,158
Legionellosis <sup>§</sup>	750	852	654	408	475	593	761	359	235	—
Leprosy	40	61	100	82	85	94	110	71	73	93
Lymphogranuloma venereum	170	335	235	263	199	250	284	348	365	363
Malaria	1,007	813	1,056	1,388	2,062	894	731	547	471	373
Measles (rubella)	2,587	1,497	1,714	3,124	13,506	13,597	26,871	57,345	41,126	24,374
Meningococcal disease	2,746	2,736	3,056	3,525	2,840	2,724	2,505	1,828	1,805	1,478
Mumps	3,021	3,355	5,270	4,941	8,576	14,225	16,817	81	69	41
Murine typhus fever	53	62	58	61	81	69	46	21,436	38,492	59,647

## HISTORICAL TABLES — 1975–1984

### Yellow fever

\* Not beginning in 1984 data reflects change in categories for tabulating encephalitis reports that were recorded by date of report to state health departments.

5 Data for 1976-1981 are from surveillance records reported by onset date.  
6 Data for 1977-1981 are from surveillance records reported by onset date.

<sup>9</sup>Beginning in 1982, data recorded by date of report to the State Auditor Department.

HISTORICAL TABLES — 1965-1974

TABLE 4. NOTIFIABLE DISEASES — summary of reported cases, United States, 1965-1974

Disease	1974	1973	1972	1971	1970	1969	1968	1967	1966	1965
Amebiasis	2,743	2,235	2,199	2,752	2,888	2,915	3,005	3,157	2,921	2,768
Anthrax	2	2	2	5	2	4	3	2	5	7
Aseptic meningitis	3,197	4,846	4,634	5,176	6,480	3,672	4,494	3,082	3,058	2,329
Botulism	28	34	22	25	12	16	7	5	9	19
Brucellosis	240	202	196	183	213	235	218	265	262	262
Chancroid	945	1,165	1,414	1,320	1,416	1,104	845	784	838	982
Cholera	—	1	—	1	—	—	—	—	—	2
Diphtheria	272	228	152	215	435	241	260	219	209	164
Encephalitis, primary	1,164	1,613	1,059	1,524	1,580	1,613	1,781	1,478	2,121	1,722
Post-infectious	218	354	243	439	370	304	502	1,060	964	981
Gonorrhea	906,121	842,621	767,215	670,268	600,072	534,872	464,543	404,836	351,738	324,925
Granuloma inguinale	47	62	81	89	124	154	156	154	148	155
Hansen disease (leprosy)	118	146	130	131	129	98	123	81	109	96
Hepatitis A (infectious)	40,358	50,749	54,074	59,606	56,797	48,416	45,893	38,909	32,859	33,856*
Hepatitis B (serum)	10,631	8,451	9,402	9,556	8,310	5,909	4,829	2,458	1,497	—
Hepatitis, unspecified	8,351	—	—	—	—	—	—	—	—	—
Leptospirosis	68	57	41	62	47	89	69	67	72	84
Lymphogranuloma venereum	394	408	756	692	612	520	485	371	308	878
Malaria	293	237	742	2,375	3,051	3,102	2,317	2,022	565	147
Measles (rubella)	22,094	26,690	32,275	75,290	47,351	25,826	22,231	62,705	204,136	261,904
Meningococcal disease	1,346	1,378	1,323	2,262	2,505	2,951	2,623	2,161	3,381	3,040
Mumps	59,128	69,612	74,215	124,939	104,953	90,918	152,209	... <sup>†</sup>	52	28
Murine typhus fever	26	32	18	23	27	36	36	33	—	—

## HISTORICAL TABLES — 1965-1974

Pertussis (whooping cough)	2,402	1,759	3,287	3,036	4,249	3,285	4,810	9,718	7,717	6,799
Plague	8	2	1	2	13	5	3	5	5	8
Poliomyelitis, total	7	8	31	21	33	20	53	41	113	72
Paralytic	7	7	29	17	31	18	53	40	106	61
Pitressin	164	33	52	32	35	57	43	41	50	60
Rabies, animal	3,151	3,640	4,369	4,310	3,224	3,490	3,591	4,481	4,178	4,574
Rabies, human	—	1	2	2	3	1	2	1	1	2
Rheumatic fever, acute	2,431	2,560	2,614	2,793	3,227	3,229	3,470	3,985	4,472	4,998
Rocky Mountain spotted fever	754	668	523	432	380	498	298	305	268	281
Rubella (German measles)	11,917	27,804	25,507	45,086	56,552	57,686	49,371	46,888	46,975	†
Rubella, congenital syndrome	45	35	42	68	77	31	14	10	11	†
Salmonellosis, excluding typhoid fever	21,980	23,818	22,151	21,928	22,096	18,419	16,514	18,120	16,841	17,161
Shigellosis	22,600	22,642	20,207	16,143	13,845	11,946	12,180	13,474	11,888	11,027
Smallpox	.....	NN	NN	NN	NN	NN	NN	NN	NN	NN
Streptococcal sore throat and scarlet fever	.....	NN	NN	NN	NN	NN	NN	NN	NN	NN
Syphilis, primary and secondary	25,385	24,825	24,429	23,783	21,982	19,130	19,019	21,053	21,414	23,388
Total, all stages	83,771	87,469	91,149	95,997	91,382	92,162	96,271	102,581	105,159	112,842
Tetanus	101	101	128	116	148	192	178	263	235	300
Trichinosis	120	102	89	103	109	215	77	66	115	199
Tuberculosis <sup>§</sup>	30,122	30,998	32,882	35,217	37,137	39,120	42,623	45,647	47,767	49,016
Tularemia	144	171	152	187	172	149	186	184	208	264
Typhoid fever	437	680	398	407	346	364	395	396	378	454
Varicella (chickenpox)	141,495	182,927	164,114	.....	.....	.....	1	.....	.....	.....
Yellow fever	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

\*Includes serum hepatitis.

†Not previously notifiable nationally.

<sup>§</sup>Case data subsequent to 1974 are not comparable to prior years because of changes in reporting criteria that became effective in 1975.

Last documented case occurred in 1949.....Last indigenous case reported 1911; last imported, 1924.....

HISTORICAL TABLES — 1955-1964

TABLE 5. NOTIFIABLE DISEASES — summary of reported cases, United States, 1955-1964

Disease	1964	1963	1962	1961	1960	1959	1958	1957	1956	1955
Amoebiasis	3,304	2,886	3,048	2,850	3,424	3,508	4,380	5,031	3,689	3,348
Anthrax	5	3	9	14	23	12	16	26	38	39
Aseptic meningitis	2,177	1,844	2,654	5,162*	1,593	.....	.....	.....	.....	.....
Botulism	23	47	10	14	12	20	6	28	17	16
Brucellosis	411	407	409	636	751	892	924	983	1,300	1,444
Chancroid	1,247	1,220	1,344	1,438	1,680	1,537	1,595	1,637	2,135	2,649
Cholera	—	—	—	—	—	—	—	—	—	—
Dengue	NN									
Diphtheria	293	314	444	617	918	934	918	1,211	1,568	1,984
Encephalitis, acute infectious (primary)	2,002	1,993	2,094	2,248	2,341	2,437	2,587	2,135	2,624	2,166
Post-infectious	1,585	.....	.....	.....	.....	.....	.....	.....	.....	.....
Gonorrhea	300,666	278,289	263,714	264,158	258,933	240,254	232,386	214,496	224,346	236,197
Granuloma inguinale	135	173	207	241	296	265	314	348	357	490
Hansen disease (leprosy)	97	103	80	63	54	44	39	36	52	75
Hepatitis <sup>§</sup>	37,740	42,974	53,016	72,651	41,666	23,574	16,294	14,922	19,234	31,961
Leptospirosis	142	89	79	71	53	83	55	47	44	24
Lymphogranuloma venereum	732	586	590	787	835	604	434	448	500	762
Malaria	93	99	118	73	72	71	85	132	234	522
Measles (rubella)	458,083	385,166	481,530	423,919	441,703	406,162	763,094	486,799	611,936	555,156
Meningococcal disease	2,826	2,470	2,150	2,232	2,259	2,180	2,581	2,691	2,735	3,455
Murine typhus fever	30	35	32	46	68	51	71	113	98	135

## HISTORICAL TABLES — 1955-1964

Pertussis (whooping cough)	13,005	17,135	17,749	11,468	14,809	40,005	32,148	28,255	31,732	62,786
Plague	-	1	-	3	2	4	-	1	1	-
Poliomyelitis, total	122	449	910	1,312	3,190	8,425	5,787	5,485	15,140	28,985
Paralytic	106	396	762	988	2,525	6,289	3,697	2,499	7,911	13,850
Psittacosis	53	76	79	102	113	147	158	278	568	334
Rabies, animal <sup>1</sup>	4,780	3,929	3,732	3,599	3,567	4,177	4,787	4,542	5,681	5,799
Rabies, human <sup>1</sup>	1	1	2	3	2	7	5	5	10	4
Rheumatic fever, acute	7,491	7,561	7,977	10,470	9,022	8,285	6,889	6,427	6,562	6,462
Rocky Mountain spotted fever	277	216	240	219	204	199	243	240	293	295
Salmonellosis, excluding typhoid fever	17,144	15,390	9,680	8,542	6,929	6,606	6,363	6,693	6,704	5,447
Shigellosis	12,984	13,009	12,443	12,571	12,487	12,888	11,861	9,822	10,306	13,912
Smallpox	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
Streptococcal sore throat and scarlet fever	402,334	342,161	315,809	338,410	315,173	334,715	264,097	226,973	176,392	147,502
Syphilis, primary and secondary	22,969	22,251	21,067	19,851	16,145	9,799	7,176	6,576	6,392	6,554
Total, all stages	114,325	124,137	126,245	124,658	122,538	120,824	113,884	123,758	130,201	122,392
Tetanus	289	325	322	379	368	445	445	447	468	462
Trichinosis	198	208	194	306	160	227	176	178	262	264
Tuberculosis**	50,874	54,042	53,315	53,726	55,494	57,535	63,534	67,149	69,895	77,368
Tularemia	342	327	328	365	390	459	587	601	522	584
Typhoid fever	501	566	608	814	816	859	1,043	1,231	1,700	1,704
Yellow fever	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

\* Includes Meningitis, other, for some states.

† Not previously notifiable nationally.

§ Data includes both infectious and serum hepatitis.

1 Registered deaths, 1955-1961.

\*\* Includes new active cases.

HISTORICAL TABLES — 1945-1954

TABLE 6. NOTIFIABLE DISEASES — summary of reported cases, United States, 1945-1954

Disease	1954	1953	1952	1951	1950	1949	1948	1947	1946	1945
Amebiasis	3,523	4,444	4,280	3,550	4,568	5,543	4,871	3,365	4,093	3,412
Anthrax	22	45	47	60	49	54	60	69	40	40
Botulism	18	18	18	33	20	24	39	44	.....NA.....	.....NA.....
Brucellosis	1,823	2,032	2,537	3,139	3,510	4,235	4,991	6,321	5,887	5,049
Chancroid*	3,003	3,338	3,738	4,233	4,977	6,707	7,661	9,515	7,091	5,515
Cholera	—	—	—	—	—	—	—	—	—	—
Dengue	6	8	5	16	26	46	24	35	40	106
Diphtheria	2,041	2,355	2,960	3,983	5,796	7,969	9,493	12,262	16,354	18,675
Encephalitis, acute infectious (primary)	2,606	1,935	1,912	1,123	1,135	903	730	785	728	785
Gonorrhea *	242,050	238,340	244,957	254,470	286,746	317,950	345,501	380,666	368,020	287,181
Granuloma inguinale*	618	667	951	1,352	1,783	2,402	2,469	2,330	2,232	1,857
Hansen disease (leprosy)	56	60	57	57	44	41	63	56	43	40
Hepatitis, infectious†	50,093	33,700	17,428	7,349	2,820	2,027	709	1,092	.....NA.....	.....NA.....
Leptospirosis	48	42	62	9	30	17	18	14	.....NA.....	.....NA.....
Lymphogranuloma venereum*	875	983	1,200	1,300	1,427	1,925	2,429	2,526	2,603	2,631
Malaria	715	1,310	7,023	5,600	2,184	4,151	9,606	15,116	48,610	62,763
Measles (rubella)	682,720	449,146	683,077	530,118	319,124	625,281	615,104	222,375	655,843	146,013
Meningococcal disease	4,436	5,077	4,884	4,164	3,788	3,519	3,376	3,420	5,693	8,208
Murine typhus fever	163	221	205	378	685	985	1,171	2,050	3,365	5,193

## HISTORICAL TABLES — 1945–1954

Pertussis (whooping cough)	60,886	37,129	45,030	68,687	120,718	69,479	74,715	156,517	109,860	133,792
Plague	—	—	—	—	—	—	—	—	—	—
Poliomyelitis, total	38,476	35,592	57,879	28,386	33,300	42,033	27,726	10,827	25,698	13,624
Psittacosis	563	169	135	25	26	35	32	27	26	27
Rabies, animal <sup>§</sup>	7,297	8,903	8,445	8,008	7,901	7,587	8,495	8,920	10,850	9,928
Rabies, human <sup>¶</sup>	13	12	24	18	18	10	24	26	34	43
Rocky Mountain spotted fever	294	313	327	347	464	570	547	596	587	472
Salmonellosis, excluding typhoid fever	5,375	3,946	2,596	1,773	1,233	1,243	882	951	723	649
Shigellosis	13,846	16,533	23,197	32,215	23,367	29,080	23,753	17,048	24,286	34,943
Smallpox	—	—	—	—	—	—	49	57	337	346
Streptococcal sore throat and scarlet fever	147,785	132,935	113,677	84,151	64,494	87,220	91,295	93,595	125,511	185,570
Syphilis, primary and secondary*	7,147	8,637	10,449	14,485	23,939	41,942	68,174	93,545	94,957	77,007
Total, all stages*	130,697	148,573	167,762	174,924	217,558	256,463	314,313	355,592	963,647	355,114
Tetanus	524	506	484	506	486	579	601	560	.....NA.....	.....NA.....
Trichinosis	277	395	367	393	327	353	487	451	.....NA.....	.....NA.....
Tuberculosis **	79,775	84,304	86,700	118,491	121,742	134,865	137,006	134,946	119,256	114,931
Tularemia	681	601	668	702	927	1,179	1,036	1,401	1,355	900
Typhoid fever	2,169	2,252	2,341	2,128	2,484	2,795	2,840	3,075	3,268	4,211
Yellow fever	—	—	—	—	—	—	—	—	—	—

\* Data were reported for fiscal years 1945–1946; data were reported by calendar year beginning in 1947.

† Data for 1953 and 1954 includes serum hepatitis.

‡ Data for 1945–1951 from Bureau of Animal Industry, U.S. Department of Agriculture, Agricultural Research Administration.

¶ Registered deaths.

\*\* Includes newly reported active and inactive cases, 1945–1951; new active cases only beginning 1952.

.....Last indigenous case reported 1911; last imported, 1924.....

## HISTORICAL TABLES — 1983-1992

**TABLE 7. NOTIFIABLE DISEASES — deaths from specified notifiable diseases, United States, 1983-1992. (Numbers in ICD column refer to the category numbers listed in the Ninth Revision of the International Classification of Diseases, 1975.)**

Cause of Death	ICD*	1992	1991	1990	1989	1988	1987	1986	1985	1984	1983
AIDS <sup>†</sup>	*042-*044	33,566	29,555	25,188	22,082	16,602	13,468	10,900	6,040	2,943	1,141
Amebiasis	006	6	5	5	4	2	1	1	1	4	21
Anthrax	022	-	-	-	-	-	-	-	-	-	-
Botulism, foodborne	005.1	1	2	4	2	2	1	1	1	4	7
Brucellosis	023	-	-	-	-	-	-	-	-	-	-
Chancroid	099.0	-	1	2	2	-	-	-	-	-	-
Cholera	001	2	2	1	1	-	-	-	-	-	-
Diphtheria	032	1	142	167	143	133	146	230	153	168	169
Encephalitis, acute infectious <sup>§</sup>	062-064,049	126	3	3	4	3	7	7	2	3	4
Gonococcal infections	098	4	3	3	3	4	3	7	2	3	4
Granuloma inguinale	099.2	-	-	-	-	-	-	-	-	-	-
<i>Haemophilus influenzae</i> , invasive	041.5	16	17	16	16	25	25	21	22	14	11
Hansen disease (leprosy)	030	2	-	3	4	1	1	2	6	6	3
Hepatitis, viral, infectious (Hep A)	070.0,070.1	82	71	76	88	70	77	65	80	77	82
Hepatitis, viral, serum (Hep B)	070.2,070.3	903	912	816	711	621	595	557	490	465	438
Hepatitis, viral, other and unsp.	070.4-070.9	1,016	857	686	717	599	510	384	372	327	343
Leptospirosis	100	2	1	2	2	2	1	-	4	-	5
Lymphogranuloma venereum	099.1	1	1	2	3	11	7	5	5	13	7
Malaria	084	8	4	3	2	3	2	5	4	1	3
Measles (rubella)	055	4	27	64	32	3	2	2	4	1	4
Meningococcal disease	036	201	198	215	273	278	258	286	257	300	299
Mumps	072	-	1	3	2	2	2	-	1	1	2
Murine typhus fever	081.0	-	-	12	1	4	1	1	6	4	7
Pertussis (whooping cough)	033	5	-	-	-	-	-	-	1	3	5
Plague	020	1	1	1	1	1	1	1	1	1	1
Poliomyelitis, total	045.0-045.9	-	-	-	-	-	-	-	-	-	-
Pituitacosis	073	4	-	2	1	1	2	-	1	1	1
Rabies, human	071	1	3	2	1	1	1	-	-	2	2
Rheumatic fever, acute	390-392	100	89	66	70	76	42	60	56	70	87
Rocky Mountain spotted fever	082.0	13	13	20	10	20	21	19	22	34	35
Rubella (German measles)	056	1	1	8	4	1	-	1	1	1	3
Salmonellosis, incl. paratyphoid fever	002.1-002.9,003	47	53	80	99	66	105	102	117	90	82
Shigellosis	004	8	10	16	8	13	4	17	8	9	9
Syphilis	090-097	91	93	106	85	98	80	80	105	121	22
Tetanus	037	9	11	9	17	16	22	23	20	23	22
Trichinosis	124	-	-	-	-	-	-	-	-	-	-
Tuberculosis (all forms)	010-018	1,705	1,713	1,810	1,970	1,921	1,755	1,782	1,752	1,729	1,779
Tularemia	021	3	2	1	1	2	4	4	2	2	1
Typhoid fever	002.0	-	1	1	1	1	2	2	2	2	3
Varicella (chickenpox)	052	100	81	120	89	83	89	47	68	53	57

\* Numbers in ICD column refer to the category numbers listed in the Ninth Revision of the International Classification of Diseases,<sup>738</sup> (The asterisks in the ICD column pertain to the ICD code, not a footnote.

† For 1983-1986, deaths are estimated from death certificates with mention of conditions coded to deficiency of cell-mediated immunity (ICD-9 No. 279.1). Includes other human immunodeficiency virus (HIV)-related deaths and other diseases classifiable as deficiency of cell-mediated immunity.

§ Arthropod-borne encephalitis and other nonarthropod-borne viral diseases of the central nervous system.

Source: National Center for Health Statistics, System, 1983-1992. Deaths are classified according to the Ninth Revision, ICD.

## State and Territorial Epidemiologists and Laboratory Directors

State and Territorial Epidemiologists and Laboratory Directors are acknowledged for their contributions to *CDC Surveillance Summaries*. The epidemiologists listed below were in the positions shown as of June 1995, and the laboratory directors listed below were in the positions shown as of June 1995.

<b>State/Territory</b>	<b>Epidemiologist</b>	<b>Laboratory Director</b>
Alabama	John P. Lofgren, MD	William J. Callan, PhD
Alaska	John P. Middaugh, MD	Gregory V. Hayes
Arizona	Lawrence Sands, DO, MPH	Barbara J. Erickson, PhD
Arkansas	Thomas C. McChesney, DVM	Michael G. Foreman
California	Duc J. Vugia, MD, MPH	Michael G. Volz, PhD
Colorado	Richard E. Hoffman, MD, MPH	Ronald L. Cada, DrPH
Connecticut	James L. Hadler, MD, MPH	Sanders F. Hawkins, PhD
Delaware	A. LeRoy Hathcock, PhD	Mahadeo P. Verma, PhD
District of Columbia	Martin E. Levy, MD, MPH	James B. Thomas, ScD
Florida	Richard S. Hopkins, MD, MSPH	E. Charles Hartwig, ScD
Georgia	Kathleen E. Toomey, MD, MPH	Elizabeth A. Franko, DrPH
Hawaii	Richard L. Vogt, MD	Vernon K. Miyamoto, PhD
Idaho	Jesse F. Greenblatt, MD, MPH	Richard H. Hudson, PhD
Illinois	Byron J. Francis, MD, MPH	David F. Carpenter, PhD
Indiana	Edmundo M. Muniz, MD, PhD, MSc	Barbara Wilder (Acting)
Iowa	M. Patricia Quintilisk, MD, MPH	W. J. Hausler, Jr, PhD
Kansas	Gianfranco Pezzino, MD, MPH	Roger H. Carlson, PhD
Kentucky	Reginald Finger, MD, MPH	Thomas E. Maxson, DrPH
Louisiana	Louise McFarland, DrPH	Henry B. Bradford, Jr, PhD
Maine	Kathleen F. Gensheimer, MD, MPH	Philip W. Haines, DrPH
Maryland	Diane M. Dwyer, MD	J. Mehseri Joseph, PhD
Massachusetts	Alfred DeMaria, Jr, MD	Ralph J. Timperi, MPH
Michigan	Kenneth R. Wilcox, Jr, MD, DrPH	Robert Martin, DrPH
Minnesota	Michael T. Osterholm, PhD, MPH	Pauline Bouchard, JD, MPH
Mississippi	Mary Currier, MD, MPH	Joe O. Graves, PhD
Missouri	H. Denny Donnell, Jr, MD, MPH	Eric C. Blank, DrPH
Montana	Todd D. Damrow, PhD, MPH	Douglas O. Abbott, PhD
Nebraska	Thomas J. Safranek, MD	John D. Blosser
Nevada	Randall L. Todd, DrPH	Arthur F. DiSalvo, MD
New Hampshire	M. Geoffrey Smith, MD, MPH	Veronica C. Malmberg, MSN
New Jersey	Kenneth C. Spitalny, MD	Shahied I. Shahied, PhD
New Mexico	C. Mack Sewell, DrPH, MS	Loris W. Hughes, PhD
New York City	Benjamin A. Mbijica, MD, MPH	Stanley Reimer
New York State	Dale L. Morse, MD, MS	Lawrence S. Sturman, MD, PhD
North Carolina	J. Michael Moser, MD, MPH	Samuel N. Merritt, DrPH
North Dakota	Larry A. Shieley, MS, MPH	James D. Anders, MPH
Ohio	Thomas J. Halpin, MD, MPH	Kathleen L. Meckstroth, DrPH
Oklahoma	Joe P. Mallonee, MPH (Acting)	Garry L. McKee, PhD
Oregon	David Fleming, MD	Michael R. Skeels, PhD, MPH
Pennsylvania	James T. Rankin, Jr, DVM, PhD, MPH	Bruce Kieger, DrPH
Rhode Island	Bela T. Matyas, MD, MPH	Walter Combs, PhD
South Carolina	James J. Gibson, MD, MPH	Harold Dowda, PhD
South Dakota	Susan E Lance, DVM, MPH	Richard S. Steece, PhD
Tennessee	William L. Moore, MD	Michael W. Kimberly, DrPH
Texas	Diane M. Simpson, MD, PhD	David L. Maserang, PhD
Utah	Craig R. Nichols, MPA	Charles D. Brokopp, DrPH
Vermont	Robert O'Grady (Acting)	Burton W. Wilke, Jr, PhD
Virginia	Grayson B. Miller, Jr, MD	James L. Pearson, DrPH
Washington	Paul Stehr-Green, DrPH, MPH	Jon M. Counts, DrPH
West Virginia	Loretta E. Haddy, MA, MS	Frank W. Lambert, Jr, DrPH
Wisconsin	Jeffrey P. Davis, MD	Ronald H. Laessig, PhD
Wyoming	Gayle L. Miller, DVM, MPH	Carl H. Blank, DrPH
American Samoa	Edgar C. Reid, DSM, MPH	—
Federated States of Micronesia	Vacant	—
Guam	Robert L. Haddock, DVM, MPH	Jeff Benjamin (Acting)
Marshall Islands	Tony de Brum	—
Northern Mariana Islands	Jose L. Chong, MD	—
Palau	Jill McCready, MS, MPH	Adolpho Firpo-Betancourt, MD
Puerto Rico	Carmen C. Deseda, MD	Norbert Mantor, PhD
Virgin Islands	Donna M. Green, MD	

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available on a paid subscription basis from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 783-3238.

The data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the succeeding Friday. Inquiries about the *MMWR* Series, including material to be considered for publication, should be directed to: Editor/*MMWR* Series, Mailstop C-08, Centers for Disease Control and Prevention, Atlanta, GA 30333; telephone (404) 332-4555.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without special permission; citation as to source, however, is appreciated.

---

☆U.S. Government Printing Office: 1995-633-175/27017 Region IV