SORPTION OF THE AIRCRAFT DEICING FLUID COMPONENT METHYL-BENZOTRIAZOLE IN SOIL

David L. Kellner, 1st Lieutenant, USAF

AFIT/GEE/ENV/99M-11

Approved for public release; distribution unlimited

20000303 174

DTIC QUALITY INSPECTED 3

.

Disclaimer Statement

The views expressed in this thesis are those of the author and do not reflect the official policy or position of the Department of Defense or the United States Government.

SORPTION OF THE AIRCRAFT DEICING FLUID COMPONENT METHYL-BENZOTRIAZOLE IN SOIL

THESIS

David L. Kellner, B. S., E.I.T. 1st Lieutenant, USAF March 1999

Presented to the faculty of the Graduate School of Engineering of the Air Force Institute of Technology

Air University

In Partial Fulfillment of the

Requirements for the Degree of

Master of Science in Environmental and Engineering Management

Idwar

Lieutenant Colonel Edward Heyse, Ph.D.

Daniel E. R fis

Dr. Charles A. Bleckmann, Ph.D. Chairman

SORPTION OF THE AIRCRAFT DEICING FLUID COMPONENT METHYL-BENZOTRIAZOLE IN SOIL

THESIS

Presented to the faculty of the Graduate School of Engineering of the Air Force Institute of Technology Air University In Partial Fulfillment of the Requirements for the Degree of Master of Science in Environmental and Engineering Management

> David L. Kellner, B.S., E.I.T. 1st Lieutenant, USAF

> > March 1999

Approved for public release, distribution unlimited.

| <u>Table of Content</u> | Table | of Content |
|-------------------------|-------|------------|
|-------------------------|-------|------------|

| Acknowledgmentiv |
|---|
| List of Figuresv |
| List of Tablesvii |
| Abstractviii |
| I. Introduction. 1-1 1.1 Overview. 1-1 1.2 Purpose for Research. 1-2 1.3 Primary Goals. 1-3 1.4 Scope. 1-3 1.5 Definitions. 1-4 |
| II. Literature Review.2-12.1 Introduction.2-12.2 Review of ADF and the Air Force.2-22.3 Composition and Structure.2-42.3.1 Glycol.2-42.3.2 4(5) Methyl-benzotriazole.2-62.4 Biodegradation Concerns.2-92.5 Sorption.2-102.5.1 Potential Hazards.2-122.6 Additional Related Research.2-16 |
| III. Methodology. 3-1 3.1 Overview. 3-1 3.2 Preliminary Experiments. 3-1 3.2.1 Pan Evaporation. 3-1 3.2.2 Toxicity Matrix. 3-2 3.3 Soil. 3-3 3.3.1 Purpose. 3-3 3.3.2 Method of Collection. 3-3 3.3.3 Soil Types. 3-4 3.3.4 Analyzed Characteristics. 3-5 3.4 Experimental Preparation. 3-5 3.4.1 Soil Preparation. 3-5 3.4.2 Solute Preparation. 3-7 3.4.3 Conditional Control for Analysis. 3-8 3.4.4 Statistical Design. 3-9 |

| 3.5 High Performance Liquid Chromatography (HPLC) | 3-11 |
|---|------|
| 3.5.1 Theory | 3-11 |
| 3.5.2 Purpose | 3-13 |
| 3.5.3 Procedure | 3-13 |
| 3.5.4 Calibration Curve | 3-14 |
| 3.6 Data Collection | 3-15 |
| 3.6.1 HPLC | 3-16 |
| 3.6.2 Quality Assurance and Quality Control (Reproducibility) | 3-17 |
| IV. Data Analysis | 4-1 |
| 4.1 Overview | 4-1 |
| 4.2 Preliminary Experiment Results | 4-2 |
| 4.2.1 Pan Evaporation | 4-2 |
| 4.2.2 Toxicity Matrix | 4-2 |
| 4.2.3 Soil Characteristics | 4-3 |
| 4.3 Statistical Quantitative/Qualitative Analysis | 4-3 |
| 4.4 Isomer Results | 4-5 |
| 4.5 Wright-Patterson Soil Results | 4-7 |
| 4.5.1 Wright-Patterson Soil – Untreated | 4-7 |
| 4.5.2 Wright-Patterson Soil – Treated | 4-9 |
| 4.6 Denver International Airport (DIA) Soil Results | 4-11 |
| 4.7 Westover ARB Soil Results | 4-13 |
| 4.8 Soil Type Differentiation | 4-15 |
| 4.9 Organic Content Differentiation | 4-17 |
| 4.10 Acclimation Differentiation | 4-18 |
| 4.11 Propylene Glycol Additive Soil Differentiation | 4-20 |
| 4.12 Adjustments | 4-22 |
| V. Conclusions and Recommendations | 5-1 |
| 5.1 Conclusions. | 5-1 |
| 5.1.1 Statistical Evaluation of Laboratory Results | 5-1 |
| 5.1.2 Significant Conclusions from Statistical Results | 5-3 |
| 5.2 Follow-on Research | 5-4 |
| 5.2.1 Analysis of Other ADF Components | 5-4 |
| 5.2.2 Breakdown of MeBT Naturally <i>in-situ</i> | 5-5 |
| 5.2.3 Environment Variation Study | 5-5 |
| 5.2.4 HPLC Identification of MeBT Degradation Product | 5-5 |
| 5.2.5 Repeat this Study | 5-5 |

.

| Appendix A: Soil Analysis Report A-1 |
|---|
| Appendix B: Calculation of Field Capacity, Solution Concentration, and Moisture Content and Individual Sample Size CalculationsB-1 |
| Appendix C: Preliminary Experimentation DataC-1 |
| Appendix D: Calibration Curve Data for MeBT IsomersD-1 |
| Appendix E: Soil Calculations & Results from Wright-Patt. AFB Soil – Untreated E-1 |
| Appendix F: Soil Calculations & Results from Wright-Patt. AFB Soil – TreatedF-1 |
| Appendix G: Soil Calculations & Results from DIA SoilG-1 |
| Appendix H: Soil Calculations & Results from Westover ARB Soil |
| Appendix I: Achieved K _{ow} Values I-1 |
| Appendix J: Statistical AnalysisJ-1 |
| BibliographyBIB-1 |
| VitaVIT-1 |

(

•

Acknowledgment

My primary word of thanks goes to Dr. Charles Bleckmann, my thesis committee chairman and advisor. Without his continued support of this important subject and his enduring patients, I may not have had the opportunity to be involved with such a technical and rewarding laboratory study.

I would like to thank Lt Col. Edward Heyse. He guided me through the error of my ways and devoted his time to this study, eventhough he has departed from the formal faculty at AFIT.

My gratitude extends to Professor Dan Reynolds for his enforcement of what is certainly the most statistical demanding work I've done to this point. Your reinforcement and encouragement are a welcomed site.

I also extend my appreciation to Major Jeff Cornell. Without his persistence and insight to involve AFIT in this endeavor, I would not have had this opportunity. He opened shortcuts that otherwise would have taken numerous hours to overcome.

Additional thanks go to my colleague, Captain Baron Burke, without whom I would not have been able to accomplish the work I did.

Lastly, and certainly not least, I want to express my deepest and most heart-felt gratitude to my wife Nancy. Though you may not understand exactly want this thesis effort encompasses in topic, you do understand what it has meant to me to have you by my side throughout the thesis process. Just stand warned that it may not be over yet. O

Dave Kellner

List of Figures

| Figure 1-1 | The structure and nomenclature of 5-MEBT and 4-MEBT1-7 | |
|-------------|---|---|
| Figure 1-2 | Propylediol (Propylene Glycol, PG)1-8 | |
| Figure 1-3 | An $[AlO_4]^{5-}$ or $[SiO_4]^{4-}$ Tetrahedra | |
| Figure 2-1 | Soil Sorption Illustration2-1 | 2 |
| Figure 2-2a | Langmuir Isotherm2-1 | 5 |
| Figure 2-2b | Freundlich Isotherm2-1 | 5 |
| Figure 3-1 | HPLC Schematic | 1 |
| Figure 3-2 | HPLC Column | 2 |
| Figure 3-3 | Retention Time Graph | 2 |
| Figure 3-4 | Isomer Calibration Curve | 4 |
| Figure 3-5 | Methodology Format 3-1 | 6 |
| Figure 3-6 | Typical HP ChemStation Data Result for MeBT | 7 |
| Figure 4-1 | Establishment of Sorption Equilibrium Point | |
| Figure 4-2 | HPLC Isomer Results | |
| Figure 4-3 | Accounting for Unrecoverable Isomer4-6 | |
| Figure 4-4 | WPAFB Untreated Sorption Coefficient vs. Concentration – 4-MeBT 4-7 | |
| Figure 4-5 | WPAFB Untreated Sorption Coefficient vs. Concentration – 5-MeBT 4-8 | |
| Figure 4-6 | WPAFB Untreated Soil Sorption Results | |
| Figure 4-7 | WPAFB Treated Sorption Coefficient vs. Concentration – 4-MeBT4-1 | 0 |
| Figure 4-8 | WPAFB Treated Sorption Coefficient vs. Concentration – 5-MeBT4-1 | 0 |
| Figure 4-9 | WPAFB Treated Soil Sorption Results 4-1 | 1 |
| Figure 4-10 | DIA Sorption Coefficient vs. Concentration – 4-MeBT4-1 | 2 |
| Figure 4-11 | DIA Sorption Coefficient vs. Concentration – 5-MeBT4-1 | 2 |

| Figure 4-12 | DIA Soil Sorption Results |
|-------------|---|
| Figure 4-13 | Westover ARB Sorption Coefficient vs. Concentration – 4-MeBT 4-14 |
| Figure 4-14 | Westover ARB Sorption Coefficient vs. Concentration – 5-MeBT 4-14 |
| Figure 4-15 | Westover ARB Soil Sorption Results 4-15 |
| Figure 4-16 | Test 1 t-test Results for 4-MeBT – Soils 3&4 4-16 |
| Figure 4-17 | Test 1 t-test Results for 4-MeBT – Soils 3&4 4-16 |
| Figure 4-18 | Average Sorption Coefficient for Soil Type Differentiation |
| Figure 4-19 | Test 2 t-test Results for 4-MeBT – Soils 3&4 4-17 |
| Figure 4-20 | Test 2 t-test Results for 4-MeBT – Soils 3&4 4-18 |
| Figure 4-21 | Average Sorption Coefficient for Organic Content Differentiation 4-18 |
| Figure 4-22 | Test 3 t-test Results for 4-MeBT – Soils 3&4 |
| Figure 4-23 | Test 3 t-test Results for 4-MeBT – Soils 3&4 4-19 |
| Figure 4-24 | Average Sorption Coefficient for Acclimation Differentiation |
| Figure 4-25 | Test 4 t-test Results for 4-MeBT – Soils 3&4 4-21 |
| Figure 4-26 | Test 4 t-test Results for 4-MeBT – Soils 3&4 |
| Figure 4-27 | Average Sorption Coefficient for PG Additive Soil Differentiation |

List of Tables

| Table 2-1 | Glycol Toxicity |
|------------|---|
| Table 2-2 | Physical and Chemical Properties of MeBT and BT 2-7 |
| Table 2-3 | Solubility of MeBT2-7 |
| Table 3-1 | Soil Characteristics |
| Table 4-1 | Soil and Concentration Identification4-1 |
| Table 4-2 | HPLC Concentration/Soil/Isomer Results4-6 |
| Table 4-3 | WPAFB Untreated Average Sorption Coefficient Results |
| Table 4-4 | WPAFB Treated Average Sorption Coefficient Results4-11 |
| Table 4-5 | DIA Soil Average Sorption Coefficient Results |
| Table 4-6 | Westover ARB Soil Average Sorption Coefficient Results |
| Table 4-7 | Average Sorption Coefficient for Soil Type Differentiation4-17 |
| Table 4-8 | Average Sorption Coefficient for Organic Content Differentiation4-19 |
| Table 4-9 | Average Sorption Coefficient for Acclimation Differentiation 4-20 |
| Table 4-10 | Average Sorption Coefficient for PG Additive Soil Differentiation4-22 |

Abstract

The air transport industry requires deicing agents to maintain flight operations in cold weather conditions. The main type of aircraft deicing fluid (ADF) being used is Type I, composed of a propylene glycol (PG)-base, water, and several additives. Among these additives are corrosion inhibitors, wetting and thickeners; some of which are known to be toxic to microorganisms. Most of the spent ADF is closely monitored and collected for treatment or recycle; however, total capture is not possible for a variety of conditions. Therefore, investigation of suitable remediation technique for used deicing fluid is appropriate.

The goal of this research was to establish if 4(5) methyl-benzotriazole (MeBT) sorbs to various soils after the application of several concentration variations. If so, then how much and in what proportions or percentage of original concentration is sorbed. Since little research has been done on this compound, I have tried to make the best of what information is closely related to MeBT. Presented here is defining information on sorption, the definition and types of sorption isotherms, and supporting, useful reports from scientific journals and one doctoral student.

Recent studies have raised questions as to the toxicity of the deicing fluid additives, especially the corrosion inhibitor MeBT. It is known that PG exerts an oxygen demand; however, it is shown that MeBT is only slightly sorbed to the soils studied and may also degrade. Having applied a MeBT solute to the several soils in varied concentrations, centrifuged the samples, washed the soil with methanol, and employed standard High Performance Liquid Chromatograph (HPLC) techniques, the percent of MeBT recovered from the soil and a sorption coefficient for each soil was determined.

viii

This study demonstrated that MeBT scarcely sorbs to soil particles and sorbs only slightly better to organic material. Portions of a specific isomer are unrecoverable at lower concentrations. In most cases, nearly 100 percent of MeBT was accounted for when an initial water and subsequent methanol extraction(s) were performed.

This work parallels that of Captain Burke and will be used as reference material for Major Cornell, a doctoral student at the University of Colorado. Previous studies have indicated that MeBT degrades along with PG; this research will more clearly define what happens to MeBT after it comes in contact with soil. Proving that MeBT sorbs to soil is the first step to define a process to deal with spent ADF.

I. Introduction

1.1 Overview

Snow and ice buildup on airframes requires the air transport industry to use deicing/anti-icing agents to maintain flight operations in cold weather conditions. Commercial airports and airbases use hundreds of millions of liters of deicing fluid, approximately 52M liters of concentrated aircraft deicing fluid (ADF) are used in North America per year in an attempt to correct this safety problem [Cancilla *et al.*, 1998:3834; Sills, 1991]. In an average winter, Chicago-O'Hare International Airport alone used more than 1.5M gallons of ADF [Mericas and Wagoner, 301]. However, without this fluid the safety of air-travel would be compromised. The main type of deicing fluid currently used is SAE Type I, consisting of numerous components including water, propylene glycol (PG), and various additives. Among these additives are corrosion inhibitors, wetting agents, flame-retardants, and thickening agents; some of these are known to be toxic to microorganisms.

Other types of deicers, Type II, III, and Type IV, have a greater viscosity than Type I. They contain polymers and other additives that allow them to stick to applied surfaces resulting in longer lasting protection against ice build-up.

The use of deicing fluid is closely monitored and attempts are made to collect spent fluid; however, spent ADF is not totally captured. Over 50% can be lost due to various conditions. A substantial percentage (49% - 80%) ADF and water solution does not make it on the aircraft due to overspray or drippage. The excess ADF enters the parking apron's storm water drainage system or is collected in the bulk snow removal [Cancilla *et al.*, 1997:430; Chesterfield *et al.*].

Various treatments of used deicing/anti-icing fluid have been suggested such as land-farming or in-situ bioremediation. An understanding of the transport of ADF and the impact of the additive 4(5) methyl-benzotriazole (MeBT) is necessary to determine if bioremediation techniques can be employed effectively.

1.2 Purpose of Research

Two principle environmental concerns exist from the use of aircraft deicing fluid. First is the high oxygen demand of propylene glycol, approximately 50% to 90% of aircraft deicing fluid [Cancilla *et al.*, 1997:430; Arco]. Oxygen consumption occurs during the decomposition by bacteria. As ADF degrades, its oxygen demand is far greater than the replenishment of oxygen to the water; therefore, disrupting the environment of oxygen using organisms. This high biochemical oxygen demand (BOD), 1.68g O₂/g PG, makes treatment by municipal wastewater treatment plants an unlikely option [Halterman-O'Malley, 2-14; Cornell, 1997:2]. The second concern is the toxic nature of MeBT, a flame retardant/corrosion inhibitor in deicing/anti-icing fluid [Cancilla et al., 1998:3834].

This study investigated the sorption characteristics of MeBT with soil. Parallel study by Burke (1999) measured biodegradation of PG and MeBT using respirometry. High Performance Liquid Chromatography (HPLC) was also used to analyze MeBT. This work supported the research of Major Jeff Cornell, a Doctoral student at the University of Colorado at Boulder through the Air Force Institute of Technology. He has examined 1) toxic nature of ADF additives, 2)

environmental and human health risk of ADFs overall, and 3) the design for future ADF formulations and treatments.

1.3 Primary Goals

The primary goal of this study was to determine the sorption characteristic of MeBT in various soils. This was accomplished by using several concentrations of MeBT solute in three specified soil types to measure variation in the sorption results. Several replicates of each soil type and solute concentrations were performed to ensure statistically sound data.

1.4 Scope

The scope of work follows the studies of Johnson (1997) and Halterman-O'Malley (1997) with additional references to the work of Cornell (1998). However, the research was centered on the sorption of MeBT in various soil types. Solutes from the procedures were evaluated using standard High Performance Liquid Chromatographic techniques to determine the amount of MeBT in each sample before and after the sorption process. This was used to model sorption characteristics of MeBT on the variety of soil types and to predict the effects of sorption on biodegradation and bioremediation.

1.5 Definitions

<u>Absolute calibration</u> – A method relating detector response to sample concentration in order to perform quantitative analysis. Standard solutions of a sample to be quantitated are prepared and equal volumes chromatographed. Peak heights or peak areas are plotted versus concentrations to produce a calibration curve.

<u>Absorption</u> – (1) The uptake, drinking in, or imbibing of a substance; the movement of substances into a cell; transfer of substances from one medium to another [Atlas and Bartha, 533; Fetter, 117]. (2) The process by which a compound in solution or attached to the surface of the solid particle moves into the interior of the solid particle; diffusion into the inner pore-spaces or lattice structure of the solid particle.

<u>Acclimated</u> – A soil that has undergone the respirometry process with exposure to MeBT and PG, the soil microcosm has shown the ability to survive in the presence of these substances. <u>Adsorption</u> – Surface phenomenon, ability of a compound to clings to a solid at the solid:liquid interface [Fetter, 117].

<u>Adsorption coefficient</u> - (a.k.a. distribution coefficient, K_d) Ratio of the sorbed phase concentration to aqueous phase concentration. Analyte retention in HPLC is proportional to its adsorption coefficient (units: mL/g).

<u>Aerobic</u> - Having molecular oxygen present; process involving electron acceptors [Schwarzenbach *et al.*, 410].

<u>Anaerobic</u> – The absence of oxygen; able to live or grow in the absence of free oxygen, low redox potential. [Atlas and Bartha, 534; Schwarzenbach *et al.*, 410]

<u>Aromatic compound</u> – Carbon skeletons containing aromatic benzene ring and compounds that resemble benzene in chemical behavior. Their ring structure and stable bonds allow them to be resistant to degradation. These molecules contain delocalized clouds of resonant π -electrons and they favor substitution rather than additional reactions, both of which contribute to their stability [Schwarzenbach *et al.*, 32; Morrison, 322] <u>Biochemical Oxygen Demand</u> – (BOD) Refers to the amount of dissolved oxygen required by aerobic and facultative microorganisms to stabilize organic matter in water (a.k.a. biological oxygen demand) [Atlas and Bartha, 535].

<u>Biodegradation</u> - The microbial mediated process of chemical breakdown of a substance to a smaller product caused by microorganisms or their enzymes [Atlas and Bartha, 535].

<u>Chemisorption</u> – Occurs when the solute is incorporated on a sediment or soil surface by a chemical reaction [Fetter, 117].

<u>Completely Mixed Batch Reactor</u> – (CMBR) Self-contained vessel in which the reactions of the sorption process will take place, conditions are the same throughout.

<u>Deicing/Anti-icing</u> - Spreading or spraying of a liquid agent to: (1) Melt already formed ice and snow; (2) Applied before ice and snow is present; provides a thin layer of protection.

Desorption - The process by which sorbed species are released to the surrounding environment.

Desorption from the particle surface proceeds more readily than desorption from the interior of a particle.

<u>Field Capacity</u> - The maximum amount of water that an unsaturated zone of soil can hold against the pull of gravity [Fetter, 639]

Hydrophilic - A compound with an affinity for water. [Schwarzenbach et al., 38]

Hydrophobic – A compound that repels or has a dislike for water. [Schwarzenbach et al., 38]

Ion Exchange – This is where cations may be attracted to the region closest to a negatively

charged (δ) clay-mineral surface and are held by electrostatic forces [Fetter, 117].

<u>Isomer</u> – Numerous arrangements of a given set of atoms to yield a different structure depending on the number and type of atoms [Schwarzenbach *et al.*, 9].

<u>Isometric</u> – *Iso*-prefix indicating equal or same. *Isometric*-Relative amount of each isomer of MeBT that composes the chemical substance; for each isomer the relative amount of each is designated by the position of the nitrogen atom on the benzene ring [Lee, 216].

<u>Isotherm, sorption</u> – Relationship between sorbed and solution concentrations of a compound of equilibrium and constant temperature. There are numerous types of isotherms; three referenced here are: Linear, Freundlich, and Langmuir (see Section 2.5 for further explanation) [Fetter, 117-123; Park, 661]. In adsorption, the dependence of an analyte concentration to solution is due to the surface interactions on the adsorbent surface on the equilibrium concentration of analyte in the bulk solution. The shape of the adsorption isotherm can predict the chromatographic behavior of the solute.

<u>Methyl-benzotriazole</u> - Compound used as a flame retardant in air transport industry, used in ADF. Pasivates the surface of metal wires and makes them less reactive to O_2 to reduce corrosion. Lower solubility in water, semi-combustible, and stable. (A.k.a. 4(5)-Methyl-benzotriazole, Tolyltriazole, MeBT, or TTZ) [Cornell et al.].

Structure:



 Numbering scheme for 5-Methyl-1H-Benzotriazole.
 Numbering scheme for 4-Methyl-1H-Benzotriazole

 Figure 1-1.
 The structure and nomenclature of 5-MEBT and 4-MEBT.

<u>Natural Attenuation</u> – Numerous processes including dilution, oxidation, and sedimentation that remove material from the environment over time.

<u>Octanol-Water Partitioning Coefficient, (K_{ow}) </u> – Ratio of concentrations of a compound in octanol and water at equilibrium, an indication of hydrophobicity, $C_{octanol}/C_{water}$ [Fetter, 132-133]. <u>Organic Carbon Partitioning Coefficient, (K_{oc}) </u> – (1) The extent to which a chemical partitions between the organic carbon and solution phases. (2) The smaller the K_{oc} value the greater the concentration of contaminant in solution. (3) K_{oc} is related to the distribution coefficient, K_d by the formula: $K_d = (K_{oc})(f_{oc})$ where K_d is an index for sorption of a given contaminant on a particular soil (at equilibrium).

(units: μg adsorbed/g OC)/($\mu g/mL$ solution)

<u>Organic Matter Partitioning Coefficient, (K_{om}) </u> - Extent that a chemical partitions between organic matter and solution. $K_d = K_{om}(f_{oc})$ Note: For "average" soil organic matter $f_{oc} = 0.58(f_{om})$ $[(\mu g/g)/(\mu g/mL)]$ [Fetter, 132].

<u>Propylene Glycol</u> – (PG) Stable, non-toxic to mammals, humectant compound that is used in the aircraft, food, and cosmetic industries [RACB, Merck Index 12th ed.].

Structure:



Figure 1-2. Propylediol (Propylene Glycol, PG)

<u>Recalcitrant</u> – A chemical that is resistant to microbial attack or degradation [Atlas and Bartha, 543].

<u>Silicate</u> – Soils made mainly of small quartz granules (a.k.a. sand or sandy soil) [Lee, 128]. <u>Solubility</u> – The maximum concentration of a contaminant that can be dissolved in water at a specific temperature. <u>Sorption</u> – It is the adherence of analyte to soil, organic, or other particles. This process involves several classifications including *absorption*, *adsorption*, *chemisorption*, and *ion exchange* (see individual definitions) [Fetter, 117; Lin *et al.*, 2178].

<u>Stationary phase</u> – This is that part of the chromatographic system which is in equilibrium with the mobile phase, refers to phase that does not move - usually a solid.

<u>Tautomers</u> - Derivative of a compound, has the same basic structure with one or more compounds added on [Cornell *et al.*].

<u>Zeolite</u> –Clay mineral with a three-dimensional framework dawn from an open framework of $[AlO_4]^{5-}$ or $[SiO_4]^{4-}$ tetrahedra linked together (a.k.a. clay or clayey soil).

Structure:



Figure 1-3: An $[AlO_4]^{5-}$ or $[SiO_4]^{4-}$ tetrahedra (primary building unit) [Lee, 128]. The tetrahedra link together leading to the formation of rings and cages (Secondary Building Units or SBU's) (a.k.a. clay or clayey soil).

II. Literature Review

2.1 Introduction

Aircraft deicing fluids may reach surface and groundwaters presenting significant long-term affects on the milieu. Only a small percentage of the ADF used at most airports and military airbases is captured and/or treated. The majority is released into the environment through surface runoff or stormwater runoff [Sills, 1991; USEPA]. The passing and subsequent enforcement of the Clean Water Act of 1987 has only recently turned public and regulatory attention to the impact that ADF has on the environment. The Environmental Protection Agency (EPA) Storm Water Discharge regulations, effective 17 December 1990, and the National Pollution Discharge Elimination System (NPDES) permit program have focused on reducing point source pollutants [Oakley and Forrest]. As a result, several large international airports, Denver IAP, Chicago O'Hare IAP, Baltimore-Washington IAP, are currently operating directly under environmental regulatory agency guidelines enforced by state or regional regulators [NRDC, Cornell et al.]. Two states, New York and Pennsylvania, have issued "zero-discharge" regulations concerning the glycol component of deicing fluids. The Air Force has even cut back on most cold weather flying operations that require ADF in attempts to lessen the problem [Cornell *et al.*].

Most concerns with ADF in the environment stem from the glycol content and the BOD burden it imparts. However, recent work [Pillard, 1993; Hartwell *et al.*, 1995; Fisher *et al.*, 1995; Cancilla *et al.*, 1997; Cornell, 1998] reveal that additives to ADF tend to create more toxic formulations than glycol alone. Of these studies, only Cancilla *et al.* [1997] isolated any specific compound that may be responsible for the increased toxicity observed. Two recent AFIT theses

dealt with environmental impacts of aircraft deicing/anti-icing fluids [Johnson; Halterman-O'Malley, 1997].

Earlier research has concluded that MeBT is found and can be detected in subsurface water from a major North American airport [Cancilla *et al.*, 1998:3834]. What is yet to be determined is the true extent and magnitude of contamination. This requires extensive testing to accurately determine how MeBT reacts once it is released into the environment. Little is known about the environmental fate and transport MeBT. The compositions of most ADFs are proprietary and compounds are modified slightly among the manufacturers. The octanol-water partitioning coefficient (K_{ow}) and metal-binding properties of MeBT suggests that sorption to soil and organic material could significantly retard the subsurface transport rate [Cornell *et al.*]. This chapter outlines the history of ADF, discusses the composition of ADF and the main components (PG and MeBT), and gives a detailed summary of the sorption process, while looking briefly at the biodegradation relationship of PG and MeBT.

2.2 Review of ADF and the Air Force

The commercial aircraft transportation industry classifies their highly used deicing fluids as Type I or Type II. The composition of these fluids includes propylene glycol and/or ethylene glycol. Commercial fluids were designed to be primary freezing point depressants (FPD) as well as meet the performance specifications of the International Standards Organization (ISO) and the Society of Automotive Engineer (SAE).

Military deicing/anti-icing fluids were developed by the Navy and adopted for Air Force use [Chesterfield *et al.*]. These military derivative specification were developed to afford maximum corrosion protection to prevent corrosion problem in steel storage tanks and deicing trucks. Secondarily, the added benefit of aircraft and other metals protection was highly desired. However, it is important to note that the time of fluid exposure to aircraft metals is of short duration and is diluted by being mixed with water before application.

There are two types of deicing fluids that have been used by the Air Force. They are MIL-A-4823D Type I - propylene glycol base with a corrosion inhibitor and MIL-A-4823D Type II ethylene and propylene glycol (independent or mix) with a corrosion inhibitor. The Navy is still the OPR on the specification, even though the USAF is the largest user of the fluid [HQAFCEE, July 1995].

The composition of Type I fluid is "unthickened," with a low viscosity, and Type II's composition is "thickened," and offers better anti-icing action than Type I fluid. However, Type I fluid can be applied with existing Air Force equipment while Type II fluid cannot, due to viscosity. In addition, in March 1992, Brigadier General James E. McCarthy, Air Force Civil Engineer, directed the immediate prohibition on the use of ethylene glycol (EG) in the USAF [HQAFCEE, July 1995]. EG is also listed under CERCLA as a hazardous substance and is therefore subject to the Emergency Planning and Community Right to Know Act (EPCRA) [HQAFCEE, July 1995]. By prohibiting the use of EG, the AF resorted to use the PG-based ADF [HQAFCEE, Dec 1995]. Type II anti-icing solutions are no longer necessary because the Type I ADF can be used just prior to flight [Chesterfield *et al.*].

The Air Force Material Command's Wright Laboratory has evaluated the SAE, Aerospace Material Specification (AMS) Type I FPD fluid with the intent of using it in the Air Force. The adoption action has been approved, authorizing Air Force units to use SAE/AMS Type I specification fluid, as long as it is propylene glycol-based. The change from military specification (MIL SPEC) fluid to SAE/AMS Type I fluid will require additional training. Air and ground crews will have to be well versed in the characteristics of both fluids until existing stocks of the military specification fluid are depleted. Military specification fluid is applied while Air Force aircraft are being serviced and is designed to remove snow and ice [Chesterfield *et al.*]. Within the Air Force's flight operations community, there exist differences in the understanding and interpretation of published holdover times after deicing/anti-icing.

2.3 Composition and Structure

Much of the ADF used in the United States is classified as PG-based Type I, which contains 85-90 percent propylene glycol (FPD), 1-2 percent surfactant and trace additives (provide adhesion), 0.5-0.6 percent MeBT (corrosion inhibitor and flame retardant), and water [SAE]. This section will review the characteristics of both PG and MeBT as major components of ADF.

2.3.1 Glycol

Glycols are straight-chained alcohols with two attached hydroxyl groups [Morrison, 456]. The length of the chain can vary greatly. Propylene glycol is the component of ADF that provides freeze-point depression (FPD). These solutions can push the freezing point temperatures down to -13°C to -59°C depending on concentrations [Sigma Products, MSDS]. Once in the soil, this component of ADF is nearly as, if not more, mobile than water in the subsurface transports

system. This phenomenon is due to PG's high miscibility in water. Other characteristics include: highly hydrophilic, low volatility (stable indefinitely at room temperature), non-toxicity, easily biodegradable in both aerobic and anaerobic conditions, and a very high oxygen demand (ca. 1.68 g O₂/g PG degraded) [EA-Engineering, Bausmith and Neufield, Gooden, Halterman-O'Malley]. In waste treatment systems or in environmental impact assessments, the BOD exerted by PG is of primary concern.

Toxicity tests performed on flathead minnows with EG and PG are summarized in Table 2-1 below [ENSR Consulting].

| MATERIAL | COMPOSITION | 48 HOUR LC ₅₀ | 96 HOUR LC ₅₀ | | |
|------------------|-------------------|--------------------------|--------------------------|--|--|
| Ethylene Glycol | Formulated in ADF | 8,541 mg/L | 8,045 mg/L | | |
| Ethylene Glycol | Pure | 81,950 mg/L | 72,860 mg/L | | |
| Propylene Glycol | Formulated in ADF | 791 mg/L | 709 mg/L | | |
| Propylene Glycol | Pure | 61,200 mg/L | 55,860 mg/L | | |

NOTE: LC₅₀ is 50 percent mortality concentration

Table 2-1. Glycol Toxicity

Further government toxicity tests on mice indicate no real toxicological threat. Results of the study concluded that there were no treatment-related effects on pup weight, health, or development. In addition, no effect was found on the fertility of adult mice [RACB]. There was one ill-effect found with dermal exposure of PG; when applied to skin, the PG tends to replace some of the important components necessary to the skin, including water. In very high doses, PG may cause liver abnormalities or kidney damage [Network Marketing].

2.3.2 Methyl-benzotriazole

MeBT is added to ADF at a concentration of ca. 0.5% w/w to reduce the flammability hazard created when a glycol solution come into contact with metal components carrying direct current [NASA, 1968]. The ADF component MeBT is actually a mix of the two isomers, 4-methyl-benzotriazole (4-MeBT) and 5-methyl-benzotriazole (5-MeBT). In this study, the mixture of the two MeBT isomers is simply referred to as MeBT.

MeBT is a weak organic acid ($pK_a = 8.8$) which is relatively hydrophobic (Log $K_{ow} \approx 2$) and reacts strongly with some metals [Matulewicz; PMC Specialties Group; Cornell, 1998]. MeBT is a member of the benzotriazole family of compounds. Benzotriazole and their derivatives (BTs) are of great importance in organic synthesis, analytical and medicinal chemistry, and industry [Cornell, *et al.*], but there is little published information on their chemistry, toxicity, and biodegradability which is useful in assessing their likely fate in the natural environment [Cornell, *et al.*]. Following is a summary of the current literature on MeBT.

MeBT is often thought of as strictly a flame retardant when it is actually classified as a corrosion inhibitor. MeBT coats the surface of conductive wire and protects it from oxidation in the presence of glycols, thus mitigating ignition hazards. Unlike other proprietary corrosion inhibitors, MeBT is manufactured from the waste product of isocyanide formation, which makes it very inexpensive. Other corrosion inhibitors are often salts of dibasic organic acids (*e.g.* triethanolamine). Industrial uses for the corrosion inhibitors include corrosion control in metal alloy cooling water towers. The chemical properties of MeBT (Table 2-2) suggest that MeBT is highly reactive in the environment. For example, the structure of 4-MeBT (specifically the proximity of the non-polar methyl group to the polar nitrogen ring) would suggest the 4-MeBT hydrophobic surface is smaller than that of 5-MeBT [Cornell, *et al.*]. In addition, 4-MeBT elutes earlier than 5-MeBT from a C-8 reverse-phase liquid chromatography column (often faster elution corresponds to less hydrophobicity). These facts seem to contradict solubility data that indicates 5-MeBT is more soluble in water. Further, the MeBT's solubility in water is much larger than the solubility of either individual isomer. There is no current explanation for this phenomenon. Notice in the following table, the melting point of MeBT is lower than the melting point of either individual isomer. This property is normally only found in metal alloy chemistry [Cornell, 1998].

| | Sol | Sol | | a la star | adian ya' | Vap. Press. | | Sp. Gr. |
|----------------------------------|---------|----------|-----|-----------|-----------|-------------|-------|---------|
| SUBSTANCE | Water | Min. Oil | | Melt. Pt. | Boil. | @ 20°C | Log | @ 20°C |
| | (wt. %) | (wt. %) | pKa | (°C) | Pt. (°C) | (mm HG) | Kow | (solid) |
| 4-Me-benzotriazole (4-MeBT) | 0.10 | N/A | 8.7 | 145-146 | N/A | N/A | N/A | N/A |
| 5-Me-benzotriazole (5-MeBT) | 0.18 | N/A | 8.7 | 82-83 | N/A | N/A | N/A | N/A |
| MeBT ; 55-60% 5-MeBT plus | 0.55 | 0.01 | 8.7 | 76-87 | >300 | 0.03 (est.) | 2.17 | 1.36 |
| 45-50% 4-MeBT | | | | | | | (est) | |
| Benzotriazole (BT) | 1.98 | 0.04 | 8.5 | 98-99 | >350 | 0.04 | 2.09 | 1.24 |

TABLE 2-2. Physical and chemical properties of MeBT and BT [Cornell, 1998]

N/A – not available

Due to the complexity of the tautometric forms of MeBT, there appears to be no simple way to describe this chemical behavior, any attempt may result in an oversimplification of the process. A possible explanation for these phenomena is that the three tautomeric forms of each isomer act differently, yielding six molecular structures in total that must be considered [Cornell 1998].

Other characteristics of MeBT that are of interest include molecular weight of 133.16 and MeBT solubility, which is summarized in Table 2-3.

| IN: WATER | 95% ETHANOL | METHANOL | ACETONE | |
|--------------------|--------------------------|---------------------|---------------------|--|
| < 0.1 mg/mL @ 18 C | \geq 100 mg/mL @ 21° C | ≥ 100 mg/mL @ 21° C | ≥ 100 mg/mL @ 21° C | |
| | | | | |

TABLE 2-3 – Solubility of MeBT

[NTP Chemical Repository (Radian Corporation) for Tolyltriazole]

There is no published information to suggest the mechanisms of MeBT toxicity to microorganisms. The Cornell (1998) on-going study suggests that MeBT may be an uncoupling agent for the electron transport chains. It is found that MeBT meets Lenhinger's definition of an uncoupling agent: a weak acid possessing hydrophobic properties. Uncouplers are known to disrupt the chemiosmotic potential of a cell or microbe by shuttling protons across the cell membrane against the natural flow set up by the cell. However, MeBT does not meet all the criteria and therefore cannot be considered in this category, but does necessitate the further need of study in this area [Cornell, *et al.*].

The toxicity of ADF is increased with the addition of MeBT. Cancilla [Cornell, 1998] identified triazole compounds as the fraction of ADF presenting the greatest toxicity as measured by the Microtox[®] assay. These Microtox[®] tests show three results: 1) the MeBT inhibited biodegradation of PG more than other additives, 2) the MeBT was significantly more toxic to microorganisms than other component (in the order of lowest to greatest toxicity 4-MeBT < MeBT < 5-MeBT), and 3) other additives were significantly more toxic to *C. dubia* and *P. promelas* than the MeBT, [Pillard, 312]. Chronic toxicity data for MeBT is unavailable. Limited acute toxicity data [PMC, 1996] indicates MeBT itself is moderately toxic to *Lepomis*

macrochirus (31 mg/L 96 hr. LC₅₀.) and *Daphnia magna* (74 mg/L 48 hr. LC₅₀) [MSDS; Cornell, 1998].

The only information available on the potential environmental fate of MeBT is from US Patent 5,503,775 (1996) as referenced by Cornell [1998], which claims that, under aerobic conditions, 5-MeBT is biodegradable while 4-MeBT is recalcitrant.

Additional toxicity results of MeBT are presented as a LD_{50} range from 675 mg/kg to 1600 mg/kg with dose given orally in mice. The MeBT component significantly decreased cell growth rates and yields and inhibited the biodegradation of propylene glycol to a greater extent than the other components. When heated to decomposition this chemical emits toxic fumes. It is also an eye and lung irritant [PMC Specialties Group].

2.4 Biodegradation Concerns

In the past, the main concern regulators had with ADF discharge to the stormwater system had been the large BOD that accompanied the mixture. New research has bought to light a different problem, the biodegradation process of ADF components. Biodegradation is the most important natural or engineered treatment method for reducing the oxygen demand exerted by these discharges [Cornell, 1998]. Recent studies have set up soils enriched with microorganisms in aerobic liquid CMBRs; they propose three hypothetical conclusions that describe the relationship of PG and MeBT biodegradation:

- 1) Propylene Glycol biodegradation rates decrease with increased MeBT concentration
- 2) Cell growth rates decrease with increased MeBT concentration
- 3) Cell yield decreases with increased MeBT concentration [Cornell, 1998]

The biodegradation process of PG can yield intermediate products such as aldehydes and organic acids (lactic, pyruvic, or acetic acids) [MSDS - 1,2 Propanidiol; RACB]. However, these compounds, if formed, are in small quantities and are rapidly reduced to carbon dioxide and water or additional organic content. Many studies show that PG is readily degradable in soil or aquatic environments.

Noticeable degradation of MeBT was not observed in these studies or after any experiment, nor did MeBT degrade during the twelve months of feeding the enrichment culture full ADF formulations [Cornell, 1997].

This thesis effort attempts to provide one piece of the MeBT puzzle; additional information about the transport of MeBT is required before any conclusions can be made about MeBT being naturally degraded [NRDC; Cornell, 1998]. All that is available now is unpublished data, and a US Patent that proposes preferential degradation of 5-MeBT over 4-MeBT [US Patent 5,503,775]. Quantifying the effects of MeBT sorption is a necessary step before effective treatment can be considered, or before the environmental impact posed by the release of MeBT into the environment can be assessed.

2.5 Sorption

Different factors effect the fate and transport of a contaminant through water. One of these factors is the adherence of contaminant to soil, organic matter, and other materials. The sorption process includes several different classifications including absorption, adsorption, chemisorption, and ion exchange. In most cases all of these processes together are referred to as sorption. The association of a particular chemical, the sorbate, with any particular solid, the sorbent, is known

as absorption (the prefix of absorption, *ab*, comes from Latin, "away from" and the suffix *sorbere* meaning, "to suck") [Fetter, 131]. The underlying principle behind this attraction results from some form of bonding between the contaminant and adsorption receptor sites on the solid. The amount of sorption that occurs in groundwater is dependent on particular characteristics of the sorbate and sorbent. It is known that the more hydrophobic the compound, the larger the fraction sorbed in soil [Maagd *et al.*, 1899]. The amount of sorption that takes place on organic matter also follows various isotherms or has varied kinetic rates.

Because of the porous nature of the soil particles, the sorbates can physically penetrate the particle and become part of the volume of the nonaqueous medium [Schwarzenbach *et al.*, 32]. Absorption is orientated in the 3-dimensional matrix. The capacity of a solid to remove a sorbate from solution is a function of the concentration of the sorbate in water (solute).



FIGURE 2-1. Soil Sorption Illustration [Ferrante, 2].

Sorption reactions generally occur over a short period of time, however if the adsorbed contaminant begins to be incorporated into the structure of the sorbent, a slower reaction, known as absorption, begins to take place. First-order rate models have been frequently used to describe sorption rate processes in transport experiments, but usually with the understanding that the firstorder coefficients are approximations for physical diffusion into often undefined regions of solute immobilization [Ball and Roberts, 1238]. The sorption coefficient (K_d) is the ratio between the concentration of the compound and that of the soil [Maagd, 1900]. Alternatively, the difference between adsorption and absorption is that adsorption is the attraction between the outer surface of a solid particle and a contaminant, whereas absorption is the uptake of the contaminant into the physical structure of the solid.

Figure 2-1 shows the primary differences between intraparticle absorption versus surface adsorption. The main difference being that some contaminant particles are attracted to the outer surface of the soil particle, while others have been incorporated into the particle's structure [Ferrante and Gallagher].

2.5.1 Potential Hazards

The process of sorption can hinder the remediation of a groundwater aquifer system. Sorption tends to cause contaminants to move more slowly than the groundwater, therefore the effects must be taken into consideration when calculating how far the contaminant will travel in a given time period. There is also the tendency to underestimate the total amount of contaminant in the system. Since the contaminant may not show up on a pump test, the assumption would be that there is no more contaminant; however, there could be upwards of 35% in the system as sorbate [Yaniga]. This can drive up remediation costs and with prolonged exposure, increase risk factors.

Several factors are key to the contribution of the contaminant properties to the sorption process. The first, the solubility of a compound is inversely proportional to the amount of sorption that the solid phase can support [Yaniga]. The second factor is polarity. A polar chemical tends to dissolve easier in water than a nonpolar chemical. Therefore a polar contaminant will absorb less to a soil particle than a nonpolar one. Another factor is the viscosity of the solute. The more viscous the solute the less that it will tend to sorb to the soil particle and may have an affinity to dissolve in water [Yaniga]. The last key factor is the octanol:water partitioning coefficient (K_{ow}) is a measure of the hydrophobicity or lack of affinity for water of an organic compound. The more hydrophobic a compound is, the less soluble in water it will be and the more likely it will be adsorbed to soil particles.

Several key soil factors affect the sorption rate, including the texture of the soil particles. Soil with high clay and high organic matter will tend to exhibit higher sorption than sandy, low organic soils. Clay soils tends to sorb more than sand due to small particle size, high surface area, and high surface charge. Organic matter provides sorption sites for hydrophobic compounds. The higher f_{oc} soils tend to have higher sorption capacities. The pH of the fluid can also have an affect. For example, organic acids adsorb best under acidic conditions.

Absorption isotherms provide a rational way of accounting for incomplete recoveries of sorbates from sorbent. These isotherms are plotted on a graph and are representations of the sorbed concentration (S) or mass of contaminant sorbed per unit dry mass of soil (K_d) or organic matter (K_{om}) versus the final concentration (C_f) of the contaminant. In order to use isotherms to estimate the mass adsorbed, an equilibrium must be reached between the sorbent and the sorbate

and the isotherm must be considered reversible, but can be used when sorption is rate-limiting and hysteric also [Ball and Roberts, 1240].

There are numerous types of isotherms; three presented here are: Langmuir isotherms, Freundlich isotherms, and Linear isotherms. The Langmuir Isotherm contains two assumptions that usually make its use difficult in real life or a heterogeneous soil case. The assumptions are that the energy of adsorption is constant and the number of binding sites is finite. The equation for the Langmuir isotherm is:

$$S = \alpha \beta C / (1 + \alpha C)$$

where α = an absorption constant related to the binding energy (L/mg)

and β = the maximum amount of solute that can be absorbed by the solid (mg/kg) [Fetter, 133]. The Langmuir isotherm is shown in Figure 2-2A. The curvature at the end of the graph marks a point at which the receptor sites on the soil particle are full and there is no more room for additional adsorption [Schwarzenbach *et al.*, 33; Ferrante and Gallagher].



FIGURE 2-2 A & B

The Freundlich Isotherm (Figure 2-2B) is an equilibrium isotherm that is used most often in practical examples. The Freundlich equation normally results in a curved graph unless the logs of both S and C are taken. By taking the log of these terms, a straight line develops making easier to obtain the slope and intercept of the line. The equation of the Freundlich isotherm is similar to the linear isotherm, but a new exponential term (1/n) is used:

$$\mathbf{S} = (\mathbf{K})\mathbf{C}^{(1/n)}$$

where K and n are constants [Ferrante and Gallagher; Fetter, 132; Schwarzenbach et al., 32].

If there is a direct, linear relationship between the amount of a solute sorbed onto a solid, C^* , and the concentration of the solute, C, the absorption isotherm of C as a function of C^* will plot as a straight line on a graph. The equation for the Linear isotherm is:

$$C^* = (K_d)C \rightarrow \text{where } n=1$$

The linear isotherm has only one fitting parameter (K_d) and if applicable the isotherm (by definition) must pass through zero and the regression through the origin is appropriate. The best estimate of the linear slope (K_d) is simply the average K_d of the samples [Ball and Roberts, 1240].

There are two limitation to this model; one, is that it does not limit the amount of solute that can be absorbed onto the solid and, two, with only a few points a curvilinear plot may be mistaken for a linear relationship [Fetter, 132].
2.6 Additional Research

There are several studies on halogenated organic chemicals, benzene, and methylbenzene in aquifer material, but very few have concentrated their efforts on 4(5) methyl-benzotriazole (tolyltriazole). The following section summarizes some of the research performed in areas close to MeBT.

Work done, by Duke and Stanford Universities, on long term sorption of halogenated organic chemicals by aquifer material explains the sorption effects on the Canadian Forces Base aquifer material in Borden, ON [Ball and Roberts, 1237; Brusseau, 1989].

The report includes details on the sorption isotherms for tetrachloroethene (PCE) and 1,2,4,5tetrachlorobenzene (TeCB). It notes that the isotherms deviated from linearity when a 4 to 5 order of magnitude range in aqueous concentration was considered. However, in the dilute range ($<50 \mu g/L$), the deviation from linearity was inconsequential. Sorption of TeCB was approximately 40 times that of PCE due to the difference in the properties of the compounds [Ball and Roberts, 1247].

Ball and Roberts (1991) makes note of several things to watch out for or concentrate efforts on. One of these points is in the time for equilibrium. The time it takes halogenated organic compound to reach full sorption equilibrium can be long. In order to help shorten that time, this study compared the difference between pulverized and unpulverized soil samples from the Borden site. The increased surface area of the pulverized soil allows the system to reach equilibrium much faster than unpulverized samples, by an approximate 20:1 ratio. Pulverized

samples yielded approximately 94.7% +/-2.8% recovery of TeCB. Results from this study show a good linear fit of data at low concentration of both PCE and TeCB (<100 μ g/L); yet, significant deviation from linear occurs at approximately 1000 μ g/L, which is less than 1% of aqueous solubility [Ball and Roberts, 1241]. Further tests with the addition of reagent-grade CaCO₃ to dissolve the inorganic carbonate show that there was no diminishing of the sorption capacity during this acid wash. This demonstrates that this compound does not sorb to inorganie carbonates, which can be significant in certain soil types.

Conclusions by Ball and Roberts pertain to MeBT in that they contrast the commonly held belief that the sorption capacity is inversely related to the particle size. Ball and Roberts states that to avoid bias in experiments of this duration, it is of paramount importance to avoid losses through volatilization or transformation. This study details background and methodology used to achieve credible results.

Weber and Huang (1993) describe a distribution reactivity model for sorption by soil and sediment using the interaction of a sorbate and sorbent. They investigated the time dependence of solute phase distribution relationships (PDRs) in completely mixed batch reactors. The experiments were conducted using a range of concentrations to obtain a time series of nonequilibrium PDRs for each sorbent-sorbate system. Freundlich isotherms were used to characterize the increasingly nonlinear results. An explicit account of the methodology used for the completely mixed batch reactor (CMBR), sorbent-sorbate system, and preparation is included in this report. Each step agrees with the basic known facts of this procedure. Weber and Huang (1993) define the changes in parameters in three stages: (1) an initiation stage, (2) a logarithmic stage, and (3) an apparent equilibrium stage. It summarizes the hypothesis that there is a boundary layer around the particle that has first-order rates associated with it [Weber and Huang, 884].

The third of the related articles is on adsorption isotherms of benzene and methylbenzene. This is closest to parallel data that I have encountered. Although the methylbenzene study used vapors on activated carbon instead of soil, it is an appropriate reference [Yun, 1997].

Since activated carbon has high surface area, high surface charge, and organic consistency it makes a good material to use in the lab for sorption studies. The report contained useful information that directly relates to MeBT. A point of interest taken from the report was that the adsorption isotherms for benzene and methylbenzene were accomplished at 30°C and 70°C, slightly higher than what was used in this study. The adsorption capacity is higher for benzene than it is for methylbenzene. However, if you look at pressure, with low pressure the amount sorbed is higher for methylbenzene than for benzene. This implies that the affinity of sorption is higher for methylbenzene [Yun, 895].

Following this line of thought, benzene can be a suitable substitute for MeBT since there are no in-depth studies with MeBT at the present time. Additionally, since the log K_{ow} values of MeBT and benzene are similar (2.17 vs. 2.13) estimations for K_{oc} will be accomplished with equations derived from benzene pesticides [Fetter, 134; Karickhoff, 1984]. This will be the basis for the solute to soil mass mixture in the CMBR.

III. <u>Methodology</u>

3.1 Overview

This chapter illustrates the reasoning and procedures used to measure the sorption coefficient of MeBT in various soil types. A high performance liquid chromatograph (HPLC) was used to analyze the MeBT extracted from soil. Mixtures of MeBT and water were added to soils, in appropriate proportions, to simulate the potential exposure in the environment. Extraction fluid, containing water and methanol, was added to the completely mixed batch reactors to recover the MeBT. All samples were prepared under the same conditions for consistency.

3.2 Preliminary Experiments

Two preliminary tests were performed to obtain a better understanding of the experimental material. Pan evaporation and nutrient matrix experiments gave insight into the characteristics of PG and a PG/MeBT mixture, respectively. The pan evaporation experiment demonstrated the hydroscopic nature of PG, while the nutrient matrix experiment demonstrated the relationship between PG and MeBT with respect to soil microorganism growth.

3.2.1 Pan Evaporation

The pan evaporation test demonstrated the hydrophilic properties of PG. The test began by adding a small amount of PG to a pre-weighed aluminum weigh boat, 11.51 grams. The PG was exposed to the atmosphere at room temperature, approximately 19° C (67° F), to observe the evaporation rate. The viscosity of PG suggested that the liquid would take several days to evaporate. Results of the test showed differently. The PG solution gained and lost mass with the changes in atmospheric conditions. If the humidity rose, the masses in the weigh boat increased

and vice versa. The liquid took some time to acclimate to the new humidity, which created a day or so lag in reaction to the change. No measurable evaporation occurred during the length of the experiment. Results of the pan evaporation are contained in Appendix C.

3.2.2 Toxicity Matrix

The second trial performed was microbial growth in a nutrient matrix. This test was to show how different concentrations of the two components in ADF under investigation, PG and MeBT, affected the growth of microbes in separate microcosms. A matrix was established with a 5-step range of each chemical, from 0 - 100 mg/L, in deionized water. Microorganisms were extracted from soil obtained from near the lab (See 3.3.2 Method of Soil Collection for details). An inorganic nutrient solution (BOD buffer) was added to the microorganisms extracted from the soil to ensure a plentiful inorganic food supply for the duration of the test. Separate 10-mL screw-top test tubes were used for each concentration intersection in the matrix. Equal volumes of PG concentration solution (4-mL) and MeBT concentration solution (4-mL) were added to the test tube. Then the same amounts (2-mL) of nutrient solute (microbes, nutrient, and water) were added to the test tubes to bring the level up to 10-mL. This mixture was then shaken to completely mix the contents. Growth of the microbes was measured with a photospectrometer. It measured the Total Suspended Solid (TSS) amount by monitoring the amount of light passed through the test tube at a wavelength of 810 nm. The more TSS or growth in the tube, the higher the count on the meter. Data from this experiment can be found in Appendix C.

3.3 Soil

Following the work of Cornell, *et al.* (1998), Johnson (1997), and Halterman-O'Malley (1997) three soil types in four configurations were used in this study. These three soils represent a range of soil types that could receive ADF applications. Soil samples were obtained from Denver International Airport (DIA), Westover Air Reserve Base (Westover), and Wright-Patterson AFB (WPAFB).

3.3.1 Purpose

Soil variations are important to this study in that a comparison matrix is constructed with the soils of interest and their mean K_d values. Four soil variations allow comparison of soil types, organic content, the exposed vs. unexposed soil difference, and the reaction to PG addition. This information allows a more complete assessment of the sorption characteristics of MeBT. This research will add to the body of knowledge for future remediation applications for ADF.

3.3.2 Method of Collection

A University of Colorado/AFIT doctoral student, Major Jeff Cornell and associates collected the DIA and Westover soils. The DIA sample was obtained from an uncontaminated site north of the C+ dam at Denver International. The soil was taken from below the root zone at a depth of approximately six inches. The soils have been analyzed in the University of Colorado's soils lab for content and consistency. A summary of the characteristics of the analysis can be found in Appendix A. The area of interest, near Bldg. 470, is one of the highest elevations on WPAFB, in a grassy open area. The sample was gathered by clearing the top 15-20 centimeters of material and collecting the next 20-30 centimeters of soil in the 0.5 square meter area. A clean metal

shovel was used to free the soil and a plastic pool filter with 6-mm openings was used to sieve out the larger organic material, rocks, and general debris. Handling and processing of the samples was kept as consistent as possible to remove any process variability. Characteristics such as surface area of the collection site, depth of the site, and size distribution were not taken into consideration between sample sites.

3.3.3 Soil Types

The DIA sample is a high clay soil with medium range organic material (\cong .0137f_{om}). The Westover sample is a sandy soil with very little organic material (< LOD). These two soils have not been exposed to ADF in the past and have not become acclimated to it. The third soil is from WPAFB; it is high clay with a medium amount of organic material (\cong .0287f_{om}). The two variations with the WPAFB soil come from the fact that the original soil is an undisturbed sample with no exposure to ADF. Refer to Table 3-1 below, for soil characteristics.

| Soil | <u>Classification</u> | Organic Content | <u>pH*</u> | Exposure to ADF |
|------------------|------------------------------|--------------------|------------|-----------------|
| | | | | |
| DIA, CO | Clay: MC=13.8 | 1.37±0.37% | 7.5 | NO |
| | | | | |
| Westover ARB, MA | Sandy: MC=6.3 | Below LOD | 8.0 | NO |
| | | assumed = 0.09% | | |
| | | | | |
| WPAFB, OH | Sandy Loam: | 2.87±0.15% | 7.8 | NO/YES** |
| | | | | |
| | MC=10.5% | | | |
| | | | | |

TABLE 3-1. Soil Characteristics

MC is moisture content and is given in percent

*NOTE: pH measurements were made using a Hach pH meter

**NOTE: The soil was part of a parallel study

The WPAFB soil was also used in a parallel experiment of respirometry with exposure to varied concentrations of PG and MeBT. The soil used in this parallel experiment was evaluated and used as a soil variation for comparison in my study.

3.3.4 Analyzed Characteristics

Analyses were performed on each of the soil samples. It is important to understand the differences between samples for comparative reasons. A summary of the soil characteristics provided by the laboratory employed has been supplied in Appendix A. In cooperation with the University of Colorado at Boulder, the attributes of interest have been narrowed down to the aforementioned characteristics. Differences in these characteristics can be identified in the responses of the 4 and 5 isomers of MeBT.

3.4 Experimental Preparation

The following sections are detailed descriptions as to how each experiment was prepared and the reasoning behind the preparation. All samples and subsets were prepared as at least triplicates for statistical consistency. It was also realized, during preliminary experimentation, that the solubility of MeBT in water was enhanced by the presence of PG. This detail was utilized while creating one solution for the study.

3.4.1 Soil Preparation

Review of related literature (Ball and Roberts, 1991) shows that a pulverized soil samples provide the maximum amount of surface area for sorption to take place. However, aside from initial sieving through a 6mm screen, the samples did not receive additional treatment. The

equilibration time for the soils used in this study were sufficient for the reaction to occur. It was also suggested that all samples be filtered prior to HPLC injection, to help avoid clogging of the column. As shown in Appendix B, HPLC results for the filtered samples did not differ significantly from the unfiltered samples. Therefore, unfiltered, centrifuged samples were used throughout the study.

It was important to use a consistent amount of soil and solution for each sample. This proportion was such that approximately half of the solvent is sorbed to the soil. If too much or too little is sorbed then accuracy will deteriorate [Heyse, 1998].

The amount of soil used was determined with the formula $M_s = V/K_d$, where M_s is the mass of the soil, V is the volume of solution and K_d is the sorption coefficient. I have estimated K_d for each soil type from the K_{ow} provided by the manufacturer on the MSDS and the relationship of K_d to K_{oc} via Karickhoff (see Section 2.6) [Fetter, 133; Karickhoff, 1984; Lyman and Reehl, 321]. The relationship is assumed to be similar to that of benzene for three reasons. One, it is the best available data on the chemical MeBT at the time of the study. Two, this benzene group studied by Karickhoff is the closest available equation. I chose the equation based on the log K_{ow} of benzene being very close to that of MeBT (2.17 vs. 2.13 respectively). Lastly, the range of K_{oc} (input parameter) values covered by this equation are similar. Thus, the equation log K_{oc} = log K_{ow} + b was used. Generally b $\approx 0.2 - 0.35$ [Karickhoff, e.1979-1982]. However, in this study the equation log $K_{oc} = 0.52 \log K_{ow} + 0.62$ was used. The conversion to K_d was accomplished by the equation $K_d = f_{oc}*K_{oc}$, where f_{oc} is the percent of organic material in the soil [Lyman and Reehl, 321]. The f_{oc} number is calculated from a f_{om} number (fraction of organic material) provide by chemical analysis $f_{oc} = 0.58*f_{om}$ [Schwarzenbach, 33]. In general, more soil was used when the organic content of the sample was lower. Specific calculations for each soil type can be found in Appendix B.

Each 40-mL screw-capped, amber glass container was rinsed with methanol and then rinsed thoroughly with distilled water. Each soil was air dried in an aluminum pan for approximately seven days. These soils were then placed in drying pans and heated to 103° for approximately 1.0 hour [Standard Methods, 2-54]. The water content for each soil type was determine by air-dried weight minus oven dried weight divided by oven dried weight. All work was accomplished using the soil's air-dried weight; however, the reports are reported on dry weight basis using the information provided in Appendix B. Therefore, in each 40-mL vial, approximately 10 grams of soil was added with a clean laboratory spoon, weighed out on an electronic scale. The appropriate amount of solution was added (see Section 3.4.2 Solute Preparation and Appendix B). Summaries of soil type and vial weights are included in Appendices E-H.

3.4.2 Solute Preparation

The range of solute concentrations for the soil study was established by the corresponding percentage of MeBT in ADF. The maximum MeBT concentration expected in the environment would be 0.5% – 0.6%. The experimental range was set at 0.01% - 0.1%, or 100ppm to 1000ppm. One producer of ADF, Octagon/Arco adds 0.05% – 0.06% MeBT to their mixture. This percentage translates into 500 mg MeBT/L deionized water obtained from dissolving laboratory grade COBRATEC TT-100[™] in deionized water. Based upon that concentration, the

concentration range of interest in this study was established. A calibration curve consisting of 10 mg/L, 100 mg/L, and 1000 mg/L (two orders of magnitude) standards were used. The limits of detection (LOD), upper and lower, of the HPLC were shown to be outside the concentration range used. Therefore, this range is effective. MeBT stock solution was added to the soil sample vials in a proportion to create slurry within the vial. Each soil had a different amount of solution added based upon the calculation explained in Section 3.4.1. The individual soil type solution calculations can be found in Appendix B.

3.4.3 Conditional Control for Analysis

All materials were stored under the same conditions unless otherwise indicated. The room temperature was regulated by air-conditioning, at approximately 19° C ($67^{\circ}F$). The sample pH was not adjusted, but was monitored with a Hach unit and stayed below 8.0. The subset samples, once prepared and treated with the MeBT solute, were placed on a rotator table for complete mixing. The samples were rotated at 20 – 25 rpm until equilibrium was reached (*e.g.* statistically, no more MeBT was sorbed to the soil). This equilibrium was established experimentally by sacrificing single vials at pre-determined time intervals. Results of equilibrium tests are in Section 4.3 and Appendix B.

Concentrations of MeBT in the solute extract were compared to the known original concentration. To account for any fraction of MeBT that might have sorbed onto sorbent without being transformed, the vial was centrifuged and the water in the CMBR was sampled along with successive methanol washes of the soil. Subsequent methanol extraction and analysis should

reveal the amount of MeBT lost (if any) from the bulk fluid. An isotherm was created from the results and plotted on the graph in Section 4.3 and in Appendix B.

3.4.4 Statistical Design

Since analytical chemistry is the science of making quantitative measurements, it is important that raw data be manipulated and reported correctly to give an accurate estimate of the certainty of the results [Tissue, 1996]. Three statistical evaluations were performed on the sorption coefficient results. Each soil was considered independent of the other soils tested. A balanced design, one where each treatment has the same number of trials compared with all other treatments, was achieved for this study.

The first evaluation was performed to show interaction between the soil and concentration results. A four by four matrix was created to include each of these variables. If significant interaction is identified (p < 0.05), the variables involved depend on each other [Ott, 906; Devore, 435]. Results of this evaluation are discussed in Section 4.3 and shown graphically in Appendix J.

Another evaluation consisted of an Analysis of Variance (Two-way ANOVA) and Pairwise Tukey analysis to look for homogeneic grouping among each independent variable. This evaluation would indicate which, if any, of the soils or concentrations acted alike in terms of their sorption coefficients. Results of these evaluations are discussed in Section 4.3 and shown in Appendix J. Lastly, a t-test or a pairwise comparison of means was performed by Statistix[®] to compare high organic content to low organic content soil, clayey soil to sandy soil, ADF on an unacclimated soil, and a soil dosed with an MeBT/PG mixture to one dosed with only MeBT. Results are discussed in throughout Chapter 4 and the tables are presented in Appendix J that also summarizes the findings. The two-sample t-test was administered with a 0.05 level of significance and populations were assumed to be normal with equal variances. The null and alternate hypotheses for each of the tests were:

 $H_o: \mu_a - \mu_b = 0$, the means are equal

H_a: $\mu_a - \mu_b \neq 0$, the means are not equal

The mean and standard deviation for each of the variables was determined in Statistix[®]. A pooled estimator, s_p^2 , was calculated, which is the estimate of the common population variance.

$$s_p^2 = (n-1)*Std Dev_a^2 + (m-1)*Std Dev_b^2$$

n + m - 2

The symbols n and m represent the sample sizes of the treatments (a and b) and Std Dev. stands for the standard deviation of the respective treatments a and b. The standard error was determined by taking the square root of the pooled estimator to yield s_p. The t statistic was then calculated from the following equation:

$$t = \frac{\overline{X}_{a} - \overline{X}_{b}}{s_{p}*(1/n + 1/m)}$$

This t value is compared to the t-critical value with the relationship $-t_{\alpha/2,n+m-2} \ge t$ where the -tcritical value is referenced from a pre-determined table [Ott, 260; Devour, 358 and 707]. Results were simply to accept or reject the null hypothesis depending upon the relationship of the t value to the t critical values. If the t value is $T \le -t$ critical or $T \ge t$ critical, then we reject the null in favor of the alternate hypothesis. If $T \ge -t$ critical or $T \le t$ critical, then we accept the null hypothesis. The full T-test is supplied in Appendix J for each set of variables.

3.5 High Performance Liquid Chromatography

Liquid chromatography (LC) relies on differences in partitioning behavior between a viscous mobile phase and a stationary phase to separate components in a solution. High-pressure pumps are used to increase efficiency of the separation. A typical schematic representation follows:



FIGURE 3-1. HPLC Schematic [Tissue].

3.5.1 Theory

Ions or molecules that are dissolved in a solvent are separated in the process. Simple liquid chromatographs consist of a column with fitted ends that holds a stationary phase. Most are made of plastic or glass that range from a few centimeters to several meters in length. HPLC columns are stainless steel tubes, typically 30-50 cm in length and 5-6 mm in diameter.



FIGURE 3-2. HPLC Column [Tissue].

The stationary phase is bonded to inert particles of 3-10 μ m diameter. A slug of the compound is injected into the column and analytes are separated as they travel through the columns due to the difference in their partitioning between the mobile or liquid phase and the stationary phase of the column. Reverse-phase partition chromatography uses a relative nonpolar stationary phase and a polar mobile phase, such as methanol or methanol/water mixture. This is the most common form due to the wide range of analytes that can be dissolved in the mobile phase. In adsorption chromatography the stationary particles are made of silica, alumina or organic material. These are used in separating isomers, which can have different sorption characteristics due to steric effects in the molecule. The retention of an analyte by a column is described by the capacity factor, k', where: $k' = t_{r2} - t_{r1}/t_m$ where t_r is the time for the analyte to pass through the column and t_m is the time for mobile phase

to pass through the column. In Figure 3-4, a typical separation of analytes is represented.



Figure 3-3. Retention Time Graph.

Usually, a relatively narrow sample band or "plug" is injected (10 - 25 μ l injection volume) into the column. During the run, the sample will be spread due to the noneven flows around and

inside the porous stationary particles, slow adsorption kinetics, longitudinal diffusion, and other factors. These processes together produce so called band broadening of the chromatographic zone. In general, the longer the component is retained on the column, the more broad its zone, or its peak on the chromatogram. Separation performance depends on both component retention and band broadening. Band broadening is a kinetic parameter that is dependent on the adsorbent particle size, porosity, pore size, column size, shape, and packing performance. An exact knowledge of the adsorbent surface area is very important, since retention is proportional to the surface area of the adsorbent.

3.5.2 Purpose

MeBT and sample solution were separated using isomeric-elution high-pressure liquid chromatography on a two successive ZORBAX RX-C8 columns and analyzed with an UV absorbence detector at $\lambda = 280 \mp 2$ nm. MeBT isomers and degradation byproducts of MeBT were separated by gradient elution HPLC on the ZORBAX RX-C8 column and detected similarly.

3.5.3 Procedure

MeBT was analyzed via direct aqueous injection into a Hewlett Packard 2170 high performance liquid chromatograph fitted with an ultra violet array detector. MeBT separation was achieved isocratically using two ZORBAX RX-C8 reverse phase, 250 x 4.6 mm columns at 35°C and a mobile phase comprised of two solvents: (1) 0.5-mL phosphoric acid (H₃PO₄) and 0.65-g potassium dihydrogen phosphate (KH₂PO₄) combined in water (1 L); along with (2) HPLC grade acetonitrile (ACN) (430-mL) and methanol. The solvent ratio for the HPLC pre-mobile phase

began at 90:10 water:methanol and moved to 30:70 water:methanol for 0.5 hours prior to analysis. Once analysis began the ratio of eluent to methanol started at 100:0 eluent:methanol and gradually moved to 50:50 eluent:methanol at the 10-minute mark. At this point the ratio was diminished to 10:90 to flush any organic material that may have accumulated. PMC Specialties Group, Cincinnati, OH, and advice from the University of Colorado have modified this technique from an established methodology [Gruden]. The peaks for each isomer of MeBT were shown to come through at approximately 7-8 minutes with a 0.25 to 0.3-minute delay between the isomers as measured with HP Chem-Station[®] for LC software. All of the samples were injected at a volume of 10 μL with a HP injection autosampler [HP ChemStation[®]].

3.6.4 Calibration Curve

Calibration standards were prepared using a background solution of deionized water in a prepared volumetric flask (0 mg/L) and solutions of 1mg/L, 5 mg/L, 10 mg/L, 50 mg/L, 100 mg/L, 500 mg/L, and 1000 mg/L of COBRATEC TT-100[™] (Tolyltriazole) in deionized water. The 1000 mg/L solution was mixed in a larger quantity (approximately 2 liters) to be diluted



Figure 3-4. Isomer Calibration Curve.

for use in the subsequent calibration solutions for the study. Past experiments with the HPLC unit have shown the upper limit of detection (LOD) to be approximately 1200-1500 mg/L for this compound. The lower LOD was established with my mixture to be approximately 5 mg/L. The calibration curve shown in Figure 3-4 passes through the origin and is linear. Results from the calibration curve can be found in Appendix D. Note that all concentration values in this study have come as a direct result of interpolation from the calibration curve. Concentrations are reported relative to the area under the HPLC trace (example found on Figure 3-7). For a given applied concentration the resulting concentrations for each isomer will not be additive to the initial concentration analyzed. For this study, the 4-MeBT isomer is relatively stable and is not assumed to degrade; therefore, the 4 isomer can be used as a normalizing factor for comparison of the two isomers.

3.6 Data Collection

In this study, all analyses were initiated on prepared soils under controlled condition. Laboratory work must be reproducible and limit the statistical error associated with the results. By controlling the variables, like temperature, and by using replicates, the data presented is statistically sound.

The diagram (Figure 3-6) shows the sample preparation process. Equilibration was enhanced through constant mixing on a rotation table. Before sampling for analysis in the HPLC, each vial was centrifuged for approximately 20 minutes. Upon completion, the solution at the top of the vial was sampled without disturbing the soil beneath (see vial "D" in the diagram).



Figure 3-5. Methodology Format.

3.6.1 HPLC

The data generated from the HPLC comes in several forms; I have chosen to use the area under a curve for my study because it allows easy correlation between the concentration and HPLC results. This working curve or plot of the analytical signal is a function of the analyte amount concentrated due to the interactions on the adsorbent surface on the equilibrium concentration of analyte in the solute. Using absolute calibration, this concentration was then back calculated, by integration, from the calibration curve generated earlier. Typical HPLC results are produced in the form showed in Figure 3-6.

HP ChemStation software integrates the area under the peak or curve to get the results shown in Appendices E-G. Note that in Figure 3-6, the separation of the two isomer peaks (4-MeBT at

7.19 and 5-MeBT at 7.38 minutes, respectively) and the peak of a possible derivative of MeBT at7.92 minutes in this diagram.



FIGURE 3-6. Typical HP ChemStation Data Result for MeBT [HP ChemStation].

Tables have been generated from all of the HPLC collected data in spreadsheet form. They can be found in Appendices E-H.

3.6.2 Quality Assurance and Quality Control (Reproducibility)

Data was generated using triplicate runs to evaluate the HPLC's performance and five replicate samples (5 bottles from the same stock) to insure that the samples were consistent within the concentration level used. Numbers are reported to at least three significant figures. Sample results were recorded and a standard deviation was taken among the five replicates to demonstrate the accuracy of the measurements. Concentrations were found by first taking the average of the individual concentration averages; then, integrating over the linear calibration curve to obtain a concentration level.

IV. Data Analysis

4.1 Overview

There were several techniques used to analyze the data generated from this study. Graphical comparisons from Excel spreadsheets were used as well as descriptive and analytical statistics from Statistix[®], a statistical software package. In order to be consistent and simplify the comparison between the different techniques, a numbering system was established for each of the data sets. The soils and concentrations from this study will be referred to in this section as the following:

| | ~ |
|--|---------|
| Wright-Patterson AFB – Untreated soil | Soil 1 |
| Wright-Patterson AFB – Treated soil | Soil 2 |
| Denver International Airport soil | Soil 3 |
| Westover ARB soil | Soil 4 |
| 1000 mg/L MeBT solution | Conc. 1 |
| 100 mg/L MeBT solution | Conc. 2 |
| 10 mg/L MeBT solution | Conc. 3 |
| Mix – approximately 100 mg/L MeBT and 1000 mg/L PG | Conc. 4 |
| | |

TABLE 4-1. Soil and Concentration Identification

Analyses of each soil's results are reviewed prior to the differentiation analysis. This allows a thorough look at each soil before interactive influences can affect judgment. An evaluation criterion for each soil type includes K_d results from each concentration level, how the addition of PG affected the sorption coefficient, and any anomalies encountered during the study. Each soil type and concentration was assumed to be independent of the others for this analysis. An Analysis of Variance and Tukey pairwise comparison of means tests were performed on each soil for both isomers of MeBT. Results from these analyses can be found in Appendix J under 4 MeBT and 5-MeBT respectively.

Differentiation analysis was performed by using four tests. These tests included:

- 1) Clayey Soil vs. Sandy Soil
- 2) Medium foc Content Soil vs. Low foc Content Soil
- 3) Acclimated Soil vs. Unacclimated Soil (on the basis of exposure to MeBT and PG)

4) Concentration Solution with PG Added vs. Solution with No PG Added The results will be presented in graphical form, based on rejecting or accepting the null hypothesis given in Section 3.4.3, and in tabular form showing a direct comparison of the sorption coefficient values. Complete test results are included separately for each isomer in Appendix J.

4.2 Results from Preliminary Experiments

4.2.1 Pan Evaporation

Results from the pan evaporation experiment are consistent with the published data that PG is a hygroscopic compound. As the humidity in the room changed with the relative humidity from the outside, the measured amount of liquid in the pan increased or decreased accordingly. The liquid gained, on average 0.386 grams of atmospheric moisture over the course of the test period. Results from the experiment can be seen graphically in Appendix C.

4.2.2 Toxicity Matrix

A spectrophotometer, used at the completion of the test, measured the amount of growth in each vial. The amount of flock or total suspended solids (TSS) in the vial is a measure of growth; the lower the light penetration through the vial, the more flock in the vial, which was an indication that more growth had taken place. Therefore, the conclusion more TSS, more growth, and less

toxicity was assumed true. Results from this toxicity experiment showed the maximum amount of growth or measured flock was realized in the 0 mg/L MeBT and 0-mg/L PG vial. Growth in a vial with only nutrient and microorganisms raises the question of what carbon source was used to produce this development. This is a question for later study.

The next several largest amounts of growth came from 100 mg/L PG row where the growth of the microbes decreased (less TSS) as the concentration of MeBT increased. Again, matching the hypothesized details of previous MeBT toxicological work [Cornell *et al.*]. This resulting matrix can be seen in Appendix C.

4.2.3 Soil Characterization

These tests consisted of a field capacity test for the WPAFB soil, a soil moisture test for each of the soils, and a sample size characterization based on published log K_{ow} value. This section includes the isotherm equilibration data for sorption and extraction. The equilibration determinations were performed on the highest f_{oc} soil and a moderate zeolite content. This soil will have the most sites available for sorption; thus, it will take the longest time to reach equilibrium (i.e. worst case).

4.3 Statistical Quantitative/Qualitative Analysis

Standards in this study were solutions containing known concentrations of analyte, MeBT. These samples provided a reference to determine unknown concentrations or to calibrate the HPLC and were used to inoculate the soils in subsequent tests. The accuracy of the measurement is how close the result comes to the true value. Determining the accuracy of a

measurement requires calibration standards. This was accomplished using several concentrations to develop a calibration or working curve. Seven known concentrations were run through the HPLC to develop the calibration seen in Figure 3-4. Results of the calibration show that linearity among concentrations from 5 mg/L to 1000 mg/L MeBT is a true assessment. The Limit of Detection (LOD) is calculated to be *approximately* 5 mg/L \pm 4 mg/L. Notwithstanding, measurements of lower concentrations were unreliable and for that fact, the 10-mg/L results have been eliminated from all further calculations. Calibration Curve data is provided in Appendix D along with LOD calculations.

Analysis of this data was performed on bottles taken from each soil type at each concentration over the equilibration period established in Appendix B. This sorption equilibrium test was accomplished as per Section 3.4.2; results are also in Figure 4-1. Statistical analyses



FIGURE 4-1. Establishment of Sorption Equilibrium Point

were performed on the bottles, 1 hour to 216 hours, to see if the K_d values of the bottles were significantly different. It was determined that equilibrium is reached in 144 hours or 7 days.

Interaction between soil types was established using Statistix[®]'s Two-way Analysis of Variance (ANOVA) table with soil and concentration being the variables in question. Results of the ANOVA reveal a p-value of 0.000. A p-value less than 0.05 is an indication of significant interaction between the variables. This interaction leads to the fact that there are similarities in each soil and how they reacted to the various concentrations. Results are provided in Appendix J for each isomer. Statistical reports were generated taking into account all of the resultant sorption coefficients.

4.4 Isomer Results

Both isomers of MeBT were evaluated over a range of concentrations and several soil variations. Table 4-2 provides a direct comparison of the concentration results for each soil at each



FIGURE 4-2. MeBT Isomer Recovery from All Soil Types Note: Error bars represent one Standard Deviation among concentration group.

concentration level of solution. Direct comparison of the isomers can be seen in Figure 4-2. For the 1000 mg/L, the relationship of 4-MeBT and 5-MeBT is fairly consistent. Except for the untreated soil at 1000 mg/L, the K_d of 5-MeBT was higher than 4-MeBT; therefore it will be the basis for comparison during each of the differentiation tests. This consistency may be due to the fact that 4-MeBT is considered recalcitrant while the 5-MeBT may be degradable, and therefore somewhat unrecoverable. This unrecoverability may appear to be excess sorption, when, in fact, it is not. Figure 4-3 shows the possible paths that can occur for the MeBT.



FIGURE 4-3. Accounting for Unrecoverable Isomer

| | Concentration | | | | |
|---|---------------|--------|-------|------|------|
| | Soil | Isomer | 1000 | 100 | Mix |
| 1 | Untreated | 4 | 734.7 | 69.0 | 68.7 |
| 2 | Untreated | 5 | 741.8 | 63.1 | 61.2 |
| 3 | Treated | 4 | 864.8 | 78.2 | 81.7 |
| 4 | Treated | 5 | 825.2 | 0.0 | 54.6 |
| 5 | Westover | 4 | 869.1 | 77.1 | 77.7 |
| 6 | Westover | 5 | 851.5 | 72.8 | 73.1 |
| 7 | DIA | 4 | 841.6 | 76.6 | 73.6 |
| 8 | DIA | 5 | 819.0 | 71.3 | 69.3 |

TABLE 4-2. HPLC Concentration/Soil/Isomer Results

Note 1: Bold numbers in Table 4-2 are addressed in the corresponding soil's anomaly section.

Note 2: Isomer concentration totals do not match applied concentration, see Section 3.6.4 for explanation.

4.5 Wright-Patterson Soil Results

Both of the WPAFB test soils came from the same sample set. The unacclimated or untreated soil had not been exposed to ADF in any way. The treated soil was exposed to MeBT and PG for a two-week period during testing by respirometry [Burke, 1999].

4.5.1 Wright-Patterson Soil - Untreated

This study found that for Soil 1, mean K_d results for the 4-MeBT isomer fell into two homogeneous groupings. The group of Conc. 2 and 4, and Conc. 1 were significantly different from one another. Conc. 1 had a mean K_d significantly lower than the other group.

The 5-MeBT isomer produced only three homogeneous groupings. The mean K_d of Conc. 2 was significantly higher than the other group. Conc. 1 and 4 comprise the lower mean K_d value homogeneous group.



FIGURE 4-4. WPAFB Untreated Sorption Coefficient vs. Concentration – 4-MeBT

Therefore, it is assumed that Soil 1 reacts relatively the same to the various concentrations applied for both isomers. Figure 4-4 and Figure 4-5 depicts the linear regression for the

concentration dependence of Kd for each isomer. Linearity is shown by the higher n value for the Freundlich isotherm: 0.859 for the 4 isomer and 0.732 for the 5 isomer. Details are in Appendix E.



FIGURE 4-5. WPAFB Untreated Sorption Coefficient vs. Concentration – 5-MeBT

The ability to separate the 4 and 5 isomer of MeBT allows for a comparison of their means at each of the concentration levels for each soil. The 1000-mg/L concentration level for the 4 and 5 isomers show similar sorption coefficients, 0.860 vs. 0.825-mL/g, respectively. At the 100-mg/L concentration level for the 4 and 5 isomers show significantly different sorption coefficients, 1.203 vs. 330-mL/g. Finally at the 100-mg/L Mix concentration, the means of the 4 and 5 isomer are similar in direction but not in magnitude (1.219 vs. 1.744-mL/g, respectively). A summary of these results is in Table 4-3.

Specifically looking at Conc.2 and Conc. 4 for Soil 1, the study reveals that there is no significant difference between the two concentrations for the 4 isomer (1.203 vs. 1.219-mL/g). This is also evident by the homogeneous grouping of the two means. The same results were not achieved for the 5-MeBT isomer. This relationship suggests that there is no difference how Soil 1 sorbs 100-mg/L of MeBT with or without the addition of PG for the 4 isomer, but there is a

difference in the 5 isomer. Results are shown in Appendix J, also refer to Table 4-3 for a summary of means sorption results and Figure 4-6 for a graphical display.

Two anomalies exist for Soil 1. First, the *high* sorption coefficients for the 10-mg/L samples, both 4 and 5-MeBT, are inconsistent with the sorption pattern established in this study. At the low end concentration level, the nonlinear portion of the calibration curve could have adversely affected the results. Secondly, the 5 isomer, in 3 of the 4 concentrations, has a higher sorption coefficient than the 4 isomer.

| Untreated | Kd-4 (mL/g) | Kd-5 (mL/g) |
|-----------|-------------|-------------|
| Conc. 1 | 0.860 | 0.825 |
| Conc. 2 | 1.203 | 1.645 |
| Conc. 4 | 1.219 | 1.735 |

TABLE 4-3. WPAFB Untreated Average Sorption Coefficient Results.



FIGURE 4-6. WPAFB Untreated Soil Sorption Results.

4.5.2 Wright-Patterson Soil – Treated

The sorption coefficient results for the 4 isomer in Soil 2 shows three homogeneous grouping of means. Conc. 2 was the highest at 0.706-mL/g, next was Conc. 4 at 0.521-mL/g, followed by Conc. 1 at 0.350-mL/g. With Soil 2 the 5-MeBT isomer produced a separate group for each concentration indicating that each concentration is sorbed differently. The means range is from

Conc. 2 at 44.5-mL/g to Conc. 1 at 0.512-mL/g. Results are presented in Table 4-4 and Figure 4-9. Figure 4-7 and Figure 4-8 depicts a linear regression for the concentration dependence of K_d , each group of coefficients is graphed on these figures. Results show a good linear fit for the 4-MeBT isomer with n = 0.708, but a poor linear relationship for the 5-MeBT isomer with n = 0.125.

Focusing attention to Conc. 2 and Conc. 4 under the 4 isomer, we find that the means are not statistically similar since they are put into two homogeneously different groups. Refer to Test 4, Section 4.11 for specific results of this comparison. For the 5 isomer, the same phenomenon exists. Refer to Table 4-4 and Figure 4-9, respectively, for a summary of mean sorption data.



FIGURE 4-7. WPAFB Treated Sorption Coefficient vs. Concentration – 4-MeBT



FIGURE 4-8. WPAFB Treated Sorption Coefficient vs. Concentration - 5-MeBT

Again, the anomaly discovered in Soil 2 was the relatively *high* sorption coefficient of the 10mg/L concentration solution for the 5 isomer (mean K_d value of 3.225-mL/g). In addition, the 5 isomer for the Conc. 2 and 3 does not show significant sorption after equilibrium is reached. This could be from the unrecoverable condition described in Section 4.4. Refer to Appendix F for HPLC results and note the appearance of an additional peak with higher retention time when the 5 isomer is unrecoverable. In this case too, each 5 isomer had a higher sorption coefficient than the 4 isomer.

| Treated | Kd-4 (mL/g) | Kd-5 (mL/g) |
|---------|-------------|-------------|
| Conc. 1 | 0.350 | 0.512 |
| Conc. 2 | 0.706 | 44.50 |
| Conc. 4 | 0.547 | 2.325 |

TABLE 4-4. WPAFB Treated Average Sorption Coefficient Results.



FIGURE 4-9. WPAFB Treated Soil Sorption Results.

4.6 Denver International Airport (DIA) Soil Results

The sorption coefficients for the 4-MeBT isomer on Soil 3 show three homogeneous groups. Conc. 3 is has the highest K_d at 2.840-mL/g, followed by Conc. 4 at 0.208-mL/g, with the lowest K_d at Conc. 1 equal to 0.044 mL/g. The 5 isomer had two homogeneous groups. Conc. 4 and 2 have the highest sorption coefficient at 0.291 and 0.255-mL/g respectively, followed by the lowest mean K_d value of Conc. 1 at 0.075-mL/g. Results are contained in Table 4-5 and Figure 4-12.



FIGURE 4-10. DIA Sorption Coefficient vs. Concentration - 4-MeBT

Figure 4-10 and Figure 4-11 depicts a linear regression for the concentration dependence of K_d , each group of coefficients is graphed on these figures. Results show an excellent linear fit.



FIGURE 4-11. DIA Sorption Coefficient vs. Concentration - 5-MeBT

The Conc. 2 and 4 result show the relationship between the addition of PG and when no PG added. With Conc. 2 and Conc. 4 being significantly different for 4-MeBT, there is a difference between PG added and not adding PG to the solution while for the 5-MeBT there is also a difference when PG is added. See Table 4-5 for a comparison.

Soil 3 with 5-MeBT has been placed into two homogeneous groups. Conc. 1, 2, and 4 are statistically the same with values of 0.0754, 0.2550, and 0.2906-mL/g respectively. Conc. 3 is higher at a K_d value of 5.0206-mL/g. The results (Table 4-5 and Figure 4-12) show that the addition of PG to the solution had little affect on the sorption coefficient of 5-MeBT.

| DIA | Kd-4 (mL/g) | Kd-5 (mL/g) |
|---------|-------------|-------------|
| Conc. 1 | 0.044 | 0.075 |
| Conc. 2 | 0.160 | 0.255 |
| Conc. 4 | 0.208 | 0.291 |
| | DIA C 1 A | <u> </u> |

Table 4-5. DIA Soil Average Sorption Coefficient Results.



Figure 4-12. DIA Soil Sorption Results.

4.7 Westover ARB Soil Results

The Tukey Comparison of Means has separated both the 4 and the 5 isomers into three homogeneous groups. The 4 isomer ranges from 0.148-mL/g to a low of 0.036-mL/g, with the grouping of Conc. 4 and Conc. 2 in the middle with values of 0.197 and 0.146-mL/g, respectively. The 5-isomer grouping has a high K_d value of 2.970 mL/g and a low of 0.056 mL/g, with the same grouping the middle. Refer to Table 4-6 for a summary of average sorption coefficient data. Figure 4-15 contains a graphical representation of the sorption data.



FIGURE 4-13. Westover ARB Sorption Coefficient vs. Concentration - 4-MeBT



FIGURE 4-14. Westover ARB Sorption Coefficient vs. Concentration - 5-MeBT

The comparison of Conc. 2 (No PG added) and Conc. 4 (PG added) show the same result for both the 4 and the 5 isomer. Both isomers have mean K_d values for Conc. 4 and Conc. 2 that are statistically equal. Details for this are shown in Section 4.11 and are presented in tabular form in Table 4-6 and graphically in Figure 4-15.

Again, the K_d value of Conc. 3 is significantly higher, for both the 4 and the 5 isomer, than the other concentrations in this soil type. Additionally, the 5 isomer is higher in sorption coefficient value than the 4 isomer, except in Conc. 3 and 4 where they are equal. Moreover, the low value of the sorption coefficient for Conc. 1 is below 0.100-mL/g. This and Soil 3's Conc. 1 are the

only ones to drop below 0.100-mL/g. In addition, the 5 isomer for the Conc. 3 does not show a significant concentration level after equilibrium is reached. This could be from the unrecoverable condition described in Section 4.4. Refer to Appendix H for HPLC results and note the appearance of an additional peak with higher retention time when the 5 isomer is unrecoverable.

| Westover | Kd-4 (mL/g) | Kd-5 (mL/g) |
|----------|-------------|-------------|
| 1000 | 0.036 | 0.051 |
| 100 | 0.148 | 0.203 |
| Mix | 0.136 | 0.193 |

TABLE 4-6. Westover ARB Soil Average Sorption Coefficient Results.



FIGURE 4-15. Westover ARB Sorption Results.

4.8 Soil Type Differentiation

The first statistical comparison was conducted to determine if there was a significant difference between the mean sorption coefficients for a sandy soil (Soil 4) and the mean K_d values of a clayey soil (Soil 3). This comparison was accomplished using a two-sample t-test as described in Section 3.4.3. As shown in Figure 4-16 (4-MeBT) and Figure 4-17 (5-MeBT), the t value obtained, indicated by the stem line and circle, fell between the t critical values ($t_{crit} = t_{.025}$ and $t_{.975}$) for the 4 isomer and outside for the 5 isomer. This indicates that the null hypothesis proposed in Section 3.4.3 should be accepted for the 4 isomer; therefore, the mean sorption coefficients for Soil 3 and Soil 4 are to be considered equal. The null is rejected for the 5 isomer and indicates that the means of the soils are not equal and the sorption is decreased with the sandy soil vs. clayey. Table 4-7 and Figure 4-18 summarizes the average sorption coefficient results for both soils.



FIGURE 4-16. Test 1 t-test Results for 4-MeBT – Soils 3 & 4.



FIGURE 4-17. Test 1 t-test Results for 5-MeBT – Soils 3 & 4.

| Soil Type Differentiation | | |
|------------------------------|-------------|-------------|
| | Kd-4 (mL/g) | Kd-5 (mL/g) |
| Soil 3 | 0.160 | 0.075 |
| Soil 4 | 0.148 | 0.051 |

TABLE 4-7. Average Sorption Coefficients for Soil Type Differentiation.


FIGURE 4-18. Average Sorption Coefficients for Soil Type Differentiation.

4.9 Organic Content Differentiation

The second statistical comparison was conducted to determine if there was a significant difference between a medium organic content soil (Soil 1) and a soil with virtually no organic content (Soil 4) with regard to MeBT sorption. A two-sample t-test was used to calculate a t value for this comparison. Figure 4-19 and Figure 4-20 depicts the results of the T-test for both the 4 and 5 isomers. The null hypothesis in Section 3.4.3 will be rejected for both isomers since the t value does not fall between the t critical values. Organic content clearly increased sorption of MeBT. Reference Table 4-8 and Figure 4-21 for a direct comparison summary of the mean K_d values.



FIGURE 4-19. Test 2 t-test Results for 4-MeBT – Soils 1 & 4.



FIGURE 4-20. Test 2 t-test Results for 5-MeBT – Soils 1 & 4.

| Organic Differentiation | | | |
|-------------------------|-------------|-------------|--|
| | Kd-4 (mL/g) | Kd-5 (mL/g) | |
| Soil 1 | 0.860 | 0.825 | |
| Soil 4 | 0.036 | 0.051 | |

TABLE 4-8. Average Sorption Coefficients for Organic Content Differentiation.



FIGURE 4-21. Average Sorption Coefficients for Organic Content Differentiation.

4.10 Acclimation Differentiation

The third statistical comparison was conducted to determine if there was a significant difference between the mean sorption coefficients for a soil that was exposed to a solution of MeBT and PG (Soil 2) and a soil that was previously unexposed to these compounds (Soil 1). To make this comparison, a two-sample t-test was employed. The result of the statistical tests determined a t value for comparison. Given the null hypothesis in Section 3.4.3 and the t critical value established from the degrees of freedom, Figure 4-22 and Figure 4-23 were created. These graphs show the t value outside the t critical values; therefore, the null hypothesis is rejected and the means of the two soils are not equal. Table 4-9 is a summary of average sorption data and Figure 4-24 is a graphical representation of the same information.



FIGURE 4-22. Test 3 t-test Results for 4-MeBT – Soils 1 & 2.



FIGURE 4-23. Test 3 t-test Results for 5-MeBT – Soils 1 & 2.

| Soil Type Differentiation | | | | | | |
|------------------------------|-------------------------|-------|--|--|--|--|
| | Kd-4 (mL/g) Kd-5 (mL/g) | | | | | |
| Soil 1 | 0.860 | 0.825 | | | | |
| Soil 2 0.350 0.512 | | | | | | |

TABLE 4-9. Average Sorption Coefficients for Acclimation Differentiation.



FIGURE 4-24. Average Sorption Coefficients for Acclimation Differentiation.

4.11 Propylene Glycol Additive Soil Differentiation

The fourth statistical comparison was conducted to determine if there was a significant difference between the mean K_d value for soils dosed with a combination of MeBT and PG (Concentration 4 – approximately 100 mg/L MeBT and 1000 mg/L PG) and the same soils dosed with only a solution of MeBT only (Concentration 2 – 100 mg/L MeBT). A two-sample t-test was conducted. Again, referencing the hypothesis in Section 3.4.3 and using the degrees of freedom for this comparison, a t value was established. Figure 4-25 and 4-26 show the results of the t-test respectively for each isomer. The results show that the null hypothesis should be accepted for the 4-MeBT because the t value is between the t critical values and rejected for the 5-MeBT isomer t value placement outside the t-critical range. This account would suggest there is no difference in the mean sorption coefficient of a solution containing PG or one without PG for the 4-MeBT and a significant difference in sorption of the added PG concentration for the 5-MeBT. Additionally, Table 4-10 and Figure 4-27 provides a direct comparison summary of average sorption data for each soil. Notice the *trend* for each soil is almost identical.



FIGURE 4-25. Test 4 t-test Results for 4-MeBT – Conc. 2 & 4.



FIGURE 4-26. Test 4 t-test Results for 5-MeBT – Conc. 2 & 4.

| Soil Ty | /pe Differen | itiation | | | | | | |
|---------|--------------|----------|----------|----------|----------|----------|----------|----------|
| | Soil 1-4 | Soil 1-5 | Soil 2-4 | Soil 2-5 | Soil 3-4 | Soil 3-5 | Soil 4-4 | Soil 4-5 |
| Conc. 2 | 1.203 | 330.000 | 0.433 | 44.500 | 0.160 | 0.255 | 0.148 | 0.203 |
| Conc. 4 | 1.219 | 1.735 | 0.377 | 2.325 | 0.208 | 0.291 | 0.136 | 0.193 |

TABLE 4-10. Average Sorption Coefficients for PG Additive Soil Differentiation.



FIGURE 4-27. Average Sorption Coefficient for PG Additive Soil Differentiation.

4.12 Adjustments

This study was based on several assumptions that have large consequences, if incorrect. The conclusions in Chapter 5 are based upon these assumptions and the best information available at the time of study.

The first assumption was that the K_{ow} or octanol:water partitioning coefficient provided by the manufacturer is accurate. This study began by using the MSDS log K_{ow} value for preliminary calculations. If the log K_{ow} provided was incorrect, calculations based upon it will be off by that proportion. Examples include the volume of solute added to soil sample and the theoretical sorption coefficient. Note that Section 5.1 touches on the differences in K_{ow} 's between theoretical and achieved and a chart of experimental K_{ow} 's is included in Appendix I. A discrepancy can be found that may be attributed to the manufacturer or the difference in reagent grade and proprietary grade product.

The next assumption I have made is in regards to the sorption equilibrium isotherm results. The isotherms were developed over a 216-hour (9-day) period at which time it appeared that the HPLC resultant areas had leveled out or reached equilibrium. It is possible that this was a temporary stagnation point and sorption of MeBT in these soils may take weeks if not years. The time frame for this study made it impractical to lengthen the time scale for sorption isotherms.

The results of most 10-mg/L concentration runs are inconsistent with the trends established by other concentrations applied. This could have arisen from the fact that the calibration curve goes

4-22

non-linear at approximately 7 mg/L. I assume that the non-linear point comes at the limit of detection (5-mg/L) and does not affect the results at the 10-mg/L point. However, inconsistent values at the 10-mg/L level may support the non-linear assumption. For this reason the 10 mg/L data was virtually eliminated from consideration in this study.

The final assumption that I have made in this study is the use of benzene as a substitute for MeBT in the calculation of the initial K_{oc} and soil mass. MeBT has an aromatic ring as part of its structure. The log K_{ow} of benzene and MeBT are similar at 2.17 and 2.13 respectively. Taking into account the second of these assumptions, this information was the best available at the time of the study. Use of toluene was considered because of the similar structure; however, this formula calculated out so little solute added to the soil that a saturated condition was not attainable for this study. A slurry or saturated condition would not have been achieved within the vial.

V. Conclusions and Recommendations

5.1 Conclusion

Results of this study help define the transport of MeBT in subsurface soils. Although limited in scope, this study describes the likely behavior of MeBT in a water-saturated soil environment. This section will begin with the major findings and continue with conclusions from each of the four tests performed during this study and finally address all other findings that have arisen.

- In the range of concentrations examined in this study (100 to 1000 mg/L), MeBT had an effective range of sorption coefficients from 0.03 to 1.45-mL/g for the 4-MeBT and 0.04 to 3.24-mL/g for the 5-MeBT.
- The two isomers, 4-MeBT and 5-MeBT, showed different sorption characteristics-the 5 isomer was consistently higher.
- Soil previously exposed to PG and MeBT, in most cases, shows approximately 50% less sorption than the unexposed, clean soil.
- The application of an additional organic material, propylene glycol, along with MeBT made no difference in the sorption of MeBT on any soil with the 4 isomer.
- Samples exposed to 5-MeBT for at least two weeks showed reduced 5-MeBT amounts and an additional peak in HPLC traces. This suggests biodegradation may occur in the 5 isomer.

5.1.1 Statistical Evaluation of Laboratory Results

The first of the four statistical tests performed for this study was a comparison of clayey soil vs. sandy soil. Samples from Denver International Airport and Westover Air Reserve Base were used. This test concluded that resultant mean sorption coefficients were equal for the 4-MeBT;

therefore, with 95% confidence, it can be said that there is no difference in how these particular clayey and sandy soils sorb 4-MeBT isomer. However, with 95% confidence, there is a significant difference in how these soils sorb 5-MeBT.

Test 2 for this study involved the comparison of a medium organic content (f_{oc}) soil vs. a soil that had virtually no organic content. Soil samples from Wright-Patterson Air Force Base and Westover ARB was used. Test 2 concluded that at all concentration levels applied, difference sorption coefficients were achieved for both the 4 and 5 isomer of MeBT. Thus, with 95% confidence, it can be said that soils with higher organic contents sorb more MeBT than those soils with low f_{oc} .

Test 3 compared the difference between a soil that had not been exposed to ADF or MeBT prior to this study and a soil that had been acclimated to MeBT and PG through a respirometry study. This test concluded that there is a large difference in how each soil sorbed the MeBT. The soil that had no exposure prior to the study had the larger of the two sorption coefficients. This phenomenon was noted for both the 4 and 5 isomer of MeBT.

The final test data results from all soils types comparing the 100-mg/L MeBT solution and 100-mg/L MeBT with 1000-mg/L propylene glycol (Mix) solution indicate that there is seemingly no difference if PG is added to the mix or if it is not added to the mix for the 4-MeBT isomer. However, the 5-MeBT indicates a significant difference in the mean sorption coefficient for the two concentrations. Reviewing the 5-MeBT, it is possible that microorganisms in the acclimated soils (Soil 2 or Treated Soil) are able to degrade the 5-MeBT isomer, thus lowering the concentration.

5-2

5.1.2 Significant Conclusions from Statistical Results

The K_d values obtained for the majority of soil types and concentrations would tend to reinforce the conclusion that MeBT does not sorb well to the soils in this study. There are certain combinations of concentrations and soil types that lead to higher sorption coefficients. These are explained in Section 4.4. These results represent low concentrations near the limit of detection and the non-linear portion of the HPLC; results may have been skewed because of this situation. Further analyses at low concentrations are necessary to support my findings.

Using the K_d values obtained during analysis, I have back-calculated log K_{ow} values for the 4 and 5 MeBT isomers. I have concluded that the published log K_{ow} values appear to be too high. Results in Appendix I reveal that few K_{ow} value approach the published value of 2.17. However, the calculation used for the log K_{ow} values is based on assumptions and cannot be relied upon; this conclusion is only speculative.

One possible reason for this discrepancy could come be differences of laboratory grade MeBT and reagent grade MeBT (see Section 4.12). It was assumed that the product used throughout my study was pure tolyltriazole; however, the laboratory grade identifier suggests that impurities may exist which could cause the difference in K_{ow} .

Another speculative conclusion for the log K_{ow} difference is the reporting process. Log K_{ow} values in this study come from isometric calculations. The K_{ow} value that has been published comes from unaltered methyl-benzotriazole.

As alluded to in Section 4.4, the 4-MeBT has a higher sorption coefficient than 5-MeBT (except for the untreated soil at 1000 mg/L). An explanation for this situation is included in Section 4.4 and reinforces areas presented in the literature review section. I would conclude that the 4-MeBT would move rather freely through the subsurface with little to slow it down, while the 5-MeBT is influenced by other factors, rendering it unrecoverable at times.

Lastly, as shown in the literature review section, the sorption coefficient is inversely related to the particle size of the soil. MeBT is a contradiction to that claim by Ball [Ball and Roberts, 1991].

5.2 Follow-on Research

This research involves the compound of interest, MeBT. The ultimate future to this subject is to find out what to do with ADF in the environment. In order to get there, numerous questions must be answered. The transport result with sorption is just one of many phases. Several areas are open to further interpretation, not only with MeBT, but ADF in general.

5.2.1 Analysis of Other ADF Components

There are numerous additives in ADF that allow it to perform in cold weather. MeBT has been the first to be studied in any depth. Further research can be done using another component of ADF and its relationship to PG and MeBT. A good working relationship with a manufacturer of ADF would be helpful to understand exactly what goes into ADF and the proportions in which they are mixed.

5.2.2 Breakdown of MeBT in-situ

An attempt can be made to isolate microorganisms that utilize MeBT aerobically as a sole substrate for growth. If the isolation is successful, experiments will then be conducted to investigate the biodegradation rates of MeBT by the isolated organisms.

5.2.3 Environment Variation Study

It is important to understand how this compound reacts under all environmental situations. The latest studies look at this compound under very specific conditions. Further research is necessary to determine how MeBT reacts under environmental changes. Such changes include temperature (-10 C - 5 C) and pH (5.0 – 9.0 or greater).

5.2.4 HPLC identification of MeBT degradation products

In several of the HPLC results, there was a small, but significant spike beyond the retention time window of interest. In most cases it was accompanied by a substantial decrease in 5-MeBT concentration. It is not known what this spike is or how to isolate it; furthermore, it is not known if it is exactly related to the decrease in the 5-MeBT isomer. Further research is necessary to determine if there is a correlation between the decrease in the 5-MeBT isomer and the area of the new spike. Ensure abiotic conditions are achieved for this portion.

5.2.5 Repeat this study

An effort can be accomplished along the same concepts as this study; however, the major emphasis would be on the metal binding properties of MeBT instead of focusing on the organic content binding. The metal binding alternative could give better information as to the difference in the K_{ow} factors.

Appendix A

Soil Analysis Report

University of Colorado at Boulder

AFIT/ENV/Charles A Bleckmann 2950 P Street Wright-Patterson AFB OH 45440

· · · · / · · · ·

and the second second and the second s

DATE RECEIVED: 12-14-1998 DATE PARTIAL REPORTED: 01-22-1999 DATE REPORTED: 02-16-1999

8-Dec-98

RESEARCH SOIL ANALYSIS

Colorado State University Soil, Water and Plant Testing Laboratory Natural & Environmental Sciences Bldg - A319 Fort Collins, CO 80523

(970) 491-5061 FAX. 491-2930

BILLING:

| | Cu | 2.93 3.15 2.89 |
|-------------|----------------|----------------------------------|
| | Mn | 2.89 2.76 2.63 |
| a ct | Fe | 50.0 51.7 50.2 |
| OTPA Extra | Zn | 2.44 2.63 2.70 |
| AB-I | K | 91.7 92.0 99.2 |
| | A | 5.3 5.6 5.6 |
| | NO3-N | 3.3 6.2 6.2 , |
| % | MO | 2.7 2.9 3.0 |
| Lime | Estimate | Medium Medium Medium |
| te | EC mmhos/cm | 1.2 0.9 0.8 |
| 889 | Hd | 7.7 7.8 7.8 |
| Sample | * | Afit # 1 Afit # 2 Afit # 3 |
| Lab | # | R3393 R3394 R3394 |

| à | \$ 0 1 | 1.6] 1.82 1.78 |
|---|--------------|----------------------------------|
| | Texture | Loam Loam Loam |
| | Clay | 16 16 |
| ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | Silt | 36 36 35 |
| | Sand | 48 48 49 |
| Sample | ID# | Afit # 1 Afit # 2 Afit # 3 |
| Lab | # | R3392 R3393 R3394 |

1

1

Appendix B

Caculation of:

Solution Concentration Moisture Content Individual Sample Sizes Sorption Equilibrium Isotherm Extraction Isotherm Filtered Sample Comparison

Field Capacity Calculations

| Mass of Soil = | Ms = | 100.01 | grams |
|---|------|--------|-------|
| Mass of Water Absorbed in Soil Achieve 100% FC | Mw = | 19.1 | grams |
| | FC = | Mw/Ms | |
| | = | 0.191 | |
| | = | 19.1% | |

.

.

Dry Soil Weight Calulations

| WPAFB | | | | | |
|----------|--------------------------|--------|--------------|-----------|-------|
| | Air- Dried Sample Weight | 100.02 | 100.01 | 100.01 g | |
| | Weight after heat/decant | 90.32 | 91.05 | 90.27 g | |
| | | | | | AVG |
| | Water Content | 10.7% | 9.8% | 10.8% | 10.5% |
| DIA | | | | | |
| | Air- Dried Sample Weight | 100.01 | 100.01 | 100.002 g | |
| | Weight after heat/decant | 88.12 | 87.56 | 87.98 g | |
| | | | | | AVG |
| | Water Content | 13.5% | 14.2% | 13.7% | 13.8% |
| Westover | | | | | |
| | Air- Dried Sample Weight | 100.00 | 100.00 | 100.01 g | |
| | Weight after heat/decant | 94.3 | 93.82 | 94.1 g | |
| | - | | | _ | AVG |
| | Water Content | 6.0% | 6 .6% | 6.3% | 6.3% |
| | | | | | |

| Solute Concentration - MeBT | Product Used: Tolyltriazole (Solid) Cobratec TT-100 Laboratory Grade |
|-----------------------------|--|
| | 1000 mg/L = <u>1 gram MeBT</u> 1 Liter Deionized Water |
| Solute Concentration - PG | Product Used: Propylene Glycol (Aqueous) Mallinckrodt, OR 1925 Laboratory Grade - 1,2 Propanediol |
| | 1000 mg/L = <u>5 grams of Liquid PG</u> 5000 mL Deionized Water |

| 1000 n | ng/L | 4 Me | 4 MeBT | | 5 MeBT | | 4 MeBT | | 5 MeBT | |
|--------|----------------|-------------|---------|-------------|----------|-------------|---------|-------------|----------|--|
| | | Retention | | Retention | | Retention | | Retention | | |
| | Starting (Hrs) | Time (Avg.) | Area | Time (Avg.) | Area | Time (Avg.) | Area | Time (Avg.) | Area | |
| | 1 | 7.020 | 7495.6 | 7.208 | 11814.4 | 6.996 | 7340.7 | 7.179 | 11505.6 | |
| | 6 | 6.969 | 7198.6 | 7.157 | 11350.1 | 6.947 | 7177.1 | 7.128 | 11294.7 | |
| | 12 | 6.954 | 7022.4 | 7.138 | 11072.9 | 6.944 | 7099.3 | 7.125 | 11152.9 | |
| 1 day | 24 | 6.948 | 7011.2 | 7.116 | 11009.5 | 6.972 | 7074.6 | 7.155 | 11083.9 | |
| 2 days | 48 | 7.019 | 7004.7 | 7.207 | 10990.5 | 6.994 | 6852.0 | 7.179 | 10740.5 | |
| 3 days | 72 | 7.063 | 7076.4 | 7.252 | 11101.3 | 6.995 | 6923.9 | 7.184 | 10836.3 | |
| 4 days | 96 | 7.021 | 6646.2 | 7.210 | 10411.4 | 7.001 | 6629.4 | 7.198 | 10396.4 | |
| 5 days | 120 | 7.087 | 6862.6 | 7.276 | 10747.1 | 7.045 | 6728.9 | 7.234 | 10509.7 | |
| 6 days | 144 | 7.065 | 6542.3 | 7.253 | T10403.8 | 7.028 | 6489.1 | 7.210 | 10374.5 | |
| 7 days | 168 | 7.036 | 6103.7 | 7.222 | 10009.1 | 6.992 | 6238.2 | 7.179 | 10126.8 | |
| 8 days | 192 | 7.042 | 6089.4 | 7.238 | 9996.6 | 6.999 | 6197.3 | 7.188 | 10001.7 | |
| 9 days | 216 | 7.027 | 6065.5 | 7.211 | 9967.0 | 6.983 | 6119.7 | 7.172 | 9982.9 | |
| | | 7.021 | 6759.88 | 7.207 | 10739.48 | 6.991 | 6739.18 | 7.178 | 10667.16 | |

Bold Numbers were adjusted for one run that was abnormal (Avg. of two runs) Example: Avg of three runs 7.351 7.249 7.228 7.276 Time 10740.4 10752.3 10748.5 10747.07 Area



| Mass of th | ne Soil:Solute Ca | lculation |
|---------------------------|------------------------------|-----------|
| MeBT K | 147 90 manufactur | er |
| log K _{ow} | 2.17 | |
| Approximatio | on from benzene | |
| $\log K_{om} = 0.5$ | 2 log K _{ow} + 0.62 | |
| log K _{om} | 1.748 | |
| K_{om} | 56.03 | |
| K _{oc} = 1.724*I | √om | |
| K_{oc} | 96.59 | |
| f _{om} for soil | 0.02867 (by percen | t) |
| $f_{oc} = .58*f_{om}$ | | |
| f _{oc} | 0.017 | |
| $Kd = f_{oc} * K_{oc}$ | | · · · |
| | | |
| Mass of soil | | |
| Ms = Vol. of | solution/Kd | |
| Vol. of soluti | on = 16.06 | mL |
| Ms ≖ | 1U.U g | |
| | | |

| Mass of t | he Soil:Solute Ca | lculation | |
|---------------------------------------|-------------------------------|-----------|---|
| MeBT K _{ow} | 147.90 | | |
| $\log K_{ow}$ | 2.17 | | |
| Approximati | on from benzene | | |
| $\log K_{om} = 0.8$ | 52 log K _{ow} + 0.62 | | |
| log K _{om} | 1.748 | | |
| K _{om} | 56.03 | | |
| K _{oc} = 1.724* | K _{om} | | |
| K _{oc} | 96.59 | | |
| f _{om} for soil | 0.0287 (by percen | t) | |
| $f_{oc} = .58 T_{om}$ | 0.017 | | |
| Kd = f _{oc} *K _{oc} | | | |
| | | | <theoretical coefficient<="" sorption="" td=""></theoretical> |
| Mass of soil | | | |
| Ms = Vol of | solution/Kd | | |
| Vol of solutio | on = 16.08 10 g | mL | |
| | | | |

| Mass of th | ne Soil: | Solute C | alculatio | - n 7 |
|---------------------------------------|-----------------------|------------|-----------|---|
| MeBT K _{ow} | 147.90 | I | | |
| log K _{ow} | 2.17 | | | |
| Approximatic | on from b | enzene | - | |
| $\log K_{om} = 0.5$ | 2 log K _{ow} | , + 0.62 | | |
| log K _{om} | 1.748 | i - | | |
| K _{om} | 56.03 | I | | |
| K _{oc} = 1.724*k | ≺ _{om} | | - | |
| K _{oc} | 96.59 | I | | |
| f _{om} for soil | 0.0137 | (by perce | _ nt) | |
| f _{oc} | 0.008 | | | |
| Kd = f _{oc} *K _{oc} | | | _ | |
| | 0.7/67/ | | | <theoretical coefficient<="" sorption="" td=""></theoretical> |
| Mass of soil | | | _ | |
| Ms = Vol of s | olution/K | d | | |
| Vol of solutio | n = | 7.67 | mL | |
| Ms 🗧 | 10 | g | | |
| Ms | 10 | g 7.07 | | |

, · · ·

.

| Mass of th | ne Soil:S | olute Calculation | |
|---------------------------|--------------------------|-------------------------|---|
| MeBT K _{ow} | 147.90 | | |
| log K _{ow} | 2.17 | | |
| Approximatio | on from ber | nzene | |
| $\log K_{om} = 0.5$ | 52 log K _{ow} + | + 0.62 | |
| log K _{om} | 1.748 | | |
| K _{om} | 56.03 | | |
| K _{oc} = 1.724*I | ≺ _{om} | | |
| K _{oc} | 96.59 | | |
| f _{om} for soil | 0.009 (| (by percent, below LOD) | |
| f _{oc} | 0.005 | | |
| $Kd = f_{oc} K_{oc}$ | | | |
| Kelmensk | 0.504 | | <theoretical coefficient<="" sorption="" td=""></theoretical> |
| Mass of soil | | | |
| Ms = Vol of s | solution/Kd | |] |
| Vol of solutio | on = [| 5.04 mL | |
| Ms = | 10 ថ្ | g | |

.

| 1000 mg/L | 0 | | 8225.7 | | 13224.4 | 4 MeBT | 5 MeBT |
|-----------|-------|-------|--------|-------|---------|-------------|-------------|
| Water | 0.5 | 7.019 | 7004.7 | 7.207 | 10990.5 | 15% | 17% |
| 1 Meth | 1 hr | 7.026 | 2069.1 | 7.215 | 3606.9 | 75% | 73% |
| 2 Meth | 6 hr | 7.025 | 562.6 | 7.211 | 1015.3 | 93 % | 92 % |
| 3 Meth | 24 hr | 7.018 | 441.7 | 7.205 | 865.9 | 95% | 93% |
| 4 Meth | 48 hr | 7.011 | 439.6 | 7.201 | 857.2 | 9 5% | 94 % |
| 5 Meth | 72 hr | 7.007 | 433.8 | 7.198 | 852.4 | 95% | 94% |

Percent MeBT Recoved 94.73% 93.55%

Bold numbers obtained from calibration curve - 10/31/98



Filtered Sample Comparison

| | Filtered | Sample | | | | | | |
|-----------|-----------|----------|---------|----------|-----------|----------|---------|---------|
| | Retention | 4 - Area | 4 | 4 Peak | Retention | 5 - Area | 5 | 5 Peak |
| 1000 mg/L | Time | (mAu*s) | Average | Std Dev | Time | (mAu*s) | Average | Std Dev |
| | 8.456 | 7390.9 | | | 8.951 | 12416.6 | | |
| | 8.345 | 7383.1 | | | 8.853 | 12385.7 | | |
| | 8.295 | 7370.7 | | | 8.742 | 12380 | | 1 |
| | 8.152 | 7361.4 | | | 8.681 | 12375.2 | | |
| | 8.108 | 7369.6 | 7375.14 | 11.73597 | 8.511 | 12379.5 | 12387.4 | 16.7447 |

Avg. Time 8.2712

.

Avg. Time 8.7476

Difference from unfiltered 1000 mg/L sample 4 MeBT

850.52

5 MeBT 836.96

Difference 10.3%

Difference 6.3%



| Sorption Coefficient - ' | | | | | | J |
|--------------------------|----------|---------|-------|---------|---------------|---------------|
| Ű | | 1.000 | | 0.800 | 0.600 | D. |
| | | | | | | "Х |
| Kd-5 | 0.825 | 0.512 | 0.075 | 0.051 | er type | |
| Kd-4 | 0.860 | 0.350 | 0.044 | 0.036 | vithin isom | alues |
| foc | 5.25% | 5.25% | 1.37% | 0.90% | % window \ | rage Kd v |
| | Untreate | Treated | DIA | Westove | s indicate 5% | s reflect ave |
| | - | 7 | ო | 4 | Brackets | Number |





| | | foc | Kd-4 | Kd-5 |
|---|----------|-------|-------|-------|
| | Untreate | 5.25% | 1.203 | 1.645 |
| ~ | Treated | 5.25% | 0.706 | 44.5 |
| ~ | DIA | 1.37% | 0.160 | 0.255 |
| + | Westove | 0.90% | 0.148 | 0.203 |
| | - | | | |

Brackets indicate 5% window within isomer type Numbers reflect average Kd values





| | 7.66 | 32.8 | %67.6 | Untreate | 4 |
|-------|-------|------|-------|----------|---|
| | 55.2 | 32.8 | 5.25% | Untreate | 4 |
| 1000 | 16860 | 0.5 | 5.25% | Treated | ო |
| | 5.02 | 2.84 | 1.37% | DIA | 7 |
| | 9401 | 3.89 | 0.90% | Westove | ~ |
| Sorpt | Kd-5 | Kd-4 | foc | | |



Brackets indicate 5% window within isomer type Numbers reflect average Kd values



| Kd-5 | 1.735 | 2.325 | 0.291 | 0.193 | |
|------|----------|---------|-------|---------|--|
| Kd-4 | 1.219 | 0.547 | 0.208 | 0.136 | |
| foc | 5.25% | 5.25% | 1.37% | 0.90% | |
| | Untreate | Treated | DIA | Westove | |
| | ~ | 2 | ę | 4 | |

Brackets indicate 5% window within isomer type Numbers reflect average Kd values



Appendix C

Preliminary Experiments

for PG and MeBT

.



Appendix C



Appendix C

Appendix D

Calibration Curve Data

for MeBT Isomers

Calibration Curve Data <u>10/31/98</u>

2 Columns, ACN, 2 Peaks

| Standard | | Retention | 4 - Area | 4 | 4 Peak | Retention | 5 - Area | 5 | 5 Peak |
|------------|-----------|-----------|----------|-------------|-----------|-----------|----------|---------|---------|
| oncentrati | Replicate | Time | (mAu*2) | Average | Std Dev | Time | (mAu*2) | Average | Std Dev |
| 1000 mg/L | 1 | 7.891 | 8237.7 | | | 8.115 | 13252.6 | | |
| - | 2 | 7.798 | 8229.2 | | | 8.031 | 13229.9 | | |
| | 3 | 7.790 | 8219.3 | | | 8.004 | 13208.3 | | |
| | 4 | 7.783 | 8221.3 | | | 7.997 | 13216.9 | | |
| | 5 | 7.777 | 8220.8 | 8225.7 | 7.76 | 7.990 | 13214.1 | 13224.4 | 17.66 |
| 500 mg/L | 1 | 7.770 | 4170.7 | | | 7.984 | 6933.8 | | |
| , i | 2 | 7.761 | 4165.5 | | | 7.974 | 6925.7 | | |
| | 3 | 7.753 | 4165.3 | | | 7.966 | 6922.6 | | |
| | 4 | 7.751 | 4165.1 | | | 7.963 | 6921.8 | | |
| | 5 | 7.750 | 4164.8 | 4166.3 | 2.48 | 7.959 | 6920.2 | 6924.8 | 5.40 |
| 100 mg/L | 1 | 7.745 | 919.0 | | | 7.959 | 1538.8 | | |
| - | 2 | 7.742 | 917.6 | | | 7.956 | 1536.9 | | |
| | 3 | 7.733 | 917.4 | 918.0 | 0.87 | 7.947 | 1535.2 | 1537.0 | 1.80 |
| 50 mg/L | 1 | 7.725 | 461.2 | | | 7.939 | 773.5 | | |
| | 2 | 7.718 | 460.6 | | | 7.932 | 771.8 | | |
| | 3 | 7.709 | 459.7 | 460.5 | 0.75 | 7.922 | 771.0 | 772.1 | 1.28 |
| 10 mg/L | 1 | 7.701 | 119.6 | | | 7.916 | 200.2 | | |
| | 2 | 7.695 | 119.8 | | | 7.908 | 201.6 | | |
| | 3 | 7.685 | 120.4 | 119.9 | 0.42 | 7.897 | 201.4 | 201.1 | 0.76 |
| 5 mg/L | 1 | 7.638 | 68.1 | | | 7.846 | 113.4 | | |
| | 2 | 7.630 | 67.6 | | | 7.837 | 113.0 | | |
| | 3 | 7.622 | 68.4 | | | 7.828 | 113.8 | | |
| | 4 | 7.614 | 68.7 | | | 7.819 | 114.2 | | |
| | 5 | 7.607 | 68.5 | 68.3 | 0.43 | 7.811 | 113.6 | 113.6 | 0.45 |
| 1 mg/L | 1 | 7.678 | 42.4 | | | 7.892 | 70.7 | | |
| | 2 | 7.668 | 41.4 | | | 7.881 | 70.0 | | |
| | 3 | 7.661 | 41.9 | | | 7.873 | 70.6 | | |
| | 4 | 7.653 | 41.7 | | | 7.862 | 70.1 | | |
| | 5 | 7.648 | 41.3 | 41.7 | 0.44 | 7.855 | 70.3 | 70.3 | 0.30 |
| 0 mg/L | 1 | | 4.8 | | | | 40.0 | | |
| | 2 | | 5.0 | | | | 5.0 | | |
| | 3 | | 4.0 | | | | 23.0 | | |
| | 4 | | 6.2 | | | | 16.5 | | |
| | 5 | | 4.1 | 4.8 | 0.9 | | 18.0 | 20.5 | 12.74 |
| Avg Retent | ion Time, | 7.714 | Avg | Retention - | Гіте, 5 = | 7.926 | | | |

.

Retention Time Difference = 0.213

| 1 | = | | 0.2 |
|---|---|--|-----|
| | | | |

| Variance at low | concentration | |
|--------------------|--------------------|------|
| Var = Std Dev | ^2 | |
| Var : | = 195.53 | |
| Std Dev : | = 13.98 | |
| Limit of Detection | on, LOD | |
| | - /105 | |
| | - 41.93 | |
| siope : | = 8.182 | |
| 4 MeBT LOD of | appro 5.127 | mg/L |

| Variance at Ic | w conce | entration | |
|---------------------------------|-------------------------|-----------|------|
| Var = Std De | ev^2 | | |
| Var | = | 519.97 | |
| Std Dev | = | 22.80 | |
| Limit of Detec LOD = 3 * Std | ction, LO Dev | D | |
| LOD | = | 68.41 | |
| slope | = | 13.19 | |
| 5 MeBT LOD o | of appro_ | 5.186 | mg/L |

| | | 4 | 4 | 5 | 5 |
|-----------|----------|--------|-----------|---------|-----------|
| | Standard | Avg | 3 Std Dev | Avg | 3 Std Dev |
| | 1000 | 8225.7 | 23.27 | 13224.4 | 52.97 |
| | 500 | 4166.3 | 7.45 | 6924.8 | 16.21 |
| | 100 | 918.0 | 2.62 | 1537.0 | 5.40 |
| | 50 | 460.5 | 2.26 | 772.1 | 3.83 |
| | 10 | 119.9 | 1.25 | 201.1 | 2.27 |
| Below LOD | 5 | 68.3 | 1.29 | 113.6 | 1.35 |
| Below LOD | 1 | 41.7 | 1.32 | 70.3 | 0.91 |
| Below LOD | 0 | 4.8 | 2.7 | 20.5 | 38.21 |







| | | | High | Low | Close | | | High | Low | Close |
|----------|---------|---------|----------|----------|--------|---------|---------|----------|----------|---------|
| | 4 | 4 | 4 | 4 | 4 | 5 | 5 | 5 | 5 | 5 |
| Standard | Std Dev | 3 sigma | +3 sigma | -3 sigma | Avg | Std Dev | 3 sigma | +3 sigma | -3 sigma | Avg |
| 1000 | 7.76 | 23.27 | 8248.9 | 8202.4 | 8225.7 | 17.66 | 52.97 | 13277.3 | 13171.4 | 13224.4 |
| 500 | 2.48 | 7.45 | 4173.7 | 4158.8 | 4166.3 | 5.40 | 16.21 | 6941.0 | 6908.6 | 6924.8 |
| 100 | 0.87 | 2.62 | 920.6 | 915.4 | 918.0 | 1.80 | 5.40 | 1542.4 | 1531.6 | 1537.0 |
| 50 | 0.75 | 2.26 | 462.8 | 458.2 | 460.5 | 1.28 | 3.83 | 775.9 | 768.3 | 772.1 |
| 10 | 0.42 | 1.25 | 121.2 | 118.7 | 119.9 | 0.76 | 2.27 | 203.3 | 198.8 | 201.1 |
| 5 | 0.43 | 1.29 | 69.6 | 67.0 | 68.3 | 0.45 | 1.35 | 115.0 | 112.3 | 113.6 |
| 1 | 0.44 | 1.32 | | | 41.7 | 0.30 | | | | 70.3 |
| 0 | 0.9 | 2.65 | | | 4.8 | 12.74 | | | | 20.5 |



.

Appendix E

Wright-Patterson AFB Soil

Untreated Sample Analysis
Appendix E

Wright-Patterson AFB Soil

4- MeBT

.

.

| 0.105 | .105 | | | | | | | | slope | all data 0.8587 | with 10 n 0.4373 | ng/L n = // - | 0.859 |
|---------|-----------|------------|--------|---------|----------|-------------|-----------|----------|------------------|--------------------|----------------------|---------------------|---------|
| d in bo | pi | | | | | | | | intercept r^2 | 0.3385 0.9946 | 5 1.3754 3 0.8883 | Kf = | 2.180 |
| | | | and | Initia | 폐 | inal | | | Initial | Sorbed | 1 point | | |
| | Vial | + Air-drie | d MeB | T Extr | ractio (| Concentrati | Soil (dry | Vol Soln | Concentrati | Concentrati | Sorption | | |
| Vial (| Only soil | Soil | mix | c | 0 | u | wt) (g) | (mL) | on | uo | Coeff. | log S I | -og Cf |
| D | ŋ | D | | | | പ് | Σ | ٧s | ö | S | Хd | | |
| | | | | | | mg/L | | | | b/brl | ml/g | | |
| 28.1 | 81 38. | 187 10.00 | 6 61.7 | 775 57. | .695 | 738.63 | 9.055 | 24.539 | 961.25 | 603.289 | 0.817 | 2.78053 | 2.86843 |
| 28.3 | 38. | 318 10.00 | 9 63.9 | 981 59 | .845 | 736.87 | 9.058 | 26.614 | 964.26 | 668.132 | 0.907 | 2.82486 | 2.86739 |
| 28.4 | 153 38. | 468 10.01 | 5 62.(| 056 59 | .129 | 735.77 | 9.063 | 24.540 | 961.22 | 610.421 | 0.830 | 2.78563 | 2.86674 |
| 28.C | 80 38. | 087 10.00 | 7 61. | 715 57 | .549 | 733.93 | 9:056 | 24.579 | 961.31 | 617.132 | 0.841 | 2.79038 | 2.86565 |
| 28.2 | 287 38. | 291 10.00 | 4 62.8 | 319 57 | .725 | 728.04 | 9.053 | 25.479 | 962.69 | 660.366 | 0.907 | 2.81978 | 2.86216 |
| 28.3 | 327 38. | 333 10.00 | 6 64. | 78 63 | .068 | 69.63 | 9.055 | 27.398 | 96.53 | 81.389 | 1.169 | 1.91056 | 1.8428 |
| 28.1 | 80 38. | 181 10.00 | 1 64. | 594 62 | .793 | 70.09 | 9.051 | 27.363 | 96.53 | 79.928 | 1.140 | 1.9027 | 1.84566 |
| 28.1 | 64 38. | 171 10.00 | 7 64.(| 544 62 | .803 | 71.07 | 9.056 | 27.424 | 96.53 | 77.106 | 1.085 | 1.88709 | 1.85169 |
| 28.3 | 316 38. | 319 10.00 | 3 64. | 504 62 | .487 | 65.11 | 9.052 | 27.136 | 96.50 | 94.085 | 1.445 | 1.97352 | 1.81365 |
| 27.9 | 323 37. | 932 10.00 | 9 64. | 135 62 | .165 | 69.29 | 9.058 | 27.154 | 96.50 | 81.563 | 1.177 | 1.91149 | 1.84067 |
| 27.4 | 121 37. | 424 10.00 | 3 63.9 | 934 61 | .603 | 0.82 | 9.052 | 27.461 | 9.65 | 26.797 | 32.680 | 1.42809 | -0.0862 |
| 27.8 | 377 37. | 878 10.00 | 1 64. | 202 62 | .449 | 0.83 | 9.051 | 27.274 | 9.65 | 26.584 | 32.029 | 1.42462 | -0.0809 |
| 27.3 | 316 37. | 324 10.00 | 8 64. | 252 62 | .428 | 0.84 | 9.057 | 27.879 | 9.66 | 27.146 | 32.317 | 1.43371 | -0.0757 |
| . 27.8 | 316 37. | 823 10.00 | 7 64. | 252 62 | 414 | 0.77 | 9.056 | 27.380 | 9.65 | 26.856 | 34.877 | 1.42904 | -0.1135 |
| 27.6 | 356 37. | 656 10.00 | 0 63.1 | 885 62 | 003 | 0.82 | 9.050 | 27.179 | 9.65 | 26.520 | 32.342 | 1.42358 | -0.0862 |
| 1 | | | | | Ļ | | | | | | | F | |
| 28.2 | 296 38. | 298 10.00 | 2 64. | 033 62 | 100. | /0.23 | 9.052 | G87.12 | 96.52 | 19.240 | 1.128 | | |
| 27.8 | 353 37. | 856 10.00 | 3 64.1 | 030 62 | .264 | 67.59 | 9.052 | 27.125 | 96.50 | 86.612 | 1.281 | | |
| 28.1 | 185 38. | 192 10.00 | 7 64.1 | 635 62 | .815 | 69.05 | 9.056 | 27.394 | 96.53 | 83.121 | 1.204 | | |
| 27.9 | 960 37. | 968 10.00 | 8 64. | 242 62 | .371 | 68.59 | 9.057 | 27.225 | 96.51 | 83.917 | 1.223 | | |
| 27.9 | 916 37. | 919 10.00 | 3 64. | 171 62 | .400 | 68.03 | 9.052 | 27.203 | 96.51 | 85.569 | 1.258 | | |
| | | | | | | Ŧ | | | | 737 737 | 727 50 | | |
| | | | | | | - 0001 | | | | 101.07 | 101.02 | | |
| | | | | | | UUUL | | | | 486.81 | 0.487 | | |

Untreated.xls





Untreated.xls

Appendix E

Wright-Patterson AFB Soil

5-MeBT

.

,

| 5-MeBT Data | | | | | | | | | | all data | with 10 m | g/L | |
|----------------------------|-----------|--------|-----------|--------|-----------|-------------|-----------|----------|-------------|-------------|-----------|---------|---------|
| Moisture Content o | f soil | 0.105 | | | | | | | slope | 0.7325 | 0.402 | Ľ | 0.732 |
| Outliers identified | in bold | | | | | | | | intercept | 0.6839 | 1.491 | ₹= | 4.829 |
| | | | | | | | | | r^2 | 0.9983 | 0.922 | | |
| | | | | and | Initial | Final | | | Initial | Sorbed | 1 point | | |
| | | Vial + | Air-dried | MeBT | Extractio | Concentrati | Soil (dry | Vol Soln | Concentrati | Concentrati | Sorption | | |
| Added Concentrati | Vial Only | ' soil | Soil | mix | L | on | wt) (g) | (mL) | uo | on | Coeff. | log S 1 | -og Cf |
| | g | ß | 6 | | | ັບ | Σ | Vs | ö | s | Кd | | |
| | | | | | | mg/L | | | | 6/6rf | ml/g | | |
| 1000 Bottle 1 | 28.181 | 38.187 | 10.006 | 61.775 | 57.695 | 744.8 | 9.055 | 24.539 | 961.25 | 586.569 | 0.788 | 2.76832 | 2.87204 |
| 1000 Bottle 2 | 28.309 | 38.318 | 10.009 | 63.981 | 59.845 | 743.81 | 9.058 | 26.614 | 964.26 | 647.740 | 0.871 | 2.8114 | 2.87146 |
| 1000 Bottle 3 | 28.453 | 38.468 | 10.015 | 62.056 | 59.129 | 742.16 | 9.063 | 24.540 | 961.22 | 593.120 | 0.799 | 2.77314 | 2.8705 |
| 1000 Bottle 4 | 28.080 | 38.087 | 10.007 | 61.715 | 57.549 | 741.46 | 9.056 | 24.579 | 961.31 | 596.695 | 0.805 | 2.77575 | 2.87009 |
| 1000 Bottle 5 | 28.287 | 38.291 | 10.004 | 62.819 | 57.725 | 736.75 | 9.053 | 25.479 | 962.69 | 635.854 | 0.863 | 2.80336 | 2.86732 |
| 100 Bottle 1 | 28.327 | 38.333 | 10.006 | 64.78 | 63.068 | 62.91 | 9.055 | 27.398 | 96.53 | 101.721 | 1.617 | 2.00741 | 1.79872 |
| 100 Bottle 2 | 28.180 | 38.181 | 10.001 | 64.594 | 62.793 | 63.48 | 9.051 | 27.363 | 96.53 | 99.913 | 1.574 | 1.99962 | 1.80264 |
| 100 Bottle 3 | 28.164 | 38.171 | 10.007 | 64.644 | 62.803 | 64.65 | 9.056 | 27.424 | 96.53 | 96.548 | 1.493 | 1.98474 | 1.81057 |
| 100 Bottle 4 | 28.316 | 38.319 | 10.003 | 64.504 | 62.487 | 62.31 | 9.052 | 27.136 | 96.50 | 102.479 | 1.645 | 2.01063 | 1.79456 |
| 100 Bottle 5 | 27.923 | 37.932 | 10.009 | 64.135 | 62.165 | 62.37 | 9.058 | 27.154 | 96.50 | 102.308 | 1.640 | 2.00991 | 1.79498 |
| 10 Bottle 1 | 27.421 | 37.424 | 10.003 | 63.934 | 61.603 | 0.53 | 9.052 | 27.461 | 9.65 | 27.677 | 52.221 | 1.44212 | -0.2757 |
| 10 Bottle 2 | 27.877 | 37.878 | 10.001 | 64.202 | 62.449 | 0.51 | 9.051 | 27.274 | 9.65 | 27.548 | 54.016 | 1.44009 | -0.2924 |
| 10 Bottle 3 | 27.316 | 37.324 | 10.008 | 64.252 | 62.428 | 0.54 | 9.057 | 27.879 | 9.66 | 28.069 | 51.980 | 1.44823 | -0.2676 |
| 10 Bottle 4 | 27.816 | 37.823 | 10.007 | 64.252 | 62.414 | 0.46 | 9.056 | 27.380 | 9.65 | 27.793 | 60.419 | 1.44393 | -0.3372 |
| 10 Bottle 5 | 27.656 | 37.656 | 10.000 | 63.885 | 62.003 | 0.48 | 9.050 | 27.179 | 9.65 | 27.541 | 57.378 | 1.43999 | -0.3188 |
| | | | | | - | | | | | | | | |
| 100 Mix 1 | 28.296 | 38.298 | 10.002 | 64.633 | 62.651 | 62.65 | 9.052 | 27.285 | 96.52 | 102.089 | 1.630 | | |
| 100 Mix 2 | 27.853 | 37.856 | 10.003 | 64.030 | 62.264 | 60.52 | 9.052 | 27.125 | 96.50 | 107.796 | 1.781 | | |
| 100 Mix 3 | 28.185 | 38.192 | 10.007 | 64.635 | 62.815 | 61.34 | 9.056 | 27.394 | 96.53 | 106.443 | 1.735 | | |
| 100 Mix 4 | 27.960 | 37.968 | 10.008 | 64.242 | 62.371 | 61.44 | 9.057 | 27.225 | 96.51 | 105.410 | 1.716 | | |
| 100 Mix 5 | 27.916 | 37.919 | 10.003 | 64.171 | 62.400 | 60.20 | 9.052 | 27.203 | 96.51 | 109.098 | 1.812 | | |
| | | | | | | | | | | | | | |

Untreated.xls

30.984 0.498

30.98 498.43

1 1000





| MeBT Isomer Extration Curve - 1000 mg/L | 12000 4 MeBT | | | 0008 | | | 4000 P | A | 2000 | | 0 2 4 6 | Number of Times Extracted | | | - | | | | | | MoBT learner Extendion Prince 400 mail | | | | | | | 1) E | | ₹ 200 | } | | | 0 2 4 0 Numbar of Times Extracted | | | | | | | |
|---|--------------|--------|--------|--------|--------|--------|---------------|-----------------|----------|--------------|---------------------|---------------------------|-----------------------|---------------|--------|--------|--------|--------|--------|--------|--|-------|--------|--------|--------|--------|--------|---------------|-----------------|-------|-------------|--------------|---------------------|--------------------------------------|----------------------|---------------|--------|--------|--------|--------|--------|
| 5 MeBT | 80% | 12% | 3% | 2% | 1% | %0 | | | | | | | _ <u>}</u> | | | | | | | | 5 MeBT | 58% | 21% | 14% | 3% | 2% | 1% | | | | | | | | | J | | | | | |
| 4 MeBT | 85% | 12% | 1% | 1% | %0 | %0 | | | | | | | | | | | | | | | 4 MeBT | 64% | 19% | 12% | 3% | 1% | 1% | | | | | | | | | | | | | | |
| 13224.4 | 10598.5 | 1523.7 | 437.1 | 261.8 | 149.6 | 62.1 | 13032.8 | 98.55% | | | | | | | | | | | | | 1537.0 | 887.4 | 318.3 | 215.7 | 52.0 | 37.1 | 17.6 | 1528.1 | 99.42% | | | | | | | | | | | | |
| _ | 7.207 | 7.215 | 7.211 | 7.205 | 7.201 | 7.198 | | | forcord, | | | | 5.094 | | 20.980 | 17.761 | 17.682 | 17.760 | 18.264 | 17.832 | | 7.264 | 7.214 | 7.439 | 7.268 | 7.299 | 7.231 | | | | fference | | | | 1.97 | | 1.794 | 1.859 | 1.968 | 1.835 | 1.869 |
| 8225.7 | 6994.7 | 1015.6 | 121.8 | 55.7 | 20.1 | 12.5 | 8220.4 | 99.94% | Diffied | | | | | 63.294 | 60.872 | 61.554 | 61.217 | 61.641 | 61.521 | | 918.0 | 591.8 | 171.9 | 105.6 | 26.4 | 13.1 | 8.5 | 917.3 | 99.92% | | Refilled Di | | | | | 64.267 | 64.512 | 64.399 | 64.133 | 64.774 | |
| | 7.019 | 7.026 | 7.025 | 7.018 | 7.011 | 7.007 | | | 1 700 0C | 38.291 | 10.004 | 62.819 | 57.725 | | 42.314 | 43.111 | 43.872 | 43.457 | 43.377 | 43.689 | | 7.072 | 7.028 | 7.223 | 7.072 | 7.148 | 7.102 | | | | 27.923 F | 37.932 | 10.009 | 64.135 | 62.165 | | 62.473 | 62.653 | 62.431 | 62.298 | 62.905 |
| 0 | 0.5 | 1 hr | 6 hr | 24 hr | 48 hr | 72 hr | | scovered | | | ied) | mix | tion | | | | | | | | 0 | 0.5 | 1 hr | 6 hr | 24 hr | 48 hr | 72 hr | | scovered | | | | ried) | mix | tion | | | | | | |
| 1000 mg/L | Water | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT Re | Emot. | ul Soil only | Soil Weight (air-dr | w/ Soil and MeBT | After Initial Extract | of water/MeBT | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | 6 Meth | 100 mg/L | Water | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT Re | | Empty | w/ Soil only | Soil Weight (air-dr | w/ Soil and MeBT | After Initial Extrac | of water/MeBT | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth |

.

Soil Calculation - Untreated.xls

MeBT Isomer Extraction Curve - 10 mg/L

| | | 20 | 3 | 25 | | 20 | | 15 | 10 | | Ω. | 0 | C | ر | | | | | | | | | | | | | | | | | |
|---------|-------|--------|--------|--------|--------|--------|---------------|----------------|------------------|---------------------|--------|------------|--------------|--------------------|------------------|-----------------------|---------------|--------|--------|--------|--------|--------|--------|-------|--------|--------|--------|--------|--------|---------------|----------------|
| | | | | | (| s*I | ١Å | ա) | 691. | A | | | | | | | | | | | | | | | | | | | | | |
| 5 MeBT | %0 | %9 | 6% | 2% | %0 | %0 | | | | | | | | | | | | | | | | | 5 MeBT | %0 | %0 | %0 | %0 | %0 | %0 | | |
| 4 MeBT | 25% | 18% | 11% | 5% | %0 | %0 | | | | | | | | | | | | | | | | | 4 MeBT | %0 | %0 | %0 | %0 | %0 | %0 | | |
| 201.1 | 0.0 | 12.6 | 12.1 | 4.1 | 0 | 0 | 28.8 | 14.32% | | | | | | | | | | | | | | | 20.5 | 0.00 | 0.00 | 0.00 | 00.0 | 0.00 | 0.00 | 00.0 | %00.0 |
| | 7.35 | 7.307 | 7.402 | 7.382 | 7.35 | 7.35 | | | | | | Difference | | | | 1.882 | | 2.061 | 1.776 | 1.911 | 1.881 | 1.944 | | 7.35 | 7.35 | 7.35 | 7.35 | 7.35 | 7.35 | | |
| 119.9 | 29.4 | 21.0 | 13.6 | 6.5 | 0 | 0 | 70.5 | 58.80% | 0 | - c 0 u | 0.0 | Refilled | | | | | 64.509 | 64.371 | 64.011 | 64.229 | 64.094 | | 4.8 | 0.00 | 0.00 | 0.00 | 00.00 | 0.00 | 0.00 | 0.00 | 0.00% |
| | 7.142 | 7.118 | 7.203 | 7.189 | 7.135 | 7.135 | | | 7 860 | 600°.1 | 1.899 | 27.656 | 37.656 | 10.000 | 63.885 | 62.003 | | 62.448 | 62.595 | 62.100 | 62.348 | 62.150 | | 7.135 | 7.135 | 7.135 | 7.135 | 7.135 | 7.135 | | |
| 0 | 0.5 | 1 hr | 6 hr | 24 hr | 48 hr | 72 hr | | ecovered | Mater | Atoth 1 | Meth 1 | | | ried) | mix | tion | | | | | | | 0 | 0.5 | 1 hr | 6 hr | 24 hr | 48 hr | 72 hr | | ecovered |
| 10 mg/L | Water | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT R | Additional Snika | אחמווחוומו שו אחווא | | Empty | w/ Soil only | Soil Weight (air-d | w/ Soil and MeBT | After Initial Extract | of water/MeB1 | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | 0 mg/L | Water | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT R |

.

ဖ

4

2

0

Number of Extractions

| | | 11 47 | 001.7 | 0.00 | nn. / | 5 |
|------------|--------------------|---------|--------|----------|------------|------|
| | 4 Meth | 48 hr | 7.135 | 0.00 | 7.35 | ö |
| | 5 Meth | 72 hr | 7.135 | 0.00 | 7.35 | ö |
| S | um Recovered | | | 0.00 | | õ |
| Ъ | rcent MeBT Re | covered | | 0.00% | | 0.00 |
| ш | pty | | 26.167 | Refilled | Difference | |
| <u>/</u> M | Soil only | | 36.172 | | | |
| S | il Weight (air-dri | ied) | 10.005 | | | |
| /M | Soil and MeBT | mix | 62.594 | | | |
| Afi | er Initial Extract | tion | 58.456 | | 4.138 | |
| | of water/MeBT | | | | | |
| | 1 Meth | | | | 0.000 | |
| | 2 Meth | | | | 0.000 | |
| | 3 Meth | | | | 0.000 | |
| | | | | | | |

| 'n |
|------------------|
| <u> </u> |
| × |
| |
| σ |
| ā |
| |
| đ |
| ň |
| ۳ |
| ₽ |
| c |
| - |
| _ |
| |
| |
| |
| c |
| S |
| <u>io</u> |
| ition |
| ation |
| ulation |
| sulation |
| Iculation |
| alculation |
| Calculation |
| Calculation |
| Calculation |
| il Calculation |
| oil Calculation |
| Soil Calculation |

| | MeBT Isomer Extraction Curve - Mix | | -+-4 MeBT | ■ -=5 MeBT | | | | | | | | | | | | - c - c - c - c | | Number of Extractions | | |
|--------|------------------------------------|--------|-----------|------------|--------|--------|----------------|--------------------|---------------|--------------------|-------------|-------|------------|--------------|---------------------|--------------------------|-----------------------|-----------------------|--------|--------|
| | | 1800 0 | | 1600.0 | 1400.0 | (| - 0.0021 s* | A 1000.0 | с ОС ш) | 6 | √ 14 | 400.0 | 0.000 | 200.0 | 0.0 | | | | | |
| 5 MeBT | 62% | 16% | 13% | 4% | 2% | %0 | | | | | | | | | | | | | | |
| 4 MeBT | %69 | 18% | 6% | 2% | 1% | %0 | | | | | | | | | | | | | | |
| 1537.0 | 949.0 | 239.3 | 198.7 | 65.4 | 23.1 | | 1475.5 | 96.00% | | | | | | | | | | | | |
| | 7.461 | 7.447 | 7.453 | 7.45 | 7.448 | | | f . | | | | | lifference | | | | 1.771 | | 2.837 | 1.914 |
| 918.0 | 637.5 | 162.5 | 56.6 | 22.3 | 8.9 | | 887.83 | 96.71% | - | | | | Refilled D | | | | | 64.248 | 64.190 | 64.236 |
| | 7.268 | 7.249 | 7.254 | 7.251 | 7.244 | | | n de compositiones | | | | | 27.916 | 37.919 | 10.003 | 64.171 | 62.400 | | 61.411 | 62.276 |
| 0 | 0.5 | 1 hr | 6 hr | 24 hr | 48 hr | 72 hr | | covered | | Water | Meth 1 | | | | ied) | mix | tion · | | | |
| Mix | Water | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT Re | | Additional Spike a | | | Empty | w/ Soil only | Soil Weight (air-dr | w/ Soil and MeBT | After Initial Extract | of water/MeBT | 1 Meth | 2 Meth |

~

2.040

62.196

3 Meth

Soil Calculation - Untreated.xls

| 1000 mg/l | | | | | | | | |
|--------------|--------------|--------------|----------|--------|--------|---------|--------|---|
| l www.mg/L | Runs | [| 4 MeBT | | | 5 MeBT | | |
| | 1 | 7.036 | 6103.7 | 738.63 | 7.222 | 10009.1 | 744.80 | |
| | 2 | 7.042 | 6089.4 | 736.87 | 7.238 | 9996.6 | 743.81 | |
| | 3 | 7.027 | 6080.5 | 735.77 | 7.211 | 9975.8 | 742.16 | |
| | 4 | 7.026 | 6065.5 | 733.93 | 7.184 | 9967.0 | 741.46 | |
| | 5 | 7.021 | 6017.7 | 728.04 | 7.157 | 9907.8 | 736.75 | |
| | avg | 7.031 | 6071.36 | | 7.202 | 9971.26 | | |
| | conc. | ······ | | 734.65 | | | 741.80 | Interpolated from calibration curve |
| | Standard D | Deviation | 33.05 | | | 39.18 | | |
| | | Average of 4 | & 5 MeBT | | 738.22 | | | |
| | | | | | | | | |
| 100 mg/L | | | | | | | | |
| | Runs | | 4 MeBT | | | 5 MeBT | | |
| 1 | 1 | 7.661 | 640.12 | 69.63 | 8.266 | 969.65 | 62.91 | |
| | 2 | 7.736 | 644.71 | 70.09 | 7.946 | 978.26 | 63.48 | |
| | 3 | 7.701 | 653.33 | 71.07 | 7.908 | 996.18 | 64.65 | |
| | 4 | 7.675 | 598.78 | 65.11 | 7.881 | 960.36 | 62.31 | |
| | 5 | 7.644 | 637.02 | 69.29 | 7.849 | 961.38 | 62.37 | |
| | avg | 7.683 | 634.79 | | 7.970 | 973.17 | | |
| | conc. | | | 69.038 | | | 63.144 | Interpolated from calibration curve |
| | Standard E | Deviation | 21.05 | | | 14.76 | | |
| | Average of | f 4 & 5 MeBT | | | 0.00 | | | |
| | <u> </u> | | | | | | | |
| 10 mg/L | _ | | | | | | | |
| | Runs | | 4 MeBT | | | 5 MeB I | 0.50 | Other |
| | 1 | 7.247 | 35.03 | 0.82 | 7.439 | 47.02 | 0.53 | 8.340 0.24 |
| | 2 | 7.232 | 35.49 | 0.83 | 7.424 | 45.99 | 0.51 | 8.337 6.39 |
| | 3 | 7.217 | 35.66 | 0.84 | 7.409 | 47.23 | 0.04 | 0.307 6.32 |
| | 4 | 7.204 | 33.10 | 0.77 | 7.395 | 43.20 | 0.40 | 0.300 7.19 8 276 7.65 |
| | 5 | 7.191 | 30.39 | 0.02 | 7.302 | 44.17 | 0.40 | 0.270 |
| | avg | 1.210 | 34.95 | 0.946 | 7.410 | 45.52 | 0 504 | Internalated from calibration curve |
| | Conc. | Deviation | 1.02 | 0.810 | | 1 77 | 0.504 | interpolated norn calibration curve |
| | Standard L | | 9 5 MODT | | 0.00 | 1.77 | | |
| L | | Average 01 4 | | | 0.00 | | | |
| 100 mg/L M | oBT / 1000 m | | | | | | | |
| live mg/c. m | Runs | | 4 MeBT | | | 5 MeBT | | Other |
| | 1 | 7 184 | 645.57 | 70.23 | 7 376 | 965.62 | 62.65 | |
| | 2 | 7 181 | 621 46 | 67.59 | 7.372 | 933.01 | 60.52 | |
| | 3 | 7.181 | 634 77 | 69.05 | 7.374 | 945.53 | 61.34 | |
| | 4 | 7.184 | 630.61 | 68.59 | 7,375 | 947.09 | 61.44 | 7.904 7.00 |
| | 5 | 7,186 | 625,49 | 68.03 | 7.377 | 928.19 | 60.20 | 7.900 7.40 |
| | avg | 7,184 | 631.58 | | 7.375 | 943.89 | | : , · · (t)(ii)/it/Part |
| | conc. | | | 68.698 | | | 61.23 | Interpolated from calibration curve |
| | Standard [| Deviation | 9.31 | | | 14.58 | | • |
| 1 | | | 0.0. | | | | | |

Below LOD

Average of 4 & 5 MeBT

.

0.00

Appendix F

Wright-Patterson AFB Soil

Treated Sample Analysis

,

•

.

Appendix F

.

Wright-Patterson AFB Soil

4-MeBT

.

. .

| 4-MeBT Data | | | | | | | | | | all data | with 10 m | g/L | |
|----------------------------|-----------|--------|-----------|--------|-----------|-------------|-----------|----------|-------------|-------------|-----------|---------|---------|
| Moisture Content o | f soil | 0.105 | | | | | | | slope | 0.7075 | 0.9055 | Ľ | 0.7075 |
| Outliers identified | in bold | | | | | | | | intercept | 0.4028 | -0.1113 | Kf= | 2.528 |
| | | | | | | | | | r^2 | 0666.0 | 0.9826 | | |
| | | | | and | Initial | Final | | | Initial | Sorbed | 1 point | | |
| | | Vial + | Air-dried | MeBT | Extractio | Concentrati | Soil (dry | Vol Soln | Concentrati | Concentrati | Sorption | | |
| Added Concentrati | Vial Only | · soil | Soil | mix | c | on | wt) (g) | (mL) | uo | on | Coeff. | log S L | og Cf |
| | g | g | g | | | ರ | Σ | Vs | ö | S | Кd | | |
| | | | | | | mg/L | | | | 6/6rl | ml/g | | |
| 1000 Bottle 1 | 28.066 | 38.068 | 10.002 | 64.183 | 62.341 | 863.67 | 9.052 | 27.065 | 964.88 | 302.644 | 0.350 | 2.48093 | 2.93635 |
| 1000 Bottle 2 | 27.799 | 37.806 | 10.007 | 64.054 | 62.239 | 866.17 | 9.056 | 27.199 | 965.04 | 296.942 | 0.343 | 2.47267 | 2.93760 |
| 1000 Bottle 3 | 28.092 | 38.092 | 10.000 | 64.564 | 62.715 | 861.52 | 9.050 | 27.422 | 965.35 | 314.616 | 0.365 | 2.49778 | 2.93527 |
| 1000 Bottle 4 | 28.285 | 38.293 | 10.008 | 64.756 | 62.849 | 862.96 | 9.057 | 27.414 | 965.31 | 309.796 | 0.359 | 2.49108 | 2.93599 |
| 1000 Bottle 5 | 28.170 | 38.177 | 10.007 | 64.652 | 62.811 | 869.78 | 9.056 | 27.426 | 965.33 | 289.364 | 0.333 | 2.46144 | 2.93941 |
| 100 Bottle 1 | 28.264 | 38.264 | 10.000 | 64.730 | 62.831 | 78.2 | 9.050 | 27.416 | 96.53 | 55.543 | 0.710 | 1.74463 | 1.89321 |
| 100 Bottle 2 | 28.057 | 38.066 | 10.009 | 64.442 | 62.639 | 76.77 | 9.058 | 27.327 | 96.52 | 55.963 | 0.718 | 1.7479 | 1.89193 |
| 100 Bottle 3 | 28.352 | 38.352 | 10.000 | 64.739 | 62.936 | 78.87 | 9.050 | 27.337 | 96.52 | 53.329 | 0.676 | 1.72696 | 1.89691 |
| 100 Bottle 4 | 27.947 | 37.949 | 10.002 | 64.330 | 62.521 | 78.2 | 9.052 | 27.331 | 96.52 | 55.325 | 0.707 | 1.74293 | 1.89321 |
| 100 Bottle 5 | 28.502 | 38.506 | 10.004 | 64.929 | 63.198 | 77.95 | 9.053 | 27.374 | 96.53 | 56.170 | 0.721 | 1.7495 | 1.89182 |
| 10 Bottle 1 | 28.176 | 38.176 | 10.000 | 65.959 | 64.114 | 8.22 | 9.050 | 28.733 | 9.67 | 4.602 | 0.560 | 0.6629 | 0.91487 |
| 10 Bottle 2 | 27.916 | 37.919 | 10.003 | 64.375 | 62.615 | 8.20 | 9.052 | 27.407 | 9.65 | 4.400 | 0.537 | 0.64341 | 0.91381 |
| 10 Bottle 3 | 28.295 | 38.295 | 10.000 | 64.356 | 62.912 | 8.25 | 9.050 | 27.011 | 9.65 | 4.173 | 0.506 | 0.62048 | 0.91645 |
| 10 Bottle 4 | 28.066 | 38.066 | 10.000 | 64.645 | 62.869 | 8.15 | 9.050 | 27.529 | 9.65 | 4.578 | 0.562 | 0.66064 | 0.91116 |
| 10 Bottle 5 | 27.041 | 37.048 | 10.007 | 63.638 | 61.805 | 8.23 | 9.056 | 27.541 | 9.65 | 4.333 | 0.526 | 0.63677 | 0.91540 |
| | | | | | | | | | | | | | |
| 100 Mix 1 | 27.748 | 37.755 | 10.007 | 64.079 | 62.321 | 81.88 | 9.056 | 27.275 | 96.51 | 44.073 | 0.538 | | |
| 100 Mix 2 | 27.557 | 37.561 | 10.004 | 63.966 | 62.181 | 81.89 | 9.053 | 27.356 | 96.53 | 44.221 | 0.540 | | |
| 100 Mix 3 | 27.852 | 37.852 | 10.000 | 64.071 | 62.281 | 81.7 | 9.050 | 27.169 | 96.50 | 44.440 | 0.544 | | |
| 100 Mix 4 | 27.665 | 37.672 | 10.007 | 64.291 | 62.492 | 82.09 | 9.056 | 27.570 | 96.55 | 44.024 | 0.536 | | |
| 100 Mix 5 | 27.305 | 37.308 | 10.003 | 63.808 | 61.980 | 81.75 | 9.052 | 27.451 | 96.54 | 44.841 | 0.549 | | |
| | | | | | | | | | | | | | |

4-MeBT

Treated-2.xls

0.774 0.403

0.7740 402.93

1 1000





Appendix F

Wright-Patterson AFB Soil

5-MeBT

.

, · · · ·

.

.

| _ |
|----|
| ш |
| Ð |
| Ş. |
| ŵ |
| |

| 5-MeBT Data | | | | | | | | | | all data | with 10 mg | ۲ ۲ | |
|----------------------------|-----------|--------|-----------|--------|-----------|-------------|-----------|----------|-------------|-------------|------------|---------|---------|
| Moisture Content of | soil | 0.105 | | | | | | | slope | 0.1253 | 0.2019 | 11 C | 0.1253 |
| Outliers identified | in bold | | | | | | | | intercept | 2.1925 | 1.8627 | Kf = | 155.79 |
| | | | | | | | | | r^2 | 0.3625 | 0.4744 | | |
| | | | | and | Initial | Final | | | Initial | Sorbed | 1 point | | |
| | | Vial + | Air-dried | MeBT | Extractio | Concentrati | Soil (dry | Vol Soln | Concentrati | Concentrati | Sorption | | |
| Added Concentrati | Vial Only | soil | Soil | mix | Ē | on | wt) (g) | (mL) | uo | on | Coeff. | log S l | .og Cf |
| | D | D | ğ | | | ç | Σ | ۷s | ö | S | Кd | | |
| | I | , | I | | | mg/L | | | | 6/6rl | ml/g | | |
| 1000 Bottle 1 | 28.066 | 38.068 | 10.002 | 64.183 | 62.341 | 821.99 | 9.052 | 27.065 | 964.88 | 427.273 | 0.520 | 2.63071 | 2.91487 |
| 1000 Bottle 2 | 27.799 | 37.806 | 10.007 | 64.054 | 62.239 | 827.7 | 9.056 | 27.199 | 965.04 | 412.482 | 0.498 | 2.6154 | 2.91787 |
| 1000 Bottle 3 | 28.092 | 38.092 | 10.000 | 64.564 | 62.715 | 820.84 | 9.050 | 27.422 | 965.35 | 437.883 | 0.533 | 2.64136 | 2.91426 |
| 1000 Bottle 4 | 28.285 | 38.293 | 10.008 | 64.756 | 62.849 | 825.02 | 9.057 | 27.414 | 965.31 | 424.634 | 0.515 | 2.62801 | 2.91646 |
| 1000 Bottle 5 | 28.170 | 38.177 | 10.007 | 64.652 | 62.811 | 830.51 | 9:056 | 27.426 | 965.33 | 408.290 | 0.492 | 2.61097 | 2.91934 |
| 100 Bottle 1 | 28.264 | 38.264 | 10.000 | 64.730 | 62.831 | 46.86 | 9.050 | 27.416 | 96.53 | 150.487 | 3.211 | 2.1775 | 1.6708 |
| 100 Bottle 2 | 28.057 | 38.066 | 10.009 | 64.442 | 62.639 | 1.52 | 9.058 | 27.327 | 96.52 | 286.607 | 188.557 | 2.45729 | 0.18184 |
| 100 Bottle 3 | 28.352 | 38.352 | 10.000 | 64.739 | 62.936 | 10.64 | 9.050 | 27.337 | 96.52 | 259.435 | 24.383 | 2.41403 | 1.02694 |
| 100 Bottle 4 | 27.947 | 37.949 | 10.002 | 64.330 | 62.521 | 46.69 | 9.052 | 27.331 | 96.52 | 150.470 | 3.223 | 2.17745 | 1.66922 |
| 100 Bottle 5 | 28.502 | 38.506 | 10.004 | 64.929 | 63.198 | 46.59 | 9.053 | 27.374 | 96.53 | 150.989 | 3.241 | 2.17895 | 1.66829 |
| 10 Bottle 1 | 28.176 | 38.176 | 10.000 | 65.959 | 64.114 | 6.08 | 9.050 | 28.733 | 9.67 | 11.396 | 1.874 | 1.05676 | 0.7839 |
| 10 Bottle 2 | 27.916 | 37.919 | 10.003 | 64.375 | 62.615 | 5.89 | 9.052 | 27.407 | 9.65 | 11.393 | 1.934 | 1.05664 | 0.77012 |
| 10 Bottle 3 | 28.295 | 38.295 | 10.000 | 64.356 | 62.912 | 0.01 | 9.050 | 27.011 | 9.65 | 28.782 | 5756 | 1.45913 | -2.301 |
| 10 Bottle 4 | 28.066 | 38.066 | 10.000 | 64.645 | 62.869 | 0.13 | 9.050 | 27.529 | 9.65 | 28.990 | 232 | 1.46224 | -0.9031 |
| 10 Bottle 5 | 27.041 | 37.048 | 10.007 | 63.638 | 61.805 | 00.0 | 9.056 | 27.541 | 9.65 | 29.361 | 195740 | 1.46777 | -3.8239 |
| | | | | | | | | | | | | | |
| 100 Mix 1 | 27.748 | 37.755 | 10.007 | 64.079 | 62.321 | 53.84 | 9:056 | 27.275 | 96.51 | 128.523 | 2.387 | | |
| 100 Mix 2 | 27.557 | 37.561 | 10.004 | 63.966 | 62.181 | 54.13 | 9.053 | 27.356 | 96.53 | 128.100 | 2.367 | | |
| 100 Mix 3 | 27.852 | 37.852 | 10.000 | 64.071 | 62.281 | 55.21 | 9.050 | 27.169 | 96.50 | 123.969 | 2.245 | | |
| 100 Mix 4 | 27.665 | 37.672 | 10.007 | 64.291 | 62.492 | 54.91 | 9.056 | 27.570 | 96.55 | 126.769 | 2.309 | | |
| 100 Mix 5 | 27.305 | 37.308 | 10.003 | 63.808 | 61.980 | 54.71 | 9.052 | 27.451 | 96.54 | 126.836 | 2.318 | | |

72.90 0.294

72.90 293.96

1000



Treated-2.xls



| 1000 mg/L 0 | | | 8225.7 | | 13224.4 | 4 MeBT | 5 MeBT | | | | |
|--------------------------|-------|----------|------------|------------|---------|----------|----------|---|---------------|-----------------------|----------|
| Water 0.4 | 9 | 6.765 | 7125.9 | 7.011 | 11036.8 | 87% | 83% | | MeBT Isome | er Extration Curve | |
| 1 Meth | L | 6.755 | 848.1 | 6.989 | 1863.5 | 10% | 14% | 1200(| | | |
| 2 Meth 61 | л | 6.739 | 372.8 | 6.968 | 543.0 | 5% | 4% | 10001 | | 4 Me | BT |
| 3 Meth 24 | hr | 6.724 | 231.4 | 6.949 | 279.4 | 3% | 2% |)) (s | | - - - 2 Me | BT |
| 4 Meth 48 | hr | 6.719 | 127.3 | 6.936 | 165.1 | 2% | 1% | 000 | | | |
| 5 Meth 72 | hr | 6.711 | 15.5 | 6.924 | 35.4 | %0 | %0 | An | | | |
| Sum Recovered | | | 8721.0 | | 13923.2 | | | 5 5 1) 1 | | | |
| Percent MeBT Reco | vered | | 106.02% | | 105.28% | | | 168 400 | | | |
| | | | - | | | | | A 200(| | | |
| | | 20.1/0 | | JITTELENCE | | | | | | | |
| w/ Soil only | | 38.177 | | | | | | - | | |] |
| Soil Weight (air-dried | | 10.007 | | | | | | | 0 2 | 4 6 | |
| w/ Soil and MeBT mix | | 64.652 | | | | | | | Numb | er of Times Extracted | |
| After Initial Extraction | | 62.811 | | 1.841 | | | | | | | |
| of water/MeBT | | | 63.294 | | | | | | | | |
| 1 Meth | , | 62.023 | 63.584 | 1.271 | | | | | | | |
| 2 Meth | | 61.435 | 63.645 | 2.149 | | | | | | | |
| 3 Meth | | 61.941 | 63.359 | 1.704 | | | | | | | |
| 4 Meth | | 61.254 | 63.698 | 2.105 | | | | | | | |
| 5 Meth | | 61.882 | 63.386 | 1.816 | | | | | | | |
| | | | 00000 | | | | | | | | |
| o Mein | | 01.342 | | 2.044 | | | | | | | |
| 100 mg/L 0 | | | 918.0 | | 1537.0 | 4 MeBT | 5 MeBT | | Extraction R | tesults - 100 mg/L | |
| Water 0. | | 7.051 | 708.4 | 7.254 | 17.4 | %11 | 1% | 800 - | | | |
| 1 Meth | | 7.035 | 166.3 | 7.243 | 9.2 | 18% | 1% | C C | • | | ł |
| 2 Meth | | 7.046 | 93.2 | 7.222 | 3.3 | 10% | %0 | - 00/ | | 4 MG | <u>-</u> |
| 3 Meth 24 | | 7 038 | 52.6 | 7.216 | 00 | %9 6% | %0 | 600 | | | BT |
| A Moth | | 1000 | 2.40 | 7 240 | | 30/2 | 200 | , (• | | | |
| 5 Meth 72 | | 7.029 | 6.7 | 7.204 | 0.0 | 1% | %0 | 000 5.,n | | | |
| Sum Recovered | | | 1055.3 | | 29.9 | | | An | | | |
| Percent MeBT Reco | vered | | 114.96% | | 1.95% | | | и) в С | | | |
| Emotv | | 28.502 F | kefilled [| Difference | | | | ere S S S S S S S S S S S S S S S S S S | | | |
| w/ Soil only | | 38.506 | | | | | | 2002 | | | |
| Soil Weight (air-dried | | 10.004 | | | | | | 100 - | | ł | |
| w/ Soil and MeBT mix | | 64.929 | | | | | | c | | | |
| After Initial Extraction | | 63.198 | | 1.731 | | | | 5 | - (| | |
| of water/MeBT | | | 63.955 | | | | | |) 2 Number | 4 6 6 | |
| 1 Meth | | 62.584 | 63.821 | 1.371 | | | | | | I OL LILLES EXHACIEN | |
| 2 Meth | | 61.960 | 63.706 | 1.861 | | | <u>,</u> | | | | |
| 3 Meth | | 62.222 | 64.029 | 1.484 | | | | | | | |
| 4 Meth | | 62.156 | 63.832 | 1.873 | | | | | | | |
| 5 Meth | | 63.010 | | 0.822 | | | | | | | |

Soil Calculation - Treated.xls

80.0 70.0 60.0

50.0

(s*uAm) ธอาA

Extraction Results - 10 mg/L

100.0 90.0

| | | ŀ | | | 1 100 | A MADT | E MADT |
|-----------------------|---------|-----------------------|----------|------------|-------|---------|--------|
| 10 mg/L | 0 | _ | 119.9 | | 1.102 | 4 MeB I | |
| Water | 0.5 | 7.135 | 0.0 | 7.350 | 0.0 | %0 | %0 |
| 1 Meth | 1 hr | 7.134 | 99.83 | 7.321 | 6.1 | 83% | 3% |
| 2 Meth | 6 hr | 7.135 | 0 | 7.35 | 0.0 | %0 | %0 |
| 3 Meth | 24 hr | 7.135 | 0 | 7.35 | 0 | %0 | %0 |
| 4 Meth | 48 hr | 7.135 | 0 | 7.35 | 0 | %0 | %0 |
| 5 Meth | 72 hr | 7.135 | 0 | 7.35 | 0 | %0 | %0 |
| Sum Recovered | | | 9.66 | | 6.1 | | |
| Percent MeBT Re | covered | night international e | 83.26% | | 3.03% | | |
| Additional snike a | Water | 7 948 | 131.9 | | | | |
| cad and at | Water | 8 803 | 7.2 | | | | |
| | Meth 1 | 8.696 | 23.4 | | | | |
| | Meth 1 | 9.613 | 13.7 | | | | |
| | Meth 2 | 7.651 | 27.2 | | | | |
| Emptv | | 27.041 | Refilled | Difference | | | |
| w/ Soil only | | 37.048 | | | | | |
| Soil Weight (air-dri | ied) | 10.007 | | | | | I |
| w/ Soil and MeBT | mix | 63.638 | | | | | |
| After Initial Extract | tion | 61.805 | | 1.833 | | | |
| of water/MeBT | | | 63.804 | | | | |
| 1 Meth | | 62.413 | 63.991 | 1.391 | | | |
| 2 Meth | | 62.288 | 64.112 | 1.703 | | | |
| 3 Meth | | 62.567 | 64.684 | 1.545 | | | |
| 4 Meth | | 62.606 | 63.863 | 2.078 | | | |
| 5 Meth | | 62.918 | 63.456 | 0.945 | | | |
| 6 Meth | | 62.451 | | 1.005 | | | |
| 0 mg/l | 0 | | 4.8 | | 20.5 | 4 MeBT | 5 MeBT |
| Water | 0.5 | 7.264 | 820.3 | 7.475 | 398.7 | 17090% | 1945% |
| 1 Meth | 1 hr | 7.264 | 254.1 | 7.474 | 211.9 | 5294% | 1034% |
| 2 Meth | 6 hr | 7.263 | 107.3 | 7.472 | 136.4 | 2235% | 665% |
| 3 Meth | 24 hr | 7.261 | 42.2 | 7.471 | 79.2 | 879% | 386% |
| 4 Meth | 48 hr | 7.261 | 20.6 | 7.469 | 34.1 | 429% | 166% |
| 5 Meth | 72 hr | | | | | %0 | %0 |

ဖ

 $\frac{2}{N}$ umber of Extractions

0

ŧ

40.0 30.0 20.0 10.0 0.0

| 3 Meth | 24 hr | 7.261 | 42.2 | |
|--------------------|----------|-------|--------|--|
| 4 Meth | 48 hr | 7.261 | 20.6 | |
| 5 Meth | 72 hr | | | |
| Sum Recovered | | | 1244.5 | |
| Percent MeBT Re | scovered | | 25927% | |
| No Additional spik | e | | | |

860.3 4197%

| .xis |
|-------|
| ated |
| Tre |
| - uoi |
| culat |
| Calc |
| Soil |

| | | | | | | | | | | | | 1537 0 | 820.4 | | |
|------------|----------------------------------|-------------------|---------------------|--------------|--------|--------|--------|--------|--------|--------|---|--------|-------|--------|--|
| Difference | | | C |) | 0000 | | 0000 | 0000 | | 0.000 | | | 7 449 | | |
| Refilled | | | | | | | | | | | i | 918.0 | 744.4 | 1 | |
| | | | | | | | | | | | | | 7.244 | 1 20 2 | |
| | 1 | -arrea) 3T mix | action | ЗТ | | | | | | | | 0 | 0.5 | 4 hr | |
| Empty | w/ Soil only Soil Moiobt (oin | w/ Soil and MeE | After Initial Extra | of water/MeE | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | 6 Meth | | Mix | Water | 1 Meth | |



 \sim

Soil Calculation - Treated.xls

HPLC Results

| | Runs | | 4 MeBT | | | 5 MeBT | | |
|-------------|---|---|--|---|---|--|--|--|
| | 1 | 7.372 | 7118.9 | 863.67 | 7.569 | 10981.6 | 821.99 | |
| | 2 | 7.356 | 7139.2 | 866.17 | 7.553 | 11053.6 | 827.70 | |
| | 3 | 7.337 | 7101.4 | 861.52 | 7.534 | 10967.2 | 820.84 | |
| | 4 | 7.301 | 7113.1 | 862.96 | 7.506 | 11019.8 | 825.02 | |
| | 5 | 7.278 | 7168.5 | 869.78 | 7.473 | 11088.9 | 830.51 | |
| | avg | 7.329 | 7128.22 | | 7.527 | 11022.22 | | |
| | conc. | | | 864.82 | | | 825.21 | Interpolated from calibration curve |
| | Standard D | Deviation | 26.35 | | | 50.28 | | |
| | | Average of 4 | & 5 MeBT | | 0.00 | | | |
| | | | | | | | | |
| 100 mg/L | | | | ····· | | | | O 11 |
| | Runs | | 4 MeBT | | | 5 MeBT | | Other |
| | 1 | 7.344 | 718.50 | 78.20 | 8.461 | 17.38 | 0.00 | 7.925 11.02 |
| | 2 | 7.323 | 716.39 | 77.97 | 8.436 | 18.06 | 0.00 | 7.923 11.16 |
| | 3 | 7.301 | 724.66 | 78.87 | 8.410 | 17.70 | 0.00 | 7.923 11.07 |
| | 4 | 7.283 | 718.57 | 78.20 | 8.387 | 18.06 | 0.00 | 7.922 11.11 |
| | 5 | 7.264 | 716.28 | 77.95 | 8.364 | 17.99 | 0.00 | 7.921 11.23 |
| | avg | 7.303 | 718.88 | | 8.412 | 17.84 | | |
| | conc. | | | 78.24 | | | 0.00 | Interpolated from calibration curve |
| | Standard D | eviation | 3.41 | | | 0.30 | | |
| | | Average of 4 | & 5 MeBT | | 0.00 | | | |
| | | | | | | | | |
| 10 mg/L | _ | | | | | | | 0# |
| | Runs | | 4 MeB I | | | 5 MeB I | | |
| | 1 | 7.149 | 101.56 | 8.22 | 7.895 | 6.31 | 0.00 | 7.896 6.13 |
| | 2 | 7.148 | 102.34 | 8.30 | 7.896 | 6.47 | 0.00 | 7.893 6.48 |
| | 3 | 7.148 | 101.79 | 8.25 | 7.892 | 6.47 | 0.00 | 7.090 |
| | | 7 4 4 0 | 100.04 | 0 4 5 | 7 000 | 0 0 4 1 | 0 00 | 1.092 |
| | 4 | 7.149 | 100.81 | 8.15 | 7.895 | 6.84 | 0.00 | 7 905 6 94 |
| | 4 5 | 7.149 | 100.81 102.63 | 8.15 8.33 | 7.895 | 6.84 6.59 | 0.00 0.00 | 7.895 |
| | 4 5 avg | 7.149 7.147 7.148 | 100.81 102.63 101.83 | 8.15 8.33 | 7.895 7.894 7.894 | 6.84 6.59 6.54 | 0.00 | 7.895 6.84 |
| | 4 5 avg conc. | 7.149 7.147 7.148 | 100.81 102.63 101.83 | 8.15 8.33 8.25 | 7.895 7.894 7.894 | 6.84 6.59 6.54 | 0.00 0.00 0.00 | 7.895 6.84 |
| | 4 5 avg conc. Standard D | 7.149 7.147 7.148 | 100.81 102.63 101.83 0.71 | 8.15 8.33 8.25 | 7.895 | 6.84 6.59 6.54 0.20 | 0.00 0.00 0.00 | 7.895 6.84 |
| | 4 5 avg conc. Standard D | 7.149 7.147 7.148 Deviation Average of 4 | 100.81 102.63 101.83 0.71 & 5 MeBT | 8.15 8.33 8.25 | 7.895 7.894 7.894 0.00 | 6.84 6.59 6.54 0.20 | 0.00 0.00 0.00 | 7.895 6.84 |
| 100 mg/L M/ | 4 5 avg conc. Standard D | 7.149 7.147 7.148 Deviation Average of 4 | 100.81 102.63 101.83 0.71 & 5 MeBT | 8.15 8.33 8.25 | 7.895 7.894 7.894 0.00 | 6.84 6.59 6.54 0.20 | 0.00 0.00 0.00 | 7.895 |
| 100 mg/L Me | 4 5 avg conc. Standard D BT / 1000 m Runs | 7.149 7.147 7.148 Deviation Average of 4 | 100.81 102.63 101.83 0.71 & 5 MeBT | 8.15 8.33 8.25 | 7.895 7.894 7.894 | 6.84 6.59 6.54 0.20 | 0.00 0.00 0.00 | 7.895 6.84 Interpolated from calibration curve Other |
| 100 mg/L Me | 4 5 conc. Standard D BT / 1000 m Runs 1 | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752 18 | 8.15 8.33 8.25 | 7.895 7.894 7.894 0.00 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 | 0.00 0.00 0.00 53.84 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 |
| 100 mg/L Me | 4 5 conc. Standard D BT / 1000 m Runs 1 2 | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 | 8.15 8.33 8.25 8.25 | 7.895 7.894 7.894 0.00 7.378 7.378 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 | 0.00 0.00 0.00 53.84 54.13 | 7.895 6.84 Interpolated from calibration curve |
| 100 mg/L Me | 4 5 conc. Standard D BT / 1000 m Runs 1 2 3 | 7.149 7.147 7.148 Deviation Average of 4 mg/L PG 7.187 7.188 7.188 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 | 8.15 8.33 8.25 8.25 81.88 81.89 81.07 | 7.895 7.894 7.894 0.00 7.378 7.378 7.379 7.379 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 | 0.00 0.00 0.00 53.84 54.13 55.21 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 |
| 100 mg/L Me | 4 5 conc. Standard D BT / 1000 m Runs 1 2 3 4 | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 7.188 7.188 7.189 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 754.11 | 8.15 8.33 8.25 8.25 81.88 81.89 81.07 82.09 | 7.895 7.894 7.894 0.00 7.378 7.379 7.379 7.379 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 847.23 | 0.00 0.00 0.00 53.84 54.13 55.21 54.91 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 7.919 12.38 |
| 100 mg/L Me | 4 5 conc. Standard D BT / 1000 m Runs 1 2 3 4 5 | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 7.188 7.188 7.189 7.189 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 754.11 751.02 | 8.15 8.33 8.25 8.25 81.88 81.89 81.07 82.09 81.75 | 7.895 7.894 7.894 0.00 7.378 7.379 7.379 7.379 7.381 7.380 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 847.23 844.21 | 0.00 0.00 0.00 53.84 54.13 55.21 54.91 54.71 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 7.919 12.38 7.918 12.53 |
| 100 mg/L Me | 4 5 2 avg conc. Standard D 2 BT / 1000 m Runs 1 2 3 4 5 5 | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 7.188 7.188 7.189 7.189 7.189 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 754.11 751.02 750.87 | 8.15 8.33 8.25 81.88 81.89 81.07 82.09 81.75 | 7.895 7.894 7.894 0.00 7.378 7.379 7.379 7.379 7.381 7.380 7.379 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 847.23 844.21 841.87 | 0.00 0.00 0.00 53.84 54.13 55.21 54.91 54.71 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 7.919 12.38 7.918 12.53 |
| 100 mg/L Me | 4 5 avg conc. Standard D BT / 1000 m Runs 1 2 3 4 5 5 avg conc | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 7.188 7.188 7.189 7.189 7.189 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 754.11 751.02 750.87 | 8.15 8.33 8.25 8.25 81.88 81.89 81.07 82.09 81.75 81.736 | 7.895 7.894 7.894 0.00 7.378 7.379 7.379 7.379 7.381 7.380 7.379 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 847.23 844.21 844.87 | 0.00 0.00 0.00 53.84 54.13 55.21 54.71 54.71 54.56 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 7.919 12.38 7.919 12.38 7.918 12.53 Interpolated from calibration curve |
| 100 mg/L Me | 4 5 avg conc. Standard D BT / 1000 m Runs 1 2 3 4 5 5 avg conc. Standard D | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 7.188 7.188 7.189 7.189 7.189 7.188 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 754.11 751.02 750.87 3.58 | 8.15 8.33 8.25 8.25 81.88 81.89 81.07 82.09 81.75 81.736 | 7.895 7.894 7.894 0.00 7.378 7.379 7.379 7.381 7.380 7.379 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 847.23 844.21 841.87 860 | 0.00 0.00 0.00 53.84 54.13 55.21 54.71 54.71 54.56 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 7.919 12.38 7.918 12.53 Interpolated from calibration curve |
| 100 mg/L Me | 4 5 conc. Standard D BT / 1000 m Runs 1 2 3 4 5 5 avg conc. Standard D | 7.149 7.147 7.148 Deviation Average of 4 ng/L PG 7.187 7.188 7.188 7.189 7.189 7.189 7.189 7.188 0.0000 7.188 | 100.81 102.63 101.83 0.71 & 5 MeBT 4 MeBT 752.18 752.27 744.78 754.11 751.02 750.87 3.58 & 5 MeBT | 8.15 8.33 8.25 8.25 81.88 81.89 81.07 82.09 81.75 81.736 | 7.895 7.894 7.894 0.00 7.378 7.379 7.379 7.381 7.380 7.379 7.381 7.380 | 6.84 6.59 6.54 0.20 5 MeBT 830.89 835.28 851.76 847.23 844.21 841.87 8.60 | 0.00 0.00 0.00 53.84 54.13 55.21 54.91 54.71 54.56 | 7.895 6.84 Interpolated from calibration curve Other 7.916 12.85 7.919 13.03 7.916 12.41 7.919 12.38 7.918 12.53 Interpolated from calibration curve |

Below LOD

.

Appendix G

Denver International Airport Soil

Sample Analysis

Appendix G

Denver International Airport Soil

4 MeBT

| 4-MeBT Data | | | | | | | | | | all data | with 10 m | g/L | |
|----------------------------|-----------|--------|-----------|---------|-----------|-------------|-----------|----------|-------------|-------------|-----------|---------|---------|
| Moisture Content c | of soil | 0.138 | | | | | | | slope | 0.4471 | 0.2703 | 11 C | 0.447 |
| Outliers identifieo | l in bold | | | | | | | | intercept | 0.2454 | 0.6944 | Kf = | 1.759 |
| | | | | | | | | | r^2 | 0.8995 | 0.8791 | | |
| | | | | w/ Soil | After | Final | | | Initial | Sorbed | 1 point | | |
| | | Vial + | Air-dried | and | Initial | Concentrati | Soil (dry | Vol Soln | Concentrati | Concentrati | Sorption | | |
| Added Concentrati | Vial Only | soil | Soil | MeBT | Extractio | on | wt) (g) | (mL) | uo | uo | Coeff. | log S | Log Cf |
| | b | D | Б | | | ų | Σ | Vs | ö | S | Кd | | |
| | | | | | | mg/L | | | | 6/6ri | ml/g | | |
| 1000 Bottle 1 | 27.983 | 37.988 | 10.005 | 46.476 | 44.697 | 854.17 | 8.792 | 9.701 | 874.94 | 22.916 | 0.027 | 1.36015 | 2.93154 |
| 1000 Bottle 2 | 28.417 | 38.418 | 10.001 | 46.826 | 45.070 | 838.69 | 8.788 | 9.621 | 873.94 | 38.592 | 0.046 | 1.58649 | 2.9236 |
| 1000 Bottle 3 | 27.933 | 37.936 | 10.003 | 46.379 | 44.542 | 835.51 | 8.790 | 9.656 | 874.38 | 42.696 | 0.051 | 1.63039 | 2.92195 |
| 1000 Bottle 4 | 27.893 | 37.901 | 10.008 | 46.429 | 44.603 | 843.82 | 8.794 | 9.742 | 875.42 | 35.002 | 0.041 | 1.5441 | 2.92625 |
| 1000 Bottle 5 | 28.047 | 38.047 | 10.000 | 46.596 | 44.815 | 835.75 | 8.787 | 9.762 | 875.77 | 44.461 | 0.053 | 1.64798 | 2.92208 |
| 100 Bottle 1 | 27.914 | 37.917 | 10.003 | 46.524 | 44.634 | 27.75 | 8.790 | 9.820 | 87.65 | 11.057 | 0.142 | 1.04365 | 1.8907 |
| 100 Bottle 2 | 28.083 | 38.088 | 10.005 | 46.589 | 44.738 | 76.40 | 8.792 | 9.714 | 87.51 | 12.276 | 0.161 | 1.08907 | 1.88309 |
| 100 Bottle 3 | 27.674 | 37.683 | 10.009 | 46.291 | 44.083 | 77.28 | 8.795 | 9.822 | 87.64 | 11.572 | 0.150 | 1.06339 | 1.88807 |
| 100 Bottle 4 | 28.018 | 38.024 | 10.006 | 46.555 | 44.656 | 75.86 | 8.793 | 9.744 | 87.55 | 12.953 | 0.171 | 1.11237 | 1.88001 |
| 100 Bottle 5 | 27.461 | 37.465 | 10.004 | 46.072 | 44.336 | 75.62 | 8.791 | 9.820 | 87.65 | 13.435 | 0.178 | 1.12822 | 1.87864 |
| 10 Bottle 1 | 28.123 | 38.128 | 10.005 | 46.875 | 45.073 | 2:57 | 8.792 | 9.960 | 8.78 | 7.038 | 2.738 | 0.84742 | 0.40993 |
| 10 Bottle 2 | 27.977 | 37.980 | 10.003 | 46.554 | 44.785 | 2.34 | 8.790 | 9.787 | 8.76 | 7.149 | 3.055 | 0.85424 | 0.36922 |
| 10 Bottle 3 | 27.591 | 37.598 | 10.007 | 46.216 | 44.372 | 2.48 | 8.793 | 9.832 | 8.77 | 7.028 | 2.834 | 0.84681 | 0.39445 |
| 10 Bottle 4 | 28.175 | 38.179 | 10.004 | 46.921 | 45.117 | 2.52 | 8.791 | 9.955 | 8.78 | 7.091 | 2.814 | 0.85069 | 0.4014 |
| 10 Bottle 5 | 27.692 | 37.695 | 10.003 | 46.337 | 44.578 | 2.52 | 8.790 | 9.855 | 8.77 | 7.006 | 2.780 | 0.84549 | 0.4014 |
| | | | | | | | | | | | | i | |
| 100 Mix 1 | 27.986 | 37.986 | 10.000 | 46.388 | 44.616 | 73.41 | 8.787 | 9.615 | 87.39 | 15.293 | 0.208 | | |
| 100 Mix 2 | 27.613 | 37.615 | 10.002 | 46.021 | 44.345 | 74.05 | 8.789 | 9.619 | 87.39 | 14.600 | 0.197 | | |
| 100 Mix 3 | 27.847 | 37.856 | 10.009 | 46.370 | 44.665 | 73.93 | 8.795 | 9.728 | 87.52 | 15.034 | 0.203 | | |
| 100 Mix 4 | 27.791 | 37.800 | 10.009 | 46.357 | 44.587 | 73.38 | 8.795 | 9.771 | 87.58 | 15.772 | 0.215 | | |
| 100 Mix 5 | 27.429 | 37.433 | 10.004 | 46.001 | 44.262 | 73.33 | 8.791 | 9.781 | 87.60 | 15.874 | 0.216 | | |
| | | | | | | Ŧ | | | | 0101 | 0101 | | |
| | | | | | | - 000 * | | | | 4.340 | 0+0.4 | | |
| | | | | | | 0001 | | | | 32.014 | 0.034 | | |

4-MeBT

DIA.xls







4-MeBT

Appendix G

Denver International Airport Soil

5 MeBT

| t of soil ed in bold | 0.138 | | | | | | | slope intercept r^2 | all data 0.4915 0.3487 0.9382 | with 10 π 0.3140 0.7943 0.9238 | ıg/L n = Kf = | 0.491 2.232 |
|-------------------------|----------|-----------|--------|-----------|---------------|-----------|----------|---------------------------|---|---|---------------------|----------------|
| | | | and | Initial | Final | | | Initial | Sorbed | 1 point | | |
| | Vial + | Air-dried | MeBT | Extractio | Concentrati 3 | Soil (dry | Vol Soln | Concentratio | Concentrati | Sorption | | |
| | ly soil | Soil | mix | L | - uo | wt) (g) | (mL) | c | on | Coeff. | log S L | og Cf |
| | D | D | | | പ് | Σ | Vs | ö | ა | Кd | | |
| | | | | | mg/L | | | | 6/6rl | b/lm | | |
| | 3 37.988 | 10.005 | 46.476 | 44.697 | 837.55 | 8.792 | 9.701 | 874.94 | 41.256 | 0.049 | 1.61548 | 2.92301 |
| | 7 38.418 | 10.001 | 46.826 | 45.070 | 819.75 | 8.788 | 9.621 | 873.94 | 59.326 | 0.072 | 1.77324 | 2.91368 |
| | 3 37.936 | 10.003 | 46.379 | 44.542 | 810.40 | 8.790 | 9.656 | 874.38 | 70.280 | 0.087 | 1.84683 | 2.9087 |
| - | 3 37.901 | 10.008 | 46.429 | 44.603 | 818.53 | 8.794 | 9.742 | 875.42 | 63.016 | 0.077 | 1.79945 | 2.91303 |
| | 7 38.047 | 10.000 | 46.596 | 44.815 | 808.96 | 8.787 | 9.762 | 875.77 | 74.222 | 0.092 | 1.87053 | 2.90793 |
| 1 N | 4 37.917 | 10.003 | 46.524 | 44.634 | 71.67 | 8.790 | 9.820 | 87.65 | 17.850 | 0.249 | 1.25163 | 1.85534 |
| ~ | 3 38.088 | 10.005 | 46.589 | 44.738 | 70.36 | 8.792 | 9.714 | 87.51 | 18.950 | 0.269 | 1.27761 | 1.84733 |
| ~ | 4 37.683 | 10.009 | 46.291 | 44.083 | 72.54 | 8.795 | 9.822 | 87.64 | 16.865 | 0.232 | 1.22698 | 1.86058 |
| ~ | 3 38.024 | 10.006 | 46.555 | 44.656 | 71.04 | 8.793 | 9.744 | 87.55 | 18.295 | 0.258 | 1.26233 | 1.8515 |
| in | 1 37.465 | 10.004 | 46.072 | 44.336 | 70.76 | 8.791 | 9.820 | 87.65 | 18.864 | 0.267 | 1.27562 | 1.84979 |
| ١Ň | 3 38.128 | 10.005 | 46.875 | 45.073 | 1.95 | 8.792 | 9.960 | 8.78 | 7.740 | 3.969 | 0.88874 | 0.29003 |
| ĸ | 7 37.980 | 10.003 | 46.554 | 44.785 | 1.39 | 8.790 | 9.787 | 8.76 | 8.207 | 5.904 | 0.91416 | 0.14301 |
| ò | 1 37.598 | 10.007 | 46.216 | 44.372 | 1.60 | 8.793 | 9.832 | 8.77 | 8.012 | 5.007 | 0.90372 | 0.20412 |
| ř | 5 38.179 | 10.004 | 46.921 | 45.117 | 1.59 | 8.791 | 9.955 | 8.78 | 8.144 | 5.122 | 0.91083 | 0.2014 |
| 9, | 2 37.695 | 10.003 | 46.337 | 44.578 | 1.58 | 8.790 | 9.855 | 8.77 | 8.060 | 5.101 | 0.90635 | 0.19866 |
| | | | | | | | | | | | | |
| ĕ | 3 37.986 | 10.000 | 46.388 | 44.616 | 69.18 | 8.787 | 9.615 | 87.39 | 19.922 | 0.288 | | |
| ÷- | 3 37.615 | 10.002 | 46.021 | 44.345 | 69.48 | 8.789 | 9.619 | 87.39 | 19.601 | 0.282 | | |
| 4 | 7 37.856 | 10.009 | 46.370 | 44.665 | 69.37 | 8.795 | 9.728 | 87.52 | 20.077 | 0.289 | | |
| ò | 1 37.800 | 10.009 | 46.357 | 44.587 | 69.08 | 8.795 | 9.771 | 87.58 | 20.549 | 0.297 | | |
| Χí | 9 37.433 | 10.004 | 46.001 | 44.262 | 69.16 | 8.791 | 9.781 | 87.60 | 20.514 | 0.297 | | |

.

5-MeBT

DIA.xls

6.227 0.054

6.227 54.499

1 1000







| 1000 ma/L | | | 8225.7 | | 13224.4 | 4 MeBT | 5 MeBT | | |
|--------------------------|--------------|-----------------------------------|------------|------------|---------|--------|--------|--------------------|--|
| Water | 5 | 6.966 | 7029.4 | 7.145 | 11161.8 | 85% | 84% | | |
| 1 Math | , 1 | 6 761 | 554 1 | 6 031 | 1147.2 | 2000 | 0% | | MeBT isomer Extraction Curve - 1000 m A MeBT |
| 2 Meth | 1 | 6.752 | 277.9 | 6.922 | 441.7 | 3% | 3% | 1 | |
| 3 Meth 2 | 4 hr | 6.750 | 142.9 | 6.919 | 209.2 | 2% | 2% | - | |
| 4 Meth | 8 hr | 6.744 | 97.8 | 6.914 | 138.2 | 1% | 1% | 5 | 0000 |
| 5 Meth 7. | 2 hr | 6.731 | 45.6 | 6.91 | 72.4 | 1% | 1% | (s | |
| Sum Recovered | | | 8147.7 | | 13170.5 | | | ມ ≩ "n ∖ | 2000 |
| Percent MeBT Reco | overed | | 99.05% | | 99.59% | | | √ɯ) | 2000 |
| Empty | | 28.047 F | Refilled D | lifference | | | | 63 | |
| w/ Soil only | | 38.047 | | | | | | ٦A | |
| Soil Weight (air-dried | ÷ | 10.000 | | | | | | •• | 2000 |
| w/ Soil and MeBT m | × | 46.596 | | | | | | | |
| After Initial Extractio | Ę | 44.815 | | 1.781 | | | | | |
| of water/MeBT | | | 46.156 | | | | | | 0 2 4 6 |
| 1 Meth | | 44.360 | 46.612 | 1.796 | | | | | Number of Extractions |
| 2 Meth | | 44.849 | 46.541 | 1.763 | | | | | |
| 3 Meth | | 44.716 | 46.083 | 1.825 | | | | | |
| 4 Meth | | 44.369 | 46.354 | 1.714 | | | | | |
| 5 Meth | | 44.455 | | 1.899 | | | | | |
| 100 mg/L | | | 918.0 | | 1537.0 | 4 MeBT | 5 MeBT | | |
| Water 0. | 2 | 7.055 | 686.3 | 7.241 | 1071.7 | 75% | 20% | | Mebi Isomer Extraction Curve - 100 mg/L |
| 1 Meth | hr | 7.045 | 105.6 | 7.229 | 214.6 | 12% | 14% | · | 1300 |
| 2 Meth | hr | 7.044 | 61.8 | 7.230 | 132.6 | 7% | %6 | | |
| 3 Meth | 4 hr | 7.040 | 32.7 | 7.223 | 67.3 | 4% | 4% | | |
| 4 Meth | 8 hr | 7.038 | 20.6 | 7.221 | 25.1 | 2% | 2% | (: | |
| 5 Meth 7. | 2 hr | 7.036 | 10.4 | 7.218 | 11.9 | 1% | 1% | s _* r | |
| Sum Recovered | | an bala and a and adams, and that | 917.4 | | 1523.2 | | | ı¥u | • · · · |
| Percent MeBT Rec | overed | | 99.93% | | 99.10% | | | u) e | |
| Empty | | 27.461 F | Sefilled D | Vifference | | | | Are | |
| w/ Soil only | | 37.465 | | | | | | , | |
| Soil Weight (air-dried | (| 10.004 | | | | | | | |
| w/ Soil and MeBT m | × | 46.072 | | | | | | | |
| After Initial Extraction | ŗ | 44.336 | | 1.736 | | | | _ | 0 2 4 6 |
| of water/MeBT | | | 46.554 | | | | | | Number of Extractions |
| 1 Meth | | 44.381 | 46.359 | 2.173 | | | | | |
| 2 Meth | | 44.414 | 46.523 | 1.945 | | | | | |
| 3 Meth | | 44.652 | 46.691 | 1.871 | | | | | |
| 4 Meth | | 44.698 | 46.578 | 1.993 | | | | | |
| 5 Meth | | 44.385 | | 2.193 | | | | | |

.

Soil Calculation - DIA.xls

| | 4 | | 0 077 | - | 1 100 | 10-11 |
|-----------------------|-----------|----------|----------|------------|--------|---------|
| 10 mg/L | 0 | | 119.9 | | 1.102 | 4 Meb I |
| Water | 0.5 | 7.156 | 46.4 | 7.346 | 63.9 | 39% |
| 1 Meth | 1 hr | 7.145 | 25.4 | 7.334 | 39.2 | 21% |
| 2 Meth | 6 hr | 7.159 | 11.3 | 7.352 | 17.5 | %6 |
| 3 Meth | 24 hr | 7.141 | 5.5 | 7.346 | 7.0 | 5% |
| 4 Meth | 48 hr | 7.135 | 0 | 7.35 | 0 | %0 |
| 5 Meth | 72 hr | 7.135 | 0 | 7.35 | 0 | %0 |
| Sum Recovered | | | 88.6 | | 127.6 | |
| Percent MeBT R | lecovered | | 73.89% | gai - | 63.45% | |
| Additional spike a | a Water | 7.894 | 6.2 | | | |
| | Meth 1 | 7.896 | 3.0 | | | |
| | | | | | | |
| Empty | | Z7.692 F | Setilled | Uitterence | | |
| w/ Soil only | | 37.695 | | | | |
| Soil Weight (air-d | Iried) | 10.003 | | | | |
| w/ Soil and MeB1 | T mix | 46.337 | | | | |
| After Initial Extract | ction | 44.578 | | 1.759 | | |
| of water/MeB1 | F | | 46.648 | | | |
| 1 Meth | | 44.857 | 45.959 | 1.791 | | |
| 2 Meth | | 44.114 | 46.223 | 1.845 | | |
| 3 Meth | | 44.452 | 46.138 | 1.771 | | |
| 4 Meth | | 44.501 | 46.161 | 1.637 | | |
| 5 Meth | | 44.499 | | 1.662 | | |
| Mix | 0 | | 918.0 | | 1537.0 | 4 MeBT |
| Water | 0.5 | 7.311 | 671.2 | 7.513 | 1057.4 | 73% |
| 1 Meth | 1 hr | 7.272 | 101.2 | 7.472 | 301.5 | 11% |
| 2 Meth | 6 hr | 7.261 | 65.3 | 7.460 | 85.2 | 7% |
| 3 Meth | 24 hr | 7.255 | 36.4 | 7.457 | 43.0 | 4% |
| 4 Meth | 48 hr | 7.251 | 12.5 | 7.453 | 24.7 | 1% |
| 5 Meth | 72 hr | | | | | %0 |
| Sum Recovered | - | | 886.6 | | 1511.8 | |
| Percent MeBT R | Recovered | | 96.58% | ; | 98.36% | |
| No Additional spi | ike | | | | | |
| Empty | | 27.429 F | Refilled | Difference | | |

| | | | | | | | | | | | | | | | ~ | | | |
|---|--------|--------|--------|--------|----------|------------------|---------|-------|------------|---------|---------|---|---------|--------|--------|--------|--------|-------|
| | 1800.0 | 1600.0 | 0.0001 | 1400.0 | 1200.0 - | | 1000.0 | 800.0 | 0000 | - 0.00a | 400.0 - | | - n.u.z | 0.0 |) | | | |
| | | | | | (s | s _* n | An | u) e | s91 | A | | | | | | | | |
| | 20% | %9 | 3% | 2% | %0 | | | | | | | | | | | | | |
| | 11% | 7% | 4% | 1% | %0 | | | | | | | | | | | | | |
| | 301.5 | 85.2 | 43.0 | 24.7 | | 1511.8 | 98.36% | | | | | | | | | | | |
| | 7.472 | 7.460 | 7.457 | 7.453 | | | ξ. , | | Difference | | | | 1.739 | | 1.861 | 2.276 | 1.835 | 1.763 |
| ; | 101.2 | 65.3 | 36.4 | 12.5 | | 886.6 | 96.58% | | Refilled | | | | | 46.152 | 46.265 | 45.932 | 46.115 | |
| ; | 272 | 261 | 255 | 251 | | | | | 29 | 33 | 4 | 5 | 82 | | 291 | 989 | 260 | 352 |

27.429 37.433 10.004 46.001 44.262

Soil Weight (air-dried) w/ Soil and MeBT mix After Initial Extraction

Empty w/ Soil only







Soil Calculation - DIA.xls

44.291 43.989 44.097 44.352

of water/MeBT 1 Meth 2 Meth 3 Meth 4 Meth
Extraction

| 0 mg/L | 0 | 4.8 | 20.5 | 4 MeBT | 5 MeBT | |
|------------------|---------------------------|-----------------------|-------|--------|--------|--|
| Water | 0.5 | 0.0 | 0.0 | %0 | %0 | |
| 1 Meth | 1 hr | 0.0 | 0.0 | %0 | %0 | |
| 2 Meth | 6 hr | 0.0 | 0.0 | %0 | %0 | |
| 3 Meth | 24 hr | | | %0 | %0 | |
| 4 Meth | 48 hr | | | %0 | %0 | |
| 5 Meth | 72 hr | | | %0 | %0 | |
| Sum Recovered | | 0 | 0 | | | |
| Percent MeBT F | tecovered | %00.0 | 0.00% | | | |
| No Additional sp | ke - Zero result after ba | ackground subtraction | | | | |

| Difference | | | | 0.00 | | 0.000 | 0.000 | 0.000 | 0000 |
|------------|--------------|-------------------------|----------------------|--------------------------|---------------|--------|--------|--------|--------|
| Refilled | | | | , | | | | | |
| Empty | w/ Soil only | Soil Weight (air-dried) | w/ Soil and MeBT mix | After Initial Extraction | of water/MeBT | 1 Meth | 2 Meth | 3 Meth | 4 Meth |

.

· ·

Soil Calculation - DIA.xls

| Runs 4 MeBT 5 MeBT 1 9,252 70417 654.17 10177 837.55 2 9,758 690.54 838.69 10.424 10942.4 819.75 3 7.497 6680.2 835.51 7.702 10933.0 818.53 3 7.444 6892.2 835.75 7.644 10942.24 819.04 avg 8.393 6897.44 8.791 10942.24 819.04 conc. 841.59 819.04 Interpolated from calibration curve Standard Deviation 64.39 143.52 Average of 4 & 5 MeBT 1 7.613 71.47 77.75 7.817 1100.56 71.67 2 7.583 702.03 76.40 7.787 1083.62 70.36 3 7.553 702.03 76.40 7.787 1093.98 71.04 4 7.526 73.66 7.267 1089.74 70.76 20.5 conc. 7.497 644.92 75.62< | 1000 mg/L | | | | | | | | |
|--|-------------|------------|---------------------------|------------------|--------|--------|----------|--------|---------------------------------------|
| 1 9.825 7041.7 864.17 10.517 11177.7 837.55 2 9.758 6900.4 836.51 7.702 10835.6 810.40 4 7.465 6830.2 835.51 7.702 10835.6 810.40 4 7.465 6830.2 835.57 7.644 10932.0 618.53 5 7.444 6830.2 835.57 7.644 10837.5 808.96 avg 8.398 6937.44 8.791 10982.4 819.04 conc. 64.39 143.52 408.79 819.04 Average of 4.8.5 MeBT 100 mg/L Runs 4 MeBT 7.756 7.817 1100.366 71.67 2 7.563 710.10 77.28 1089.74 70.76 70.36 3 7.554 703.76 7.766 1093.98 71.42 10.57 2.64 4 7.525 697.26 7.366 1087.47 1.07.75 10.54 1.303 Average of 4.8.5 MeBT 7.362 7.658 | - | Runs | | 4 MeBT | | | 5 MeBT | | |
| 2 9.758 6905.4 833.69 10.424 10942.4 819.75 3 7.497 68902 835.51 7.702 10935.6 810.40 4 7.465 6957.7 843.82 7.667 10938.0 818.53 5 7.444 6892.2 835.75 7.644 10937.6 606.96 avg 6.398 6937.4 841.59 819.04 toom.c. 843 9 143.52 Average of 4.8.5 MeBT 830.31 100 mg/L Runs 7.653 702.03 76.40 7.767 1083.82 70.36 3 7.553 702.03 76.40 7.767 1083.82 70.36 3 7.552 697.26 77.86 1099.74 70.76 avg 7.554 703.76 77.56 1099.74 70.76 avg 7.554 703.76 77.56 1097.57 conc. 75.54 703.76 77.56 1097.57 conc. 75.54 70.376 71.27 10 mg/L Runs 1 7.167 52.14 2.57 7.367 80.541 1.95 5 7.128 51.78 80.541 1.95 5 7.128 51.78 80.542 1.60 Average of 4.8.5 MeBT 73.93 10 mg/L Runs 2 7.157 50.68 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.335 77.06 1.58 avg 7.146 51.59 7.335 77.06 1.59 5 7.128 51.78 2.57 7.367 80.541 1.95 5 7.128 51.78 2.57 73.35 77.06 1.59 3 7.146 51.59 7.335 77.06 1.59 5 7.128 51.78 2.57 7.367 80.541 1.95 5 7.128 51.78 2.57 7.367 80.541 1.95 5 7.128 51.78 2.57 7.367 80.541 1.95 7.924 1.335 77.06 100 mg/L MeBT / 1000 mg/L PG Runs 1 7.189 674.677 73.41 7.380 1065.51 69.18 3 7.146 51.59 7.337 77.06 100 mg/L MeBT / 1000 mg/L PG Runs 1 7.189 674.677 73.41 7.380 1065.51 69.18 7.924 13.35 7.925 13.03 3 7.146 674.677 73.41 7.380 1065.51 69.18 7.924 13.35 7.924 13.35 7. | | 1 | 9.825 | 7041.7 | 854.17 | 10.517 | 11177.7 | 837.55 | |
| 3 7.467 690.2 835.51 7.702 10935.6 810.40 4 7.465 6957.7 843.82 7.667 10938.0 818.53 5 7.444 6892.2 835.75 7.644 10817.5 808.96 avg 8.398 6937.44 8.791 10942.24 conc. 841.59 819.04 Interpolated from calibration curve Standard Deviation 64.39 143.52 Average of 4 & 5 MeBT 830.31 100 mg/L Runs 4 MeBT 77.75 7.817 1103.56 71.67 2 7.583 702.03 76.40 7.787 1083.62 71.03 5 7.553 710.10 77.28 7.756 1116.93 72.54 4 7.525 697.26 75.86 7.726 1093.98 71.04 5 7.749 609.92 75.68 7.726 1093.98 71.04 5 7.7457 609.92 75.68 77.756 1097.57 conc. 76.58 71.27 standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 73.93 10 mg/L Runs 4 MeBT 73.93 10 mg/L 7146 617.67 73.41 7.380 1065.51 69.18 7.126 51.76 73.94 73.36 7.306 10.68 7.126 51.76 4.07 73.93 77.06 8 7.127 1000 mg/L PG Runs 4 MeBT 2.05 100 mg/L MeBT 71000 mg/L PG Runs 7.146 676.63 7.379 1066.65 7.924 13.33 7.924 13.33 7.924 13.33 7.924 15.75 7.924 | | 2 | 9.758 | 6905.4 | 838.69 | 10.424 | 10942.4 | 819.75 | |
| 4 7.465 6957.7 843.82 7.667 10938.0 815.35 5 7.444 6892.2 835.75 7.644 10942.24 803.96 avg 6.398 6937.44 6.791 10942.24 803.96 conc. 841.59 819.04 Interpolated from calibration curve Standard Deviation 64.39 143.52 Average of 4 & 5 MeBT 830.31 100 mg/L 7.653 702.03 76.40 7.787 108.56 71.67 2 7.583 702.03 76.64 7.786 1169.39 72.54 3 7.555 707.26 7.766 1169.39 72.54 4 7.525 694.92 75.62 7.696 1089.74 70.76 conc. 7.654 703.76 71.27 interpolated from calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 7.393 10 mg/L 1 7.167 52.41 2.57 7.337 80.54 1.59 2 7.157 50.63 2.34 7.345 <th></th> <th>3</th> <th>7.497</th> <th>6890.2</th> <th>835.51</th> <th>7.702</th> <th>10835.6</th> <th>810.40</th> <th></th> | | 3 | 7.497 | 6890.2 | 835.51 | 7.702 | 10835.6 | 810.40 | |
| 5 7.444 6992.21 835.76 7.644 10842.24 80.96 avg 8.398 6937.44 8.791 10942.24 819.04 conc. 841.59 819.04 1043.52 819.04 Average of 4 & 5 MeBT 830.31 100 mg/L Runs 4 MeBT 7.613 714.471 77.75 7.817 1103.56 71.67 1 7.613 714.471 77.75 7.817 1103.56 71.67 2 7.553 710.10 77.28 7.756 1116.93 72.54 3 7.554 703.76 7.726 1093.96 71.04 5 7.497 694.92 75.58 70.57 1097.57 conc. 7.554 703.76 7.756 1097.57 100 avg 7.145 51.57 2.48 7.333 7.642 1.95 2 7.157 50.63 2.34 7.345 1.95 1.95 3 7.145 51.57 2. | | 4 | 7.465 | 6957.7 | 843.82 | 7.667 | 10938.0 | 818.53 | |
| avg 8.398 6937.44 8.791 10942.24 etherpolated from calibration curve Standard Deviation 64.39 143.52 Average of 4 & 5 MeBT 830.31 100 mg/L Runs 4 MeBT 7.613 714.47 77.755 7.817 1103.56 71.67 2 7.633 702.03 76.40 7.787 1003.62 70.36 3 7.553 710.10 77.28 7.786 1116.93 72.54 4 7.525 597.26 75.62 7.786 1099.74 70.76 avg 7.554 703.76 1099.77 10.07 10 10 8tandard Deviation 8.34 1 13.03 Average of 4 & 5 MeBT 13.03 Average of 4 & 5 MeBT 73.93 76.52 1.93 1 1.167 1 7.167 52.47 7.357 80.54 1.95 3 7.145 51.57 2.48 7.333 76.82 1.60 4 7.143< | | 5 | 7.444 | 6892.2 | 835.75 | 7.644 | 10817.5 | 808.96 | |
| conc. 841.59 819.04 Interpolated from calibration curve Standard Deviation 64.39 143.52 Average of 4 & 5 MeBT 830.31 100 mg/L Runs 4 MeBT 5 MeBT 1 7.613 714.47 77.75 2 7.553 702.03 76.40 7.787 2 7.553 710.10 77.28 77.56 3 7.552 708.40 7.087 1083.62 70.36 5 7.497 694.92 75.56 77.26 1093.96 71.04 5 7.497 694.92 75.56 71.27 Interpolated from calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 73.93 10 mg/L Runs 4 MeBT 5 MeBT 73.93 74.55 7.367 80.54 1.95 2 7.157 50.63 2.34 7.345 74.54 1.39 3 7.145 51.87 2.48 7.33 77.66 | | avg | 8.398 | 6937.44 | | 8.791 | 10942.24 | | |
| Standard Deviation 64.39 143.52 Average of 4 & 5 MeBT 830.31 100 mg/L Runs 4 MeBT 5 MeBT 1 7.613 714.47 77.75 7.817 1003.62 70.36 3 7.553 70.010 77.728 7.756 116.39 72.54 4 7.552 697.26 75.68 77.756 1108.32 70.76 avg 7.554 70.376 77.56 1089.74 70.76 conc. 76.58 77.26 10.39.74 70.76 conc. 76.58 71.27 Interpolated from calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 1.303 Average of 4 & 5 MeBT 73.93 76.62 1.60 4 7.137 3 7.145 51.67 2.48 7.333 76.62 1.60 4 7.137 50.63 2.23 7.345 77.56 1.65 5 7.128 51.77 | | conc. | | | 841.59 | | | 819.04 | Interpolated from calibration curve |
| Average of 4 & 5 MeBT 830.31 100 mg/L Runs 4 MeBT 5 MeBT 1 7.613 714.47 77.75 7.817 1103.56 71.67 2 7.583 702.03 76.40 7.787 1083.62 70.36 3 7.553 710.10 77.26 7.756 1116.93 72.54 5 7.497 694.92 75.62 7.696 1089.74 70.76 conc. 75.54 703.76 7.756 107.756 71.27 Interpolated from calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 1.303 Average of 4 & 5 MeBT 7.167 50.63 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.333 76.82 1.60 1.58 2 7.157 50.63 2.34 7.345 74.54 1.39 1.59 3 7.146 51.59 7.322 76.73 1.59 1.58 | | Standard I | Deviation | 64.39 | | | 143.52 | | |
| 100 mg/L Runs 4 MeBT 5 MeBT 1 7.613 714.47 77.75 7.817 1103.56 71.67 2 7.583 702.03 76.40 7.787 1083.62 70.36 3 7.553 701.01 77.28 7.756 1116.93 72.54 4 7.525 697.26 75.86 70.86 1093.98 71.04 5 7.497 694.92 75.62 7.696 1099.74 70.76 conc. 76.58 71.27 Interpolated from calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 73.93 10 mg/L 1 7.167 52.14 2.57 7.357 80.54 1.99 2 7.157 50.63 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.332 76.62 1.69 4 7.134 51.59 2.34 7.317 76.68 1.59 | | | Average of 4 | & 5 MeBT | | 830.31 | | | |
| Runs 4 MeBT 5 MeBT 1 7.613 714.47 77.75 71.817 1103.56 71.67 2 7.583 702.03 76.40 7.787 1083.62 70.36 3 7.553 710.10 77.28 7.756 1116.93 72.54 4 7.525 697.26 75.58 77.276 1093.98 71.04 5 7.497 694.92 75.62 7.696 1089.74 70.76 avg 7.554 703.76 7.755 1097.57 Tococ. 76.58 71.27 Interpolated from calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 73.93 Tomg/L Runs 4 MeBT 5 MeBT 1.95 2 7.167 52.14 2.57 7.327 70.68 1.58 3 7.145 51.78 2.52 7.327 70.68 1.58 4 <t< td=""><td>100 mg/L</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<> | 100 mg/L | | | | | | | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | Runs | | 4 MeBT | | | 5 MeBT | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | 1 | 7.613 | 714.47 | 77.75 | 7.817 | 1103.56 | 71.67 | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | 2 | 7.583 | 702.03 | 76.40 | 7.787 | 1083.62 | 70.36 | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | 3 | 7.553 | 710.10 | 77.28 | 7.756 | 1116.93 | 72.54 | |
| $ \begin{array}{ c c c c c c c c c c c c c c c c c c c$ | | 4 | 7.525 | 697.26 | 75.86 | 7.726 | 1093.98 | 71.04 | |
| $\begin{array}{ c c c c c c c c c c c c c c c c c c c$ | | 5 | 7.497 | 694.92 | 75.62 | 7.696 | 1089.74 | 70.76 | |
| Conc. 76.58 71.27 Interpolated from Calibration curve Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 73.93 10 mg/L Runs 4 MeBT 5 MeBT 1 7.167 52.14 2.57 7.357 80.54 1.95 2 7.157 50.63 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.333 76.82 1.60 4 7.134 51.83 2.52 7.327 76.68 1.58 2 7.146 51.59 7.335 77.06 1.62 conc. 2.49 1.62 1.62 1.62 1.62 Standard Deviation 0.57 2.17 2.17 Average of 4 & 5 MeBT 2.05 7.189 680.59 7.380 1065.51 69.18 7.924 13.35 7.924 13.35 7.924 13.35 7.924 13.25 5 7.189 679.46 73.33 7.379 1066.45 <td></td> <td>avg</td> <td>7.554</td> <td>703.76</td> <td></td> <td>7.756</td> <td>1097.57</td> <td></td> <td></td> | | avg | 7.554 | 703.76 | | 7.756 | 1097.57 | | |
| Standard Deviation 8.34 13.03 Average of 4 & 5 MeBT 73.93 10 mg/L Runs 4 MeBT 5 MeBT 1 7.167 52.14 2.57 7.357 80.54 1.95 2 7.157 50.63 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.333 76.82 1.60 4 7.134 51.83 2.52 7.322 76.73 1.59 5 7.128 51.77 2.48 7.335 77.06 conc. 2.49 1.62 1.62 Standard Deviation 0.57 2.17 2.17 Average of 4 & 5 MeBT 2.05 Other Other 1 7.189 674.67 7.3.41 7.380 1065.51 69.18 7.924 13.35 7.924 13.35 2 7.189 674.67 7.3.31 7.380 1066.45 69.37 2 7.189 674.67 7.3.31 7.380 1066.45 69.37 3 7.189 676.63 7.379 1066.65 7.924 15.75 5 7.189 | | conc. | - | | 76.58 | | | /1.2/ | Interpolated from calibration curve |
| Average of 4 & 5 MeB1 73.93 10 mg/L Runs 4 MeBT 5 MeBT 1 7.167 52.14 2.57 7.357 80.54 1.95 2 7.157 50.63 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.333 76.82 1.60 4 7.134 51.83 2.52 7.317 76.68 1.58 avg 7.146 51.59 7.335 77.06 1.62 Interpolated from calibration curve Standard Devlation 0.57 2.17 Average of 4 & 5 MeBT 2.05 1.62 Interpolated from calibration curve 1 7.189 674.67 73.41 7.380 1065.51 69.18 7.924 13.35 2 7.189 674.67 73.41 7.380 1065.51 69.37 7.924 13.35 3 7.189 674.67 73.33 7.379 1068.45 69.37 7.924 13.35 3 7.189 674.67 73.33 7.379 1066.65 7.924 15.75 | | Standard I | Deviation | 8.34 | | | 13.03 | | |
| $ \begin{array}{ c c c c c c c c c c c c c c c c c c c$ | | | Average of 4 | & 5 MeB I | | 73.93 | | | |
| Runs 4 MeBT 5 MeBT 1 7.167 52.14 2.57 7.357 80.54 1.95 2 7.157 50.63 2.34 7.345 74.54 1.39 3 7.145 51.57 2.48 7.333 76.82 1.60 4 7.134 51.83 2.52 7.317 76.68 1.58 avg 7.146 51.59 7.335 77.06 1.62 1.62 conc. 2.49 1.62 1.62 1.62 1.62 1.62 Standard Deviation 0.57 2.17 2.17 Average of 4 & 5 MeBT 2.05 1.62 100 mg/L MeBT / 1000 mg/L PG 7.189 674.67 73.41 7.380 1065.51 69.18 7.924 13.35 2 7.189 674.67 73.33 7.379 1068.45 69.37 7.925 12.89 3 7.189 674.44 73.38 7.380 1063.99 69.08 7.924 15.75 | 10 mg/L | | | | ······ | ····· | | | |
| $ \begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | Runs | L | 4 MeBT | | | 5 MeBT | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | 1 | 7.167 | 52.14 | 2.57 | 7.357 | 80.54 | 1.95 | |
| 3 7.145 51.57 2.48 7.333 76.82 1.60 4 7.134 51.83 2.52 7.322 76.73 1.59 5 7.128 51.78 2.52 7.317 76.68 1.58 avg 7.146 51.59 7.335 77.06 1.59 conc. 2.49 1.62 Interpolated from calibration curve Standard Deviation 0.57 2.17 1.62 Average of 4 & 5 MeBT 2.05 1.62 Interpolated from calibration curve Other 1 7.189 674.67 73.41 7.380 1065.51 69.18 7.924 13.35 2 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 3 7.189 674.44 73.38 7.380 1063.99 69.08 7.924 15.75 5 7.187 673.99 73.33 7.379 1066.65 7.924 15.75 5 7.189 676.63 7.379 1066.65 69.25 Interpolated from | | 2 | 7.157 | 50.63 | 2.34 | 7.345 | 74.54 | 1.39 | |
| $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | | 3 | 7.145 | 51.57 | 2.48 | 7.333 | 76.82 | 1.60 | |
| $ \begin{array}{c c c c c c c c c c c c c c c c c c c $ | | 4 | 7.134 | 51.83 | 2.52 | 7.322 | 76.73 | 1.59 | |
| avg 7.146 51.59 7.335 77.06 conc. 2.49 1.62 Interpolated from calibration curve Standard Deviation 0.57 2.17 Average of 4 & 5 MeBT 2.05 100 mg/L MeBT / 1000 mg/L PG Conc. 5 MeBT 1 7.189 674.67 73.41 7.380 1065.51 69.18 2 7.189 680.59 74.05 7.380 1070.15 69.48 3 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 4 7.189 674.44 73.38 7.380 1065.14 69.16 7.924 13.35 3 7.189 674.44 73.38 7.379 1068.45 69.37 7.925 12.89 4 7.189 676.63 7.379 1066.65 7.924 15.75 5 7.187 673.99 7.333 7.379 1066.65 7.924 16.23 avg 7.189 676 | | 5 | 7.128 | 51.78 | 2.52 | 7.317 | 76.68 | 1.58 | |
| Conc. 2.49 1.62 Interpolated from calibration curve Standard Deviation 0.57 2.17 Average of 4 & 5 MeBT 2.05 100 mg/L MeBT / 1000 mg/L PG 0ther 1 7.189 674.67 73.41 7.380 1065.51 69.18 2 7.189 680.59 74.05 7.380 1070.15 69.48 7.924 13.35 3 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 4 7.189 674.44 73.38 7.380 1063.99 69.08 7.924 15.75 5 7.187 673.99 73.33 7.379 1066.65 7.924 16.23 avg 7.189 676.63 7.379 1066.65 7.924 16.23 conc. 73.62 69.25 69.25 10.12 10.23 10.23 Average of 4 & 5 MeBT 71.44 71.44 71.44 10.23 10.23 10.23 | | avg | 7.146 | 51.59 | | 7.335 | //.06 | 4:00 | Internalistad from collibration or mo |
| Standard Deviation 0.57 2.17 Average of 4 & 5 MeBT 2.05 100 mg/L MeBT / 1000 mg/L PG 0ther 1 7.189 674.67 2 7.189 674.67 3 7.189 679.46 3 7.189 679.46 4 7.189 679.46 5 7.189 679.46 679.46 73.93 7.379 1068.45 69.37 7.924 13.35 7.925 12.89 3 7.189 679.46 73.93 4 7.189 674.44 73.38 7.380 1065.14 69.08 7.924 5 7.187 673.99 7.333 7.362 69.25 Interpolated from calibration curve Standard Deviation 3.13 2.56 | | conc. | <u> </u> | 0.57 | 2.49 | | 0.47 | 1.02 | |
| Ioo mg/L MeBT / 1000 mg/L PG SMeBT Other 1 7.189 674.67 73.41 7.380 1065.51 69.18 7.924 13.35 2 7.189 680.59 74.05 7.380 1070.15 69.48 7.928 13.03 3 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 4 7.189 674.44 73.38 7.380 1065.14 69.08 7.924 15.75 5 7.187 673.99 73.33 7.379 1066.65 7.924 16.23 avg 7.189 676.63 7.379 1066.65 7.924 16.23 conc. 73.62 69.25 Interpolated from calibration curve Standard Deviation 3.13 2.56 2.56 | | Standard I | Jeviation Average of 4 | 0.57 & 5 MeBT | | 2.05 | 2.17 | | |
| Runs 4 MeBT 5 MeBT Other 1 7.189 674.67 73.41 7.380 1065.51 69.18 7.924 13.35 2 7.189 680.59 74.05 7.380 1070.15 69.48 7.924 13.35 3 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 4 7.189 674.44 73.38 7.380 1065.14 69.18 7.924 15.75 5 7.187 673.99 73.33 7.379 1066.65 7.924 16.23 avg 7.189 676.63 7.379 1066.65 69.25 Interpolated from calibration curve Standard Deviation 3.13 2.56 2.56 4.5 5.6 71.44 | 400 | DT / 4000 | | | | | | | |
| Mills 4 MeB 1 5 MeB 1 0 MeB 1 1 7.189 674.67 73.41 7.380 1065.51 69.18 7.924 13.35 2 7.189 680.59 74.05 7.380 1070.15 69.48 7.928 13.03 3 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 4 7.189 674.44 73.38 7.380 1063.99 69.08 7.924 15.75 5 7.187 673.99 73.33 7.378 1065.14 69.16 7.924 16.23 avg 7.189 676.63 7.379 1066.65 7.924 16.23 conc. 73.62 69.25 Interpolated from calibration curve 16.23 Standard Deviation 3.13 2.56 2.56 11.44 11.44 | 100 mg/L Me | Bune | | | | | 5 MoBT | | Other |
| 1 1.165 0.14.07 10.41 1.300 1000.31 031.10 2 7.189 680.59 74.05 7.380 1070.15 69.48 7.928 13.03 3 7.189 679.46 73.93 7.379 1068.45 69.37 7.925 12.89 4 7.189 674.44 73.38 7.380 1063.99 69.08 7.924 15.75 5 7.187 673.99 73.33 7.379 1066.65 7.924 16.23 avg 7.189 676.63 7.379 1066.65 7.924 16.23 conc. 73.62 69.25 Interpolated from calibration curve Standard Deviation 3.13 2.56 Average of 4 & 5 MeBT 71.44 71.44 | | 1 | 7 190 | 674 67 | 73.41 | 7 380 | 1065 51 | 60.18 | 7 024 13 35 |
| 2 7.165 000.35 74.03 7.300 1010.15 00.40 1.320 | | י 2 | 7.109 | 680.59 | 74.05 | 7 380 | 1070 15 | 69.10 | 7 928 13 03 |
| 4 7.189 674.44 73.38 7.380 1063.99 69.08 7.924 15.75 5 7.187 673.99 73.33 7.378 1065.14 69.16 7.924 16.23 avg 7.189 676.63 7.379 1066.65 1000000000000000000000000000000000000 | | 2 | 7.109 | 679.46 | 73.03 | 7.300 | 1068.45 | 69 37 | 7 925 12 89 |
| | | 4 | 7.109 | 674 44 | 73 38 | 7 380 | 1063 99 | 69.08 | 7 924 15 75 |
| avg 7.189 676.63 7.379 1066.65 conc. 73.62 69.25 Interpolated from calibration curve Standard Deviation 3.13 2.56 Average of 4 & 5 MeBT 71.44 | | 5 | 7 187 | 673.99 | 73.33 | 7.378 | 1065 14 | 69 16 | 7.924 16.23 |
| conc. 73.62 69.25 Standard Deviation 3.13 2.56 Average of 4 & 5 MeBT 71.44 | | ava | 7 189 | 676.63 | 10.00 | 7,379 | 1066.65 | 00.10 | |
| Standard Deviation 3.13 2.56 Average of 4 & 5 MeBT 71.44 | | conc | | | 73.62 | | | 69.25 | Interpolated from calibration curve |
| Average of 4 & 5 MeBT 71.44 | | Standard I | Deviation | 3 13 | | | 2 56 | | |
| | | | Average of 4 | & 5 MeBT | | 71.44 | 1.00 | | |

Below LOD

.

Appendix H

.

.

.

Westover ARB Soil

Sample Analysis

Appendix H

Westover ARB Soil

4-MeBT

. . .

. .

| 5 | |
|---|--|
| 뿺 | |
| ž | |
| Ī | |
| ব | |

| 4-MeBT Data | | | | | | | | | | all data | with 10 m | g/L | |
|-------------------------------|-----------|--------|-----------|--------|-----------|-------------|-----------|----------|---------------|--------------|-----------|---------|---------|
| Moisture Content of | fsoil | 0.063 | | | | | | | slope | 0.4099 | 0.2274 | II C | 0.410 |
| Outliers identified | in bold | | | | | | | | intercept | 0.2836 | 0.7470 | Kf = | 1.921 |
| | | | | | | | | | r^2 | 0.9554 | 0.8750 | | |
| | | | | and | Initial | Final | | | Initial | Sorbed | 1 point | | |
| | | Vial + | Air-dried | MeBT | Extractio | Concentrati | Soil (dry | Vol Soln | Concentrati (| Concentratic | Sorption | | |
| Added Concentrati | Vial Only | soil | Soil | mix | c | uo | wt) (g) | (mL) | uo | c | Coeff. | log S l | og Cf |
| | D | g | D | | | ç | Σ | Vs | ö | S | Кd | | |
| | | | | | | mg/L | | | | 6/6rl | ml/g | | |
| 1000 Bottle 1 | 28.279 | 38.281 | 10.002 | 44.738 | 43.035 | 868.82 | 9.409 | 7.050 | 915.92 | 35.286 | 0.041 | 1.5476 | 2.93893 |
| 1000 Bottle 2 | 28.508 | 38.511 | 10.003 | 44.545 | 42.822 | 870.29 | 9.410 | 6.627 | 910.54 | 28.345 | 0.033 | 1.45247 | 2.93966 |
| 1000 Bottle 3 | 28.426 | 38.428 | 10.002 | 44.661 | 42.904 | 865.53 | 9.409 | 6.826 | 913.16 | 34.549 | 0.040 | 1.53844 | 2.93728 |
| 1000 Bottle 4 | 28.462 | 38.462 | 10.000 | 44.266 | 43.002 | 871.83 | 9.407 | 6.397 | 907.35 | 24.151 | 0.028 | 1.38294 | 2.94043 |
| 1000 Bottle 5 | 27.735 | 37.737 | 10.002 | 44.041 | 42.279 | 868.85 | 9.409 | 6.897 | 914.05 | 33.130 | 0.038 | 1.52023 | 2.93894 |
| 100 Bottle 1 | 27.882 | 37.891 | 10.009 | 44.411 | 42.622 | 76.67 | 9.416 | 7.113 | 91.66 | 11.325 | 0.148 | 1.05403 | 1.88463 |
| 100 Bottle 2 | 28.161 | 38.168 | 10.007 | 44.971 | 43.132 | 77.6 | 9.414 | 7.396 | 91.98 | 11.299 | 0.146 | 1.05303 | 1.88986 |
| 100 Bottle 3 | 28.052 | 38.056 | 10.004 | 44.577 | 42.767 | 76.89 | 9.411 | 7.114 | 91.67 | 11.169 | 0.145 | 1.04801 | 1.88587 |
| 100 Bottle 4 | 27.757 | 37.762 | 10.005 | 44.578 | 42.703 | 76.26 | 9.412 | 7.409 | 92.00 | 12.388 | 0.162 | 1.09299 | 1.8823 |
| 100 Bottle 5 | 27.834 | 37.839 | 10.005 | 44.648 | 42.827 | 78.15 | 9.412 | 7.402 | 91.99 | 10.884 | 0.139 | 1.03677 | 1.89293 |
| 10 Bottle 1 | 28.020 | 38.021 | 10.001 | 46.43 | 44.535 | 1.98 | 9.408 | 9.002 | 9.34 | 7.043 | 3.557 | 0.84778 | 0.29667 |
| 10 Bottle 2 | 28.234 | 38.240 | 10.006 | 46.643 | 44.808 | 1.53 | 9.413 | 8.996 | 9.34 | 7.465 | 4.879 | 0.87302 | 0.18469 |
| 10 Bottle 3 | 28.460 | 38.467 | 10.007 | 46.886 | 45.014 | 1.75 | 9.414 | 9.012 | 9.34 | 7.268 | 4.153 | 0.86141 | 0.24304 |
| 10 Bottle 4 | 28.267 | 38.267 | 10.000 | 46.729 | 44.905 | 1.82 | 9.407 | 9.055 | 9.35 | 7.243 | 3.980 | 0.85994 | 0.26007 |
| 10 Bottle 5 | 28.479 | 38.480 | 10.001 | 46.991 | 45.128 | 2.33 | 9.408 | 9.104 | 9.35 | 6.792 | 2.915 | 0.83198 | 0.36736 |
| | | | | | | | | | | | | | |
| 100 Mix 1 | 27.433 | 37.433 | 10.000 | 44.008 | 42.257 | 78.19 | 9.407 | 7.168 | 91.73 | 10.318 | 0.132 | | |
| 100 Mix 2 | 27.847 | 37.855 | 10.008 | 44.443 | 42.674 | 78.51 | 9.415 | 7.181 | 91.74 | 10.091 | 0.129 | | |
| 100 Mix 3 | 27.556 | 37.560 | 10.004 | 44.045 | 42.270 | 77.42 | 9.411 | 7.078 | 91.62 | 10.682 | 0.138 | | |
| 100 Mix 4 | 27.956 | 37.963 | 10.007 | 44.506 | 42.642 | 77.4 | 9.414 | 7.136 | 91.69 | 10.832 | 0.140 | | |
| 100 Mix 5 | 27.681 | 37.681 | 10.000 | 44.161 | 42.341 | 77.04 | 9.407 | 7.073 | 91.62 | 10.962 | 0.142 | | |

5.585 0.027

5.585 26.872

1 1000

Westover-2.xls





Appendix H

· ·

Westover ARB Soil

5-MeBT

.

| ıta | | | | | | | | | all data | with 10 m | g/L | |
|-----------|--------|-----------|--------|-----------|-------------|-----------|----------|-------------|-------------|-----------|---------|----------|
| ÷. | 0.063 | | | | | | | slope | 0.4389 | 0.2384 | Ľ | 0.4389 |
| bold | | | | | | | | intercept | 0.3514 | 0.8405 | Кf = | 2.246 |
| | | | | | | | | r^2 | 0.9735 | 0.8321 | | |
| | | | and | Initial | Final | | | Initial | Sorbed | 1 point | | |
| | Vial + | Air-dried | MeBT | Extractio | Concentrati | Soil (dry | Vol Soln | Concentrati | Concentrati | Sorption | | |
| Vial Only | r soil | Soil | mix | c | uo | wt) (g) | (mL) | uo | uo | Coeff. | log S L | og Cf |
| ۵ | D | b | | | ŭ | Σ | ۷s | ö | S | Кd | | |
| | | | | | mg/L | | | | 6/6rl | ml/g | | |
| 28.279 | 38.281 | 10.002 | 44.738 | 43.035 | 852.71 | 9.409 | 7.050 | 915.92 | 47.356 | 0.056 | 1.67537 | 2.930801 |
| 28.508 | 38.511 | 10.003 | 44.545 | 42.822 | 851.32 | 9.410 | 6.627 | 910.54 | 41.704 | 0.049 | 1.62017 | 2.930093 |
| 28.426 | 38.428 | 10.002 | 44.661 | 42.904 | 846.42 | 9.409 | 6.826 | 913.16 | 48.412 | 0.057 | 1.68496 | 2.927586 |
| 28.462 | 38.462 | 10.000 | 44.266 | 43.002 | 855.74 | 9.407 | 6.397 | 907.35 | 35.092 | 0.041 | 1.54521 | 2.932342 |
| 27.735 | 37.737 | 10.002 | 44.041 | 42.279 | 851.33 | 9.409 | 6.897 | 914.05 | 45.972 | 0.054 | 1.6625 | 2.930098 |
| 27.882 | 37.891 | 10.009 | 44.411 | 42.622 | 72.34 | 9.416 | 7.113 | 91.66 | 14.596 | 0.202 | 1.16423 | 1.859379 |
| 28.161 | 38.168 | 10.007 | 44.971 | 43.132 | 73.5 | 9.414 | 7.396 | 91.98 | 14.520 | 0.198 | 1.16196 | 1.866287 |
| 28.052 | 38.056 | 10.004 | 44.577 | 42.767 | 72.38 | 9.411 | 7.114 | 91.67 | 14.578 | 0.201 | 1.1637 | 1.859619 |
| 27.757 | 37.762 | 10.005 | 44.578 | 42.703 | 71.86 | 9.412 | 7.409 | 92.00 | 15.851 | 0.221 | 1.20006 | 1.856487 |
| 27.834 | 37.839 | 10.005 | 44.648 | 42.827 | 73.9 | 9.412 | 7.402 | 91.99 | 14.226 | 0.193 | 1.15308 | 1.868644 |
| 28.020 | 38.021 | 10.001 | 46.43 | 44.535 | 2.28 | 9.408 | 9.002 | 9.34 | 6.756 | 2.963 | 0.82972 | 0.357935 |
| 28.234 | 38.240 | 10.006 | 46.643 | 44.808 | 0.14 | 9.413 | 8.996 | 9.34 | 8.795 | 63.733 | 0.94424 | -0.86012 |
| 28.460 | 38.467 | 10.007 | 46.886 | 45.014 | 2.25 | 9.414 | 9.012 | 9.34 | 6.789 | 3.017 | 0.83182 | 0.352183 |
| 28.267 | 38.267 | 10.000 | 46.729 | 44.905 | 2.31 | 9.407 | 9.055 | 9.35 | 6.772 | 2.931 | 0.8307 | 0.363612 |
| 28.479 | 38.480 | 10.001 | 46.991 | 45.128 | 1.03 | 9.408 | 9.104 | 9.35 | 8.050 | 7.815 | 0.90578 | 0.012837 |
| | | | | | | | | | | | | |
| 27.433 | 37.433 | 10.000 | 44.008 | 42.257 | 73.6 | 9.407 | 7.168 | 91.73 | 13.815 | 0.188 | | |
| 27.847 | 37.855 | 10.008 | 44.443 | 42.674 | 73.91 | 9.415 | 7.181 | 91.74 | 13.600 | 0.184 | | |
| 27.556 | 37.560 | 10.004 | 44.045 | 42.270 | 72.65 | 9.411 | 7.078 | 91.62 | 14.269 | 0.196 | | |
| 27.956 | 37.963 | 10.007 | 44.506 | 42.642 | 72.74 | 9.414 | 7.136 | 91.69 | 14.364 | 0.197 | | |
| 27.681 | 37.681 | 10.000 | 44.161 | 42.341 | 72.47 | 9.407 | 7.073 | 91.62 | 14.398 | 0.199 | | |

Westover-2.xls

6.927 0.036

6.927 35.944

1 1000





| | | | | | | | | • | 0000 | MeBT Isomer Extr | action Curve - 1000 mg/L |
|--------------------------|-------|------------------|------------|------------|---------|--------|--------|------------------|--------|------------------|--------------------------|
| 1000 mg/L | 0 | | 8225.7 | | 13224.4 | 4 MeBT | 5 MeBT | - | | | |
| Water | 0.5 | 7.019 | 6699.9 | 7.207 | 10702.8 | 81% | 81% | | | | |
| 1 Meth | 1 hr | 7.026 | 1059.7 | 7.215 | 1626.9 | 13% | 12% | - | 0000 | | |
| 2 Meth | 6 hr | 7.025 | 241.4 | 7.211 | 645.9 | 3% | 5% | | | | |
| 3 Meth | 24 hr | 7.018 | 124.8 | 7.205 | 137.5 | 2% | 1% | (1 | 8000 - | | |
| 4 Meth | 48 hr | 7.011 | 71.8 | 7.201 | 71.6 | 1% | 1% | s*u/ | | • | |
| 5 Meth | 72 hr | 7.007 | 21.4 | 7.198 | 33.9 | %0 | %0 | Am | 6000 - | | |
| Sum Recovered | | | 8219.0 | | 13218.6 | | |) ea | | | |
| Percent MeBT Recover | 'ed | n e e e entre Ju | 99.92% | : | 99.96% | | | чA | 4000 | | |
| Empty | | 27.735 F | Sefilled [| Difference | - | | | | 2001 | _ | |
| w/ Soil only | | 37.737 | | | | | | | 0000 | | |
| Soil Weight (air-dried) | | 10.002 | | | | | | | | | |
| w/ Soil and MeBT mix | | 44.041 | | | | | | | • | <i>)</i> | |
| After Initial Extraction | | 42.279 | | 1.762 | | | | | 5 | | |
| of water/MeBT | | | 44.221 | | | | | | 5 | | .3 4 5 6 |
| 1 Meth | | 42.355 | 44.395 | 1.866 | | | | | | Ζ | umber of Extractions |
| 2 Meth | | 42.113 | 44.249 | 2.282 | | | J | | | | |
| 3 Meth | | 42.343 | 44.340 | 1.906 | | | | | | | |
| 4 Meth | | 42.813 | 44.199 | 1.527 | | | | | | | |
| 5 Meth | | 42 746 | | 1.453 | | | | | | | |
| | | | | | | | | | | | |
| 100 mg/L | 0 | | 918.0 | | 1537.0 | 4 MeBT | 5 MeBT | | | MeBT Isomer Ext | raction Curve - 100 mg/L |
| Water | 0.5 | 7.056 | 665.2 | 7.241 | 920.6 | 72% | 60% | - | 000 | | |
| 1 Meth | 1 hr | 7.041 | 134.7 | 7.226 | 367.5 | 15% | 24% | | - 006 | | -+-4 MeBT |
| 2 Meth | 6 hr | 7.04 | 65.1 | 7.224 | 155.4 | %2 | 10% | | | / | |
| 3 Meth | 24 hr | 7.037 | 25.5 | 7.224 | 45.9 | 3% | 3% | (| | | |
| 4 Meth | 48 hr | 7.035 | 12.2 | 7.222 | 26.7 | 1% | 2% | S _* I | | • | |
| 5 Meth | 72 hr | 7.031 | 4.2 | 7.216 | 7.8 | %0 | 1% | n∀ | - 009 | | |
| Sum Recovered | | | 906.9 | | 1523.9 | | | ա) | 200 - | | |
| Percent MeBT Recover | red | Araidhidhean | 98.79% | | 99.15% | | | ee | 400 - | | |
| No additional spikes | | | | | | | | ЪЧ | 300 - | | |
| Emotv | | 27.834 F | Sefilled 1 | Difference | | | | | 200 | | |
| w/ Soil only | | 37.839 | | | | | | | | | - |
| Soil Weight (air-dried) | | 10.005 | | | | | | | ò | | |
| w/ Soil and MeBT mix | | 44.648 | | | | | | | - | 0 | 4 6 |
| After Initial Extraction | | 42.827 | | 1.821 | | | | | | NUI | mber of Extractions |
| of water/MeBT | | | 44.722 | | | |] | | | | |
| 1 Meth | | 42.864 | 44.608 | 1.858 | | | | | | | |
| 2 Meth | | 42.620 | 45.006 | 1.988 | | | | | | | |
| 3 Meth | | 42.955 | 44.638 | 2.051 | | | | | | | |
| 4 Meth | | 42 994 | 44 875 | 1644 | | | | | | | |

Soil Calculation - Westover.xls

1.694

43.181

5 Meth

Extraction

| 10 mg/L | | | | | | | | | | | | | | | i | | 0 | | | | | Curve - Mix | | → 4 MeBT | | | | | | | | | | | | 2 4 5 6 | Extractions | | | | |
|--------------------------------|-------|--------|--------|--------|--------|--------|---------------|----------------------|---------------------|----|------------|--------------|-------------------------|----------------------|--------------------------|-----------------|--------|---------------------|--------|--------|--------|------------------------|-----------------|----------|--------|--------|---------|------------------|---------------|----------------------|---------------------|------------|--------------|-------------------------|----------------------|--------------------------|---------------|--------|--------|--------|--------|
| MeBT Isomer Extraction Curve - | | | | 2 | 0 | | | | .2 .2 | 0 | к | | 0 | 5 | | | 0 2 4 | Nimbor of Extractio | | | | MeBT Isomer Extraction | 1800.0 | | 1600.0 | 1400 0 | | 1200.0 | 1000.0 | | 600.00 | 400.0 | | Z00.0 | 0.0 | с с | Number of | | | | |
| | | ц | · · · | 4 | 4 | |) (s, | n n | າພ) | 69 | ٩٢ | _ | ~ | | | | | | | | | 1 | _ | - | - | | (s | s _≁ r | ١Ą | ա) | rea | A | | | | | | | | | |
| 5 MeBT | %0 | %0 | %0 | %0 | %0 | %0 | | | | | | | | | | | | | | | 5 МеВТ | 100000 | 900% 00% | 0/ 77 | γ 2 | 4% | %0 0 | %0 | | | | | | | | | | | | | |
| MeBT | 40% | 22% | 8% | %0 | %0 | %0 | | | | | | | | | | | | | | | 1 MART | 1000 | 66% 18% | 201 | %8 | 3% | %0 | %0 | | | | | | | | | | | | | |
| 201.1 4 | 0.0 | 0.0 | 0 | 0 | 0 | 0 | 0 | 0.00% | | | | | | | | | | | | | 1537 0 | 0.1001 | 1002.3 336.1 | | 124.1 | 63.6 | | | 1526.1 | 99.29% | | | | | | | | | | | |
| | 7.35 | 7.35 | 7.35 | 7.35 | 7.35 | 7.35 | | | | | ference | | | | 1.863 | | 1.623 | 1.438 | 1.719 | 1.738 | | - | 7.468 | 1.100 | (.455 | 7.451 | | | | 1.2 2 | | ference | | | | 1.820 | | 1.622 | 1.921 | 1.706 | 1.811 |
| 119.9 | 47.9 | 26.3 | 9.8 | 0 | 0 | 0 | 84 | 70.06% | | | Tilled Dim | | | | | 45.880 | 46.084 | 45.896 | 46.293 | | 018.0 | | 602.4 164 7 | 104.7 | (6.6 | 31.1 | | | 874.8 | 95.29% | | filled Dif | | | | | 44.483 | 44.530 | 44.216 | 44.453 | |
| | 7.139 | 7.140 | 7.164 | 7.135 | 7.135 | 7.135 | | | | | 28.479 Kei | 38.480 | 10.001 | 46.991 | 45.128 | | 44.257 | 44.646 | 44.177 | 44.555 | | | 7.278 | E07.1 | 7.256 | 7.252 | | | | | - | 27.681 Re | 37.681 | 10.000 | 44.161 | 42.341 | | 42.861 | 42.609 | 42.510 | 42.642 |
| 0 | 0.5 | 1 hr | 6 hr | 24 hr | 48 hr | 72 hr | | 'ed | | | | | | | | | | | | | c | , , | 0.5 1 hr | = | 6 hr | 24 hr | 48 hr | 72 hr | | red | | | | | | | | | | | |
| 10 mg/L | Water | 1 Meth | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT Recover | No additional spike | | Empty | w/ Soil only | Soil Weight (air-dried) | w/ Soil and MeBT mix | After Initial Extraction | of water/MeBT · | 1 Meth | 2 Meth | 3 Meth | 4 Meth | Wic | | Water | | 2 Meth | 3 Meth | 4 Meth | 5 Meth | Sum Recovered | Percent MeBT Recover | No additional spike | Emoty | w/ Soil only | Soil Weight (air-dried) | w/ Soil and MeBT mix | After Initial Extraction | of water/MeBT | 1 Meth | 2 Meth | 3 Meth | 4 Meth |

Soil Calculation - Westover.xls

Extraction

| 0 ma/i | | 4.8 | 20.5 | 4 MeBT | 5 MeBT |
|---------------------|-------|-------|-------|--------|--------|
| | 2 | 2.12 | | | |
| Water | 0.5 | 0.0 | 0.0 | %0 | %0 |
| 1 Meth | 1 hr | 0.0 | 0.0 | %0 | %0 |
| 2 Meth | 6 hr | 0.0 | 0.0 | %0 | %0 |
| 3 Meth | 24 hr | | | %0 | %0 |
| 4 Meth | 48 hr | | | %0 | %0 |
| 5 Meth | 72 hr | | | %0 | %0 |
| Sum Recovered | | 0 | 0 | | |
| Percent MeBT Recove | red | %00.0 | 0.00% | | |

No additional spike - Zero result after background subtraction

| Refilled Difference | | 0.000 | | · 0.000 | 0.000 | 0.000 | 0.000 | |
|-----------------------|---|--------------------------|---------------|---------|--------|--------|--------|--|
| Empty w/ Soil only | Soil Weight (air-dried) w/ Soil and MeBT mix | After Initial Extraction | of water/MeBT | 1 Meth | 2 Meth | 3 Meth | 4 Meth | |

,

.

Soil Calculation - Westover.xls

| 1000 mg/L | | | · · · · · · · · · · · · · · · · · · · | | | | | · ····· |
|------------|---------------------|--------------|---------------------------------------|---------------|-------|----------|-----------|--|
| | Runs | | 4 MeBT | | | 5 MeBT | | |
| | 1 | 7.432 | 7160.7 | 868.82 | 7.631 | 11368.7 | 852.71 | |
| | 2 | 7.422 | 7172.6 | 870.29 | 7.621 | 11351.1 | 851.32 | |
| | 3 | 7.409 | 7134.0 | 865.53 | 7.608 | 11289.4 | 846.42 | |
| | 4 | 7.427 | 7185.1 | 871.83 | 7.631 | 11406.9 | 855.74 | |
| | 5 | 7.386 | 7160.9 | 868.85 | 7.584 | 11351.3 | 851.33 | |
| | avg | 7.415 | 7162.66 | | 7.615 | 11353.48 | | |
| | conc. | | | 869.064 | | | 851.50 | Interpolated from calib |
| | Standard E | Deviation | 18.91 | | | 42.43 | | |
| | | Average of 4 | & 5 MeBT | | 0.00 | | | |
| 400 | | | | | | | | |
| 100 mg/L | Dune | | 4 MoBT | | | 5 MeBT | | 1 |
| | 1 | 7.467 | 704 57 | 76.67 | 7 667 | 1113.88 | 72 34 | |
| | 2 | 7.407 | 713.08 | 77.60 | 7 640 | 1131 53 | 73.50 | |
| | 2 | 7.441 | 706.54 | 76.89 | 7.606 | 1114 38 | 72.38 | |
| | 4 | 7 389 | 700.78 | 76.26 | 7.587 | 1106.49 | 71.86 | |
| | 5 | 7.367 | 718.06 | 78.15 | 7.564 | 1137.67 | 73.90 | |
| | avo | 7.416 | 708.61 | | 7.613 | 1120.79 | · · · · · | |
| | conc. | | | 77.114 | | | 72.80 | Interpolated from calib |
| | Standard D | Deviation | 6.91 | | | 13.17 | | |
| | | Average of 4 | & 5 MeBT | | 0.00 | | | |
| | | | | | | | | |
| 10 mg/L | | | | | | | | 1 |
| | Runs | | 4 MeBT | | | 5 MeBT | | |
| | 1 | 7.147 | 48.21 | 1.98 | 7.343 | 21.73 | 0.03 | |
| | 2 | 7.142 | 45.21 | 1.53 | 7.350 | 20.67 | 0.01 | |
| | 3 | 7.143 | 46.66 | 1.75 | 7.335 | 20.51 | 0.00 | |
| | 4 | 7.145 | 47.14 | 1.82 | 7.335 | 27.43 | 0.14 | |
| | C | 7.146 | 00.03 47.55 | 2.33 | 7.337 | 23.47 | 0.00 | |
| | avg | 7.145 | 47.00 | 4 000 | 7.340 | 22.70 | 0.05 | Internelated from calib |
| | Stondard F | Deviation | 1.09 | 1.002 | | 2.86 | 0.00 | |
| | Standard L | | 1.90 & 5 MoBT | | 0.00 | 2.00 | | |
| | | Average of 4 | a o meo i | | 0.00 | | | |
| 100 mg/L M | eBT / 1000 r | ng/L PG | | | | | | ····· |
| | Runs | Г . | 4 MeBT | | | 5 MeBT | | |
| | 1 | 7.189 | 718.48 | 78.19 | 7.379 | 1133.13 | 73.60 | |
| | 2 | 7.189 | 721.39 | 78.51 | 7.380 | 1137.91 | 73.91 | |
| | 3 | 7.190 | 711.39 | 77.42 | 7.381 | 1118.58 | 72.65 | |
| | 4 | 7.187 | 711.25 | 77.40 | 7.379 | 1119.97 | 72.74 | |
| | 5 | 7.190 | 707.88 | 77.04 | 7.38 | 1115.83 | 72.47 | |
| | avg | 7.189 | 714.08 | | 7.380 | 1125.08 | | |
| | | | | NO 140 | | | 70.07 | the second state of free second states |
| | conc. | | | 77.712 | | | 73.07 | interpolated from callo |
| | conc. Standard E | Deviation | 5.62 | 77.712 | | 9.79 | 73.07 | interpolated from callb |

Below LOD

Appendix I

Achieved Kow Values

Calculations

.

.

| 1000 mg/ | Untre | eated | Trea | ated | DI | A | West | tover |
|-----------|--------|--------|--------|--------|-------|-------|-------|--------|
| ľ | 4 | 5 | 4 | 5 | 4 | 5 | 4 | 5 |
| erage Kd | 0.860 | 0.825 | 0.350 | 0.512 | 0.044 | 0.075 | 0.036 | 0.051 |
| foc | 0.030 | 0.030 | 0.030 | 0.030 | 0.008 | 0.008 | 0.005 | 0.005 |
| Koc | 28.680 | 27.507 | 11.667 | 17.053 | 5.450 | 9.425 | 7.200 | 10.280 |
| Kom | 16.636 | 15.955 | 6.767 | 9.892 | 3.161 | 5.467 | 4.176 | 5.963 |
| log Kow | 1.16 | 1.12 | 0.40 | 0.72 | -0.23 | 0.23 | 0.00 | 0.30 |
| log Kow | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 |
| ifference | 1.01 | 1.05 | 1.77 | 1.45 | 2.40 | 1.94 | 2.17 | 1.87 |

| 100 mg/L | Untreated | | Treated | | DIA | | Westover | |
|-----------|-----------|----------|---------|---------|--------|--------|----------|--------|
| | 4 | 5 | 4 | 5 | 4 | 5 | 4 | 5 |
| erage Kd | 1.203 | 330.260 | 0.706 | 44.523 | 0.160 | 0.255 | 0.148 | 0.203 |
| foc | 0.030 | 0.030 | 0.030 | 0.030 | 0.008 | 0.008 | 0.005 | 0.005 |
| Koc | 40.107 | 11008.67 | 23.547 | 1484.10 | 20.050 | 31.875 | 29.600 | 40.600 |
| Kom | 23.264 | 6385.538 | 13.658 | 860.847 | 11.630 | 18.489 | 17.169 | 23.550 |
| log Kow | 1.44 | 6.13 | 0.99 | 4.45 | 0.86 | 1.24 | 1.18 | 1.45 |
| log Kow | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 |
| ifference | 0.73 | -3.96 | 1.18 | -2.28 | 1.31 | 0.93 | 0,99 | 0.72 |

| 10 mg/L | Untreated T | | Trea | Treated DI/ | | A Wes | | over |
|-----------|-------------|--------|-------|-------------|-------|-------|-------|--------|
| | 4 | 5 | 4 | 5 | 4 | 5 | 4 | 5 |
| erage Kd | 32.850 | 55.203 | 0.521 | 400.000 | 2.844 | 5.021 | 3.897 | 16.092 |
| foc | 0.030 | 0.030 | 0.030 | 0.030 | 0.008 | 0.008 | 0.005 | 0.005 |
| Koc | 1095.0 | 1840.1 | 17.4 | 13333 | 355.5 | 627.6 | 779.4 | 3218.4 |
| Kom | 635.2 | 1067.3 | 10.1 | 7734.0 | 206.2 | 364.0 | 452.1 | 1866.8 |
| log Kow | 4.20 | 4.63 | 0.74 | 6.29 | 3.26 | 3.73 | 3.91 | 5.10 |
| log Kow | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 |
| ifference | -2.03 | -2.46 | 1.43 | -4.12 | -1.09 | -1.56 | -1.74 | -2.93 |

| 100 Mix | Untreated | | Treated | | DIA | | Westover | |
|-----------|-----------|-------|---------|-------|-------|-------|----------|-------|
| | 4 | 5 | 4 | 5 | 4 | 5 | 4 | 5 |
| erage Kd | 1.219 | 1.735 | 0.547 | 2.325 | 0.208 | 0.291 | 0.136 | 0.193 |
| foc | 0.030 | 0.030 | 0.030 | 0.030 | 0.008 | 0.008 | 0.005 | 0.005 |
| Koc | 40.6 | 57.8 | 18.2 | 77.5 | 26.0 | 36.4 | 27.2 | 38.6 |
| Kom | 23.6 | 33.5 | 10.6 | 45.0 | 15.1 | 21.1 | 15.8 | 22.4 |
| log Kow | 1.45 | 1.74 | 0.78 | 1.99 | 1.07 | 1.35 | 1.11 | 1.40 |
| log Kow | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 | 2.17 |
| ifference | 0.72 | 0.43 | 1.39 | 0.18 | 1.10 | 0.82 | 1.06 | 0.77 |

Back-calculated log Kow values Differences < or = 1.00 (acceptable) Differences < 0 (not good)

Appendix J

Statistical Analysis

Calculations

.

.

Appendix J

Statistical Analysis

4-MeBT

, ·

.

.

4 MeBT

| | 1000 | 100 | 10 | Mix |
|----------|-------|-------|--------|-------|
| Untreate | 0.860 | 1.203 | 32.849 | 1.219 |
| Treated | 0.350 | 0.706 | 0.521 | 0.547 |
| DIA | 0.044 | 0.160 | 2.844 | 0.208 |
| Westove | 0.036 | 0.148 | 3.897 | 0.136 |



| CASE | SOIL | CONC | OBSERVE | YHAT | RESID |
|-----------|--------|--------|---------|------------------|---------|
| 1 | 1 | 1 | 0.817 | 0.8604 | -0.0434 |
| 2 | 1 | 1 | 0.907 | 0.8604 | 0.0466 |
| 3 | 1 | 1 | 0.83 | 0.8604 | -0.0304 |
| 4 | 1 | 1 | 0.841 | 0.8604 | -0.0194 |
| 5 | 1 | 1 | 0.907 | 0.8604 | 0.0466 |
| 6 | 1 | 2 | 1.169 | 1.2032 | -0.0342 |
| 7 | 1 | 2 | 1.14 | 1.2032 | -0.0632 |
| 8 | 1 | 2 | 1.085 | 1.2032 | -0.1182 |
| 9 | 1 | 2 | 1.445 | 1.2032 | 0.2418 |
| 10 | 1 | 2 | 1.177 | 1.2032 | -0.0262 |
| 11 | 1 | 3 | 32.68 | 32.849 | -0.169 |
| 12 | 1 | 3 | 32.029 | 32.849 | -0.82 |
| 13 | 1 | 3 | 32.317 | 32.849 | -0.532 |
| 14 | 1 | 3 | 34.877 | 32.849 | 2.028 |
| 15 | 1 | 3 | 32.342 | 32.849 | -0.507 |
| 16 | 1 | 4 | 1.128 | 1.2188 | -0.0908 |
| 17 | 1 | 4 | 1.281 | 1.2188 | 0.0622 |
| 18 | 1 | 4 | 1.204 | 1.2188 | -0.0148 |
| 19 | 1 | 4 | 1.223 | 1.2188 | 0.0042 |
| 20 | 1 | 4 | 1.258 | 1.2188 | 0.0392 |
| 21 | 2 | 1 | 0.35 | 0.35 | -0.0 |
| 22 | 2 | 1 | 0.343 | 0.35 | -0.007 |
| 23 | 2 | 1 | 0.365 | 0.35 | 0.015 |
| 24 | 2 | 1 | 0.359 | 0.35 | 0.009 |
| 25 | 2 | 1 | 0.333 | 0.35 | -0.017 |
| 26 | 2 | 2 | 0.71 | 0.7064 | 0.0036 |
| 27 | 2 | 2 | 0.718 | 0.7064 | 0.0116 |
| 28 | 2 | 2 | 0.676 | 0.7064 | -0.0304 |
| 29 | 2 | 2 | 0.707 | 0.7064 | 0.0006 |
| 30 | 2 | 2 | 0.721 | 0.7064 | 0.0146 |
| 31 | 2 | 3 | 0.56 | 0.5212 | 0.0388 |
| 32 | 2 | 3 | 0.494 | 0.5212 | -0.0272 |
| 33 | 2 | 3 | 0.506 | 0.5212 | -0.0152 |
| 34 | 2 | 3 | 0.562 | 0.5212 | 0.0408 |
| 35 | 2 | 3 | 0.484 | 0.5212 | -0.0372 |
| 36 | 2 | 4 | 0.538 | 0.547 | -0.009 |
| 37 | 2 | 4 | 0.54 | 0.547 | -0.007 |
| 38 | 2 | 4 | 0.572 | 0.54/ | 0.025 |
| 39 | 2 | 4 | 0.536 | 0.54/ | -0.011 |
| 40 | 2 | 4 | 0.549 | 0.54/ | 0.002 |
| 41 | 3 | 1 | 0.027 | 0.0436 | -0.0166 |
| 42 | 3 | 1 | 0.046 | 0.0436 | 0.0024 |
| 43 | 3 | 1 | 0.051 | 0.0436 | 0.0074 |
| 44 | 3 | 1 | 0.041 | 0.0436 | -0.0026 |
| 45 | 3 | 1 | 0.053 | 0.0436 | 0.0094 |
| 40 | ວ ວ | 2 | 0.142 | 0.1604 | -0.0184 |
| 47 | ວ ວ | 2 | 0.15 | 0.1604 | -0.0104 |
| 40 | с С | 2 | 0.15 | 0.1604 | -0.0104 |
| 49 | с С | 2 | 0.171 | 0.1604 | 0.0106 |
| 50 | ວ ວ | 2 | 0.170 | 0.1604 | 0.0176 |
| 51 | 3 | 3 | 2.738 | 2.8442 | -0.1062 |
| 52 52 | ა ი | с r | 3.000 | 2.0442 9 0119 | -0.0102 |
| 55 E A | ა ი | 3 | 2.034 | 2.044Z | -0.0102 |
| 54 | د د | 3 | ∠.0⊥4 | 2.0442 | -0.0302 |
| 55 | 3 2 | د ۸ | 2./ð | 2.0442 | ~0.0042 |
| 30 - 7 | 3 | 4 | 0.208 | 0.2078 | 0.0002 |
| 5/ E0 | 3 | 4 | 0.19/ | 0.2078 | -0.0108 |
| 58 50 | د م | 4 | 0.203 | 0.2078 | -0.0048 |
| 59 | 3 | 4 | 0.215 | 0.2078 | 0.0072 |
| 60 | ک ک | 4 | U.216 | 0.20/8 | 0.0082 |

| CASE | SOIL | CONC | OBSERVE | YHAT | RESID |
|------|------|------|---------|--------|---------|
| 61 | 4 | 1 | 0.041 | 0.036 | 0.005 |
| 62 | 4 | 1 | 0.033 | 0.036 | -0.003 |
| 63 | 4 | 1 | 0.04 | 0.036 | 0.004 |
| 64 | 4 | 1 | 0.028 | 0.036 | -0.008 |
| 65 | 4 | 1 | 0.038 | 0.036 | 0.002 |
| 66 | 4 | 2 | 0.148 | 0.148 | 0.0 |
| 67 | 4 | 2 | 0.146 | 0.148 | -0.002 |
| 68 | 4 | 2 | 0.145 | 0.148 | -0.003 |
| 69 | 4 | 2 | 0.162 | 0.148 | 0.014 |
| 70 | 4 | 2 | 0.139 | 0.148 | -0.009 |
| 71 | 4 | 3 | 3.557 | 3.8968 | -0.3398 |
| 72 | 4 | 3 | 4.879 | 3.8968 | 0.9822 |
| 73 | 4 | 3 | 4.153 | 3.8968 | 0.2562 |
| 74 | 4 | 3 | 3.98 | 3.8968 | 0.0832 |
| 75 | 4 | 3 | 2.915 | 3.8968 | -0.9818 |
| 76 | 4 | 4 | 0.132 | 0.1362 | -0.0042 |
| 77 | 4 | 4 | 0.129 | 0.1362 | -0.0072 |
| 78 | 4 | 4 | 0.138 | 0.1362 | 0.0018 |
| 79 | 4 | 4 | 0.14 | 0.1362 | 0.0038 |
| 80 | 4 | 4 | 0.142 | 0.1362 | 0.0058 |

4MEBT, 03/02/99, 9:50

ANALYSIS OF VARIANCE TABLE FOR OBSERVE

| SOURCE | DF | SS | MS | F | P |
|----------|----|---------|---------|---------|--------|
| SOIL (A) | 3 | 1019.49 | 339.830 | 2846.19 | 0.000 |
| CONC (B) | 3 | 1371.45 | 457.149 | 3828.79 | 0.0000 |
| A*B | 9 | 2492.13 | 276.903 | 2319.16 | 0.0000 |
| RESIDUAL | 64 | 7.64147 | 0.11940 | | |
| TOTAL | 79 | 4890.71 | | | |

GRAND MEAN 2.8581 SE 0.0386

MEANS OF OBSERVE FOR SOIL

| SOIL | MEAN | SS (MEAN) |
|------|--------|-----------|
| 1 | 9.0328 | 3787.3 |
| 2 | 0.5312 | 0.3277 |
| 3 | 0.8140 | 27.612 |
| 4 | 1.0542 | 56.022 |

OBSERVATIONS PER CELL20STD ERROR OF AN AVERAGE0.0773STD ERROR (DIFF OF 2 AVE'S)0.1093ERROR TERM USED: RESIDUAL, 64 DF

MEANS OF OBSERVE FOR CONC

| CONC | MEAN | SS (MEAN) |
|------|--------|-----------|
| 1 | 0.3225 | 2.2585 |
| 2 | 0.5545 | 3.9029 |
| 3 | 10.028 | 3509.4 |
| 4 | 0.5274 | 3.6831 |
| | | |

OBSERVATIONS PER CELL20STD ERROR OF AN AVERAGE0.0773STD ERROR (DIFF OF 2 AVE'S)0.1093ERROR TERM USED: RESIDUAL, 64 DF

MEANS OF OBSERVE FOR SOIL*CONC

| SOIL | CONC | MEAN | SS (MEAN) |
|-------------|--------------|------------|-----------|
| 1 | 1 | 0.8604 | 7.53E-03 |
| 1 | 2 | 1.2032 | 0.0783 |
| 1 | 3 | 32.849 | 5.3538 |
| 1 | 4 | 1.2188 | 0.0139 |
| 2 | 1 | 0.3500 | 6.44E-04 |
| 2 | 2 | 0.7064 | 1.29E-03 |
| 2 | 3 | 0.5212 | 5.52E-03 |
| 2 | 4 | 0.5470 | 8.80E-04 |
| 3 | 1 | 0.0436 | 4.31E-04 |
| 3 | 2 | 0.1604 | 8.69E-04 |
| 3 | 3 | 2.8442 | 0.0609 |
| 3 | 4 | 0.2078 | 2.59E-04 |
| 4 | 1 | 0.0360 | 1.18E-04 |
| 4 | 2 | 0.1480 | 2.90E-04 |
| 4 | 3 | 3.8968 | 2.1167 |
| 4 | 4 | 0.1362 | 1.21E-04 |
| OBSERVATION | S PER CELL | | 5 |
| STD ERROR O | F AN AVERAGE | E 0.1 | 545 |
| STD ERROR (| DIFF OF 2 AV | VE'S) 0.2 | 185 |
| ERROR TERM | USED: RESIDU | JAL, 64 DF | |

4MEBT, 03/02/99, 10:23

.....

4MEBT, 03/02/99, 10:28

TUKEY (HSD) COMPARISON OF MEANS OF OBSERVE BY SOIL

| MEAN | HOMOGENEOUS GROUPS |
|--------|--|
| 9.0328 | I |
| 1.0543 | I |
| 0.8140 | I I |
| 0.5312 | I |
| | MEAN 9.0328 1.0543 0.8140 0.5312 |

THERE ARE 3 GROUPS IN WHICH THE MEANS ARE NOT SIGNIFICANTLY DIFFERENT FROM ONE ANOTHER.

| CRITICAL | Q VALU | JE | | 3.731 | REJECTION | LEVEL | 0.050 |
|----------|--------|-----|------------|--------|-----------|-------|-------|
| CRITICAL | VALUE | FOR | COMPARISON | 0.2883 | | | |
| STANDARD | ERROR | FOR | COMPARISON | 0.1093 | | | |

ERROR TERM USED: RESIDUAL, 64 DF

4MEBT, 03/02/99, 10:28

TUKEY (HSD) COMPARISON OF MEANS OF OBSERVE BY CONC

| CONC | MEAN | HOMOGENEOUS GROUPS |
|------|--------|-----------------------|
| 3 | 10.028 | I |
| 2 | 0.5545 | I |
| 4 | 0.5274 | I |
| 1 | 0.3225 | I |

THERE ARE 2 GROUPS IN WHICH THE MEANS ARE NOT SIGNIFICANTLY DIFFERENT FROM ONE ANOTHER.

| CRITICAL | Q VALU | JE | | 3.731 | REJECTION | LEVEL | 0 050 |
|----------|--------|-----|------------|--------|-----------|-------|-------|
| CRITICAL | VALUE | FOR | COMPARISON | 0.2883 | | 50,50 | 0.000 |
| STANDARD | ERROR | FOR | COMPARISON | 0.1093 | | | |

ERROR TERM USED: RESIDUAL, 64 DF

Box and Whisker Plot



· · · · ·

4MEBT, 03/02/99, 10:04

| SOIL | MEAN | SAME SIZ | PLE ZE | S.D. | S.E. | |
|------------------------------|---------------------------|------------------------|-----------|------------------|----------------------------|--------------------|
| 3 4 DIFFERENCE | 0.043 0.030 7.60E-0 | 36 5 50 5 03 | 5 (| 0.0104 43E-03 | 4.64E-03 2.43E-03 | |
| NULL HYPOTHE ALTERNATIVE | SIS: DIF HYP: DIF | FERENCE = FERENCE < | 0 > 0 | D | OF& CT FOP D | τϝϝϝϗͺ |
| ASSUMPTION | | Т | DE | P | 95% CI FOR D | |
| EQUAL VARIAN UNEQUAL VARI | CES ANCES | 1.45 1.45 | 8 6.0 | 0.1850 0.1969 | (-4.48E-03, (-5.20E-03, | 0.0197) 0.0204) |
| | | F | NUM DF | DEN DF | Р | |
| TESTS FOR EQ OF VAR | UALITY LIANCES | 3.65 | 4 | 4 | 0.1187 | |
| CASES INCLUE | DED 10 | MISSING | CASES 0 | | | |

4MEBT, 03/02/99, 10:06

| SOI | L | MEAI | SA N S | MP: IZI | LE E | s. | D. | S.I | c. | |
|------------------|----------------------------|--------------|--------------|------------|----------|----------|--------------|----------------|------------------|--------------------|
| | 3 | 0.16 | 04 | 5 | | 0.0 |)147 | 6.59E- | -03 | |
| 4 | Į | 0.148 | 30 | 5 | 8.5 | 51E | 2-03 | 3.81E- | -03 | |
| DIFFEF | RENCE | 0.012 | 24 | | | | | | | |
| NULL H ALTERN | IYPOTHESIS: IATIVE HYP: | DIFI DIFI | FERENCE | = <> | 0 0 | | | | | |
| ASSUMP | PTION | | Т | | DF | | Р | 95% CI | FOR D | IFFERENCE |
| EQUAL UNEQUA | VARIANCES L VARIANCE | S | 1.63 1.63 | | 8 6.4 | 0. 0. | 1420 1514 | (-5.1 (-5.9 | 6E-03, 5E-03, | 0.0300) 0.0307) |
| TESTS | FOR FOULT.T | ͲV | F | | NUM DF | | DEN DF | E |) | |
| | OF VARIANC | ES | 3.00 | | 4 | | 4 | 0.15 | 64 | |
| CASES | INCLUDED 1 | 0 | MISSING | CÆ | ASES 0 | | | | | |

Test 1 for 4-MeBT looks at the clayey vs. sandy soil comparison.

t _{star} = 1.637 From Statistix

 $t_{025} < t_{star} < t_{975}$

Table A.5, Devore, 8 degree of freedom

t .025 :=- 2.306 t .975 := 2.306

Do Not Reject H_o

ORGIN≡0

$$t := -5, -4.9..5$$



4MEBT, 03/02/99, 10:09

| SOIL | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|--|-------------------------------|-------------------------|-----------|------------|------------|
| 1 | 0.8604 | 5 | 0.0434 | 0.0194 | |
| 4 | 0.0360 | 5 | 5.43E-03 | 2.43E-03 | |
| DIFFERENCE | 0.8244 | | | | |
| NULL HYPOTHESIS: ALTERNATIVE HYP: ASSUMPTION | : DIFFEREN(DIFFEREN(T | CE = 0 CE <> 0 DF | Ρ | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES | 42.1 | - <u>-</u> | 0.0000 | (0 7793 | 0 8695) |
| UNEQUAL VARIANCE | ES 42.1 | 7 4.1 | 0.0000 | (0.7708, | 0.8780) |
| TESTS FOR FOILT | F | NUM | DF DEN DF | Р | |
| OF VARIANC | CES 63. | 79 4 | 4 | 0.0007 | |
| CASES INCLUDED 1 | .0 MISS | ING CASES | 0 | | |

•

.

4MEBT, 03/02/99, 10:10

| SOIL | MEAN | SAMI SIZ | PLE ZE | S.D. | S.E. | |
|--|-------------------------|------------------------------|----------------|------------------|----------------------|--------------------|
| 1 4 DIFFEBENCE | 1.203 0.148 1.055 | 2 5 | 5 8.5 | 0.1399 51E-03 | 0.0626 3.81E-03 | |
| NULL HYPOTHESI ALTERNATIVE HY ASSUMPTION | S: DIFE P: DIFE | 'ERENCE = 'ERENCE <> T | 0 > 0 DF | Ρ | 95% CI FOR 3 | DIFFERENCE |
| EQUAL VARIANCE UNEQUAL VARIAN | S CES | 16.83 16.83 | 8 4.0 | 0.0000 0.0001 | (0.9107, (0.8817, | 1.1997) 1.2287) |
| TESTS FOR EQUA OF VARIA | LITY NCES | F 269.96 | NUM DF | DEN DF 4 | P 0.0000 | |
| CASES INCLUDED | 10 | MISSING (| CASES 0 | | | |

Test 2 for 4-MeBT looks at the higher $\,f_{oc}^{}\,$ vs. low $f_{oc}^{}\,$ soil comparison.

t _{star} = 42.495 From Statistix

Table A.5, devour, 22 degree of freedom

t .025 :=- 2.306 t .975 := 2.306

Do Reject $\rm H_{o}$

ORGIN≡0



4MEBT, 03/02/99, 10:12

| SOTI | MEAN | SAMPLE | S D | 9 F | |
|------------------------------------|--------------------------|-------------------|---------|------------|------------|
| | | <u> </u> | | ۰ü. | |
| 1 | 0.8604 | 5 | 0.0434 | 0.0194 | |
| 2 | 0.3500 | 5 | 0.0127 | 5.67E - 03 | |
| DIFFERENCE | 0.5104 | | | | |
| NULL HYPOTHESIS ALTERNATIVE HYP | : DIFFEREN : DIFFEREN | CE = 0 CE <> 0 | | | |
| ASSUMPTION | Т | DF | P | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES | 25.2 | 5 8 | 0.0000 | (0.4638. | 0.5570) |
| UNEQUAL VARIANC | ES 25.2 | 5 4.7 | 0.0000 | (0.4574, | 0.5634) |
| | F | NUM DF | DEN DF | Р | |
| TESTS FOR EQUAL | ITY | | | | |
| OF VARIAN | CES 11. | 69 4 | 4 | 0.0177 | |
| CASES INCLUDED | 10 MISS | ING CASES 0 | | | |

. . .

4MEBT, 03/02/99, 10:14

| SOIL | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|--------------------------------------|------------------------|-----------------|------------------|--------------------|------------------------|
| 1 | 1.2032 | 5 | 0.1399 | 0.0626 | |
| 2 DIFFERENCE | 0.7064 0.4968 | 5 | 0.0179 | 8.02E-03 | |
| NULL HYPOTHESIS: ALTERNATIVE HYP: | DIFFERENC DIFFERENC | E = 0 E <> 0 | | | |
| ASSUMPTION | т | DF | P | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCE | 7.88 S 7.88 | 8 4.1 | 0.0000 0.0012 | (0.3513 (0.3238 | , 0.6423) , 0.6698) |
| TESTS FOR EQUALI | F TY ——— | NUM D | F DEN DF | P | |
| OF VARIANC | ES 60.9 | 2 4 | 4 | 0.0008 | |
| CASES INCLUDED 1 | 0 MISSI | NG CASES 0 | | | |

Test 3 for 4-MeBT looks at an acclimated vs. unacclimated soil comparison.

t _{star} = 26.309 From Statisitx

 $t_{025} < t_{star} < t_{975}$

Table A.5, devour, 22 degree of freedom

t .025 :=-2.306 t .975 := 2.306

Do Reject H_o

ORGIN≡0



4MEBT, 03/02/99, 10:16

| CONC | MEA | N SI | IPLE ZE | S.D. | S.E. | |
|------------------------------|----------------------|------------------------|-------------|--------|------------------------|--------------------|
| 2 | 1.20 | 32 | 5 | 0.1399 | 0.0626 | |
| 4 | 1.21 | 88 | 5 | 0.0589 | 0.0264 | |
| DIFFERENCE | -0.01 | 56 | | | | |
| NULL HYPOTHE ALTERNATIVE | SIS: DIF HYP: DIF | FERENCE = FERENCE < | = 0 :> 0 | | | |
| ASSUMPTION | | Т | DF | Р | 95% CI FOR | DIFFERENCE |
| EQUAL VARIAN UNEQUAL VARI | CES ANCES | -0.23 -0.23 | 8 5.4 | 0.8240 | (-0.1721, (-0.1865, | 0.1409) 0.1553) |
| TECTS FOD FO | יזאד דיזע | F | NUM DF | DEN DF | Р | |
| OF VAR | IANCES | 5.64 | 4 | 4 | 0.0613 | |
| CASES INCLUD | ED 10 | MISSING | CASES 0 | | | |

Test 4 for 4-MeBT looks at a PG additive vs. no PG added concentration mix comparison.

 $t_{star} = -0.249$

$$t_{025} < t_{star} < t_{975}$$

Table A.5, devour, 22 degree of freedom

t .025 := - 2.306 t .975 := 2.306

Do Not Reject H_o

ORGIN≡0



.

4MEBT, 03/02/99, 10:17

| CONC | MEA | N SA | MP: IZI | LE E | S.D. | S.E. | |
|----------------------------------|--------------------|--------------------|------------|------------|------------------|------------------------|--------------------|
| 2 | 0.43 | 34 | 10 | | 0.2882 | 0.0911 | |
| 4 DIFFERENCE | 0.37 0.05 | 74 60 | 10 | l l | 0.1791 | 0.0566 | |
| NULL HYPOTHESI ALTERNATIVE HY | S: DIF: P: DIF: | FERENCE FERENCE | = <> | 0 0 | | | |
| ASSUMPTION | | Т | | DF | Р | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCE UNEQUAL VARIAN | S CES | 0.52 | | 18 15.1 | 0.6081 0.6093 | (-0.1694, (-0.1726, | 0.2814) 0.2846) |
| TESTS FOR EQUA | LITY | F | | NUM DF | DEN DF | P | |
| OF VARIA | NCES | 2.59 | | 9 | 9 | 0.0864 | |
| CASES INCLUDED | 20 | MISSING | CF | ASES 0 | | | |

4MEBT, 03/02/99, 10:19

| | | SAMPLE | | | |
|------------------------------------|----------------------|-------------------|----------|------------|------------|
| CONC | MEAN | SIZE | S.D. | S.E. | |
| 2 | 0.1604 | 5 | 0.0147 | 6.59E-03 | |
| 4 | 0.2078 | 5 | 8.04E-03 | 3.60E-03 | |
| DIFFERENCE - | -0.0474 | | | | |
| NULL HYPOTHESIS ALTERNATIVE HYP | DIFFEREN DIFFEREN | CE = 0 CE <> 0 | | | |
| ASSUMPTION | Т | DF | Р | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES | -6.3 | 1 8 | 0.0002 | (-0.0647, | -0.0301) |
| UNEQUAL VARIANCH | ES -6.3 | 1 6.2 | 0.0007 | (-0.0656, | -0.0292) |
| | F | NUM | DF DEN D | F P | |
| TESTS FOR EQUAL | ΙТΥ | | | | |
| OF VARIAN | CES 3.3 | 36 4 | 4 | 0.1338 | |
| CASES INCLUDED | 10 MISS | ING CASES | 0 | | |

4MEBT, 03/02/99, 10:20

| CONC | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|-------------------------------------|------------------------|-----------------|------------------|--------------------------|--------------------|
| 2 | 0.1480 | 5 8 | .51E-03 | 3.81E-03 | |
| JIFFERENCE | 0.1382 | 5 5 | .50E-03 | 2.46E-03 | |
| NULL HYPOTHESIS ALTERNATIVE HYP | DIFFERENC DIFFERENC | E = 0 E <> 0 | | | |
| ASSUMPTION | т | DF | Р | 95% CI FOR I | IFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCE | 2.60 2.60 | 8 6.8 | 0.0314 0.0360 | (1.35E-03, (1.03E-03, | 0.0223) 0.0226) |
| TESTS FOR EQUAL | F | NUM D | F DEN DF | P | |
| OF VARIANO | CES 2.40 | 0 4 | 4 | 0.2086 | |
| CASES INCLUDED | 0 MISSI | NG CASES 0 | | | |

Appendix J

Statistical Analysis

5-MeBT

.

.

5 MeBT

| | 1000 | 100 | 10 | Mix |
|----------|-------|---------|--------|-------|
| Untreate | 0.825 | 330.000 | 55.200 | 1.735 |
| Treated | 0.512 | 44.500 | 400 | 2.325 |
| DIA | 0.075 | 0.255 | 5.021 | 0.291 |
| Westove | 0.051 | 0.203 | 16.092 | 0.193 |
| | | | | |



| CASE | SOIL | CONC | OBSERVE | YHAT | RESID |
|----------|--------|--------|------------------|-----------|-------------------|
| 1 | 1 | 1 | 0.788 | 0.8252 | -0.0372 |
| 2 | 1 | 1 | 0.871 | 0.8252 | 0.0458 |
| 3 | 1 | 1 | 0.799 | 0.8252 | -0 0262 |
| 4 | 1 | 1 | 0.805 | 0 8252 | -0.0202 |
| 5 | 1 | 1 | 0.863 | 0.8252 | 0.0202 |
| 6 | 1 | 2 | 1.617 | 330 2648 | -328 6478 |
| 7 | 1 | 2 | 1 574 | 330 2648 | -328 6908 |
| 8 | 1 | 2 | 1 493 | 330 2648 | -328 7710 |
| 9 | 1 | 2 | 1645 | 330 2648 | 131/ 7352 |
| 10 | 1 | 2 | 1 64 | 330 2648 | -328 6240 |
| 11 | - | 3 | 52 221 | 55 2028 | |
| 12 | 1 | 3 | 54 016 | 55 2028 | 2.9010 _1 1060 |
| 13 | 1 | 3 | 51 98 | 55 2020 | -3 2220 |
| 14 | 1 | 3 | 60 419 | 55 2020 | -J.ZZZO 5 0160 |
| 15 | 1 | 3 | 57 378 | 55 2028 | 2 1752 |
| 16 | 1 | 4 | 1 63 | 1 7349 | 2.1752 |
| 17 | 1 | 4 | 1 781 | 1 7340 | -0.1048 |
| 18 | 1 | 4 | 1 735 | 1 7240 | 0.0462 |
| 19 | 1 | 4 | 1 716 | 1 7340 | 0.0002 |
| 20 | 1 | 4 | 1.710 | 1 7240 | -0.0188 |
| 21 | 2 | | 1.012 | 1./348 | 0.0772 |
| 22 | 2 | 1 | 0.52 | 0.5116 | 0.0084 |
| 23 | 2 | 1 | 0.490 | 0.5116 | -0.0136 |
| 24 | 2 | 1 | 0.555 | 0.5116 | 0.0214 |
| 25 | 2 | 1 | 0.010 | 0.5116 | 0.0034 |
| 26 | 2 | ⊥ 2 | 0.492 | U.5116 | -0.0196 |
| 27 | 2 | 2 | J.ZII 100 EE7 | 44.523 | -41.312 |
| 28 | 2 | 2 | 100.00/ | 44.523 | 144.034 |
| 29 | 2 | 2 | 24.303 | 44.523 | -20.14 |
| 30 | 2 | 2 | 3.223 | 44.523 | -41.3 |
| 31 | 2 | 4 | J.241 1 071 | 44.523 | -41.282 |
| 32 | 2 | с г | 1 024 | 40346.361 | -40344.48 |
| 33 | 2 | 2 | 1.934 | 40346.361 | -40344.42 |
| 34 | 2 | 2 | 5756 | 40346.361 | -34590.36 |
| 35 | 2 | ່ ວ | 105740 | 40346.361 | -40114.36 |
| 36 | 2 | 3 | 195740 | 40346.361 | 155393.63 |
| 37 | 2 | 4 | 2.307 | 2.3252 | 0.0618 |
| 38 | 2 | 4 | 2.307 | 2.3252 | 0.0418 |
| 39 | 2 | 4 | 2.240 | 2.3252 | -0.0802 |
| 40 | 2 | 4 | 2.309 | 2.3252 | -0.0162 |
| 41 | 2 | 4 | 2.318 | 2.3252 | -0.0072 |
| 42 | 3 | 1 | 0.049 | 0.0754 | -0.0264 |
| 43 | 2 | 1 | 0.072 | 0.0754 | -0.0034 |
| 44 | 3 | 1 | 0.087 | 0.0754 | 0.0116 |
| 45 | 3 | ⊥ 1 | 0.077 | 0.0754 | 0.0016 |
| 46 | 3 | 1 2 | 0.092 | 0.0754 | 0.0166 |
| 47 | 3 | 2 | 0.249 | 0.255 | -0.006 |
| 48 | 3 | 2 | 0.269 | 0.255 | 0.014 |
| 49 | 3 | 2 | 0.232 | 0.255 | -0.023 |
| 50 | 3 | 2 | 0.258 | 0.255 | 0.003 |
| 51 | 3 | 2 | 0.267 | 0.255 | 0.012 |
| 52 | 3 | 2 | 5.969 | 5.0206 | -1.0516 |
| 53 | ר ר | ン っ | 5.904 | 5.0206 | 0.8834 |
| 54 | ר ר | ა ი | 5.UU/ E 100 | 5.0206 | -0.0136 |
| 55 | ン ? | 3 | 5.122 E 101 | 5.0206 | 0.1014 |
| 56 | ン ? | 3 | 5.101 | 5.0206 | 0.0804 |
| 57 | ວ ຈ | 4 | 0.288 | 0.2906 | -0.0026 |
| 52 | ວ ຈ | 4 | 0.282 | U.2906 | -0.0086 |
| 50 59 | ン っ | 4 | 0.289 | 0.2906 | -0.0016 |
| 59 60 | 3 | 4 | 0.297 | 0.2906 | 0.0064 |
| 00 | 3 | 4 | U.297 | 0.2906 | 0.0064 |

| CASE | SOIL | CONC | OBSERVE | YHAT | RESID |
|------|------|------|---------|---------|----------|
| 61 | 4 | 1 | 0.056 | 0.0514 | 0.0046 |
| 62 | 4 | 1 | 0.049 | 0.0514 | -0.0024 |
| 63 | 4 | 1 | 0.057 | 0.0514 | 0.0056 |
| 64 | 4 | 1 | 0.041 | 0.0514 | -0.0104 |
| 65 | 4 | 1 | 0.054 | 0.0514 | 0.0026 |
| 66 | 4 | 2 | 0.202 | 0.203 | -0.001 |
| 67 | 4 | 2 | 0.198 | 0.203 | -0.005 |
| 68 | 4 | 2 | 0.201 | 0.203 | -0.002 |
| 69 | 4 | 2 | 0.221 | 0.203 | 0.018 |
| 70 | 4 | 2 | 0.193 | 0.203 | -0.01 |
| 71 | 4 | 3 | 2.963 | 16.0918 | -13.1288 |
| 72 | 4 | 3 | 63.733 | 16.0918 | 47.6412 |
| 73 | 4 | 3 | 3.017 | 16.0918 | -13.0748 |
| 74 | 4 | 3 | 2.931 | 16.0918 | -13.1608 |
| 75 | 4 | 3 | 7.815 | 16.0918 | -8.2768 |
| 76 | 4 | 4 | 0.188 | 0.1928 | -0.0048 |
| 77 | 4 | 4 | 0.184 | 0.1928 | -0.0088 |
| 78 | 4 | 4 | 0.196 | 0.1928 | 0.0032 |
| 79 | 4 | 4 | 0.197 | 0.1928 | 0.0042 |
| 80 | 4 | 4 | 0.199 | 0.1928 | 0.0062 |

5MEBT, 03/02/99, 10:50

ANALYSIS OF VARIANCE TABLE FOR OBSERVE

| SOURCE | DF | SS | MS | F | P |
|---|-------------------|--|--|----------------------|----------------------------|
| SOIL (A) CONC (B) A*B RESIDUAL | 3 3 9 64 | 1.519E+09 1.522E+09 4.578E+09 3.021E+10 | 5.065E+08 5.075E+08 5.086E+08 4.720E+08 | 1.07 1.08 1.08 | 0.3676 0.3667 0.3916 |
| TOTAL | 79 | 3.783E+10 | | | |

GRAND MEAN 2550.2 SE 2429.1

MEANS OF OBSERVE FOR SOIL

| SOIL | MEAN | SS (MEAN) |
|------|--------|-----------|
| 1 | 97.007 | 2.53E+06 |
| 2 | 10098 | 3.63E+10 |
| 3 | 1.4104 | 88.929 |
| 4 | 4.1348 | 3807.9 |

OBSERVATIONS PER CELL20STD ERROR OF AN AVERAGE4858.2STD ERROR (DIFF OF 2 AVE'S)6870.5ERROR TERM USED: RESIDUAL, 64 DF

MEANS OF OBSERVE FOR CONC

| CONC | MEAN | SS (MEAN) |
|------|--------|-----------|
| 1 | 0.3659 | 2.0858 |
| 2 | 93.811 | 2.57E+06 |
| 3 | 10106 | 3.63E+10 |
| 4 | 1.1359 | 16.918 |

OBSERVATIONS PER CELL20STD ERROR OF AN AVERAGE4858.2STD ERROR (DIFF OF 2 AVE'S)6870.5ERROR TERM USED: RESIDUAL, 64 DF

MEANS OF OBSERVE FOR SOIL*CONC

| SOIL | CONC | MEAN | SS (MEAN) | | |
|--|------|--------|------------|--|--|
| 1 | 1 | 0.8252 | 6.00E-03 | | |
| 1 | 2 | 330.26 | 2.16E+06 | | |
| 1 | 3 | 55.203 | 52.626 | | |
| 1 | 4 | 1.7348 | 0.0194 | | |
| 2 | 1 | 0.5116 | 1.11E - 03 | | |
| 2 | 2 | 44.523 | 26268 | | |
| 2 | 3 | 40346 | 3.02E+10 | | |
| 2 | 4 | 2.3252 | 0.0123 | | |
| 3 | 1 | 0.0754 | 1.12E-03 | | |
| 3 | 2 | 0.2550 | 9.14E-04 | | |
| 3 | 3 | 5.0206 | 1 9032 | | |
| 3 | 4 | 0.2906 | 1 65E-04 | | |
| 4 | 1 | 0.0514 | 1 73E-04 | | |
| 4 | 2 | 0.2030 | 4 54E-04 | | |
| 4 | 3 | 16.092 | 2854 7 | | |
| 4 | 4 | 0.1928 | 1.67E-04 | | |
| OBSERVATIONS PER CELL 5 STD ERROR OF AN AVERAGE 9716.3 STD ERROR (DIFF OF 2 AVE'S) 13741 ERROR TERM USED: RESIDUAL, 64 DF | | | | | |

5MEBT, 03/02/99, 10:51

GRAND MEAN 31.771 SE 27.556

MEANS OF OBSERVE FOR SOIL

| SOIL | MEAN | SS (MEAN) |
|------|--------|-----------|
| 1 | 110.94 | 2.52E+06 |
| 2 | 15.787 | 32470 |
| 3 | 0.2070 | 0.1353 |
| 4 | 0.1491 | 0.0726 |

OBSERVATIONS PER CELL15STD ERROR OF AN AVERAGE55.113STD ERROR (DIFF OF 2 AVE'S)77.941ERROR TERM USED: RESIDUAL, 48 DF

MEANS OF OBSERVE FOR CONC

| CONC | MEAN | SS (MEAN) |
|------|--------|-----------|
| 1 | 0.3659 | 2.0858 |
| 2 | 93.811 | 2.57E+06 |
| 4 | 1.1359 | 16.918 |

OBSERVATIONS PER CELL20STD ERROR OF AN AVERAGE47.729STD ERROR (DIFF OF 2 AVE'S)67.499ERROR TERM USED: RESIDUAL, 48 DF

MEANS OF OBSERVE FOR SOIL*CONC

| SOIL | CONC | MEAN | SS (MEAN) |
|--------------|------------|--------|-----------|
| 1 | 1 | 0.8252 | 6.00E-03 |
| 1 | 2 | 330.26 | 2.16E+06 |
| 1 | 4 | 1.7348 | 0.0194 |
| 2 | 1 | 0.5116 | 1.11E-03 |
| 2 | 2 | 44.523 | 26268 |
| 2 | 4 | 2.3252 | 0.0123 |
| 3 | 1 | 0.0754 | 1.12E-03 |
| 3 | 2 | 0.2550 | 9.14E-04 |
| 3 | 4 | 0.2906 | 1.65E-04 |
| 4 | 1 | 0.0514 | 1.73E-04 |
| 4 | 2 | 0.2030 | 4.54E-04 |
| 4 | 4 | 0.1928 | 1.67E-04 |
| OBSERVATIONS | PER CELL | | 5 |
| STD ERROR OF | AN AVERAGE | 95.4 | 58 |

| STD | ERROR | OF AN | AVER | AGE | | 95.458 |
|------|--------|-------|------|--------|----|--------|
| STD | ERROR | (DIFF | OF 2 | AVE'S) |) | 135.00 |
| ERRO | R TERM | USED | RES | IDUAL, | 48 | DF |

5MEBT, 03/02/99, 10:52

5MEBT, 03/02/99, 10:55

BONFERRONI COMPARISON OF MEANS OF OBSERVE BY SOIL

| SOIL | MEAN | HOMOGENEOUS GROUPS |
|------|--------|-----------------------|
| 2 | 10098 | I |
| 1 | 97.007 | I |
| 4 | 4.1347 | I |
| 3 | 1.4104 | I |

THERE ARE NO SIGNIFICANT PAIRWISE DIFFERENCES AMONG THE MEANS.

| CRITICAL | T VALU | JE | | 2.723 | REJECTION | LEVEL | 0.050 |
|----------|--------|-----|------------|--------|-----------|-------|-------|
| CRITICAL | VALUE | FOR | COMPARISON | 18707 | | | |
| STANDARD | ERROR | FOR | COMPARISON | 6870.5 | | | |

ERROR TERM USED: RESIDUAL, 64 DF

.

BONFERRONI COMPARISON OF MEANS OF OBSERVE BY CONC

| CONC | MEAN | HOMOGENEOUS GROUPS |
|------|--------|-----------------------|
| 3 | 10106 | I |
| 2 | 93.811 | I |
| 4 | 1.1359 | I |
| 1 | 0.3659 | I |

THERE ARE NO SIGNIFICANT PAIRWISE DIFFERENCES AMONG THE MEANS.

| CRITICAL | T VALU | JE | | 2.723 | REJECTION | LEVEL | 0.050 |
|----------|--------|-----|------------|--------|-----------|-------|-------|
| CRITICAL | VALUE | FOR | COMPARISON | 18707 | | | |
| STANDARD | ERROR | FOR | COMPARISON | 6870.5 | | | |

ERROR TERM USED: RESIDUAL, 64 DF

Box and Whisker Plot



.....

5MEBT, 03/02/99, 10:59

| SOIL | MEAN | SAMPLE SIZE | S | .D. | S.E. | |
|--------------------------------------|----------------------------|-------------------|--------------|--------------|--------------------------|--------------------|
| 3 4 DIFFERENCE | 0.0754 0.0514 0.0240 | 5 5 | 0.0 6.581 | 0167 E-03 | 7.49E-03 2.94E-03 | |
| NULL HYPOTHESIS: ALTERNATIVE HYP: | DIFFEREN DIFFEREN | CE = 0 CE <> 0 | | | | |
| ASSUMPTION | Т | DF | | P | 95% CI FOR D | IFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCE | 2.98 S 2.98 | 8 8 8 5. | 0. 2 0. | 0175 0292 | (5.45E-03, (3.56E-03, | 0.0426) 0.0444) |
| TESTS FOR FOULT. | F ۳۷ — | NUM | I DF | DEN DF | Р | |
| OF VARIANC | ES 6.4 | 17 | 4 | 4 | 0.0489 | |
| CASES INCLUDED 1 | 0 MISSI | ING CASES | 0 | | | |

| SOIL | MEA | N S | MPLE IZE | S.D. | S.E. | |
|----------------------------------|--------------|--------------|-------------|--------|--------------------|------------------------|
| 3 | 0.25 | 50 | 5 | 0.0151 | 6.76E-03 | |
| 4 DIFFERENCE | 0.20 0.05 | 30 20 | 5 | 0.0107 | 4.76E-03 | |
| NULL HYPOTHESI ALTERNATIVE HY | S: DIF | FERENCE | = 0 <> 0 | | | |
| ASSUMPTION | | Т | DF | Р | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCE UNEQUAL VARIAN | IS ICES | 6.29 6.29 | 8 7.2 | 0.0002 | (0.0329 (0.0325 | , 0.0711) , 0.0715) |
| TESTS FOR EQUA | TTTY | F | NUM DF | DEN DF | Р | |
| OF VARIA | NCES | 2.01 | 4 | 4 | 0.2573 | |
| CASES INCLUDED | 0 10 | MISSING | CASES 0 | | | |

Test 1 for 5-MeBT looks at the clayey vs. sandy soil comparison.

t _{star} := 2.98 From Statistix

$$t_{025} < t_{star} < t_{975}$$

Table A.5, devour, 22 degree of freedom

t .025 == 2.306 t .975 = 2.306

Do Reject H_o

ORGIN≡0





| SOIL | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|------------------------------------|--------------------------|---------------------|----------|----------------------|--------------------------|
| 1 4 | 0.8252 | 5 | 0.0387 | 0.0173 | - |
| DIFFERENCE | 0.7738 | 5 | 0.001-05 | 2.946-05 | |
| NULL HYPOTHESIS ALTERNATIVE HYP | : DIFFEREN : DIFFEREN | ICE = 0 ICE <> 0 | | | |
| ASSUMPTION | Т | DF | Р | 95% CI FC | R DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANC | 44.0 ES 44.0 | 93 8 93 4. | 0.0000 | 0 (0.733 0 (0.726 | 3, 0.8143) 0, 0.8216) |
| TECTC FOR FOUNT | F | ' NUM | DF DEN | DF P | |
| OF VARIAN | CES 34. | 67 | 4 4 | 4 0.0023 | |
| CASES INCLUDED | 10 MISS | ING CASES | 0 | | |

5MEBT, 03/02/99, 11:02

| SOIL | MEAN | SAMPLE SIZE | S.D. | S.E. |
|--------------------------------------|----------------------------|----------------|------------------|--|
| 1 4 DIFFERENCE | 330.26 0.2030 330.06 | 5 5 | 734.96 0.0107 | 328.68 4.76E-03 |
| NULL HYPOTHESIS: ALTERNATIVE HYP: | DIFFERENC DIFFERENC | E = 0 $E <> 0$ | | |
| ASSUMPTION | Т | DF | Р | 95% CI FOR DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCE | 1.00 1.00 | 8 4.0 | 0.3447 0.3721 | (-427.88, 1088.0) (-582.51, 1242.6) |
| TESTS FOR EQUALI OF VARIANC | F TY ES47591648 | NUM DE | F DEN DF | P 4 0,0000 |
| CASES INCLUDED 1 | 0 MISSI | NG CASES 0 | | |

Test 2 for 5-MeBT looks at the higher f_{oc} vs. low f_{oc} soil comparison.

t _{star} = 44.658 From Statistix

t_{.025} < t_{star} < t_{.975}

Table A.5, devour, 22 degree of freedom

t .025 :=- 2.306 t .975 := 2.306

Do Reject H_o

ORGIN≡0



.

| SOIL | SA MEAN S | AMPLE SIZE | S.D. | S.E. | |
|--------------------------------------|--------------------------|---------------|--------|----------------------|--------------------|
| 1 | 0.8252 | 5 | 0.0387 | 0.0173 | |
| 2 DIFFERENCE | 0.5116 0.3136 | 5 | 0.0167 | 7.45E-03 | |
| NULL HYPOTHESIS: ALTERNATIVE HYP: | DIFFERENCE DIFFERENCE | = 0 <> 0 | | | |
| ASSUMPTION | Т | DF | P | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCE | 16.63 S 16.63 | 8 5.4 | 0.0000 | (0.2701, (0.2662, | 0.3571) 0.3610) |
| TRANC FOR FOURI | F | NUM DF | DEN DF | Р | |
| OF VARIANC | CES 5.41 | 4 | 4 | 0.0654 | |
| CASES INCLUDED 1 | 0 MISSING | G CASES 0 | | | |

5MEBT, 03/02/99, 11:04

| SOIL | SAN MEAN SI | APLE [ZE | S.D. | S.E. | |
|--|-----------------------------------|-------------------|------------------|------------------------|--------------------|
| 1 2 DIFFERENCE | 330.26 14.523 | 5 5 | 734.96 81.037 | 328.68 36.241 | |
| NULL HYPOTHESIS: ALTERNATIVE HYP: ASSUMPTION | DIFFERENCE = DIFFERENCE < T | = 0 <> 0 DF | Ρ | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCES | 0.86 | 8 4.1 | 0.4127 0.4352 | (-476.80, (-623.83, | 1048.3) 1195.3) |
| TESTS FOR EQUALIT OF VARIANCE | F S 82.25 | NUM DF 4 | DEN DF | P 0.0004 | |
| CASES INCLUDED 10 | MISSING | CASES 0 | | | |

Test 3 for 5-MeBT looks at an acclimated vs. unacclimated soil comparison.

t _{star} = 18.098

Table A.5, devour, 22 degree of freedom

t .025 :=-2.306 t .975 := 2.306

Do Reject H_o

ORGIN≡0



5MEBT, 03/02/99, 11:06

| CONC | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|------------------------------------|------------------------------|-------------------|------------------|------------------------|--------------------|
| 2 4 DIFFERENCE | 330.26 1.7348 328.53 | 5 5 | 734.96 0.0697 | 328.68 0.0312 | |
| NULL HYPOTHESIS ALTERNATIVE HYP | S: DIFFEREN(P: DIFFEREN(| CE = 0 CE <> 0 | | | |
| ASSUMPTION | Т | DF | P | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANC | 5 1.00 CES 1.00 | 8 9 4.0 | 0.3468 0.3741 | (-429.42, (-584.04, | 1086.5) 1241.1) |
| TESTS FOR EQUAL | F | | OF DEN DF | P | |
| CASES INCLUDED | 10 MISSI | NG CASES 0 | 4 4 | £ 0.0000 | 1 |

Test 4 for 5-MeBT looks at a PG additive vs. no PG added concentration mix comparison.

 $t_{star} = -5.009$

$$t_{025} < t_{star} < t_{975}$$

Table A.5, devour, 22 degree of freedom

t .025 == 2.306 t .975 = 2.306

Do Reject H_o

ORGIN≡0





TWO-SAMPLE T TESTS FOR OBSERVE BY CONC

| CONC | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|------------------------------------|------------------|----------------|----------|------------------------|--------------------|
| 2 | 44.523 | 5 | 81.037 | 36.241 | |
| 4 DIFFERENCE | 2.3252 42.198 | 5 | 0.0555 | 0.0248 | |
| NULL HYPOTHESIS ALTERNATIVE HYP | DIFFERENC | E = 0 $E <> 0$ | | | |
| ASSUMPTION | Т | DF | Р | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANC | 1.16 ES 1.16 | 8 4.0 | 0.2778 | (-41.374, (-58.423, | 125.77) 142.82) |
| | F | NUM DI | F DEN DF | Р | |
| TESTS FOR EQUAL OF VARIAN | CES2133388. | 64 4 | 4 | 0.0000 | |

CASES INCLUDED 10 MISSING CASES 0

5MEBT, 03/02/99, 11:09

| CONC M | | SAM SI | PLE ZE | S.D. | S.E. | |
|--------------------------------|----------------------|-----------|-----------|--------|------------|------------|
| 2 | 0.255 | 0 | 5 (| 0.0151 | 6.76E-03 | |
| 4 | 0.290 | 6 | 5 6.4 | 43E-03 | 2.87E-03 | |
| DIFFERENCE | -0.035 | 6 | | | | |
| NULL HYPOTHES ALTERNATIVE H | IS: DIFF YP: DIFF | ERENCE = | 0 > 0 | | | |
| ASSUMPTION | | Т | DF | Р | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANC | ES | -4.85 | 8 | 0.0013 | (-0.0525, | -0.0187) |
| UNEQUAL VARIA | NCES | -4.85 | 5.4 | 0.0038 | (-0.0541, | -0.0171) |
| TESTS FOR FOU | ልፐ.ፐጥሃ | F | NUM DF | DEN DF | P | |
| OF VARI | ANCES | 5.53 | 4 | 4 | 0.0631 | |
| CASES INCLUDE | D 10 | MISSING (| CASES 0 | | | |

5MEBT, 03/02/99, 11:11

| CONC | MEAN | SAMPLE SIZE | S.D. | S.E. | |
|--------------------------------------|-------------------------|----------------|--------------------|--------------------------|------------------------|
| 2 0 4 0 DIFFERENCE 0 | .2030 .1928 .0102 | 5 5 | 0.0107 6.46E-03 | 4.76E-03 2.89E-03 | |
| NULL HYPOTHESIS: ALTERNATIVE HYP: | DIFFERENC DIFFERENC | E = 0 $E <> 0$ | | | |
| ASSUMPTION | T | DF | P | 95% CI FOR | DIFFERENCE |
| EQUAL VARIANCES UNEQUAL VARIANCES | 1.83 1.83 | 8 6.6 | 0.1045 0.1125 | (-2.65E-03 (-3.14E-03 | , 0.0230) , 0.0235) |
| TESTS FOR EQUALITY | F | NUM | DF DEN DF | P | |
| OF VARIANCE: | 5 2.72 | 2 4 | 4 | 0.1778 | |
| CASES INCLUDED 10 | MISSIN | NG CASES | C | | |

BIBLIOGRAPHY

Arco Chemical Company, "Biodegradation and Toxicity of Glycols. Newton Square: Arco Chemical Company Report, May 1990.

Atlas, Ronald M., Bartha, Richard, <u>Microbial Ecology: Fundamentals and Applications</u>. Benjamin Cummings: Redwood City, 1993.

Ball, William P., Roberts, Paul V., "Long-term Sorption of Halogenated Organic Chemicals by Aquifer Material. 2. Intraparticle Diffusion", <u>Environmental Science Technology</u>, 25: 1238. 1996.

Bausmith, D.S., Neufeld, R.D., "Land Treatment of Aircraft Deicing Fluids." WEFTEC '96, Dallas, TX. 1996.

Brusseau, M.L., Rao, P.S.C., Jessup, R.E., Davidson, J.M., "Borden, ON. Site." Contaminant Hydrology. 4: 223-240. 1989.

Burke, Baron, Capt., "Biodegradation of Aircraft Deicing Fluid Components in Soil.", Air Force Institute of Technology, WPAFB, OH, Air University, 1999.

Cancilla, D.A., A. Holtkamp, L. Matassa and X. Fang, "Isolation and Characterization of Microtox[®] Active Components from Aircraft Deicing/Anti-icing Fluids.", <u>Environmental Toxicology and Chemistry</u>, March 1997. 16(3): 430-434.

Cancilla, D.A., Martinez, J., Van Aggelen, G.C., "Detection of Aircraft Deicing/Antiicing Fluid Additives in a Perched Water Monitoring Well at an International Airport." <u>Environmental</u> <u>Science and Technology</u>. 32(28):3834-3835. Dec 1, 1998.

Chesterfield, B. Major, Bean, Jim C., and Dawson Tony, "Air Force Deicing Fluid." Electronic Publication: <u>http://www-afsc.saia.af.mil/magazine/htdocs/Octmag96</u>. 1997.

Cornell, Jeff S., Pollard, David A., Hernandez, Mark T., "Chemical Components of Aircraft Deicer Fluid: How They Affect Propylene Glycol Degradation Rates and Deicing Waste Stream Toxicity". Water Environment Federation Technical Conference, Orlando, FL, 1998.

., "Aspects of the Chemistry, Toxicity, and Biodegradability of Benzotriazole and It's Derivatives." 1997.

., University of Colorado – Boulder, Ph.D. Student, Boulder, Colorado, Telephone interviews and e-mail, 1998.

<u>Cross-Talk</u>, "Exploring Available De-icing Technologies.", Pro-Act: San Antonio, TX., 24 May 1996.

Devore, Jay L., <u>Probability and Statistics for Engineering and the Sciences</u>. 4th ed., Duxbury Press, 1995.

EA-Engineering, Science and Technology, R.S.a.M.G., Comparative Toxicity and Environmental Impact of Ethylene Glycol and Propylene Glycol: A Review. Silver Spring, MD. EA-Engineering, Science and Technology. 1996.

ENSR Consulting . Evaluation of the Biotic Communities and Chemistry of the Water and Sediments in Sand Creek Near Stapleton IA, Denver, CO Stapleton IA Authority. 1993.

Ferrante, Dan A., Gallagher, Daniel L., "Groundwater Pollution Primer", Class Notes, Civil Engineering Department, Virginia Tech University, 1998.

Fetter C.W., Contaminant Hydrogeology. Prentice Hall, Upper Saddle River, NJ, 1993.

Gooden, W., "Aircraft Deicing Fluid Biodegradation." M.S. Thesis, ChEng, Ft. Collins, Colorado State University. 1997.

Gruden, Cyndee L., Experimental Protocol – Sorption Experiments, University of Colorado – Boulder. 1998.

Fisher, D.J., Knott, M.H., "The Acute Whole Effluent Toxicity of Storm Water from an International Airport." <u>Environmental Toxicology and Chemistry</u>. 14(6): 1103-1111. 1995.

Hach DR/2000 Spectrophotometer, "Measure Suspended Solids (0-750 mg/L)." <u>Hach Co.</u> <u>Procedures Manual</u>. Advanced Water Quality Laboratory. 1993, 1994.

Halterman-O'Malley, AM,. 2Lt., "Biodegradation of Deicing Agents in Various Soil Types." Air Force Institute of Technology, WPAFB, OH, Air University. 1997.

Heyse, Edward Lt Col., AF Center for Environmental Excellence, Personal interviews and emails, 1998-1999.

Hartwell, S.I., Jordahl, D.M., "Toxicity of Aircraft Deicer and Anti-icer Solutions to Aquatic Organisms." <u>Environmental Toxicology and Chemistry</u>. 14(8): 1375-1386. 1995.

HP ChemStation, Hewlett Packard. 3rd ed. HP Part No. G2070-90104, 1994, 1995.

HQ Air Force Center for Environmental Excellence, PRO-ACT Fact Sheet, TI6159, July 1995.

HQ Air Force Center for Environmental Excellence, PRO-ACT Fact Sheet, Dec 1995.

Johnson, Laura M., Capt., "Evaluation of the Natural Biodegradation of Aircraft Deicing Fluid Components in Soils." Air Force Institute of Technology, WPAFB, OH, Air University. 1997. Karickhoff, S. W., "Organic Pollutant Sorption in Aquatic Systems." Journal of Hydraulic Engineering, 110(6): 707-35. 1984.

Lee, C.C., Environmental Engineering Dictionary. 2nd ed. Government Institute, CA. pp. 216. 1992.

Lee, C-K., Chiang, A.S.T., "Lattice Model for the Adsorption of Benzene in Silicalite I." <u>AIChE</u> Journal. 38(1): 128-135. 1992.

Lin, T-F., Van Loy, M.D., Nazaroff, W.W., "Gas-Phase Transport and Sorption of Benzene in Soil." <u>Environmental Science and Technology</u>. 30(7): 2178-2186. 1996.

Lyman, W.J., Reehl, W.F., <u>Handbook of Chemical Property Estimation Methods, Environmental</u> <u>Behavior of Organic Compounds</u>. Arthur D. Little, Inc., Cambridge, MA., McGraw-Hill, Inc., 1982.

Maagd, P. G-J., Sinnige, Theo L., Schrap, S. Marca, Opperhuizen, A., Sijm, D., "Sorption Coefficients of Polycyclic Aromatic Hydrocarbons for Two Lake Sediments: Influence of the Bactericide Sodium Azide.", <u>Environmental Toxicology and Chemistry</u>, 17(10): 1899-1907. 1998.

Matulewicz, W., Degradation Studies of Cobratec 99 and Cobratec 100. 1997.

Merck Index, 12th ed., #8040, 1996.

Mericas, DD., Wagoner, B., "Balancing Safety and the Environment." <u>Water Pollution</u>. 16: 301-315. 1994.

Morrison, Robert T., Boyd, Robert N., Organic Chemistry. 2nd ed., Boston: Allyn and Bacon, Inc., 1971.

NASA, Downs, R.W., "Chemically Inducted Ignition in Aircraft and Spacecraft Electrical Circuitry by Glycol/Water Solutions." NASA Manned Spacecraft Center, Houston, TX. 1968.

Network Marketing, Propylene Glycol: Antifreeze and Brake Fluid, Electronic reference, <u>http://www.successteams.com</u>. 1998.

NRDC, "Environmental Groups to Sue BWI and O'Hare Airports." Press Release: <u>http://www.nrdc.org/news</u>, New York. 1998.

NTP Chemical Repository (Radian Corporation), Tolyltriazole, Catalog ID Number: 001439. http://ntp-do.niehs.nih.gov. August 29, 1991.

Oakley, Monica M., Forrest, Carol L., "The Clock is Ticking to Comply with New Stormwater Regulations." <u>Water Environment and Technology</u>. 51-66, March 1991.

Ott, R. Lyman, <u>An Introduction to Statistical Methods and Data Analysis</u>. 4th ed., Duxbury Press, 1993.

Park, I., Knaebel, K.S., "Adsorption Breakthrough Behavior: Unusual Effects and Possible Causes." <u>AIChE Journal</u>. 38(5): 660-670. 1992.

Pillard, D.A., "Comparative Toxicity of Formulated Glycol Deicers and Pure Ethylene and Propylene Glycol to *Ceriodaphnia dubia* and *Pimephales promelas*." <u>Environmental Toxicology</u> and <u>Chemistry</u>. 14(2): 311-315. 1995.

PMC Specialties Group, Inc., Cobratec[®] 99/100 Technical Bulletin, Rocky River, OH. PMC Specialties Group, Inc. 1996.

., Materials Safety Data Sheet (MSDS), Cobratec® TT-100, 1996.

RACB. 1,2-Propanediol (Propylene Glycol), CAS# 57-55-6, Study Number RACB84068, Electronic Reference, <u>http://www.ntp-server.niehs.nih.gov</u>. 1998.

SAE. "Specification for Deicing/Antiicing Fluid, Aircraft, SAE Type I." AMS 1424, Warrendale, PA. 1992.

Schwarzenbach, Philip M., Gschwend, and Dieter, Imboden M. Environmental Organic Chemistry, Wiley & Sons, Inc., New York, NY, 1993.

Sigma Products Information Sheet. 1,2,Propanediol, Electronic reference. <u>http://www.sigma.sial.com</u>. 1998.

Sills, Robert D., Blakeslee, Paul A., <u>The Environmental Impact of Deicers in Airport Stormwater</u> <u>Runoff</u>. Michigan Department of Natural Resources-Environmental Impact Study, Lansing, MI: Surface Water Quality Division, March 1991.

Standard Methods for the Examination of Water and Wastewater, "2540B. Total Solids Dried at 103-105°C." 18th ed. pp. 2-54. 1992.

Tissue, B.M., Analytical Chemistry, SCIMEDIA. Electronic Publication, <u>http://www.scimedia.com</u>. 1996.

United States Environmental Protection Agency (USEPA). <u>Overview of the Storm Water</u> <u>Program</u>. Washington: Government Printing Office, 13 December, 1991.

United States Environmental Protection Agency (USEPA)."Emerging Technology Report: Preliminary Status of Airplane Deicing Fluid Recovery Systems." EPA 832-B-95-005, Washington DC. 1995. United States Patent 5,503,775. Rao, N.M., Lu, F.F., Johnson, D.A., Nghiem, N.P., "Method of Preventing Yellow Metal Corrosion in Aqueous Systems with Superior Corrosion Performance in Reduced Environmental Impact."

Yaniga, Paul; Bioremediation Specialist, Entrepreneur, Personal interview, 25 Aug 98.

Weber, Walter J., Weilin, Haung, "A Distributed Reactivity Model for Sorption by Soils and Sediments. 4. Intraparticle Heterogeneity and Phase-Distribution Relationships under Nonequilibrium Conditions", <u>Environmental Science Technology</u>, 30: 881-888. 1993.

Yun, Jeong-Ho, "Adsorption isotherms of Benzene and Methylbenzene vapors on activated carbon", Journal of Chemical and Engineering Data, 42: 894-896 (Sept/Oct '97).

<u>Vita</u>

First Lieutenant David L. Kellner was born in Buffalo, New York on February 11, 1968. After high school graduation he obtained an Associate Degree of Applied Science in Accounting in 1989. He then joined the United States Air Force in November 1989 as a Nondestructive Inspection Specialist in the 28th Maintenance Squadron at Ellsworth AFB, South Dakota after a 3 month technical school at Chanute AFB, IL. He was married on April 21, 1990 to Nancy Sunick. While at Ellsworth AFB, he attended the South Dakota School of Mines and Technology and obtained his Bachelor of Science in Civil Engineering in only 4 and ¹/₂ years. Upon his graduation from Officers Training School (OTS) 22 September 1995 at Maxwell AFB, Alabama: he was assigned to the 437th Civil Engineer Squadron, Charleston AFB, South Carolina from 28 September 95 to 13 August 1997. While at Charleston AFB, he served as a Project Officer (32E1G), Section Commander (C36P1), and Readiness Flight Commander (32E3B). He was selected to study for his Masters of Science in Engineering and Environmental Management (GEEM) at the Air Force Institute of Technology from August 1997 to March 1999. Upon completion of the GEEM program, he will be assigned to the 7th Civil Engineer Squadron, Dyess AFB, Texas.

| REPORT DOCUMENTATION PAGE | | | | | Form Approved OMB No. 0704-0188 | | |
|--|---------|-----------------------------|----------------------------|---------------|---|--|--|
| Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of the Office and Budget, Paperwork Reduction Project (0704-0188), Washinaton. DC 20503. | | | | | | | |
| 1. AGENCY USE ONLY (Leave b | lank) | 2. REPORT DATE | 3. REPORT TYPE AM | ND DATE | S COVERED | | |
| | | March 1999 | | Maste | er Thesis | | |
| 4. IIILE AND SUBITILE | o Fluid | Component Methyl Ben | rotriazola in Sail | 5. FUN | DING NUMBERS | | |
| Solption of the Anterant Determ | g Fluit | r component memyi-ben | | | | | |
| 6. AUTHOR(S) | | | | | | | |
| DAVID L. KELLNER, 1Lt, U | JSAF | | | | | | |
| 7. PERFORMING ORGANIZATIO | | E(S) AND ADDRESS(ES) | | 8. PER | FORMING ORGANIZATION | | |
| All Force institute of Technolo | ogy | | | | | | |
| Wright-Patterson AFB, OH 45 | 433-77 | 65 | | A | AFIT/GEE/ENV/99M-11 | | |
| | | | | | | | |
| Capt Steven M. Anderson | AGENC | Y NAME(S) AND ADDRESS | ES) | 10. SPC AG | DNSORING/MONITORING ENCY REPORT NUMBER | | |
| 130 Barnes Drive Suite 2 | | | | | | | |
| Tyndal AFB, FL 32403-5323 | | | | | | | |
| 11. SUPPLEMENTARY NOTES | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| 12a. DISTRIBUTION AVAILABILIT | Y STA | TEMENT | | 12b. Di | STRIBUTION CODE | | |
| | | | | | | | |
| Approved for Public Release, Distribution Unlimited | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| 13. ABSTRACT (Maximum 200 w | ords) | | | | | | |
| The air transportation industry | require | es deicing/anti-icing (ADA | F) fluids to maintain fli | ght oper | ations in cold weather | | |
| conditions. Additives to the Al | DAF ai | d in the performance of the | nese agents. Approxima | tely 52N | 1 liters of concentrated | | |
| aircraft deicing fluid are used in | n Nortl | n America per year in an | attempt to correct this sa | fety prol | blem. Among the additives to | | |
| be toxic to microorganisms Id | / Hame | retardants, wetting agent | s, and thickening agents | . Some | of these agents are known to | | |
| step to understanding how to re | mediat | e ADAF Methyl-Benzot | riazole (MeBT) is a corr | me envir | bibitor/flome retendent torio | | |
| additive in ADAF. This study investigates the sorption characteristics of MeRT in various soils. Three soil times more used | | | | | | | |
| to evaluate the sorption characteristics of MeBT at 1000 mg/L. 100 mg/L. and a mixture of MeBT and provulene given | | | | | | | |
| (PG). A High Pressure Liquid Chromatograph (HPLC) was used to conduct the recovery experimentation after the samples | | | | | | | |
| had reached equilibrium. Resulting sorption coefficients revealed that MeBT did not sorb well to any of the soils types used. | | | | | | | |
| The organic carbon content affects the sorption coefficient by slightly raising the sorbed amount of MeBT. | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| 14. SUBJECT TERMS | | | | | | | |
| Adsorption, Sorption Characteristics, Sorption Isotherm, ADF, Tolyltriazole, MeBT, Hi | | | | | | | |
| Performance Liquid Chromatography | | | | | 16. PRICE CODE | | |
| 17. SECURITY CLASSIFICATION | 18.5 | CURITY CLASSIFICATION | 19 SECURITY OF ASSIST | CATION | | | |
| OF REPORT | 0 | F THIS PAGE | OF ABSTRACT | | 20. LINITATION OF ABSTRACT | | |
| Unclassified | | Unclassified | Unclassified | | UL | | |

Standard Form 298 (Rev. 2-89) (EG) Prescribed by ANSI Std. 239.18 Designed using Perform Pro, WHS/DIOR, Oct 94