

AD _____

Award Number: DAMD17-96-C-6105

TITLE: Treatment of Metastatic Breast Carcinoma Refactory to
Doxorubicin with Liposomal-Annamycin

PRINCIPAL INVESTIGATOR: Aysegul A. Sahin, M.D.

CONTRACTING ORGANIZATION: University of Texas
MD Anderson Cancer Center
Houston, Texas 77030

REPORT DATE: October 1999

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 1999	3. REPORT TYPE AND DATES COVERED Annual (23 Sep 98 - 22 Sep 99)
----------------------------------	--------------------------------	--

4. TITLE AND SUBTITLE Treatment of Metastatic Breast Carcinoma Refractory to Doxorubicin with Liposomal-Annamycin	5. FUNDING NUMBERS DAMD17-96-C-6105
--	--

6. AUTHOR(S)
Aysegul A. Sahin, M.D.

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)
University of Texas MD Anderson Cancer Center
Houston, Texas 77030

E-MAIL:
perezr01@gcrc.med.nyu.edu

8. PERFORMING ORGANIZATION REPORT NUMBER

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)
U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT
Approved for public release; distribution unlimited

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 Words)

Fourteen patients with metastatic breast carcinoma refractory to doxorubicin were entered in a Phase II study of liposomal-annamycin, a new anthracycline antibiotic which has shown ability to circumvent multidrug resistance. A total of 25 doses of liposomal-annamycin were given at three different dose levels. Toxicity was mild and limited to granulocytopenia in less than 50% of patients. No antitumor responses were seen.

14. SUBJECT TERMS
Breast Cancer

20010228 073

15. NUMBER OF PAGES 7
16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT
Unclassified

18. SECURITY CLASSIFICATION OF THIS PAGE
Unclassified

19. SECURITY CLASSIFICATION OF ABSTRACT
Unclassified

20. LIMITATION OF ABSTRACT
Unlimited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.



PI - Signature

10-19-99

Date

TABLE OF CONTENTS

Front cover	1
SF 298	2
Foreword	3
Table of Contents	4
Introduction	5
Report	5
Conclusions	7
References	7

INTRODUCTION

Liposomal-Annamycin is a liposome entrapped new anthracycline antibiotic which has shown lack of cross-resistance in vitro and in vivo in different cell lines that express P-glycoprotein and MRP. In a Phase I study conducted in patients with solid tumors, the dose limiting toxicity was myelosuppression. No alopecia, mucositis, cardiac, skin, nor gastrointestinal toxicities were observed. The maximum tolerated dose was 210 mg/m² administered intravenously every 3 weeks. Because the multidrug resistance phenotype has been associated with some human malignancies, particularly acute leukemia and breast carcinoma, when they become refractory to standard chemotherapy, we proposed and initiated a Phase II study of liposomal-Annamycin in patients with metastatic breast carcinoma refractory to doxorubicin. This report summarizes the status of this study.

REPORT

Because of a transfer of the PI to New York University Medical Center, work during the last period of support was very limited. An extension of one year will be needed and requested to complete this project.

ELIGIBILITY CRITERIA

1. Metastatic breast carcinoma
2. Anthracycline-resistant
3. Measurable disease
4. Life expectancy >12 weeks
5. Prior anthracycline <350 mg/m² of doxorubicin equivalent by bolus, <450 mg/m² by prolonged infusion
6. Adequate bone marrow function
7. Ejection fraction >55%

PATIENT CHARACTERISTICS

A total of fourteen patients have been entered in the study and a total of 25 doses of liposomal-annamycin have been given. The following Table summarizes the characteristics of the patients entered.

Number of patients entered	14
Number of patients evaluable	14
Age median (range)	50 (34-73)
Performance status	

1	12
2	2
Sex: female	14
Race:	
Black	7
Hispanic	3
White	4
Histology	
Ductal carcinoma, invasive	14
Prior therapy	
Chemotherapy	14
Hormonal therapy	3
Radiation therapy	6
Surgical therapy	7
Prior chemotherapy: number of regimens	
1	2
2	2
3	8
4	2
number of agents	
3	1
>3	13

TOXICITY

Toxicity has been mild and mostly limited to granulocytopenia as indicated below:

1. Myelosuppression (first 9 patients)

Dose	No courses	Granulocyte nadir (range) (day)	Platelet nadir (range) (day)
190mg/m ²	4	1.2 (0.7-1.8) 11	192 (134-251) 14
210	8	1.5 (0.5-3.0) 15	218 (145-288) 10

250	2	0.8 (0.3-1.3) 11	169 (146-192) 8
-----	---	---------------------	--------------------

2. Non-hematological toxicities

Nausea and vomiting was observed after 6 courses (grade 3 in one, grade 2 in two, and grade 1 in three). Diarrhea grade 1 was observed after one course. Stomatitis was observed after 4 courses (grade 2 in two, and grade 1 in two). One grade 1 allergic reaction was observed.

ANTITUMOR ACTIVITY

Fourteen patients progressed after 1 or 2 cycles of liposomal-annamycin. One patients remained stable after 2 cycles of liposomal-annamycin. No tumor responses have been seen so far.

CORRELATIVE TISSUE STUDIES

Three tissue specimens were obtained pre-therapy for MDR analysis. These samples are kept frozen in Dr. Sahin's laboratory and will be assayed when the study is completed and/or responses are seen.

CONCLUSIONS

Results obtained to date suggest that liposomal-annamycin is very well tolerated with grade 3 granulocytopenia being observed in a minority of patients. Non-hematological toxicity is minimal.

No tumor responses have been observed so far.

The study will continue to complete fourteen fully evaluable patients. Baseline tumor biopsies will be obtained to analyze MDR status. Because no responses have been seen in this cohort of 14 patients, further accrual is restricted to patients with MDR ⊕ tumors.

REFERENCES

1. Booser D, Zou Y, Priebe W, Perez-Soler R. Phase I clinical and pharmacology study of liposomal-annamycin. In preparation for submission to Clinical Cancer Research.
2. Booser D, Esparza-Guerra L, Zou Y, Priebe W, Perez-Soler R. Liposomal-annamycin. Phase I clinical and pharmacological study. Proceed. ASCO 16:762 p217a, 1997.