

REPORT DOCUMENTATION PAGE

Form Approved
OMB NO. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 15 Dec 00	3. REPORT TYPE AND DATES COVERED Final 1 Mar 99-31 Aug 00	
4. TITLE AND SUBTITLE Monitoring Low Temperature Physiology in Hibernating Mammals			5. FUNDING NUMBERS DAAD19-99-1-0071	
6. AUTHOR(S) Matthew T. Andrews				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) North Carolina State University Department of Genetics Box 7614 Raleigh, NC 27695-7614			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 39467.1-LS-RIP	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12 b. DISTRIBUTION CODE 20010227 133	
13. ABSTRACT (Maximum 200 words) We have successfully identified genes controlling the physiological characteristics of hibernation in the thirteen-lined ground squirrel (<i>Spermophilus tridecemlineatus</i>). Identification of these specific genetic markers has allowed us to study the function of their gene products under highly controlled conditions of temperature, light and humidity. To accomplish this goal specific instrumentation was purchased with this DURIP grant. Measurements of low-temperature physiology in hibernating mammals under controlled conditions required an environmental chamber that allows us to maintain a variety of ambient temperatures ranging from -5 to 20°C. The response and adaptation of our ground squirrel species to environmental change has been monitored by precisely measuring activity, core body temperature and heart rate using miniature, battery-free, implantable transmitters that allow uninterrupted lifetime studies. Tissues collected from these animals were quick-frozen in liquid N ₂ and placed in a -86°C freezer where they are stored until we analyze patterns of gene expression controlling the physiological characteristics of hibernation in mammals.				
14. SUBJECT TERMS hibernation, gene expression, physiological monitoring			15. NUMBER OF PAGES 7	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

MASTER COPY: PLEASE KEEP THIS "MEMORANDUM OF TRANSMITTAL" BLANK FOR REPRODUCTION PURPOSES. WHEN REPORTS ARE GENERATED UNDER THE ARO SPONSORSHIP, FORWARD A COMPLETED COPY OF THIS FORM WITH EACH REPORT SHIPMENT TO THE ARO. THIS WILL ASSURE PROPER IDENTIFICATION. NOT TO BE USED FOR INTERIM PROGRESS REPORTS; SEE PAGE 2 FOR INTERIM PROGRESS REPORT INSTRUCTIONS.

MEMORANDUM OF TRANSMITTAL

U.S. Army Research Office
ATTN: AMSRL-RO-BI (TR)
P.O. Box 12211
Research Triangle Park, NC 27709-2211


- | | |
|--|---|
| <input type="checkbox"/> Reprint (Orig + 2 copies) | <input type="checkbox"/> Technical Report (Orig + 2 copies) |
| <input type="checkbox"/> Manuscript (1 copy) | <input checked="" type="checkbox"/> Final Progress Report (Orig + 2 copies) |
| | <input type="checkbox"/> Related Materials, Abstracts, Theses (1 copy) |

CONTRACT/GRANT NUMBER: DAAD19-99-1-0071

REPORT TITLE: **DURIP: Monitoring Low Temperature Physiology in Hibernating Mammals**

is forwarded for your information.

SUBMITTED FOR PUBLICATION TO (applicable only if report is manuscript):



Sincerely,

Matthew T. Andrews

Matthew T. Andrews
Professor, Dept. of Biochemistry & Molecular Biology
University of Minnesota School of Medicine
10 University Drive
Duluth, MN 55812
Telephone: 1-218-726-7271
e-mail: mandrews@d.umn.edu

Enclosure 3

b. FINAL PROGRESS REPORT

DURIP: Monitoring Low Temperature Physiology in Hibernating Mammals

Matthew T. Andrews, P. I.

Statement of the problem studied

Mammals in deep hibernation have an observed respiratory quotient (RQ) of 0.7 (Lyman, 1982) RQ is a unit-less value representing the moles of CO₂ respired per moles of O₂ consumed. A value of 1.0 indicates combustion of carbohydrate; however, an RQ of 0.7 indicates that fat is the major substrate for energy production. This fuel selection of fat over carbohydrate seen during hibernation also occurs in starvation, when conservation of glucose for utilization by the brain is essential for survival and is controlled by the phosphorylation of pyruvate dehydrogenase (Randle, 1986) Since hibernating ground squirrels can survive 5-6 months without feeding, mechanistic insight into the activation of genes such as PDK-4 could provide a better understanding of how nutrient-gene interactions influence an animal's capacity to cope with the effects of starvation. Recent experiments with rats have shown that starvation for a period of 12 to 24 hours results in a sharp increase in heart PDK-4 mRNA (Wu, 1998) As seen with ground squirrel hearts at spring arousal, this up-regulation of PDK-4 in rats is reversible as heart PDK-4 mRNA returns to control levels upon re-feeding.

It has been known for over 30 years that diabetes, like starvation, causes an overall reduction in cardiac glucose oxidation (Bleehen, 1954; Morgan, 1961; Morgan, 1961) An examination of working hearts in diabetic pigs has shown that the major defect in carbohydrate catabolism was not in glycolysis, but in the oxidation of pyruvate by pyruvate dehydrogenase (Hall, 1996). Experiments with rat hearts (Wu, 1998) have shown that high levels of PDK-4 expression were induced in hearts of animals rendered diabetic by treatment with streptozotocin - an antibiotic that destroys the insulin producing capacity of the pancreas. As with starvation and spring arousal, this induction is reversible as high levels of both PDK-4 protein and mRNA in the hearts of diabetic animals are reduced to control levels upon treatment with insulin (Table 1). As a positive regulator of glucose uptake and utilization, it has been a long-standing observation that insulin inhibits heart PDK activity (Hutson, 1978) Only now is it evident that this effect is mediated by regulating the expression of the gene encoding PDK-4.

Table 1. Conditions affecting PDK-4 expression in the heart

PDK- 4 ↑ - stable increase in PDK-4 mRNA levels

1. Hibernation (ground squirrel; Andrews *et al.*, 1998)
2. Starvation (rat; Wu *et al.*, 1998)
3. Diabetes (rat; Wu *et al.*, 1998)

PDK- 4 ↓ - reversal of PDK-4 mRNA to control levels (refs. for each numbered item same as above)

1. Spring arousal
 2. Starvation + re-feeding
 3. Diabetes + insulin
-

Summary of the most important results

The thirteen-lined ground squirrel, a characteristic deep hibernator, slows its metabolism and drops its body temperature to survive the harsh winter months when food is scarce. To survive without eating, meticulous regulation of energy resources is essential during hibernation. In the heart of the thirteen-lined ground squirrel, this regulation is accomplished by changes in gene expression that result in a switch from carbohydrate catabolism to fatty acid oxidation. Using a PCR-based subtractive hybridization scheme, three genes encoding mitochondrial proteins were found to be differentially expressed in the hearts of hibernating ground squirrels. One gene, pyruvate dehydrogenase kinase isozyme 4 (PDK-4), is nuclear encoded and is upregulated during hibernation. PDK-4 acts by phosphorylating pyruvate dehydrogenase (PDH) and attenuating its activity. PDH catalyzes the oxidation of pyruvate to acetyl CoA, the first irreversible step in glycolysis. Northern blot analysis shows that PDK4 message is upregulated 20-fold in heart and 5-fold in skeletal muscle during hibernation.

Unlike PDK-4, the two remaining differentially expressed genes are located in the mitochondrial genome and are down regulated during hibernation in heart and *ndh6* is down in skeletal muscle. These genes encode NADH dehydrogenase 5 (NDH5) and NADH dehydrogenase 6 (NDH6), two subunits of NADH ubiquinone-oxidoreductase (complex I). Complex I is a large multi-protein component of the mitochondrial electron transport chain containing more than 30 subunits and is the first carrier of electrons in the system. NDH5 is transcribed from the mitochondrial H-strand along with 11 other protein coding genes. Although NDH5 is down

regulated, genes flanking either side of it on the H-strand are not. This difference in expression suggests a specific mechanism for reduction of NDH5 message. NDH6, on the other hand, is the only L-strand encoded protein. Its open reading frame partially overlaps with NDH5. Down-regulation of NDH5 and NDH6 in combination with the upregulation of PDK-4 is concordant with the observed reduction in mitochondrial state 3 respiration during hibernation in mammals.

In the mitochondrial genome, the bi-directional promoter region regulates transcriptional control for all genes in the mitochondria. Because *ndh6* is the only protein encoding gene on the L strand, a reduction in L strand initiation would curtail *ndh6* mRNA. Transcriptional initiation, however, cannot explain the reduction in *ndh5* mRNA. *Ndh5* is down-regulated, but flanking genes are not. The mechanism for the down-regulation of a single gene on the H strand is still unknown.

Differences in tissue-specific expression patterns for *ndh5* and *ndh6* imply distinct mechanisms for control. Decreased expression of *ndh6* is not muscle-type specific. In an individual animal, *ndh6* is down-regulated in both heart and skeletal muscle suggesting an organismal signal or factor causing the abated *ndh6* expression during hibernation. *Ndh5*, on the other hand, is down-regulated in heart but not skeletal muscle. This divergence implies a tissue-specific signal or factor controlling its expression.

List of all publications and technical reports

Published articles:

Squire, T.L. and Andrews, M.T. (2000) Genetic control of carbon utilization during hibernation: mechanistic considerations, in *"Life in the Cold"* (G. Heldmaier, S. Klaus, M. Klingenspor, Eds.) pp. 325-337, Springer-Verlag, Berlin.

Abstracts:

Buck, M.J. and Andrews, M.T. (2000) Coordinated regulation of PDK-4 in cardiac, skeletal muscle, and adipose tissue during hibernation in the thirteen-lined ground squirrel. Eleventh International Hibernation Symposium, Jungholz, Austria, August 13-18, 2000.

Squire, T.L., Bauer, V.W., Lowe, M.E., and Andrews, M.T. (2000) Genetic regulation of lipolysis during hibernation. Eleventh International Hibernation Symposium, Jungholz, Austria, August 13-18, 2000.

Participating scientific personnel

GRADUATE STUDENTS

Teresa Squire - Ph.D. in Genetics, expected May 2001
Michael Buck
Jayita Guhaniyogi

UNDERGRADUATE RESEARCH PROJECTS DIRECTED

Luis Miguel Gonzalez, 1999-2000

Report of inventions

None

Bibliography

- Andrews, M. T., Squire, T. L., Bowen, C. M., and Rollins, M. B. (1998). Low-temperature carbon utilization is regulated by novel gene activity in the heart of a hibernating mammal. *Proc Natl Acad Sci U S A* 95, 8392-7.
- Bleehen, N. M., and Fisher, R. B. (1954). The action of insulin in the isolated rat heart. *J. Physiol. (Lond.)* 123, 260-276.
- Hall, J. L., Stanley, W. C., Lopaschuk, G. D., Wisneski, J. A., Pizzurro, R. D., Hamilton, C. D., and McCormack, J. G. (1996). Impaired pyruvate oxidation but normal glucose uptake in diabetic pig heart during dobutamine-induced work. *Am J Physiol* 271, H2320-9.
- Hutson, N. J., Kerbey, A. L., Randle, P. J., and Sugden, P. H. (1978). Conversion of inactive (phosphorylated) pyruvate dehydrogenase complex into active complex by the phosphatase reaction in heart mitochondria is inhibited by alloxan-diabetes or starvation in the rat. *Biochem. J.* 173, 669-680.
- Lyman, C. P., Willis, J. S., Malan, A., and Wang, L. C. H. (1982). Hibernation and Torpor in Mammals and Birds, T. T. Kozlowski, ed. (New York, NY: Academic Press, Inc.).
- Morgan, H. E., Cadenas, E., Regen, D. M., and Park, C. R. (1961). Regulation of glucose uptake in muscle. II. Rate-limiting steps and effects of insulin and anoxia in heart muscle from diabetic rats. *J. Biol. Chem.* 236, 262-268.

Morgan, H. E., Henderson, M. J., Regen, D. M., and Park, C. R. (1961). Regulation of glucose uptake in muscle. I. The effects of insulin and anoxia on glucose transport and phosphorylation in the isolated perfused heart of normal rats. *J. Biol. Chem.* 236, 253-261.

Randle, P. J. (1986). Fuel selection in animals. *Biochem. Soc. Trans.* 14, 799-806.

Wu, P., Sato, J., Zhao, Y., Jaskiewicz, J., Popov, K. M., and Harris, R. A. (1998). Starvation and diabetes increase the amount of pyruvate dehydrogenase kinase isoenzyme 4 in rat heart. *Biochem J* 329, 197-201.