7 🔼	
AD	

Award Number: DAMD17-99-2-9043

TITLE: Drug Development and Conservation in West and Central Africa/Performance of Neurochemical and Radio Receptor

Assays of Plant Extracts

PRINCIPAL INVESTIGATOR: Deborah C. Mash, Ph.D.

CONTRACTING ORGANIZATION: University of Miami

Miami, Florida 33136

REPORT DATE: September 2002

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

Form Approved

REPORT DOCUMENTATION PAGE

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE 3. REPORT TYPE AND DATES COVERED						
	September 2002	Annual (1 Sep	p 01 - 31 Aug 02) 5. FUNDING NUMBERS			
4. TITLE AND SUBTITLE Drug Development and Confidence of Development and D	Neurochemical and Radi		5. FUNDING N			
7. PERFORMING ORGANIZATION N	AME(S) AND ADDRESS(ES)			G ORGANIZATION		
University of Miami Miami, Florida 33136		:	REPORT NUMBER			
E-Mail: dmash@med.miami.edu						
9. SPONSORING / MONITORING A	GENCY NAME(S) AND ADDRESS(E	S)	10. SPONSORI	NG / MONITORING		
U.S. Army Medical Research and Fort Detrick, Maryland 21702-50	Materiel Command			EPORT NUMBER		
11. SUPPLEMENTARY NOTES						
12a. DISTRIBUTION / AVAILABILIT				12b. DISTRIBUTION CODE		
Approved for Public Re		limited.				
Approved for Public Re		limited.				
		limited.				
13. ABSTRACT (Maximum 200 Wo.		limited.		15. NUMBER OF PAGES		
13. ABSTRACT (Maximum 200 Wo		limited.	L	15. NUMBER OF PAGES  6  16. PRICE CODE		
13. ABSTRACT (Maximum 200 Wo		19. SECURITY CLASSIF OF ABSTRACT Unclassif:	ICATION	6		

# **Table of Contents**

Cover
SF 298
ntroduction4
3ody4
Key Research Accomplishments
Reportable Outcomes
Conclusions
deferences6
nnendices

### **ICBG PROGRESS REPORT 6/7/2002**

#### Specific Aims

The main objective of these studies was to study the CNS activity of these plant fractions using specific radio ligand binding assays to identify their selectivity profiles as well as to determine their functional activities as either a full or partial agonist or antagonist.

#### Materials

The following table gives a list of the samples provided to us thus far.

Table 1: ICBG PLANT SAMPLES

Table 1. ICBG I LANT SAWIFLES						
In House Sample#	House Sample# SU-Lab Number		Wt given (mg)			
ICBG 1	SU1904	Whole Plnt	25			
ICBG 2	SU1905	Sd Pulp	25			
ICBG 3	SU1906	Pn polar	25			
	~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	i ii potai	23			
ICBG 4	SU1907	Lf/Stem	25			
ICBG 5	SU1908	Lf/stem				
10000	301708	Listem	25			
ICBG 6	SU1909	Stbk	25			
TODO	501707	Siok	25			
ICBG 7	SU1910	C41.1.	25			
ICDU /	301910	Stbk	25			
ICBG 8	SU1911	041.1.	0.5			
ICDG 6	201911	Stbk	25			
ICBG 9	CL11010	TYPE I DI				
<b>}</b> }	SU1912	Whole Plnt	25			
ICBG 10	SU1913	Whole Plnt	25			
100011	GY 7. 4.1.					
ICBG 11	SU1914	Ft pulp	25			

## Methods

*Preparation of Extracts*: Stocks of the solid extracts were prepared by weighing 4 mg of sample and diluting to a total volume of 5 ml with 50% ethanol solvent for a final concentration of 0.8 mg/ml.

Opioid Recetor(kappa-1) Ligand Binding<sup>a</sup>: Two-point binding screens using [ $^3$ H]U69593 were performed as per metheds within the grant and previously (Izenwasser et al,1999). Briefly, mokey insular cortex was homogenized in ice-cold buffer (1:10 w/w, 50 mM Tris, 176  $\mu$ M MnCl<sub>2</sub>, 0.1% BSA pH 7.4), centrifuged at 32,000 x g for 10 minutes, and the supernatant was discarded. The pellet was resuspended (1:10, w/w) with buffer, and 100  $\mu$ l was added to each tube for a final concentration of 10.0 mg/ml (original tissue wet weight), in a final assay tube volume of 1.0 ml. Screens were performed by incubating 2 concentrations of extract (0.8 and 80

 $\mu$ g/tube) in the presence 1.0 nM of radioligand for 1 hr at 25°C, to determine their inhibitory potency (% inhibition). Nonspecific binding was determined by binding in the presence of 10  $\mu$ M naloxone. Incubations were terminated by vacuum filtration through glass fiber filters (Whatman 934-AH) presoaked in 1% polyethylenimine (PEI) followed by washing with ice-cold buffer (2 x 5 ml).

Norepinephrine Transporter(NET) Ligand Binding<sup>b</sup>: Two-point binding screens using [³H]Nisoxetine were performed as per metheds within the grant and previously (Tejani-Butt et al,1990). Briefly, mokey hypothulamus was homogenized in ice-cold buffer (1:10 w/w, 50 mM Tris, 300 mM NaCl, 5 mM KCl pH 7.4), centrifuged at 32,000 x g for 10 minutes, and the supernatant was discarded. The pellet was resuspended (1:10, w/w) with buffer, and 100 μl was added to each tube for a final concentration of 10.0 mg/ml (original tissue wet weight), in a final assay tube volume of 1.0 ml. Screens were performed by incubating 2 concentrations of extract (0.8 and 80 μg/tube) in the presence 3.0 nM of radioligand for 4 hr at 4°C, to determine their inhibitory potency (% inhibition). Nonspecific binding was determined by binding in the presence of 10 μM Mazindol. Incubations were terminated by adding 4ml of ice-cold buffer and vacuum filtration through glass fiber filters (Whatman 934-AH) presoaked in 0.5% polyethylenimine (PEI) followed by washing with ice-cold buffer (2 x 4 ml).

Data Analysis: Data are analyzed using the non-linear regression algorithms found in EBDA/LIGAND<sup>TM</sup> and GraphPad Prism<sup>TM</sup> computer software programs.

Results.

Table 2: Plant Extract Activity (% inhibition) on DAT, SERT, Mμ, Kappa-1 & NET

	DAT	75 (20)	SERT		Мμ	Иμ		Kappa-1		Карра-1	
1	$(f^3H)WIN$		([ <sup>125</sup> I]RTI-55)		([³H]DAMGO)		([ <sup>3</sup> H]U69593)		([ <sup>3</sup> H]U69593)		
In House	@ 0.8	@ 80	@ 0.8	@ 80	@ 0.8	@ 80	@ 0.8	@ 80	IC50	nН	
Sample#	μg/tube	μg/tube	μg/tube	μg/tube	μg/tube	μg/tube	μg/tube	μg/tube	Ug/ml		
ICBG 1		58.7		42.0	1.8	60.4		84.0	37.4	1.21	
ICBG 2		46.3		90.1	16.7	60.2		91.1	15.9	0.93	
ICBG 3		18.8		46.7	31.8	39.4		78.8	51.6	1.32	
ICBG 4		77. <b>7</b>			33.0	41.0		82.1	72.6	1.25	
ICBG 5		22.9				33.3			N/A	1.23	
ICBG 6		27.2		20.5		33.4		51.6	68.6	1.49	
ICBG 7						29.6		9.9	N/A	1.47	
ICBG 8		50.6		31.0	10.3	29.2		64.0	44.9	1.04	
ICBG 9		82.0				32.8		64.0	81.8	1.13	
ICBG 10		74.9	3.9	76.1	28.6	32.9		74.5	47.8	1.13	
ICBG 11				87.2	32,4	47.1		41.1	47.8 N/A	1,11	

-- denotes no inhibitory activity

As can be seen in Table 2, a number of extracts inhibited greater than 50% of the binding at DAT, SERT and  $M\mu$ . Therefore, full characterization of these compounds will be conducted.

Future Aims

Future studies will be to further characterize those extracts found to be active with full competition curves. In addition we plan to expand the screens with other ligands specific to other neuroreceptors and neurotransporters, and adding new compounds as made available.

### References

<sup>a</sup>Sari Izenwasser, Julie K. Staley, Stephanie Cohn, and Deborah C. Mash. Characterization of Kappa-Opioid Receptor Binding in Human Insular Cortex. *Life Sciences* 1999; 65(9):857-862.

<sup>b</sup>Shanaz M, Tejani-Butt, David J, Brunswick and Alan Frazer. [<sup>3</sup>H]Nisoxetine: a new radioligand for norepinephrine uptake sites in brain. *European Journal of Pharmacology* 1990; 191:239-243.