

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

|                             |                                    |                              |
|-----------------------------|------------------------------------|------------------------------|
| 1. REPORT DATE (DD-MM-YYYY) | 2. REPORT TYPE<br>Technical Papers | 3. DATES COVERED (From - To) |
|-----------------------------|------------------------------------|------------------------------|

|                       |                            |
|-----------------------|----------------------------|
| 4. TITLE AND SUBTITLE | 5a. CONTRACT NUMBER        |
|                       | 5b. GRANT NUMBER           |
|                       | 5c. PROGRAM ELEMENT NUMBER |

|  |                                |
|--|--------------------------------|
| 6. AUTHOR(S)<br><i>Please see attached</i> | 5d. PROJECT NUMBER<br>1011     |
|  | 5e. TASK NUMBER<br>0046        |
|  | 5f. WORK UNIT NUMBER<br>346204 |

|   |                                   |
|---|-----------------------------------|
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br>Air Force Research Laboratory (AFMC)<br>AFRL/PRS<br>5 Pollux Drive<br>Edwards AFB CA 93524-7048 | 8. PERFORMING ORGANIZATION REPORT |
|---|-----------------------------------|

|  |   |
|--|---|
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)<br>Air Force Research Laboratory (AFMC)<br>AFRL/PRS<br>5 Pollux Drive<br>Edwards AFB CA 93524-7048 | 10. SPONSOR/MONITOR'S ACRONYM(S)                              |
|  | 11. SPONSOR/MONITOR'S NUMBER(S)<br><i>Please see attached</i> |

12. DISTRIBUTION / AVAILABILITY STATEMENT  
Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

20030116 049

15. SUBJECT TERMS

|                                 |                             |                              |                                     |                     |  |
|---------------------------------|-----------------------------|------------------------------|-------------------------------------|---------------------|--|
| 16. SECURITY CLASSIFICATION OF: |                             |                              | 17. LIMITATION OF ABSTRACT<br><br>A | 18. NUMBER OF PAGES | 19a. NAME OF RESPONSIBLE PERSON<br>Leilani Richardson          |
| a. REPORT<br>Unclassified       | b. ABSTRACT<br>Unclassified | c. THIS PAGE<br>Unclassified |                                     |                     | 19b. TELEPHONE NUMBER<br>(include area code)<br>(661) 275-5015 |

MEMORANDUM FOR PR (In-House/Contractor Publication)  
FROM: PROI (TI) (STINFO)

15 March 2000

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2000-048**  
Larson, C. William; Harper, Jessica; Presilla-Marquez, J.D. (Schafer Corp.), "Matrix Isolation of Boron and Carbon Vapor. Control of Cluster Formation During Preparation and Annealing"

**10<sup>th</sup> Internat'l IUPAC Conference on High Temperature Materials Chemistry (Statement A)**  
**(Papers, oral presentation, poster, Proceedings)**  
**(Juelich, Germany, 10-14 April 2000)(Deadline: 31 March 2000)**

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Signature \_\_\_\_\_ Date \_\_\_\_\_

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review.

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Signature \_\_\_\_\_ Date \_\_\_\_\_

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b.) appropriateness of distribution statement, c.) military/national critical technology, d.) economic sensitivity, e.) parallel review completed if required, and f.) format and completion of meeting clearance form if required

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Signature \_\_\_\_\_ Date \_\_\_\_\_

4. This request has been reviewed by PR for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/national critical technology, and f.) data rights and patentability

Comments: \_\_\_\_\_  
\_\_\_\_\_

APPROVED/APPROVED AS AMENDED/DISAPPROVED

\_\_\_\_\_  
ROBERT C. CORLEY (Date)  
Senior Scientist (Propulsion)  
Propulsion Directorate

# MATRIX ISOLATION OF BORON AND CARBON VAPOR. CONTROL OF CLUSTER FORMATION DURING PREPARATION AND ANNEALING

C. W. Larson, J. Harper and J. D. Presilla-Márquez<sup>a</sup>

Air Force Research Laboratory  
Edwards AFB, CA 93524-7680  
E-mail: carl.larson@ple.af.mil

## Abstract

The formation and recombination of small boron/carbon clusters were studied by quantitative Fourier transform infrared (FTIR) spectroscopy. Samples were prepared by evaporation of powder mixtures from a resistively heated oven (3000 K) and isolated at 10 K in solid argon. C<sub>3</sub> and BC<sub>2</sub> disappeared entirely over the course of several annealings. Yields of all the larger clusters (B<sub>J</sub>C<sub>n-J</sub>, n ≥ 4, J = 0, 1, 2) increased during annealing. Clusters with n ≥ 4 are linear, with the boron atoms of the J = 1 and 2 clusters capping the ends of carbon chains; B<sub>2</sub>C<sub>n-2</sub> clusters are inert. During annealing, C<sub>n</sub> and BC<sub>n-1</sub> clusters grow by adding a mobile atom, dimer, or trimer to the carbon end of the chain. Dramatic growth of B<sub>2</sub>C<sub>2</sub> during the first annealing is consistent with the presence of B atoms and/or BC molecules in the initial matrices. (do not react to form larger clusters w/in matrix)

## Introduction

Under the auspices of the U. S. Air Force High Energy Density Matter (HEDM) Program [1], we have pursued the goal of production of a cryogenic matrix with an atom density of ~ 1 × 10<sup>21</sup> atoms cm<sup>-3</sup>, which amounts to ~ 5 mole percent atoms in an inert solid, or an equivalent atom gas pressure of ~ 60 atm at 298 K. This and our companion paper [2] describe qualitative and quantitative analysis of B<sub>J</sub>C<sub>n-J</sub> clusters in prototypical HEDM. Previously, the first conclusive identifications of small boron clusters, BC<sub>2</sub>, B<sub>2</sub>C, BC<sub>3</sub>, and B<sub>2</sub>C<sub>2</sub> were reported [3-5]. Observed FTIR fundamentals and isotope shifts were matched with predictions of *ab-initio* theory in those studies.

Verhaegen, Stafford, and Drowart [6] published the first experimental evidence of B<sub>J</sub>C<sub>n-J</sub> 35-years ago. Mass spectrometry was used to measure boron and boron carbide vaporization from graphite Knudsen cells heated by electron bombardment to ~ 2500 K. The vapor pressure of boron at 2200 K was ~ 0.008 torr, 94% atoms, 5% BC<sub>2</sub>, 1% B<sub>2</sub>C, and 0.07% BC. At the same temperature, boron carbide had about one-half the vapor pressure with 96% atoms, 4% BC<sub>2</sub>, 0.1% B<sub>2</sub>C. Distributions of anions and cations of B<sub>x</sub>C<sub>y</sub> with up to 17-atoms have been measured in plumes of laser vaporized boron carbide targets by Becker and Dietze [7]. Their mass spectrometric analysis showed that B<sub>2</sub>C<sup>-</sup>, B<sub>2</sub>C<sup>+</sup>, BC<sub>2</sub><sup>-</sup>, and BC<sub>2</sub><sup>+</sup> were major components of the plume. Matrix isolated BC<sub>2</sub> has been produced by oven and laser vaporization of carbon/boron mixtures and targets and measured by FTIR [3,8] and ESR [9,10] spectroscopy. Most recently, Wyss, Grutter and Maier [11] isolated BC<sub>2</sub><sup>-</sup> and BC<sup>-</sup> in neon matrices by sputtering a boron carbide target with Cs<sup>+</sup> and trapping the mass-selected anions in neon matrices. Neutrals were produced by electron photodetachment with uv radiation. Electronic transitions of BC<sub>2</sub> and BC and their anions were studied. The FTIR

<sup>a</sup> Present address: Schafer Corporation, 26565 W. Agoura Rd., Calabasas, California 93102.

spectrum of the previously identified [3]  $\nu_2(a_1)$  fundamental of  $BC_2$  was also observed. The infrared spectrum of BC has never been measured, but a CCSD(T)/cc-pVTZ calculation with quartic force field predicted [12] its vibrational fundamental at  $1148\text{ cm}^{-1}$  with intensity of  $43\text{ km mol}^{-1}$ . Of the pure boron clusters, only  $B_3$  has been experimentally observed. Li, Van Zee and Weltner [13] identified all four isotopomers of  $B_3$  in argon matrices by FTIR spectroscopy and found that experimental frequencies matched *ab-initio* frequencies of its cyclic geometry. Although *ab-initio* studies of  $B_n$ ,  $n \geq 4$ , have been carried out, only geometries and energies have been reported [14,15]. Unlike  $C_n$ ,  $B_n$  for  $n \geq 5$  are predicted to have stable 3-dimensional geometries.

Carbon clusters were recently reviewed by Van Orden and Saykally [16]. Cyclic  $C_6$  [17, 18], cyclic  $C_8$  [19, 20], linear  $C_{11}$  [21], have recently been identified by their FTIR isotopomer spectroscopy. Most recently, electronic spectra of linear  $C_{17}$ ,  $C_{19}$ , and  $C_{21}$  [22] and cyclic  $C_{10}$  and cyclic  $C_{12}$  [23] isolated in neon matrices were reported, and Raman spectra of linear  $C_{16}$ ,  $C_{18}$ , and  $C_{20}$  were reported [24].

Figure 1 shows a Pascal's triangle representation of the boron/carbon system. In the limit of statistical recombination of well mixed atoms, the relative yields of  $B_J C_{n-J}$  clusters of given  $n$  is given by

$$\rho(B_J C_{n-J})/\rho(C_n) = n!/J!(n-J)! [B/C]^J, \quad (1)$$

where  $[B/C]$  is the molar ratio of boron to carbon, and  $\rho(i)$  is the column density of species  $i$ .

## Experimental

Mixtures of boron and carbon powders were evaporated with a resistively heated oven to temperatures  $\sim 3000\text{ K}$ , and vapors were co-condensed with  $\sim 1000:1$  excess argon at  $10\text{ K}$  to form  $\sim 60\ \mu$  thick matrices in about one-hour. Beer's Law enabled quantitative analysis of the various species,  $\rho(i) = 2.303 A_i(\text{experiment})/I_i(\text{theory})$ , where  $\rho(i)$  is the average column density, which is the average value of the product of the species density and matrix thickness.  $A_i(\text{experiment})$  is the integrated intensity of the infrared absorbance of species  $i$ , equal to  $-\log_{10} \tau_i(\nu)$ , where  $\tau_i(\nu)$  is the transmittance at absorption frequency  $\nu$ .  $I_i(\text{theory})$  is the infrared absorption intensity of species  $i$ , which by necessity must be obtained by *ab-initio* calculation because experimental values are not available at this time. Fig. 1 indicates values of  $I_i(\text{theory})$  calculated by DFT/cc-pVDZ [25-28].

boron  
cannon"  
by it here  
ince it is  
ntioned, but  
t defined, in  
a conclusion

## Results and Discussion

Annealing of pure carbon matrices led to disappearance of atoms, dimers and trimers from the initial deposits. The identified products of recombination included cyclic  $C_6$  and cyclic  $C_8$ , and linear  $C_n$  molecules,  $4 \leq n \leq 13$ . After annealing to constant composition, the total measurable carbon was found to be as much as twice the initial measurable carbon, indicating that about 50 % of the initially deposited carbon was isolated as atoms and dimers. The cyclic molecules, the dominant condensation products, amounted to as much as  $\sim 60\%$  of total carbon in the fully annealed matrices.

In matrices containing boron, linear  $C_3$  and cyclic  $BC_2$ ,  $B_2C$ , and  $B_3$ , constituted about 80% of the total observable boron and carbon in the initially deposited matrix. The observation of approximately statistical distributions [Equation (1)] in clusters with  $n = 3, 4$ , and 5 in the initially deposited matrices implies that these smaller clusters form by random condensation of well-mixed atoms, uninfluenced by their relative energies, the energies of their precursors, or preferential kinetics pathways that could otherwise distort the statistics.




Figure 1. Representation of the  $B_n C_{n-J}$  ( $n = 1 - 13$ ,  $J = 0 - n$ ) system by Pascal's triangle. Numbers below molecular formulas denote the *ab-initio* intensities of the most intense infrared stretching fundamentals, Refs. [25] - [28]. Cyclic carbon clusters are removed from the triangle to the left.




Figure 2. Survey spectrum of matrix containing carbon and boron at natural abundance with  $[B/C] \sim 1/3$  after three annealings. All of the peaks indicated grow upon annealing except  $C_3$ . Fundamentals of  $BC_{n-1}$  for  $n = 5, 6, 7$ , and  $9$  are similarly red-shifted from fundamentals of linear  $C_n$  and their experimental absorbances are all slightly greater in this matrix. Two fundamentals of  $BC_6$  are observed at  $2112$  and  $1866 \text{ cm}^{-1}$ , red-shifted from the two fundamentals of linear  $C_7$ .

$B_2C_{n-2}$  clusters, once formed, were inert to further condensation upon annealing. Thus, energy loss by recombination during preparation of high energy density matrices (HEDM) is minimized in matrices with large B/C ratios. For example, when B/C  $\sim$  2, any recombination that does occur during co-deposition would tend to produce  $B_2C$ , a high-energy molecule that is inert in the cryogenic environment. In pure boron HEDM, the cyclic  $B_3$  molecule may be similarly inert, which would suggest that energy loss by recombination will also stop at a  $B_3$  "island of stability." Continuing experimental work to produce five-mole percent pure boron HEDM is concentrating on development of a resistively heated high-flux boron atom source, the "boron cannon", and production of higher density matrices.

## References

1. M. R. Berman, Editor, *Proceedings of the High Energy Density Matter (HEDM) Contractors' Conference* (Monterey, California, May 1998), Air Force Office of Scientific Research, Bolling AFB, DC 20332-8050, and annual proceedings dating from 1987.
2. C. W. Larson, J. Harper, J. D. Presilla-Márquez, Poster Paper, **This Volume**.
3. J. D. Presilla-Márquez, C. W. Larson, P. G. Carrick, and C. M. L. Rittby, *J. Chem. Phys.* **105**, 3398 (1997).
4. J. D. Presilla-Márquez, P. G. Carrick, and C. W. Larson, *J. Chem. Phys.* **110**, 5702 (1999).
5. C. W. Larson and J. D. Presilla-Márquez, *J. Chem. Phys.* **111**, 1988 (1999).
6. G. Verhaegen, F. E. Stafford, and J. Drowart, *J. Chem. Phys.* **40**, 1622 (1964).
7. S. Becker, and H. J. Dietze, *Intl. J. Mass Spectrometry and Ion Proc.* **82**, 287 (1988).
8. J. M. L. Martin, P. R. Taylor, J. T. Yustein, T. R. Burkholder, and L. Andrews, *J. Chem. Phys.* **99**, 12 (1993).
9. W. C. Easley and W. Weltner, Jr., *J. Chem. Phys.* **52**, 1489 (1970).
10. L. B. Knight, Jr., S. Cobranchi, E. Earl, and A. J. McKinley, *J. Chem. Phys.* **104**, 4927 (1996).
11. M. Wyss, M. Grutter, and J. P. Maier, *J. Phys. Chem. A* **102**, 9106 (1998).
12. J. M. L. Martin and P. R. Taylor, *J. Chem. Phys.* **100**, 9002 (1994).
13. S. Li, R. J. Van Zee, and W. Weltner, Jr., *Chem. Phys. Lett.* **262**, 298 (1996).
14. J. Niu, B. K. Rao and P. Jena, *J. Chem. Phys.* **107** 132 (1997).
15. I. Boustani, *Chem. Phys. Lett.* **233** 273 (1995), **240** 135 (1995).
16. A. Van Orden, and R. J. Saykally, *Chem. Rev.* **98**, 2313 (1998).
17. J. D. Presilla-Márquez, J. A. Sheehy, J. D. Mills, P. G. Carrick, and C. W. Larson, *Chem. Phys. Lett.* **274**, 439 (1997).
18. S. L. Wang, C. M. L. Rittby, and W. R. M. Graham, *J. Chem. Phys.* **107**, 6032 (1997).
19. J. D. Presilla-Márquez, J. Harper, J. A. Sheehy, P. G. Carrick, and C. W. Larson, *Chem. Phys. Lett.* **300**, 719 (1999).
20. S. L. Wang, C. M. L. Rittby, and W. R. M. Graham, *J. Chem. Phys.* **107**, 7025 (1997), **112**, 1457 (2000), and X. D. Ding, et al., *J. Chem. Phys.* **112**, 5113 (2000).
21. L. Lapinski and M. Vala, *Chem. Phys. Lett* **300**, 195 (1999).
22. M. Wyss, M. Grutter, and J. P. Maier, *Chem. Phys. Lett* **304**, 35 (1999)
23. M. Grutter, M. Wyss, E. Riaplov, and J. P. Maier, *J. Chem. Phys.* **111**, 7397 (1999).
24. A. K. Ott, G. A. Rechtsteiner, C. Felix, O. Hampe, M. F. Jarrold, R. P. Van Duyne, and K. Raghavachari, *J. Chem. Phys.* **109**, 9652 (1998).
25. C. M. L. Rittby, J. A. Sheehy, and M. Vala, M. Zerner (personal communications).
26. J. M. L. Martin, J. P. François, and R. Gijbels, *J. Chem. Phys.* **90**, 6469 (1989).
27. J. M. L. Martin, J. El-Yazal, and J. P. François, *Chem. Phys. Lett.* **242**, 570 (1995).
28. J. M. L. Martin and P. R. Taylor, *J. Chem. Phys.* **100**, 9002 (1994).

# MATRIX ISOLATION OF BORON AND CARBON VAPOR.

C. W. Larson, J. Harper and J. D. Presilla-Márquez

*Air Force Research Laboratory*

*Edwards AFB, CA 93524-7680*

*E-mail: carl.larson@ple.af.mil*

## Abstract

This report supplements the companion paper published in this volume [1]. Details about the FTIR quantitative analysis of the disappearance and formation of clusters during annealing of solid argon matrices containing boron and carbon species are presented.

## Introduction

The companion paper [1] summarized the literature of small boron clusters, about  $B_J C_{n-J}$  ( $n = 2$  to 10,  $J = 0, 1, 2, 3$ ) and described the formation of clusters during preparation and annealing of solid argon. This report presents additional details.

## Experimental

A precision matched pair of matrices was prepared as described in Ref. [1] with two different boron isotopic compositions and natural abundance carbon. Matrix (a) contained natural abundance boron,  $^{11}B/^{10}B = 4.0$ , and matrix (b) contained boron with an inverted isotope ratio,  $^{11}B/^{10}B = 0.37$ . Each matrix had a [C/B] molar ratio of  $\sim 3.0$ , a thickness of  $\sim 60$  m, and very similar absolute amounts of each element. Each matrix was annealed repeatedly according to the same annealing protocol given in the caption of Figure 1.

## Results and Discussion

Figure 1 shows the annealing behaviors of the FTIR bands belonging to  $BC_2$ ,  $BC_3$  and  $B_2C_2$  for each of the matrices. Cyclic  $BC_3$  disappears entirely when one of its carbon atoms is attacked by a radical species. One of the B-C bonds is broken, and the  $BC_3$  ring opens. The dramatic appearance of  $B_2C_2$  during the first annealing is consistent with a mechanism involving recombination of BC, whose density lies below the limit of detection, or condensation of B atoms with  $BC_2$ .

Figure 2 shows the annealing behaviors of the  $^{11}B_J^{12}C_{n-1}$  clusters,  $n = 3$  to 9 and  $J = 0, 1, 2$  that were measured in matrix (a). Loss of clusters during the sixth annealing by matrix sublimation is generally greatest for  $C_n$  and smallest for  $B_2C_{n-2}$ , which is consistent with a stratified matrix with a higher [C/B] ratio near the subliming surface. Apparently, boron evaporates more rapidly than carbon during the early stages of oven evaporation, resulting in boron-rich layers near the substrate.

Column densities of the trimer clusters in the initially deposited matrices are in the ratio  $\rho(C_3):\rho(BC_2):\rho(B_2C):\rho(B_3) \approx 1.0:1.5:0.5: < 0.05$ , where only the upper limit for  $\rho(B_3)$  may be determined because it was not observed. This is consistent with formation of the trimers by statistical condensation from a matrix with  $[B/C] \sim 1/3$ , which would produce a 1.0:1.0:0.33:0.05 distribution,  $\rho(B_J C_{n-J})/\rho(C_n) = n!/J!(n-J)! [B/C]^J$ , Eq. (1), Ref. [1].

Agreement between distributions implies trimers form by random condensation of well-mixed atoms, uninfluenced by the relative energies of the trimers, the energies of their precursors, or preferential kinetics pathways that could otherwise distort the statistics. Linear  $C_3$  and cyclic  $BC_2$ , disappeared entirely when the matrices were repeatedly annealed to




Figure 1. Absorbance spectra of matrix (a) and matrix (b) showing disappearance of  $BC_2$ ,  $BC_3$  and  $C_4$  and appearance of  $B_2C_2$  upon annealing. Annealing temperatures and times were (1) 27.5 K/150 s, (2) 30.0 K/75 s, (3) 32.5 K/45 s, (4) 35.0 K/30 s, (5) 37.5 K/20 s, (6) 40.0 K/20 s. The decrease in the peaks in the fifth and sixth annealing is due to matrix sublimation. The red shift of the  $BC_3$  absorption from the  $C_4$  absorption is similar to the red-shifts observed for all the other  $BC_{n-1}$  clusters that occur when a boron atom is substituted for a carbon atom in a  $C_n$  cluster.

temperatures between 25 K and 35 K, but cyclic  $B_2C$  was inert. Linear  $C_4$  and  $BC_3$  (BCCC) disappeared more slowly, and linear  $B_2C_2$  (BCCB) grew to  $\sim 95\%$  of its final value during the first annealing. Once formed,  $B_2C_2$ , like  $B_2C$ , was also inert to further reaction.


Figure 3 shows the absolute column densities of the analyzed clusters that was obtained by transformation of the Fig. 2 data with Beer's law and theoretical intensities shown in Fig. 1, Ref [1]. The  $BC$  molecule was not observed, but its upper limit column density was established to be somewhat larger than that of  $B_2C_2$ .

The sources of  $B_2C_2$  are from condensation of atom plus trimer ( $B + BC_2$  but not  $C + B_2C$ ) or dimer + dimer ( $BC + BC$  but not  $B_2 + C_2$ ). Although  $BC$  was not observed, the upper limit of  $\rho(BC)$  is larger than  $\rho(B_2C_2)$  so that  $BC$  cannot be ruled out as a source of  $B_2C_2$ . The growth of  $B_2C_2$  is consistent with the presence of  $BC$  and/or  $B$  in the originally deposited matrix in an amount at least as great as the growth of  $B_2C_2$ .

Figures 1 and 2 show that  $C_4$  and  $BC_3$  do not change during the first annealing but begin to disappear in subsequent annealings. This indicates that a source for these tetramers, atom + trimer and/or dimer + dimer, operates during the first annealing.


Figure 2 shows that linear  $C_5$ ,  $BC_4$  (BCCCC) and  $B_2C_3$  (BCCCB) and larger linear clusters all grew upon annealing. The sources of  $B_2C_3$  are dimer + trimer ( $BC + BC_2$  but not  $B_2 + C_3$ ) and atom + tetramer ( $B + BC_3$  but not  $C + B_2C_2$ ). Since  $\rho(BC_2) \sim 5\rho(BC_3)$  in the initially deposited matrix, the  $BC + BC_2$  source is dominant. Growth of  $B_2C_3$  is consistent with the presence of  $BC$  in the initially deposited matrix in an amount at least as great as the amount by which  $B_2C_3$  grows.





2

Figure 1. Annealing behaviors of  $^{11}\text{B}^{12}\text{C}_{n-J}$  species in matrix (a), which contained natural abundance carbon and boron,  $^{11}\text{B}/^{10}\text{B}=80/20$ ,  $^{12}\text{C}/^{13}\text{C}=99/1$ , with  $[\text{C}/\text{B}] \sim 3$ . Spectra labeled '0' were obtained from the originally deposited matrix, and spectra labeled '1' to '6' were obtained after successive annealings. During the 6th annealing (at 40 K for 20 seconds), almost half the matrix evaporates.



3. Figure 2. Distributions of  $B_J C_{n-J}$  clusters,  $J = 0, 1, 2$ ;  $n = 3 - 11$ . Circles, triangles and squares represent  $BC_{n-1}$ ,  $C_n$  and  $BC_{n-2}$  clusters, respectively. Large symbols denote upper and lower limits of measurement, based on a nominal minimum measurable absorbance of  $0.0023 \text{ cm}^{-1}$ , and a maximum absorbance for linearity of Beer's law of  $2.3 \text{ cm}^{-1}$  (1% transmittance). Small symbols denote measured quantities in the initial matrix, and in six annealed matrices. Some of the larger clusters ( $n = 8, 10, 11$ ) have not been identified,  $BC_7, B_2C_6, BC_9, B_2C_8, BC_{10}, B_2C_9$ .

Growth of  $BC_4$  occurs primarily by  $BC + C_3$  rather than  $B + C_4$  or  $C + BC_3$  because  $\rho(C_3) \sim 10\rho(C_4)$  and  $\rho(C_3) \sim 2\rho(BC_3)$ . Growth of  $C_5$  occurs by  $C + C_4$  and  $C_2 + C_3$ , which establishes the presence of  $C$  and/or  $C_2$  in the original matrix in an amount at least as great as  $C_5$  growth.

### Conclusions

Disappearance of triangular  $BC_2$  requires breaking of one of its B-C bonds when one of its carbon atoms is attacked. The major reorganization of electronic energy involved in opening the ring appears to occur with little ( $< \sim 3 \text{ kcal mol}^{-1}$ ) or no energy barrier, which makes this small molecule a candidate for an interesting *ab-initio* study of unusual reactivity at low temperature. Annealing kinetics of disappearance of  $C_3$  and  $BC_2$ , and of appearance of  $B_2C, C_4, BC_3, B_2C_2, C_5, BC_4,$  and  $B_2C_3$  unequivocally establishes the presence of atoms and dimers in the originally deposited matrix. About 80% or more of the initially deposited HEDM existed as atoms, dimers and trimers. Molecules with two boron atoms are immune from radical attack and condensation during annealing.

### Reference

1. C. W. Larson, J. Harper, J. D. Presilla-Márquez, Poster Paper, This Volume.

(high eng data substrate)