

AD \_\_\_\_\_

Award Number: DAMD17-01-1-0680

TITLE: Role of the 5HT<sub>3</sub> Receptor in Alcohol Drinking and Aggression Using a Transgenic Mouse Model

PRINCIPAL INVESTIGATOR: Andrea M. Allan, Ph.D.

CONTRACTING ORGANIZATION: University of New Mexico  
Albuquerque, New Mexico 87131-5041

REPORT DATE: September 2002

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE September 2002	3. REPORT TYPE AND DATES COVERED Annual (1 Sep 01 - 31 Aug 02)
----------------------------------	----------------------------------	---

4. TITLE AND SUBTITLE Role of the 5HT <sub>3</sub> Receptor in Alcohol Drinking and Aggression Using a Transgenic Mouse Model	5. FUNDING NUMBERS DAMD17-01-1-0680
--	--

6. AUTHOR(S):  
Andrea M. Allan, Ph.D.

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  
University of New Mexico  
Albuquerque, New Mexico 87131-5041  
  
E-Mail: aallan@salud.unm.edu

8. PERFORMING ORGANIZATION REPORT NUMBER

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  
U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

20030502 110

12a. DISTRIBUTION / AVAILABILITY STATEMENT  
Approved for Public Release; Distribution Unlimited

12b. DISTRIBUTION CODE

13. Abstract (Maximum 200 Words) *(abstract should contain no proprietary or confidential information)*  
Alcohol use has been identified as an important factor in aggressive, or violent, behavior in humans. Alcohol not only increases the incidence but also the severity of violent attacks. Several clinical studies have reported the observation that highly aggressive individuals display a serotonin-deficient trait. A number of studies indicate that the 5HT<sub>3</sub> receptor system mediates alcohol consumption and the subjective effects of alcohol. The 5HT<sub>3</sub> receptor is unique in the serotonin receptor family in that it is a cation channel and modulates the release of other neurotransmitters, including GABA and dopamine. Thus, the 5-HT<sub>3</sub> receptor is likely to play a critical role influencing alcohol consumption, which appears to involve dopamine and influencing both natural and alcohol-heightened aggression through the GABA<sub>A</sub> receptor system. We have developed a 5-HT<sub>3</sub> receptor over-expressing mouse to study the role of this receptor in alcohol drinking and aggression. We hypothesize that 5HT<sub>3</sub> receptor over-expression decreases alcohol preference and aggressive activity through a 5HT<sub>3</sub> receptor sensitive mechanism increasing the release of GABA. We will evaluate the transgene on 3 different inbred strains (C57, DBA, 129) who differ in alcohol preference and alcohol-induced aggression. We will measure the impact of 5HT<sub>3</sub> receptor over-expression on alcohol preference in a two-bottle choice design and aggression using intruder-aggression test. We will test the ability of 5HT<sub>3</sub> receptor antagonists to block and GABA receptor agonist to mimic the phenotypic effects of over-expression. These mice should prove useful in testing hypothesis regarding the role 5-HT<sub>3</sub> receptors play in alcohol abuse and alcohol-heightened aggression.

14. SUBJECT TERMS  
alcohol, brain, serotonin, genetics, drinking, aggression

15. NUMBER OF PAGES  
11  
16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT  
Unclassified

18. SECURITY CLASSIFICATION OF THIS PAGE  
Unclassified

19. SECURITY CLASSIFICATION OF ABSTRACT  
Unclassified

20. LIMITATION OF ABSTRACT  
Unlimited

## Table of Contents

Cover.....	1
SF 298.....	2
Table of contents.....	3
Introduction.....	4
Body.....	5
Key Research Accomplishments.....	9
Reportable Outcomes.....	9
Conclusions.....	9
References.....	10
Appendices.....	N/A

## Introduction

Alcohol use has been identified as an important factor in aggressive, or violent, behavior in humans. In its 10th Special Report to the U.S. Congress on Alcohol and Health (June 2000), the Public Health Service stated that one in four victims of violent crime described their attacker as having consumed alcohol prior to committing the crime. When the victim was a current or former intimate partner, the incidence increased to two out of three offenders having been drinking prior to the attack. Alcohol not only increases the incidence but also the severity of the attack. While these statistics may be correlational, a causal relationship between alcohol use and physical violence needs to be explored.

Experimentally, aggression in mice can be brought about by a isolating male for some time, then, subsequently, to introduce a group-housed male to the isolated male's home cage (*isolation-induced aggression or intruder-aggression*), with heightened attacks seen when the resident mouse is an actively breeding male (Malick, 1979; Miczek, 1999). Many environmental and experiential variables factor into aggressive behavior, but the administration of alcohol is an important societal variable in aggressive behavior (Miczek et al., 1998). Several lines of research on the neurochemical basis for aggression have implicated the serotonin system (Korte et al., 1996; Fish et al., 1999; DeAlmedia et al., in press). Alterations in the serotonin system, using knock out technology, have produced lines of mice that display altered aggressive behavior (Saudou et al., 1994; Cases et al., 1995). Pharmacologic studies have implicated the involvement of 5-HT<sub>1a</sub> (deBoer et al., 2000), 5-HT<sub>1b</sub> (Fish et al., 1999) and 5-HT<sub>3</sub> (Rudissaar et al., 1999) receptors. Several clinical studies have reported the observation that highly aggressive individuals display a serotonin-deficient trait, as measured by either low levels of cerebral spinal fluid (CSF) 5-hydroxyindole acetic acid (5-HIAA) or a flattened prolactin response to serotonin activation (Brown et al., 1982; Linnoila et al., 1983; Coccaro et al., 1990, Mann, 1999). While these data provide support for a role for serotonin in aggression, there is strong support for gamma amino butyric acid (GABA) as well (Soderpan and Svensson, 1999; Glaysheva et al., 1998; Guillot et al., 1998; Navarro and Pedraza, 1996; Miczek et al., 1995; 1994,1993; Weerts et al., 1993).

Besides the suggestion that serotonin plays a role in aggression, several lines of research have suggested a role for serotonin in the regulation of alcohol consumption. A number of studies indicate that the 5HT<sub>3</sub> receptor system mediates alcohol consumption and the subjective effects of alcohol. 5HT<sub>3</sub> receptor antagonists decrease alcohol intake in laboratory animals (Jankowska et al., 1995, Jankowska et al. 1994, Tomkins et al. 1995, Knapp and Pohorecky 1992, Hodge et al. 1993) and humans (Johnson et al., 1993; Sellers et al., 1994). In addition, 5HT<sub>3</sub> receptor antagonists increase the subjective feeling of alcohol intoxication in humans (Swift et al., 1996), suggesting that 5HT<sub>3</sub> receptors are involved in alcohol sensitivity. Lovinger and his colleagues have shown that alcohol directly potentiates the 5HT<sub>3</sub> receptor (Lovinger, 1991; Lovinger and White, 1991; Lovinger and Zhou, 1994). The 5HT<sub>3</sub> receptor is unique in the serotonin receptor family in that it is a cation channel and modulates the release of a number of other neurotransmitters, including GABA and dopamine. Thus, the 5-HT<sub>3</sub> receptor is likely to play a critical role influencing alcohol consumption, which appears to involve dopamine and influencing both natural and alcohol-heightened aggression through the GABA<sub>A</sub> receptor system.

We hypothesized that over-expression of 5-HT<sub>3</sub> receptors decreases alcohol consumption because the presence of an increased number of 5-HT<sub>3</sub> receptors increased the potentiation of dopamine release at lower alcohol concentrations. Thus, the animal requires less alcohol to obtain the same behavioral effect. Alternatively, it is possible that the over-expression of 5-HT<sub>3</sub> receptors increases the release of gamma aminobutyric acid (GABA), an inhibitory neurotransmitter. GABA plays an important role in the inhibitory circuit from the prefrontal cortex to the amygdala and ventral striatal areas. This pathway is thought to play a role in moderating impulsive and compulsive behaviors. Thus, the lower level of alcohol consumption seen in the 5-HT<sub>3</sub> receptor over-expressing mice may be the result of increased inhibitory control over alcohol consumption. That is, these transgenic mice know "when to say when". To add support for this hypothesis, we found that the 5-HT<sub>3</sub> receptor over-expressing mice fail to behave aggressively in an intruder aggression test. The wild type mice clearly displayed high levels of aggression by attacking the intruder within seconds of introduction and continuing the attacks until the test was halted. This lower level of aggression in the 5-HT<sub>3</sub> receptor over-expressing mice may be due to an increase in GABA activity. The proposed studies will explore the neurochemical relationship between the lower alcohol consumption and lower levels of aggression seen in 5-HT<sub>3</sub> receptor over-expressing transgenic mice. The first goal will examine the impact of 5HT<sub>3</sub> receptor over-expression on alcohol preference using a two-bottle free choice test. The second goal will examine the impact of 5HT<sub>3</sub> receptor over-expression on natural aggressive behavior. Lastly, the impact of 5HT<sub>3</sub> receptor over-expression on alcohol-heightened aggressive behavior will be measured.


### **Body**

Progress made towards our goals are as follows: Breeding onto the three different backgrounds was initiated. Generations N1, N3 and N5 are the tested generations planned. We have completed generation N1 testing and have begun generation N3 testing.


#### ***Lower Alcohol Drinking Phenotype Conserved On Various Genotypic Backgrounds in the first generation of cross breeding***

Reduced alcohol preference as a result of 5HT<sub>3</sub> receptor over-expression was conserved even when expressed on a genetic background known for high alcohol preference, C57BL/6J (Figure 1). Using a two-bottle free choice test, C57BL/6J mice drank an average of 14 g/kg/day, while the Line 13 x C57BL cross transgenic mice drank an average of 8 g/kg/day, while the 129 mice drank an average of 8 g/kg/day the line13X129 drank 3.75 g/kg/day. The transgene did not influence the DBA drinking which remained below 1g/kg/day.. **Thus, the presence of the transgene was able to influence alcohol preference on a different genetic background even though these mice are heterozygous for the transgene.**

As depicted below in Figure 1 below, the influence on the transgene to reduce alcohol drinking in high alcohol consuming strains is maintained on the first generation background. However, preliminary data from the N3 generation suggests that this influence is lost, at least on the C57BL/6 background. This data is still being analyzed and evaluated.



**Figure 1** Ethanol consumption averaged over 10 days in a two bottle free choice test in C57BL/6 xLine13 cross compared to C57BL/6 mice (left panel), DBA/2J vs. DBA/2J x line 13 cross mice (center panel) and 129/J xLine13 cross compared to 129/J mice (right panel). Data are mean ± SEM, n=10. Asterisk indicates significance at p<0.05.



**Figure 2:** Intruder aggression test. Latency to first attack in seconds. Data are mean ± SEM, n=5 mice. Male group housed A/J mice served as intruders. Residents housed singly 2 week prior. Testing was performed in the resident's home cage. Transgenic mice failed to attack during the 10 minute test period, thus no data is presented for them.

**5-HT<sub>3</sub> Receptor Over-Expression Significantly Lowers Intruder Aggression**

Using a standard intruder test, we evaluated the level of aggressive behavior in the transgenic 5-HT<sub>3</sub> receptor over-expressing mice. Resident mice were isolated for




one week prior to the testing. A/J male group housed mice were used as the intruders. The intruder mouse was placed bite. If no biting took place the test was terminated after 5 minutes. Thus, the longest the test was run would have been 10 minutes. In this way we avoid any bias against aggressive behavior that may begin later in the testing period but yet can still be very intense in nature. Further measurements of latency alone may distort the measures of actual fighting once it is initiated.

Transgenic mice failed to engage in biting or other forms of attack (Figure 2). These were single tests and these were not breeding males. Repeated testing and breeding male tests reveal a different picture. In repeated engagements, transgenic mice did attack the intruder with a frequency nearly indistinguishable from that of the wild type males. We have measured alcohol-heightened aggression in preliminary tests. While the transgenics do show an increase in aggressive behavior with alcohol, aggression is still significantly lower than in the control mice, with an average of 3.5 attack bites for transgenics and 11 attack bites for controls after 1 g/kg alcohol injected i.p. 10 minutes prior to testing. Data on aggression following free access alcohol are preliminary and additional animals need to be studied. Further we are measuring the effect of repeated encounters in the resident's cage.

#### ***5-HT<sub>3</sub> Receptor Over-Expression Significantly Improves Learning***

As an additional measure of impulse control we have been examining learning using a fear conditioned freezing measure with these lines. Fear conditioning is a classical Pavlovian learning measure where the mouse learns to anticipate a foot shock through the association of a preceding tone and the learning context itself. Learning is indicated by freezing. Over-expression of the 5HT<sub>3</sub> receptor significantly improves learning in this system (see figure below). These data suggest that the over-expression of the 5HT<sub>3</sub> receptor permits better stimuli association and information processing. Additionally, these results indicate that over-expression of the 5-HT<sub>3</sub> receptor in mouse forebrain results in enhanced hippocampal-dependent learning and attention.

Freezing scores for 5-HT3 receptor transgenic and control mice



### Key Research Accomplishments

- ❖ Initiate moving the transgene on to the three inbred strain backgrounds through breeding.
- ❖ Test the two bottle choice drinking on the wild type and transgenic N1 generations of the three strains.
- ❖ Test initial and repeated aggressive behavior on the wild type and transgenic N1 strains.
- ❖ Characterize unique learning and stress responding in the mice which impacts consumption and aggression.
- ❖ Began alcohol induced aggressive behavior testing on the N1 strains.
- ❖ Breed forward to the N3 generations.

### Reportable Outcomes

Harrell, A.V., Caldwell, K.K, **Allan, A.M.** "Transgenic mice over-expressing the 5-HT3 receptor have enhanced learning in latent inhibition and contextual fear conditioning paradigms" Society for Neurosciences, 2001

Carta, M. **Allan, A.M.** Partridge, L.D., Valenzuela, C.F. Effect of cocaine on 5-HT3 receptor mediated currents in hippocampal neurons from transgenic mice over expressing the receptor. Society for Neurosciences, 2001

**Allan, AM**, Chynoweth, J and Caldwell, K K "A Moderate exposure fetal alcohol mouse model using saccharin fading technique." Research Society on Alcoholism 26:531, 2002

Carta, M. **Allan, A.M.** Partridge, L.D., Valenzuela, C.F. Effect of cocaine on 5-HT3 receptor mediated currents in hippocampal neurons from transgenic mice over expressing the receptor. Submitted to Eur. J. Pharmacology, 2002

Harrell, A. V. and **Allan, A.M.** Improvements in Hippocampal-Dependent Learning and Decremental Attention in 5-HT3 Receptor Over-Expressing Mice. Submitted to Learning & Memory, 2002

### Conclusions

At least a generation 1 of the cross breeding the original impact of the transgene appears stable. However, as the background becomes more inbred the influence seems to be diluted. Further, the impact of the transgene on reducing aggressive behavior is upon initial exposure to the intruder but is overwhelmed but repeated presentations of the threat. This suggests a learned component or plasticity involved in aggressive responding that has not been noted in the literature at this time. Can aggressive responding be interfered with pharmacologically? We plan to look into these questions in future studies.

## References

- Brown GL, Ebert MH, Goyer PF, Jimerson DC, Klein WJ, Bunney WE (1982). Aggression, suicide, and serotonin-relationships to CSF amine metabolites. *Am J Psychiatry*;139:741-6.
- Cases O, Seif I, Grimsby J, Gaspar P, Chen K, Pournin S, (1995) Aggressive behavior and altered amounts of brain serotonin and norepinephrine in mice lacking MAOA. *Science* 268:1763-6.
- Coccaro EF, Astill JL, Herbert JL, and Schut AG. Fluoxetine Treatment of Impulsive Aggression in DSM-III-R Personality Disorder Patients. *J Clin Psychopharmacol*, 1990;10:373-5.
- DeAlmeida RMM, Nikulina E, Faccidomo S, Fish E, and Miczek KA. 5-HT1B/D receptors and aggression: specific reductions of alcohol-heightened aggression in male mice by CP-94,253 and zolmitriptan. *Psychopharmacology*, 2000;(submitted)
- DeBoer, SF, Lesourd, M, Mocaer, E and Koolhaas, JM (2000) Somatodentritic 5-HT1a autoreceptors mediate anti-aggressive actions of 5-HT1a receptor agonists in rats: An ethopharmacological study with S-15535, alnespirone and WAY-100635. *Neuropsychopharmacol.* 23:20-33.
- Fish E, Faccidomo S, and Miczek KA. (1999) Aggression heightened by alcohol or social instigation in mice: reduction by the 5-HT1B receptor agonist CP-94,253. *Psychopharmacology*, 146:391-9.
- Gladysheva, OS, Troistakaya, VT and Raevskii, KS (1998) Effect of intranasal administration of gamma-aminobutyric acid on intraspecies aggression of male mice. *Bull. Exper. Biol. & Med.* 125:353-354.
- Jankowska E., Bidzinski A., Kostowski W. (1994) Alcohol-Drinking In Rats Treated With 5,7-Dihydroxytryptamine :Effect Of 8-Oh-Dpat And Tropicsetron (Ics-205-930). *Alcohol* , 11:283-288 .
- Jankowska E, Kostowski W (1995) The effect of tropisetron injected into the nucleus-accumbens-septi on ethanol-consumption in rats *Alcohol* 12: 195-198
- Jankowska E., Krutki P., Lackberg Z.S., Hammar I. (1995) Effects Of Serotonin On Dorsal Horn Dorsal Spinocerebellartract Neurons. *Neuroscience* , 67:489-495.
- Knapp, D.J., Pohorecky L.A. (1992) Zaccopride, a 5-HT3 receptor antagonist, reduces voluntary ethanol consumption in rats. *Pharm. Biochem. Behav.* 41:847-850.
- Korte, SM, Meijer, OC, Kloet, ER, Buwalda, B Keijser, J, Sluyter, FS, Oortmerssen, G, Bohus, B (1996) Enhanced 5-HT1a receptor expression in forebrain regions of aggressive house mice. *Brain res.* 736:338-343.
- Linnoila M, Virkkunen M, Scheinin M, Nuutila A, Rimon R, and Goodwin FK. Low cerebrospinal fluid 5-hydroxyindoleacetic acid concentration differentiates impulsive from nonimpulsive violent behavior. *Life Sci*, 1983;33:2609-14.
- Lovinger, D.M. (1991) Ethanol potentiation of 5HT3 receptor-mediated ion current in NCB-20 neuroblastoma cells. *Neurosci. Lett.* 122:57-60.
- Lovinger, D.M. White G. (1991) Ethanol Potentiation of 5-Hydroxytryptamine3 Receptor-Mediated Ion Current in Neuroblastoma Cells and Isolated Adult Mammalian Neurons. *Mol. Pharm.* 40:263-270.

- Lovinger, D.M., Zhou Q. (1994) Alcohols Potentiate Ion Current Mediated by Recombinant 5-HT<sub>3</sub>RA Receptors Expressed in a Mammalian Cell Line. *Neuropharmacol.* 33:1567-1572.
- Malick JB. The pharmacology of isolation-induced aggressive behavior in mice. In: Essman WB, Valzelli L, editors. *Current Developments in Psychopharmacology*, vol. 5, New York: SP Medical and Scientific Books, 1979, 1-27.
- Mann JJ. Role of the serotonergic system in the pathogenesis of major depression and suicidal behavior. *Neuropsychopharmacology*, 1999;21:99S-105S.
- Miczek, KA, Barros, HM, Sakoda, L, Weerts, EM (1998) Alcohol and heightened aggression in individual mice. *Alc;Clin Exper. Res.* 22:1698-1705.
- Miczek, KA, Debold, JF, Vanerp, AMM (1994) Neuropharmacological characteristics of individual differences in alcohol effects on aggression in rodents and primates. *Behav. Pharmacol.* 5:407-421.
- Miczek, KA, Weerts, EM, Debold, JF (1993) Alcohol benzodiazepine GABA-A receptor complex and aggression: Ethological analysis of individual-differences in rodents and primates. *J. Stud. On Alcoholism* S11:170-179.
- Miczek, KA, Weerts, EM, Vivian, JA and Barros, HM (1995) Aggression anxiety and vocalization in animals: GABA(A) and 5-HT anxiolytics. *Psychopharmacology* 121:38-56.
- Miczek KA. (1999) Aggressive and social stress responses in genetically modified mice: from horizontal to vertical strategy. *Psychopharmacology* 147:17-19.
- Rudissaar, R, Pruus, K, Skrebuhhova, T, Allikmets, L and Matto, V (1999) Modulatory role of 5-HT<sub>3</sub> receptors in mediation of apomorphine-induced aggressive behavior in male rats. *Behav. Brain Res.* 106:91-96.
- Soderpam, B and Svensson, AI (1999) Naloxone reverses disinhibitory/ aggressive behavior in 5,7-DHT-lesioned rats: involvement of GABA(A) receptor blockade. *Neuropharmacol.* 38:1851-1859.
- Swift, R.M., Davidson, D., Whelihan, W. and Kuznetsov O. (1996) Ondansetron alters human alcohol intoxication. *Biol. Psychiatry* 40:514-521.
- Tomkins, D.M., Le, A.D., Sellers E.M. (1995) Effect of the 5-HT<sub>3</sub> antagonist ondansetron on voluntary ethanol intake in rats and mice maintained on a limited access procedure. *Psychopharm.* 117:479-485.