REPORT DOCUMENTATION PAGE	. <b>А</b>	FRL-SR-AR-TR-03-
The public reporting burden for this collection of information is estimated to average 1 hour per rescrict- gathering and maintaining the data needed, and completing and reviewing the collection of information. Sen of information, including suggestions for reducing the burden, to Department of Defanse, Washington (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Artington, VA 22202-4302. Respondents shou subject to any penalty for failing to comply with a collection of information if it does not display a currently to DEFACE DO NOT DEFILIPATION FORM TO THE AROVE ADDRESS.	, including the d comments n Headquartu ld be aware valid OMB cou	0109
1 REPORT DATE (DD-MM-YYYY) 2. REPORT TYPE	······	3. DATES COVERED (From - To)
31-03-2003 FINAL		01-02-2002-31-12-2002
4. TITLE AND SUBTITLE	5a. CO	NTRACT NUMBER
Workshop on Dielectric Properties and Electrical Breakdown of Biological		
Membranes	5b. GR	ANT NUMBER
		AFOSR F49620-02-1-0110
	5c PR	OGRAM ELEMENT NUMBER
		O POT WINDER
6. AUTHOR(S)	5d. PR	OJECI NUMBER
Karl H. Schoenbach		
	5e. TA	SK NUMBER
	5f. W0	DRK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(LS)		REPORT NUMBER
Old Dominion University Research Foundation		ODURF 221101
Hughes Hall		
NOTIOIK, VA 25529		ж.
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)
Dr. Robert I Barker		
AFOSR		
4015 Wilson Boulevard, Room 713		NUMBER(S)
Arlington, VA 22203-1954		
40 DISTRIBUTION (AVAILADILITY STATEMENT		
	n e estat Alexander	
		1117100 ANTACA
13. SUPPLEMENTARY NOTES		
	l	FAAAAAA MAL
14. ABSTRACT		
"Dielectric Properties and Electrical Breakdown of Biological Membranes"	was the top	ic of a workshop at the Conference on orkshop, supported by the IEEE Dielectrics
Electrical Insulation and Dielectric Phenomena on October 20, 2002, in Mer	hv Karl H.	Schoenbach. The presentations focused on
and Electrical insulation overery, the US ALCON, and ULIN, was organized	he interest i	n this topic was stimulated by recent
nonthermal biological effects caused by high power pulsed electric fields. T		•
nonthermal biological effects caused by high power pulsed electric fields. The observations of modifications in cell functions and induction of apoptosis in	cells when	exposed to ultrashort electrical pulses.
nonthermal biological effects caused by high power pulsed electric fields. The observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on members.	cells when brane structu	exposed to ultrashort electrical pulses. rres and functions, dielectric properties,
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memb electroporation, and applications. The funding was utilized to support the tra	cells when brane structu avel expense	exposed to ultrashort electrical pulses. area and functions, dielectric properties, es of invited speakers. A special issue on a UTD and K H. Schearberth (ODD) will
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on membrane electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes"	cells when brane structu avel expense with J. Wea	exposed to ultrashort electrical pulses. rres and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will negative
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on membre electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and	cells when brane structu avel expense with J. Wea Electrical I	exposed to ultrashort electrical pulses. ares and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will nsulation.
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memb electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and 15. SUBJECT TERMS	cells when brane structu avel expense with J. Wea Electrical I	exposed to ultrashort electrical pulses. Ires and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will insulation.
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memb electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and <b>15. SUBJECT TERMS</b> membrane structures, membrane functions, dielectric properties, electropora	cells when brane structu avel expense with J. Wea Electrical I	exposed to ultrashort electrical pulses. Ires and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will insulation. ermal biological effects, high power pulsed
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on meml electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and <b>15. SUBJECT TERMS</b> membrane structures, membrane functions, dielectric properties, electropora electric fields, ultrashort electrical pulses, apoptosis, gene therapy, biofoulir	a cells when brane structu avel expense with J. Wea Electrical I ation, nonth- ng, bacterial	exposed to ultrashort electrical pulses. ares and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will nsulation. ermal biological effects, high power pulsed decontamination
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memi- electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and <b>15. SUBJECT TERMS</b> membrane structures, membrane functions, dielectric properties, electropora electric fields, ultrashort electrical pulses, apoptosis, gene therapy, biofoulir	a cells when brane structu avel expense with J. Wea Electrical I ation, nonth- ng, bacterial	exposed to ultrashort electrical pulses. Ires and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will nsulation. ermal biological effects, high power pulsed decontamination AME OF RESPONSIBLE PERSON
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memi- electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and <b>15. SUBJECT TERMS</b> membrane structures, membrane functions, dielectric properties, electropora electric fields, ultrashort electrical pulses, apoptosis, gene therapy, biofoulir <b>16. SECURITY CLASSIFICATION OF:</b> <b>17. LIMITATION OF</b> <b>18. NUM</b> <b>OF</b>	cells when brane structu avel expense with J. Wea Electrical I ation, nonthong, bacterial	exposed to ultrashort electrical pulses. Ires and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will insulation. ermal biological effects, high power pulsed decontamination AME OF RESPONSIBLE PERSON I. Schoenbach
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memi- electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and <b>15. SUBJECT TERMS</b> membrane structures, membrane functions, dielectric properties, electropora electric fields, ultrashort electrical pulses, apoptosis, gene therapy, biofoulir <b>16. SECURITY CLASSIFICATION OF:</b> <b>a. REPORT b. ABSTRACT c. THIS PAGE</b> <b>17. LIMITATION OF</b> <b>ABSTRACT</b> <b>C. THIS PAGE</b>	a cells when brane structu avel expense with J. Wea Electrical I ation, nonth- ng, bacterial IBER 19a. N Karl H 19b. T	exposed to ultrashort electrical pulses. Ires and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will nsulation. ermal biological effects, high power pulsed decontamination AME OF RESPONSIBLE PERSON I. Schoenbach ELEPHONE NUMBER (Include area code)
nonthermal biological effects caused by high power pulsed electric fields. T observations of modifications in cell functions and induction of apoptosis in Prominent scientists from various countries presented invited talks on memil electroporation, and applications. The funding was utilized to support the tra "Dielectric Properties and Electrical Breakdown of Biological Membranes" be published (in October 2003) in the IEEE Transactions on Dielectrics and 15. SUBJECT TERMS membrane structures, membrane functions, dielectric properties, electropora electric fields, ultrashort electrical pulses, apoptosis, gene therapy, biofoulir 16. SECURITY CLASSIFICATION OF: 	a cells when brane structu avel expense with J. Wea Electrical I ation, nonth- ng, bacterial IBER 19a. N Karl H 19b. T	exposed to ultrashort electrical pulses. Ires and functions, dielectric properties, es of invited speakers. A special issue on aver (MIT) and K.H. Schoenbach (ODU) will insulation. ermal biological effects, high power pulsed decontamination AME OF RESPONSIBLE PERSON I. Schoenbach ELEPHONE NUMBER (Include area code) 757-683-4625

e

e

4. 4. 1000000

Mount for the second

News.compost v S.A.

NAME OF A LOCAL OF

.

8 1.1.1.1.1.1.1.1.1.1.1.1.1 Final Report on the Project

# "Workshop on Dielectric Properties and Electrical Breakdown of Biological Membranes"

Grant No: AFOSR F49620-02-1-0110

February 1, 2002 to December 31, 2002

submitted to:

### Dr. Robert J. Barker Air Force Office of Scientific Research

4015 Wilson Boulevard, Room 713 Arlington VA 22203-1954 Phone: (703) 696-8574 FAX: (703) 696-8481 e-mail: robert.barker@afosr.af.mil

by:

### Karl H. Schoenbach

Center for Bioelectrics Old Dominion University, Norfolk VA 23529 Phone: (757) 683-4625 FAX: (757) 683-3220 e-mail: schoenbach@ece.odu.edu

## DISTRIBUTION STATEMENT A

Approved for Public Release Distribution Unlimited

### Abstract

"Dielectric Properties and Electrical Breakdown of Biological Membranes" was the topic of a one-day workshop at the Conference on Electrical Insulation and Dielectric Phenomena on October 20, 2002, in Cancun, Mexico. The workshop, which was supported by the IEEE Dielectrics and Electrical Insulation Society, the US Air Force Office of Scientific Research, and the Office of Naval Research, was organized by Karl H. Schoenbach, director of the Center for Bioelectrics at Old Dominion University. The presentations and discussions focused on nonthermal biological effects caused by high power pulsed electric fields. The interest in this topic was stimulated by recent observations of modifications in cell functions and induction of apoptosis in cells when exposed to ultrashort electrical pulses. Prominent scientists from the United States, Japan, Germany and Israel presented invited talks on membrane structures, membrane functions, dielectric properties, and electroporation. In addition, an overview of applications of both reversible and irreversible electroporation in medicine and biology was presented. This included gene therapy, cancer treatment, biofouling prevention, and bacterial decontamination. The workshop concluded with a panel discussion. The funding provided by AFOSR was utilized to support the travel expenses of invited speakers. As a follow-up to the workshop, a special issue on "Dielectric Properties and Electrical Breakdown of Biological Membranes" with J. Weaver (MIT) and K.H. Schoenbach (ODU) will be published in October 2003 in the IEEE Transactions on Dielectrics and Electrical Insulation.

### Introduction

The membrane surrounding biological cells consists of a lipid bilayer with a thickness of approximately 5 nm. The dielectric constant is about 5, and its membrane resistivity is  $2x10^9 \ \Omega \text{cm}$  [1]. The function of the cell membrane is to isolate regions of different materials, but also to facilitate the flow of selected types of ions and molecules from one region to the other. This flow of ions passes through membrane channels, consisting of proteins, which are embedded in the membrane. These ion channels are either chemically (ligand) or electrically (voltage) controlled.

In the resting state, a constant voltage of approximately 70 mV (for mammalian cells) is applied across the membrane. The potential gradient is caused by differences in sodium and potassium ion concentrations inside and outside the cell. For excitable cells, such as nerve and muscle cells, relatively small depolarizing changes in the membrane voltage, on the order of 10 to 20 mV, lead to strong nonlinear changes in the membrane conductance, an effect which allows electrical signal propagation along the axons of neurons. It was studied and modeled by Hodgkin and Huxley in the 1950s [2], who received the Nobel prize for their research in 1962.

A transmembrane voltage of 70 mV corresponds to an electric field of 140 kV/cm. In experiments with artificial membranes free of proteins (channels), 180 mV could be applied before changes in the conductance were observed [3]. This voltage corresponds to a dielectric strength of 360 kV/cm. Higher dielectric strengths can be achieved when pulsed voltages are applied. Generally, for electroporation to occur, which is a kind of electrical breakdown of the membrane, voltages of 1 V need to be applied across the membrane [4] corresponding to dielectric strengths of 2 MV/cm. This holds true for a pulse duration in the microsecond range. For pulses with submicrosecond duration, this value is even higher.

An understanding of the dielectric properties of plasma membranes, and the changes in conductance with increasing voltage, are not only important for biologists, but might offer opportunities for physicists and engineers interested in novel dielectric materials. The facts that artificial membranes can easily be generated, and features such as self-healing effects (reversible electroporation), make them an interesting object of study for the dielectrics and electrical insulation community.

{In addition to effects on the outer membrane, new research results indicate that it is possible to affect subcellular structures [5]. Although this topic was only being presented in one of the ten invited talks, it generated so much interest in the audience that almost all of the discussion centered on this new topic of "Intracellular Electromanipulation".}

### References

¥

[1]. I. Ermolina, Y. Polevaya, Y. Feldman, B.Z. Ginzburg, and M. Schlesinger, "Study of Normal and Malignant White Blood Cells by Time Domain Dielectric Spectroscopy," IEEE Trans. Dielectrics and Electrical Insulation 8, 253 (2001).

[2]. A.L. Hodgkin and A.F. Huxley, "A Quantitative Description of Membrane Current and its Application to Conduction and Excitation in Nerve," J. Physiol. <u>117</u>, 500 (1952).

[3]. T.J. Lewis, "Electronic Processes in Biology," Phys. Med. Biol. 27, 335 (1982).

[4]. J.C. Weaver, "Electroporation of Cells and Tissues," IEEE Trans. Plasma Science <u>28</u>, 24 (2000).

[5]. Karl H. Schoenbach, Stephen J. Beebe, and E. Stephen Buescher, "Intracellular Effect of Ultrashort Pulses," J. Bioelectromagnetics <u>22</u>, 440 (2001).

Û

### **Technical Program**

The workshop encompassed invited talks, contributed papers (poster-papers), and a discussion session. The program included invited talks on the structure of biological membranes, electrical breakdown (electroporation), measurements of dielectric properties of membranes, modeling of electroporation, subcellular effects caused by nanosecond electrical pulses, and medical and environmental applications of electroporation and intracellular electromanipulation. The presentations were followed by a discussion session, where the invited speakers and representatives of funding agencies served as panelists. Much of the discussion had focused on the effect of ultrashort electrical pulses on biological cells, where the most recent research results were presented by E. Stephen Buescher. A poster session, as part of the general DEIS program on Monday, October 21 with contributed papers, complemented the invited talks (attachment).

### **Speakers and Topics**

- 1. Hidenori Akiyama and Sunao Katsuki, Kumamoto University, Japan, "Environmental Applications of Biological Inactivation by Pulsed Electric Field and/or Pulsed Discharges in Water"
- 2. E. Stephen Buescher, Eastern Virginia Medical School, USA, "Intracellular Electromanipulation of Mammalian Cells with Submicrosecond Pulsed Electric Fields"
- 3. Yuri Feldman, Hebrew University, Jerusalem, Israel, "Dielectric Properties of Biological Systems; from Amino Acids to Cells"
- 4. Ravindra. P. Joshi, Old Dominion University, USA, "Dynamical Modeling of Cellular Response to Short High-Intensity Electric Fields"
- 5. **Raphael C. Lee,** University of Chicago, USA "Trauma and Injury Resulting From Strong Electric Field Exposure"
- 6. Dietmar P. Rabussay, Genetronics, Inc, San Diego, USA, "Medical Applications of Electropermeabilization of Cell Membranes"
- 7. Christoph Schultheiss, Forschungszentrum Karlsruhe GmbH, Germany, "Industrial Applications of Electroporation"
- 8. HT Tien and A. Ottova, Michigan State University, USA, "Electrical and Dielectric Properties of Experimental Lipid Bilayers in Relation to Biotechnology"
- 9. James C. Weaver, Harvard-MIT Division of Health Sciences and Technology, USA, "Electroporation of Biological Membranes from Multicellular to Nano Scales"
- 10. Ahmed E. Yousef, C. Patrick Dunne, Ragip Unal and Beatrice H. Lado, Ohio State University, USA, "Damage of Bacterial Cell Membranes by Pulsed Electric Field Treatment."

## Workshop on

# **"Dielectric Properties and Electrical Breakdown of Biological Membranes"**

7.30 am	Registration
8.15 am	Introduction (Karl H. Schoenbach)
8.20 am	James C. Weaver, Harvard-MIT Division of Health Sciences
	and Technology, USA, "Electroporation of Biological
	Membranes from Multicellular to Nano Scales"
9.00 am	HT Tien, Michigan State University, USA, "Electrical and
	Dielectric Properties of Experimental Lipid Bilayers in Relation
	to Biotechnology"
9.30 am	Yuri Feldman, Hebrew University, Israel, "Dielectric
	Properties of Biological Systems; from Amino Acids to Cells"
10.00 am	Break
10.30 am	Raphael C. Lee, University of Chicago, USA, "Trauma and
	Injury Resulting from Strong Electric Field Exposure"
11.00 am	Ravindra. P. Joshi, Old Dominion University, USA,
	"Dynamical Modeling of Cellular Response to Short High-
	Intensity Electric Fields"
11.30 am	E. Stephen Buescher, Eastern Virginia Medical School, USA,
	"Intracellular Electromanipulation of Mammalian Cells with
	Submicrosecond Pulsed Electric Fields"
12.00 am	Lunch-Break
1.30 pm	Dietmar P. Rabussay, Genetronics, Inc, San Diego, USA,
	"Medical Applications of Electropermeabilization of Cell
	Membranes"
2.00 pm	Ahmed E. Yousef, Ohio State University, USA, "Damage of
	Bacterial Cell Membranes by Pulsed Electric Field Treatment."
2.30 pm	Break
2.45 pm	Christoph Schultheiss, Forschungszentrum Karlsruhe GmbH,
<b>•</b> • • •	Germany, "Industrial Applications of Electroporation"
3.15 pm	Hidenori Akiyama, Kumamoto University, Japan,
	"Environmental Applications of Biological Inactivation by
0.45	Pulsed Electric Field and/or Pulsed Discharges in Water"
3.45 pm	Break
4.15 pm	Panel Discussion

5.30 pm Adjourn **Abstracts of Invited Talks** 

## Environmental Applications of Biological Inactivation by Pulsed Electric Field and/or Pulsed Discharges in Water

Hidenori Akiyama and Sunao Katsuki Kumamoto University, Kumamoto 860-8555, Japan

The phosphorus and nitrogen compounds flow into lakes and marshes near urban areas. In these nourishing surroundings, algae plankton increases and covers the water surface. Lakes often become the sources of tap water, and a kind of algae is poisonous. The protection of the environment at lakes and marshes is one of important subjects. Also, the sewage without enough treatment flows into a river and the seaside, and the destruction of water environments is caused. The sewage treatment is also one of important subjects. The protection of water environment is our obligation for descendants. The biological inactivation using pulsed electric field and/or pulsed streamer discharges in water is a potential method to clean the water. Here, some challenging trials to protect the water environment with pulsed electric field and/or pulsed streamer discharges in water are described and discussed.

## Intracellular Electromanipulation of Mammalian Cells with Submicrosecond Pulsed Electric Fields

E. Stephen Buescher, Pamela S. Hair, Stephen J. Beebe, and Karl H. Schoenbach\* Eastern Virginia Medical School and \*Old Dominion University, Norfolk, VA

The effects of high power submicrosecond pulsed electric fields on living cells have been examined. Theoretically, reducing the duration of high electric field (MV/m) pulses from microseconds to nanoseconds should increasingly affect intracellular rather than surface membranes of living cells. Experimentally, square-wave, 60 ns duration, 3.6 - 5.3 MV/m electric field pulses applied in trains of 1-10 pulses result in progressive increases in the numbers of permeabilized intracellular granules in a human eosinophil cell model without large surface membrane effects. Electron micrographic examination of cells treated in this way demonstrates alteration of intracellular granule morphology consistent with permeabilization of granule membrane, i.e., intracellular electromanipulation. Continuous microscopic examination of individual living cells exposed to long or short duration pulsed electric field applications shows that rapid permeabilization of surface membrane (median 5 minutes) with anodic preference (electroporation) and prompt cellular swelling follow a single, long duration (100 ms) pulse. In contrast, after a single short duration (60 ns) pulse, onset of surface membrane permeability is delayed (median 17 minutes), the increased permeability shows no anodic preference, and cellular swelling is absent suggesting that these effects are due to intracellular electromanipulation rather than direct effects on the surface membrane. Submicrosecond, high power pulsed electric fields applied to living cells achieve preferential effects on intracellular rather than surface membranes, potentially providing new approaches for selective/generalized cell or tissue ablation, growth stimulation and tissue remodeling.

## Dielectric Properties Of Biological Systems; From Amino Acids To Cells

Yu.Feldman, The Department of Applied Physics, The Hebrew University of Jerusalem, Jerusalem, 91904, Israel

An interesting and important subject in biophysics is the investigation of the dielectric properties of complex biological systems like cells and their structural components (amino acids, proteins, membranes, cytoplasm etc.). These can bring valuable knowledge about the structure and dynamic behavior of proteins in solutions and different cell structures and their functions and metabolic mechanism in suspensions.

The aim of this contribution is to present a comprehensive theoretical and experimental study by means of time domain dielectric spectroscopy (TDDS) of static and dynamic dielectric properties of different biological systems including amino acids, regular and membrane proteins, normal and malignant blood cells of different types.

The dipole correlation functions for different globular proteins, such as myoglobin, lysozyme and RNase A solutions at different concentration, pH and temperature were obtained. The processes of the intramolecular interactions of myoglobin molecules, the pH-induced dimerization of lysozyme and thermal denaturation of RNase A are discussed. The dynamic structure study of native glucose oxidase and glucose oxidase, which was hydrophobized by covalent attachment of palmitoyl chains, is presented. We report also the application of (TDDS) to bacteriorhodopsin (bR) containing purple membrane films. The results of these measurements unexpectedly show that oriented purple membrane has unique liquid crystal like ferroelectric behavior. The dielectric behavior can be considered as a soft mode relaxation processes in ferro-electric liquid crystals near smectic-C\* - smectic-A phase transition.

Suspensions of normal lymphocytes of T and B types and leukaemia lymphocytes (Raji, Daudi, Bjab, Peer and HDMar) were investigated at the low volume fractions (3-10%) where the cellular interactions are negligible small. The measurements were performed at a wide frequency range from 200 KHz to 3 GHz at room temperature 25 °C and physiological temperature 37 °C. The effect of electrode polarization in TDDS experiment was corrected. Cell radii and volume fraction were measured by independent methods.

It was shown that dielectric permittivity; capacitance and conductivity values of cell membrane are higher for normal lymphocytes in comparison with malignant ones. The difference of the same parameters between normal-B cells and normal-T is also discussed.

## Dynamical Modeling of Cellular Response to Short High-Intensity Electric Fields

### R. P. Joshi, Q. Hu and K. H. Schoenbach Department of Electrical and Computer Engineering Old Dominion University, Norfolk, VA 23529

The application of short, high-intensity electric pulses provides a mechanism for the non-thermal manipulation of biological cells. Some of the direct cellular effects include electroporation-induced variations in conductivity, ionic transport, shape deformation, localized energy absorption, and the onset of electro-chemical processes. It is already well known that the overall cellular outcome can either be reversible or irreversible depending on the characteristics of the external electrical excitation. An important first step for achieving a desired cellular response with as much control as possible, is to develop a quantitative understanding of electromagnetic field-cell interactions for such pulsed exposures. Previous simulation work pertaining to electric field effects on biological systems has generally been based on quasi-static models or invoked Laplace equation for the field distribution. In this presentation the dynamical aspects will be discussed. A comprehensive model at the cellular level that includes electrodynamics, changes in cell geometry, and electroporative variations associated with factors such as surface tension and pore area will be emphasized. A dynamical model allows for a better physical representation, and better agreement with observed experimental trends emerge. Implications for nanosecond, high voltage pulses will also be discussed.

## Trauma and Injury Resulting From Strong Electric Field Exposure

### Raphael C. Lee, MD, ScD, PhD (Hon) University of Chicago, USA

Human use of electrical power is increasing. Along with that are increased opportunities for electrical injury. Victims of contact with high-energy electrical systems manifest a wide spectrum of injury depending on the electrical power capacity, current path through the body, duration of contact and healing response. Primary mechanisms of damage include membrane electroporation, electroconformational denaturation of membrane proteins and Joule heating. If arcing occurs, then acoustic blast injury can result. Skeletal muscle and nerves, tissues designed to be very sensitive to electrical signals, are particularly vulnerable to injury by the supraphysiological electric forces that are often associated with human contact with high voltage industrial power sources. Neurological and muscular damage often manifest despite lack of evidence of thermal burn. Severely injured victims present enormous medical management challenges due to release of inflammatory cytokines and altered cardiovascular behavior.

This paper will discuss the field exposure-tissue injury relationship and clinical manifestations of electrical injury. Approaches to rapid diagnostic imaging and therapy will be reviewed. In addition, the mechanism of action of promising new therapeutic amphipolar copolymers, which mimic behavior of natural chaperones, that have been shown to reduce tissue damage will be mentioned.

#### **References:**

- 1. Chen, W. and Lee, R.C. "Altered Ion Channel Conductance and Ionic Selectivity Induced by Large Imposed Membrane Potential Pulse" Biophys. J. 67(2): 603-612, 1994
- 2. Mankani, M., Abramov, G., Boddie, A., and Lee, R.C. "Detection of Peripheral Nerve Injury in Electrical Shock Patients", Annals New York Academy of Science, Vol. 720, pp. 206-212, 1994
- 3. Marks, J., Pan, C., Bushell, T., Cromie, WJ and Lee RC, "Amphiphilic, tri-block Copolymers Provide Potent, Membrane-targeted Neuroprotection" FASEB J. 15(6): 1107-9, 2001
- 4. Maskarinec, S.A., Hannig, J., Lee, R.C., and Lee, K-Y. "Direct Observation of Poloxamer 188 Insertion Into Lipid Monolayers" Biophys. J 2002 82(3):1453-9
- 5. <u>Occupational Electrical Injury and Safety</u>. Chen, C.-T., Lee, R.C., Shih, J.-X., and Zhong, M.-H. Eds., Annals of the New York Acad. of Science, Vol. 888, 1999

## Medical Applications of Electropermeabilization of Cell Membranes

Dietmar P. Rabussay

Genetronics, Inc., 11199 Sorrento Valley Road, San Diego, CA 92121

Electropermeabilization (EP, also referred to as "electroporation") of outer cell membranes can be achieved by exposing cells to electrical pulses of field strengths between 100 and 2000 V/cm and pulse durations of 10<sup>-6</sup> to 10<sup>-1</sup> sec. EP is applicable to cells in suspension ("in vitro") as well as to cells in live tissue ("in vivo"). The optimal conditions for EP depend on many factors, including the properties of the target cells and their environment, the pulse parameters, and the geometry of the electrodes and their contact with the cells to be electroporated. EP has been used for about 20 years to deliver DNA and other molecules into cells in vitro, but its potential for medical applications was not realized until about a decade ago. Since then, a variety of medical applications have been explored in pre-clinical and clinical studies. The most advanced application, now in a Phase III clinical trial, is the treatment of malignant tumors. In this application, EP enhances the intracellular uptake of anti-tumor agents by tumor cells up to several thousand-fold compared to the uptake observed in conventional chemotherapy, and causes significantly improved tumor shrinkage or elimination. This cancer treatment modality has shown excellent potential for many different types of solid tumors. Another application of EP, the effective delivery of DNA into cells of several target tissues opens up the possibility of breakthroughs in gene therapy and DNA vaccination. EP appears to combine high efficiency, safety and economical feasibility, a combination that other competing DNA delivery methods have not been able to achieve. Experiments in large animals have confirmed the potential for clinical applications of EP in gene therapy and DNA vaccinations and the first clinical study is now in progress. Other in vivo medical applications of EP include many forms of vascular therapies (both drug and gene-based), treatment of glaucoma, and the use of EP-facilitated intradermal or transdermal delivery of drugs and vaccines to treat local and systemic diseases. Finally, EP has been explored as a technique to enable or enhance ex vivo therapeutic approaches. In ex vivo therapies, cells or organs are treated outside the patient's body before they are infused or implanted Specific applications include the modification of blood clotting into the patient. behavior, cellular immune therapies of cancer and other diseases, and prevention of organ rejection after transplant.

In summary, EP has opened up new approaches to solving numerous medical problems whose solutions were hampered by the difficulty of introducing drugs and genetic material across the barrier of the cell membrane. Some applications have progressed to advanced clinical trials and seem to fulfill the promise of this technology. Many other potential applications are still in early stages of development or have not yet been explored at all.

## Industrial Applications of Electroporation

C. Schultheiss, H.-J. Bluhm and H.-G. Mayer\*

Forschungszentrum Karlsruhe GmbH, Institute for Pulsed Power and Microwave

Technology,

\*Technology Transfer and Marketing, Postfach 3640, D-76021 Karlsruhe, Germany M. Kern, Fa. Lutz & Kern, Saarstrasse 4, D-68753 Waghäusel-Kirrlach, Germany

Electroporation (EP) of biological cells has become a rapidly developing area of biotechnology. If external electric fields are applied across biological cells in a water suspension the cytoplasm becomes polarized. The charge displacement in the cytoplasm and in the suspension leads to high transmembrane potentials in the order of several volts. As a result large irreversible openings can built-up in the cell membrane, the cytoplasm can flow-out of and the cell can perish.

The subject of this talk is firstly to report on actual experiments at the mobile industrial electroporation pilot system KEA (Karlsruhe Elektroporation Anlage) with whole plants[1]. The maximum throughput of this device is 2 tons/hour. The reactor is a 16 cm-tube with integrated electrode pairs, where pulses with amplitudes up to 300 kV are applied at a repetition rate of 7 Hz. An overview of results from treated plants like beets, apples, grapes etc. will be given.

Secondly, design studies, development and test of an industrial EP production device will be presented. Since plants, as for example beets, are large in diameter and nearly weightless in water (suspension), the size of an electroporation reactor gap has to be large and the plants must be transported continuously by means of a forced feedthrough. In the test facility ZAR (Zell Aufschluss Reaktor) a throughput in the order of 1 ton/min has been demonstrated. The operation of such a powerful device with a reactor spacing of 30 cm needs an assembly of Marx generators with high pulse repetition rates to establish fields at the reactor site as well as to fulfill power demands.

At Forschungszentrum Karlsruhe different types of Marx generators with gas spark gaps have been developed and tested. They are able to run with pulse frequencies in the order of 20-30 Hz (test Marx generator with 100 kV pulses). The pulse amplitude needed is 350 kV, the pulse length is 1 µs and the pulse energy fed into a 20 Ohm load is about 1 kJ. The life time of the electric components is designed to withstand a minimum of up to 200 million discharge cycles (3 month operation). The complete system consisting of electroporation reactor plus conveyance is designed to be mainly built from dielectric materials to avoid undesired shielding effects of the electric field in the area of the reactor chamber. However high voltage strength dielectric materials like polyethylene or polypropylene underlie enhanced mechanical wear which must be taken into account.

1. C. Schultheiss, H.-J. Bluhm, H.-G. Mayer and M. Kern, "Industrial-Scale Electroporation of Plant Material Using High Repetition Rate Marx Generators", IEEE Pulsed Power Plasma Science 2001, Las Vegas June 17-22, 2001

## Electrical and dielectric properties of experimental lipid bilayers in relation to biotechnology

HT Tien and A Ottova

Physiology Department, Biomedical & Physical Sciences Building Michigan State University, East Lansing, MI 48824

Transmembrane voltage pulses between 200 mV to 1 V are known to create transient 'pores' in the lipid bilayer, with a diameter about 2 nm. This phenomenon, termed electroporation, has been extensively investigated [1]. Electroporation occurs following electric pulses up to  $10^6$  V/cm with duration between µsec and msec to membranes in close contact and is believed to initiate primarily in the lipid bilayer. This talk will begin with a brief summary of the origin of lipid bilayer research. One of practical applications of electroporation is cell transfection for gene expression. Other applications include encapsulation of drugs in controlled-release and insertion of proteins in living cells. For instance, the effects of the channel-forming peptide gramicidin D (gD) on the conductance and electroporation thresholds of BLMs have been reported [2]. It seems likely that the presence of membrane proteins affect the electroporation of the lipid bilayer by changing its mechanical properties. Transport of ions such as Na<sup>+</sup>, K<sup>+</sup>, Cl<sup>-</sup> through membrane pores discharge the membrane potential, and at times an external pulse of sufficient amplitude and duration tends to cause dielectric breakdown of the lipid bilayer. Molecular transport through primary pores and pores enlarged by secondary processes provides the basis for transporting molecules into and out of cells. Tissue electroporation, by longer, large pulses, is involved in electrocution injury. Tissue electroporation by shorter, smaller pulses is under investigation for biomedical engineering applications of medical therapy aimed at cancer treatment, gene therapy, and transdermal drug delivery [3].

#### References

- 1. H. T Tien and A.L. Ottova, Membrane Biophysics: As viewed from experimental bilayer lipid membranes (planar lipid bilayers and spherical liposomes), Elsevier, Amsterdam, 2000, 648 pp.
- 2. H.T. Tien and A.L. Ottova, The Lipid Bilayer Concept and Its Experimental Realization: From Soap Bubbles, Kitchen Sink, to Bilayer Lipid Membranes J. Memb. Sci., 189 (2001) 83
- 3. A. Ottova, V. Tvarozek and H.T. Tien, Supported Planar Lipid Bilayers (s-BLMs) in Planar lipid bilayers (BLMs) and their applications, Elsevier, 2002, to be published

## Electroporation of Biological Membranes from Multicellular to Nano Scales

### James C. Weaver Harvard-MIT Division of Health Sciences and Technology Massachusetts Institute of Technology Cambridge, MA 02139

Several decades ago both cell and artificial planar bilayer membranes were found to exhibit dramatic changes if the transmembrane voltage was raised to 0.2 to 1 V by short pulses. Ionic and molecular transport were found to be increased by orders of magnitude, with both reversible and irreversible outcomes possible. Initially the term "breakdown" was applied. However, mechanistic interpretations presently exclude ion pair generation of classic dielectric breakdown. Instead, a transient aqueous pore hypothesis can account for many features of electroporation in bilayer membranes comprised of low dielectric constant constituents. My presentation will consider transient aqueous pore-based phenoemena in planar bilayers, single cell systems often encountered in vitro, multicellular systems relevant to in vivo applications, and the expectation for significantly modified behavior for systems with very small spatial extent.

## Damage of Bacterial Cell Membranes by Pulsed Electric Field Treatment

Ahmed E. Yousef<sup>1</sup>, C. Patrick Dunne<sup>2</sup>, Ragip Unal<sup>3</sup> and Beatrice H. Lado<sup>1</sup> <sup>1</sup> Ohio State University, USA; <sup>2</sup> US Army Natick Soldier Center, Natick, Massachusetts, USA; <sup>3</sup> Department of Food Engineering, Mersin University, Turkey

High-intensity electric pulses damage membranes of the microbial cell, leading to its injury and death. Based on this concept, pulsed electric field (PEF) emerged as a preservation technology to inactivate disease-causing microorganisms in food (Lado and Yousef, 2002). PEF treatment effectively inactivates bacteria and yeast in orange juice, milk, skim milk, yogurt, liquid egg and pea soup (Vega\_Mercado et al., 1997); however, PEF is ineffective against bacterial and mold spores. This emerging food processing technology is developed with the promise to produce safe food and preserve its sensory qualities.

Lethality of PEF probably results from the electroporation of cell membranes. Formation of pores alters membranes' barrier properties, leading to loss of cell homeostasis. Membrane damage may be measured using fluorescence probes that gain entry only in cells with dysfunctional membranes. A fluorescent nucleotide-binding probe, Propidium Iodide (PI), was used recently to measure cell membrane damage during PEF treatment (Unal et al., 2002). According to this study, membrane porosity correlates well with cells' inactivation by PEF treatment. Therefore, measurement of membrane damage resulting from pore formation may serve as an index of PEF effectiveness against foodborne pathogens. Fluorescence staining technique showed that pre-treatment of gram-negative bacterial cells with lysozyme and EDTA increased the PI uptake by the PEF-treated cells. Therefore, it is possible to use the fluorescence staining technique to measure PEF-caused cell injury that is not detectable by cultural techniques.

#### **References.**

Lado, B. H. and Yousef, A.E. 2002. Alternative food preservation technologies: Efficacy and mechanisms. Microbes and Infection 4: 433\_440

Vega\_Mercado, H., O. Martin\_Belloso, B.L. Qin, F.J. Chang, M.M. Gongora\_Nieto, G.V. Barbosa\_Canovas, and B.G. Swanson. 1997. Non\_thermal food preservation: Pulsed electric fields. Trends Food Sci. 8:151\_157.

Unal, R., Yousef, A.E., and Dunne, C.P. 2002. Spectrofluorometric assessment of bacterial cell membrane damage by pulsed electric field. Innovative Food Science & Emerging Technologies (In press).

**Poster Papers** related to Dielectrics can be found in the CEIDP, 2002 Annual Report Conference on Electrical Insulation and Dielectric Phenomena, 02CH37372.

M. Stacey, J. Stickley, P. Fox, C. O'Donnel, K. Schoenbach, S. Beebe, and S. Buescher. "Increased cell killing and DNA damage in cells exposed to ultra-short pulsed electric fields" pp. 79-82

T. J. Lewis. "Electromechanical effects in biological membranes" pp. 87-90

N. Hassan, I. Chatterjee, N. G. Publicover, and G. L. Craviso "A combined experimental and computational analysis of membrane potential variations in excitable cells in response to DC electric fields" pp. 91-94

### **2002 IEEE/DEIS conference**

The workshop was part of the 2002 IEEE/DEIS conference, which was held in Cancun, Mexico, October 20–24, 2002. The Conference on Electrical Insulation and Dielectric Phenomena (CEIDP) is sponsored by the IEEE Dielectrics and Electrical Insulation Society to provide an international forum for the discussion of work in progress on research on dielectric phenomena and measurements. The conference provides an opportunity for specialists from around the world to meet and exchange their findings.

#### Workshop Fee:

A conference fee of \$ 50 was charged to participants. This fee was used to cover expenses of documentation and beverages at coffee breaks.

### Documentation

A book with the abstracts of the invited talks and copies of relevant publications has been provided to the participants. In addition, a special issue on "Biodielectrics" in the IEEE Transactions on Dielectrics and Electrical Insulation with J.C. Weaver and K.H. Schoenbach as Guest Editors will be published. The special issue will contain invited papers based on invited talks at the workshop and contributed papers.

# Publicity



# Workshop on "Dielectric Properties and Electrical Breakdown of Biological Membranes"

A one-day workshop on "Dielectric Properties and Electrical Breakdown of Biological Membranes" was held on 20 October during the 2002 CEIDP in Cancun, Mexico. The workshop was supported by the IEEE Dielectrics and Electrical Insulation Society, the US Air Force Office of Scientific Research, and the Office of Naval Research.

The membrane, which encompasses biological cells, consists of a lipid bilayer with a thickness of approximately 5 nm. The dielectric constant is about 5, and its membrane resistivity is  $2 \times 10^9 \Omega$ cm. The function of the cell membrane is to isolate regions of different materials, but also to facilitate the flow of selected types of ions and molecules from one region to the other. Electrical breakdown of the membrane is defined as the onset of reversible or irreversible changes in the cell membrane (electroporation)

ogy was presented. This included gene therapy, cancer treatment, biofouling tion, an overview of applications of electroporation in medicine and bioland occurs at membrane voltages of greater than 1 V, which corresponds to electric fields of greater than  $2 \times 10^6$ V/cm. A group of leading scientists in the field of biodielectrics presented invited talks on membrane structures, irreversible prevention, and bacterial decontamimembrane functions, dielectric properties, and electroporation. In addireversible and both



Karl Schoenbach, summarizing the Workshop on Biological Membranes.

nation. The workshop was concluded with a panel discussion.

biodielectrics subcommittee to offer a follow-up workshop in biodielectrics workshop in the IEEE Transactions on Guest editors are J. Weaver (MIT) and line for submission of manuscripts is on the part of the DEIS community in engineering and biology. This was shop participants when asked about shop, and by the decision of the at the 2003 CEIDP in Albuquerque, NM. Another opportunity to learn about, and to participate in, biodielectrics is the publication of a special issue on the topic of the 2002 K.H. Schoenbach (ODU). The dead-Obviously, there is a strong interest this research area at the interface of manifested by the very positive feedback of the approximately 50 workthe content and format of the work-Dielectrics and Electrical Insulation. 15 January 2003.

—This report was written by Karl H. Schoenbach, Chairperson of the Workshop. Announcing a Special Issue on

# Dielectric Properties and Electrical Breakdown of Biological Membranes

## in the

# IEEE Transactions on Dielectrics and Electrical Insulation

## with

J.C. Weaver and K.H. Schoenbach as Guest Editors.

Deadline for Submission: January 15, 2003

Information for authors can be found in any recent issue of the IEEE Transactions on Dielectrics and Electrical Insulation.

Please mail manuscripts to either

Dr. James C. Weaver

MIT Bldg #16-319 MIT Cambridge, MA 02139 Phone: 617-253-4194 FAX: 617-253-2514 e-mail: jcw@mit.edu or

Dr. Karl H. Schoenbach Old Dominion University 231 Kaufman Hall Norfolk Virginia 23529-0246 *Phone:* 757-683-4625 *FAX:* 757-683-3220 *e-mail:* schoenbach@ece.odu.edu

# List of Attendees

. . .

Mahmoud Abou-Dakka 1500 Montreal Rd INMS M-50, R 225 Ottawa, Ontario K1J 6W1 Canada Tel: 613-990-3619 Fax: 613-952-9366 Mahmoud.abou-dakka@nrc.ca

Hidenori Akiyama Kurokami 2-39-1 Kumamoto, 860-8555 Tel: 81-96-42-3618 Fax: 81-96-342-3618 Japan akiyama@eecs.kumamoto-u.ac.jp

Mike Arnold IRL Communications & Sensors P.O. Box 31-310 Lower Hutt New Zealand Tel: 64-4-569-0233 <u>m.Arnold@irl.cri.nz</u>

Soli Bamji 1200 Montreal Rd, Bldg M-50 Ottawa, Ontario K1A OR6 Canada Tel: 613-990-4021 Soli.bamji@nrc.ca

Yves Bertrand Sites des Renardieres Moret sur Loing, 77250 France Tel: 33-1-60736994 Fax: 33-1-60737477 Yves.Bertrand@edf.fr

Stephen Buescher 855 W Brambleton Ave Norfolk, VA 23510 Tel: 757-668-6464 Fax: 757-668-6476 sbuesche@chkd.com Indira Chatterjee Dept of ElectricalEngineering MS 260, University of Nevada Reno, NV 89557 USA Tel: 775-784-1346 Fax: 775-784-6627 indira@ee.unr.edu

्यत्र मोनः

Ladji Cisse 1500 Montreal Rd INMS M-50 Room 225 Ottawa, Ontario K1A 0R6 Canada Tel: 613-990-3619 Fax: 613-952-9366 Ladji cisse@nrc.ca

Gale L. Craviso Department of Pharmacology Howard Building, Room 219 University of Nevada, Mail Stop 318 Reno, NV 89557 USA Tel: 775-784-4118 Fax: 775-784-1620 gcraviso@unr.edu

Eric David 1100 Notre-Dam Ouest Montreal, Quebec H3C 1K3 Canada Tel: 514-396-8528 Fax: 514-396-8624 Eric.david@etsmtl.ca

Luigi Egiziano Dept of Information & Electrical Eng University of Salerno Via Ponte don Melillo Fisciano, I-84084 Italy Tel: 39-089-964293 Fax: 39-089-964218 egiziano@diiie.unisa.it Yuri Feldman Department of Applied Physics The Hebrew University of Jerusalem 91904 Jerusalem, Israel yurig@vms.huji.ac.il

R A Fouracre Dept of Electronic and Electrical Engineering University of Strathclyde Glasgow G1 1XW t.fouracre@eee.strath.ac.uk

Michel Frechette 1800 boul.Lionel-Boulet Varennes, Quebec J3X 151 Canada Tel: 450-652-8216 Fax: 450-652-8962 frechett@ireg.ca

Kaori Fukunaga 4-2-1, Nukui-Kita Koganei Tokyo, 184-8795 Japan Tel: 81-42-327-6259 Fax: 81-42-327-6675 kaori@crl.go.jp

Allen Garner Center For Bioelectrics Old Dominion University 231 Kaufman Hall Norfolk, VA 23529 agarner@odu.edu

Reimund Gerhard-Multhaupt Dept of Physics University of Potsdam Am Neuen Palais 10 D-14469 Potsdam Germany Tel: 49-331-977-1615 Fax: 49-331-977-1577 rgm@rx.uni-potsdam.de Ron Gilgenbach Bioelectromagnetism Lab Nuclear Engineering & Radiological Sciences Dept University of Michigan Ann Arbor, MI 48109-2104 USA Tel: 734-763-1261 Fax: 734-763-4540 rongilg@umich.edu

Francisco Camacho Gonzalez Hasenheide 57 Berlin Germany Tel: 4930-6981-8388 fcamacho@canopus.physik.uni-potsdam.de

Erling Ildstad Tronohem, 7491 Norway Tel: 47-7394226 Fax: 47-73394279 Erling.ildstad@elkraft.nthu.no

D. Randy James Oak Ridge National Lab P.O. Box 2008 4500 S MS-6122 Oak Ridge, TN 37831-6122 USA Tel: 865-574-6213 Fax: 865-574-6210 jamesdr@ornl.gov

Tom Johnson USU HS/PMB 4301 Jones Bridge Rd Bethesda, MD 20814 Tel: 301-319-6953 tjohnson@usuhs.mil

Ravindra Joshi Old Dominion University 231 Kaufman Hall Norfolk, VA 23529 USA Tel: 757-683-4827 Fax: 757-683-3220 rjoshi@odu.edu

۶,

Karan V. Kaler Dept of Electrical & Computer Eng 2500 University Dr, N.W. Calgary, Alberta T2N-1N4 Canada Tel: 403-220-5809 Fax: 403-282-6855 kaler@enel.ucalgary.ca

Sunao Katsuki Dept Electrical & Computer Eng Kumamoto University Kurokami 2-39-1 Kumamoto, Japan 860-8555 katsuki@eecs.kumamoto-u.ac.jp

Yue-Ying lau 2355 Bonisteel Blvd Ann Arbor, MI 48109-2104 USA Tel: 734-764-5122 Fax: 734-763-4540 yylau@umich.edu

Raphael C. Lee Plastic & Reconstructive Surgery 5841 South Maryland Ave MC 6035/suite J-641 Chicago, Il 60637 USA Tel: 773-702-6302 Fax: 773-702-1634 rlee@surgery.bsd.uchicago.edu

T. John Lewis School of Informatics University of Wales, Bangor Gwynedd LL57 1UT. U.K. Tel: 44-1248-382684 Fax: 44-1248-361429 lewis@informatics.bangor.ac.uk

Youlan Li Dept of Electrical & Computer Eng University of Calgary 2500 University Dr N.W. Calgary, Alberta T2N 1N4 Canada Tel: 1-403-289-3498 liy@enel.ucalgary.ca John Mathes Honeywell FMIT/New Mexico Box 5250 Albuquerque, NM 87185 Tel: 505-844-2625 Fax: 505-844-3751 jmathes@kcp.com

Gavita Mugala Teknikringen 33 Royal Institute of Technology (KTH) Dept of Electrical Eng Stockholm, 100 44 Sweden Tel: 46-8-7906258 Gavita.mugala@ekc.kth.se

Masayuki Nagao Dept of Electrical & Electronic Eng Toyohashi University of Technology Hibarigaoka 1-1 Tempaku Toyohashi, 441-8580 Japan Tel: 81-532-44-6725 Fax: 81-532-48-7634 Nagao@eee.tut.ac.jp

Angelica Ottova Department of Physiology Biomedical & Physical Sciences Bldg East Lansing, MI 48824 USA Tel: 517-355-6475 ext 1347 Fax: 517-355-5125 ottova@msu.edu

Dietmar Rabussay 1119 Sorrento Valley Rd San Diego, CA 92075 USA Tel: 858-410-3183 Fax: 858-410-3397 dietmarr@genetronics.com Govinda Raju Electrical Engineering Dept Essex Hall, Sunset Ave Windsor, Ontario N9B3P4 USA Tel: 519-253-3000 raju@uwindsor.ca

Clive W Reed 116 Woodhaven Dr Scotia, NY 12302 USA reedcw@nycap.rr.com

Gerhard Salge Segelhof Baden-Daettwil, CH-5405 Switzerland Tel: 41-58-5868113 Gerhard.salge@ch.abb.com

Karl H. Schoenbach Center For Bioelectrics Old Dominion University 231 Kaufman Hall Norfolk, VA 23529 Tel: 757-683-4625 Fax: 757-683-3220 schoenbach@ece.odu.edu

Christoph Schultheiss Forschungszentrum karlsruhe GmbH POB 3640 Karlsruhe, D-76021 Germany Tel: 49-7247-82-4384 Fax: 49-7247-82-6126 Christoph.schultheiss@ihm.fzk.de

Raji Sundararajan 7001 E Williams Field Rd, Bldg 50 Mesa, AZ 83212 USA Tel: 480-727-1507 Fax: 480-727-1723 raji@asu.edu H T Tien Physiology (BPSB) Michigan State University East Lansing, MI 48824 USA Tel: 517-355-6475 Fax: 517-432-1967 tien@msu.edu

Vincento Tucci Dept of Information & Electrical Eng University of Salerno Vio Point don Melillo Fisciano Italy Tel: 39-089-964270 Fax: 39-089-964218 vtucci@unisa.it

P. Thomas Vernier MOSIS, USC-ISI 4676 Admiralty Way Marina Del Rey, CA 90292 USA Tel: 310-448-8752 vernier@ieee.org

Massimo Vitelli Via Roma Aversa (CE) 81031 Italy Tel: 39-081-5010240 Fax: 39-081-5037042 vitelli@unina.it

James Weaver Room 16-319 Harvard – MIT, Division of Health Sci & Tech Cambridge, MA 02139 USA Tel: 617-253-4194 Fax: 617-253-2514 jcw@mit.edu

Timothy J. Whelan PO Box 419159 FC39 Kansas City, MO 64141 USA Tel: 816-997-7188 Fax: 816-997-7169 twhelan@kcp.com Ahmed Yousef 2015 Fyffe Rd Parker Building Columbus, OH 43210 USA Tel: 614-292-7814 Fax: 614-292-0218 Yousef.l@osu.edu