

AD \_\_\_\_\_

Award Number: DAMD17-98-1-8353

TITLE: Modulation of Growth and Differentiation in Breast Cancer  
by Soy Isoflavones

PRINCIPAL INVESTIGATOR: Omer Kucuk, M.D.

CONTRACTING ORGANIZATION: Wayne State University  
Detroit, Michigan 48202

REPORT DATE: November 2002

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20030513 094

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE November 2002	3. REPORT TYPE AND DATES COVERED Annual (1 Oct 01 - 1 Oct 02)	
4. TITLE AND SUBTITLE Modulation of Growth and Differentiation in Breast Cancer by Soy Isoflavones			5. FUNDING NUMBERS DAMD17-98-1-8353	
6. AUTHOR(S): Omer Kucuk, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Wayne State University Detroit, Michigan 48202  E-MAIL: kucuko@karmanos.org			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words)  Our studies investigate the <i>in vivo</i> effects of soy isoflavones on human breast tissues obtained from lumpectomy/mastectomy specimens. We investigate the effect of increased tissue concentration of isoflavones for a period of three weeks on breast cell proliferation, differentiation and cell cycle regulatory proteins. Patients with ductal carcinoma <i>in situ</i> (DCIS) or invasive breast cancer are randomly assigned to take 100 mg soy isoflavone (Novasoy™, Archer Daniels Midland Company, Decatur, Illinois) or placebo daily for three weeks prior to surgery. Plasma isoflavone levels are measured at baseline and after three weeks in both groups. Tissue isoflavone levels are measured on samples from benign breast tissues in both groups. Biomarker studies are performed on surgical specimens by immunohistochemistry and Western blot. These studies will enable us to determine if a short duration of exposure to increased tissue levels of isoflavones will modulate biomarkers of cell differentiation (Cx43), adhesion (E-cadherin), proliferation (MIB-1), and cell growth and apoptosis (bcl-2, bax, p53, p21, Rb, EGF-R, cyclin D1, CDK5, CDK6) in benign, pre-malignant and malignant areas of breast epithelial tissues. Biomarker studies on the patients randomized on this study will be completed in early 2003.  We have entered 68 patients on study and 4 patients will be entered before the study is closed. We have started to do immunohistochemical and blood biomarker studies including soy isoflavone levels and NF-kappaB and Akt levels. The results will be fully analyzed after the accrual is completed over the next several months.				
14. SUBJECT TERMS CTR, breast cancer			15. NUMBER OF PAGES 7	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

## Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	5
Conclusions.....	7
References.....	7
Appendices.....	7

#### 4. INTRODUCTION

Epidemiological studies have shown an inverse association between dietary intake of fruits and vegetables and carcinoma of the breast. One group of major micronutrients in vegetables and fruits, which have been postulated to prevent breast cancer, are soy isoflavones. The mechanism by which isoflavones may prevent breast cancer is not known. Based on our preliminary studies, we hypothesize that isoflavones inhibit cell proliferation, upregulate the expression of gap junctional protein connexin 43 (cx43) and alter the expression of cell cycle regulatory proteins. Our studies will investigate the *in vivo* effects of isoflavones on human breast tissues obtained from lumpectomy/mastectomy specimens. We will investigate the effect of increased tissue concentration of isoflavones for a period of three weeks on breast cell proliferation, differentiation and cell cycle regulatory proteins. Sixty-four patients with ductal carcinoma *in situ* (DCIS) or invasive breast cancer scheduled to have surgery will be randomly assigned to supplement their diet with 100 mg soy isoflavone or placebo daily for three weeks. Plasma isoflavone levels will be measured at baseline and after three weeks in both groups. Tissue isoflavone levels will be measured on samples from surgical specimens, in benign and malignant areas of the epithelia, in both groups. Biomarker studies will be done on surgical specimens by immunohistochemistry and Western blot analysis. Comparisons will be made between areas of comparable microscopic characteristics [malignant, DCIS, lobular carcinoma *in situ* (LCIS), dysplasia, hyperplasia and benign] on breast tissues of patients from intervention and control groups. These studies will enable us to determine if a short duration of exposure to increased tissue levels of isoflavones will modulate biomarkers of cell differentiation (cx43), adhesion (E-cadherin), proliferation (MIB-1), and cell growth and apoptosis (bcl-2, bax, p53, p21, Rb, EGF-R, cyclin D1, CDK5, CDK6) in benign, pre-malignant and malignant areas of breast epithelial tissues. In addition, since baseline biopsy samples are available, a limited number of the marker studies (prioritized in the order cx43, bcl-2, p21, CDK5) will be performed on pre-intervention biopsy samples of patients in the intervention group, giving us an opportunity to compare pre- and post-intervention marker levels in the same patient.

#### 5. BODY


During the first year of the study, there was a delay in getting the study started because of difficulty in hiring study personnel and change of study personnel. An additional delay in starting the study was due to changes made in the study design by introducing isoflavone and placebo tablets and making patients with invasive cancer also eligible for entry. The study intervention was changed from soy protein isolate to soy isoflavone tablet, in order to make the study intervention easier to take and to improve the compliance with the study intervention. The change also improved the study design by introducing a placebo arm instead of a no intervention arm. The study design is now better scientifically and it is easier for the patients to accept a placebo controlled study compared to one with no intervention arm. However, these changes resulted in additional delays in starting the study because of resubmission to the IRB.

The study is currently accruing at a slower rate, however 68 patients have been randomized and remaining 4 patients anticipated to be randomized over the next 2-3 months. The no cost extension we received for the study will allow us to finish the accrual and all the biomarker studies over the next year before November 2003.

Below please find the Statement of Work copied from the original application, which outlines the work to be completed over the period of the grant.

#### STATEMENT OF WORK

##### Timeline:



It is estimated that over a period of 45 months, 64 eligible and evaluable patients will be randomized on the study. We will evaluate study compliance by returned remaining pill counts and questioning the subjects with regard to their intake of study or non-study soy supplements. Since the duration of study is only three weeks and soy isoflavone is a non-toxic nutritional supplement, non-compliance with the study intervention should not be a problem in this study. However, to accommodate a non-compliance rate of about 4 patients on each arm we will randomize a total of 72 patients over 45 months, i.e. an accrual rate of 1.6 patient/month.

During the last three months of the study, we will complete the final three-week intervention/non-intervention period and surgery. This will leave approximately 9-10 weeks for completion of the biomarker and micronutrient analyses on the most recent patients and overall biostatistical analyses of the results. Biomarker studies will be performed throughout the grant period between the first month and the 46th month.

## 6. KEY RESEARCH ACCOMPLISHMENTS

As the study is double blind, randomized in design, no findings can be attributed to study intervention until the end of the study period when the code will be broken and the identity of the study compound taken by each patient will be known. This will be accomplished within the next 2-3 months as study accrual has reached 68.

## 7. REPORTABLE OUTCOMES

Because of the study design, no reportable outcomes can be provided until the end of the study period when the code will be broken and the identity of the study compounds will be known.

However, we reported the results of a pilot study with a poster presentation at the 2000 DOD Era of Hope Meeting in Atlanta in June 2000 and 2002 Era of Hope Meeting September 2002. This study was conducted as a part of this application. Please find below the two abstracts presented at the two DOD Meetings.

### **EFFECTS OF SOY ISOFLAVONES ON BIOMARKERS OF OXIDATIVE STRESS AND CELL GROWTH IN PATIENTS WITH BREAST CANCER**

**Omer Kucuk, MD, Fazlul Sarkar, PhD, Zora Djuric, PhD, David Bouwman, MD,  
Mary Ann Kosir, MD, Michael White, MD, Lynn Hryniuk, William Hryniuk, MD**

Ecological studies have shown an inverse association between dietary intake of soy isoflavones and development of breast cancer. The mechanism by which soy isoflavones may prevent breast cancer is not known. Based on our preliminary studies, we hypothesize that soy isoflavones inhibit cell proliferation,

upregulate the expression of gap junctional protein connexin 43 (Cx43) and alter the expression of cell cycle regulatory proteins. Our studies investigate the *in vivo* effects of soy isoflavones on human breast tissues obtained from lumpectomy/mastectomy specimens. We investigate the effect of increased tissue concentration of isoflavones for a period of three weeks on breast cell proliferation, differentiation and cell cycle regulatory proteins. Patients with ductal carcinoma *in situ* (DCIS) or invasive breast cancer are randomly assigned to take 100 mg soy isoflavone (Novasoy™, Archer Daniels Midland Company, Decatur, Illinois) or placebo daily for three weeks prior to surgery. Plasma isoflavone levels are measured at baseline and after three weeks in both groups. Tissue isoflavone levels are measured on samples from benign breast tissues in both groups. Biomarker studies are performed on surgical specimens by immunohistochemistry and Western blot. These studies will enable us to determine if a short duration of exposure to increased tissue levels of isoflavones will modulate biomarkers of cell differentiation (Cx43), adhesion (E-cadherin), proliferation (MIB-1), and cell growth and apoptosis (bcl-2, bax, p53, p21, Rb, EGF-R, cyclin D1, CDK5, CDK6) in benign, pre-malignant and malignant areas of breast epithelial tissues. Biomarker studies on the patients randomized on this study will be completed in early 2003.

We have conducted a pilot study in six women, who took 50 mg of soy isoflavones (Novasoy™) daily for three weeks. DNA was isolated from the nuclei of peripheral blood lymphocytes and analyzed for levels of 5-hydroxy-methyl-2'-deoxyuridine (5-OHmdU) by gas chromatography-mass spectrometry. The mean level of 5-OHmdU was decreased by 35% (relative to baseline) after 1 week and by about 50% after 2 weeks and 3 weeks of supplementation. Mean plasma levels of 8-isoprostanes also decreased somewhat after supplementation.

### **CLINICAL TRIAL OF SOY ISOFLAVONES PRIOR TO BREAST CANCER SURGERY**

**Omer Kucuk, Fazlul Sarkar, Lisa Newman, David Bouwman, Volkan Adsay, Zora Djuric, Daniel Doerge, Ralph Parchment, Mousumi Banerjee, Adhip Majumdar, Lynn Hryniuk, and William Hryniuk**

Our studies investigate the *in vivo* effects of soy isoflavones on human breast tissues obtained from lumpectomy/mastectomy specimens. We investigate the effect of increased tissue concentration of isoflavones for a period of three weeks on breast cell proliferation, differentiation and cell cycle regulatory proteins. Patients with ductal carcinoma *in situ* (DCIS) or invasive breast cancer are randomly assigned to take 100 mg soy isoflavone (Novasoy™, Archer Daniels Midland Company, Decatur, Illinois) or placebo daily for three weeks prior to surgery. Plasma isoflavone levels are measured at baseline and after three weeks in both groups. Tissue isoflavone levels are measured on samples from benign breast tissues in both groups. Biomarker studies are performed on surgical specimens by immunohistochemistry and Western blot. These studies will enable us to determine if a short duration of exposure to increased tissue levels of isoflavones will modulate biomarkers of cell differentiation (Cx43), adhesion (E-cadherin), proliferation (MIB-1), and cell growth and apoptosis (bcl-2, bax, p21, EGFR, cyclin D1, CDK5, CDK6) in benign, pre-malignant and malignant areas of breast epithelial tissues. We will also measure plasma equol, a metabolite of soy isoflavones, and lipid peroxidation products. DNA oxidation products will be measured on peripheral blood lymphocytes. Metabolites of 5-lipoxygenase (5-HETE, LTB4) and cyclooxygenase (PEG2) pathways will also be measured in the plasma. We will also assess ERRP (EGFR Related Protein) levels in tissues by immunohistochemistry and by Western blot. Biomarker studies on the patients randomized on this study will be presented.

We have conducted a pilot study in six women, who took soy isoflavones 50 mg (Novasoy™) daily for three weeks. DNA was isolated from the nuclei of peripheral blood lymphocytes and analyzed for levels of 5-hydroxy-methyl-2'-deoxyuridine (5-OHmdU) by gas chromatography-mass spectrometry. The mean level of 5-OHmdU was decreased by 35% (relative to baseline) after 1 week and by about 50% after 2

weeks and 3 weeks of supplementation. Mean plasma levels of 8-isoprostane also decreased after supplementation.

## 8. CONCLUSIONS

Because of the study design, again, no conclusions can be drawn until the end of the study period when the code will be broken and the identity of the study compounds taken by the patients will be known.

## 9. REFERENCES

Since the study is still accruing patients, no publications can be reported until the end of the study when the code will be broken and the identity of the study compounds taken by the patients will be known. However, we have published the following papers based on data from our pilot study with soy isoflavones.

Davis JN, Kucuk O, Sarkar FH. Effect of soy isoflavone supplementation on NF-kappaB activation in healthy human subjects. *Free Rad. Biol. Med.* 30:1293-1302, 2001.

Djuric Z, Chen G, Doerge D, Heilbrun L, Kucuk O. Effect of soy supplementation on oxidative stress in men and women. *Cancer Lett.* 172:1-6, 2001.

## 10. APPENDICES

No appendices are available for submission at this time.