Preparation and Characterization of the First Binary Titanium Azides, $Ti(N_3)_4$, $[P(C_6H_5)_4][Ti(N_3)_5]$ and $[P(C_6H_5)_4]_2[Ti(N_3)_6]$ and on Linear Ti-N-NN Coordination^{**,†}

Ralf Haiges^{*}, Jerry A. Boatz, Stefan Schneider, Thorsten Schroer, Muhammed Yousufuddin, and Karl O. Christe^{*}

Whereas numerous partially azide-substituted titanium compounds had previously been reported,^[1-7] no binary titanium azides were known. In a recent theoretical study, the group 4 metal tetrazides $M(N_3)_4$ (M = Ti, Zr, Hf, Th) were predicted^[8] to be vibrationally stable, exhibiting tetrahedral structures with unique linear M-N-NN bond angles (see Figure 1). All previously characterized covalent binary azide species possess bent M-N-NN angles.^[9] In this paper, we wish to report the synthesis, isolation and characterization of the first binary titanium azide species Ti(N₃)₄, [Ti(N₃)₅]⁻ and [Ti(N₃)₆]²⁻, and provide explanations for the observed and predicted Ti-N-NN bond angles.

[*] Prof. Dr. K. O. Christe, Dr. R. Haiges, Dr. S. Schneider, Dr. T. Schroer Loker Research Institute, University of Southern California Los Angeles, CA 90089-1661 (USA), Fax: (+1) 213-740-6679 E-mail: <u>kchriste@usc.edu</u>

Prof. Dr. K. O. Christe, Dr. J. A. Boatz ERC, Inc. and Space and Missile Propulsion Division, Air Force Research Laboratory (AFRL/PRSP) 10 East Saturn Boulevard, Bldg 8451 Edwards Air Force Base, CA 93524 (USA)

- [**] This work was funded by the Defense Advanced Research Projects Agency, with additional support from the Air Force Office of Scientific Research and the National Science Foundation. We thank Prof. G. A. Olah, and Drs. A. Morrish, D. Woodbury, and M. Berman, for their steady support, and Prof. R. Bau and Dr. R. Wagner for their help and stimulating discussions.
- [†] Dedicated to Dr. Robert Corley on the occasion of his 70th birthday.

DISTRIBUTION A. Approved for public release; distribution unlimited.

Report Documentation Page				Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.						
1. REPORT DATE 12 MAR 2004		2. REPORT TYPE		3. DATES COVE	RED	
4. TITLE AND SUBTITLE Preparation and C	the First Binary Tit	anium Azides	5a. CONTRACT NUMBER F04611-99-C-0025			
			5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER				
6. AUTHOR(S) Ralf Haiges; Jerry Boatz; Stefan Schneider; Thorsten Schroer;				5d. PROJECT NUMBER DARP		
Muhammed Yousi			5e. TASK NUMBER A205			
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) ERC Incorporated,555 Sparkman Drive,Huntsville,AL,35816-0000				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITC		10. SPONSOR/MONITOR'S ACRONYM(S)				
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT Where as numerous partially azide-substituted titanium compounds had previously been reported,[1-7] no binary titanium azides were known. In a recent theoretical study, the group 4 metal tetrazides M(N3)4 (M = Ti, Zr, Hf, Th) were predicted[8] to be vibrationally stable, exhibiting tetrahedral structures with unique linear M-N-NN bond angles (see Figure 1). All previously characterized covalent binary azide species possess bent M-N-NN angles.[9] In this paper, we wish to report the synthesis, isolation and characterization of the first binary titanium azide species Ti(N3)4, [Ti(N3)5]- and [Ti(N3)6]2-, and provide explanations for the observed and predicted Ti-N-NN bond angles.						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC		17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON		
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified		20		

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std Z39-18 The reaction of TiF_4 with $(CH_3)_3SiN_3$ in acetonitrile solution at room temperature results within minutes in complete fluoride-azide exchange and yields a clear orange solution of $Ti(N_3)_4$ according to Equation (1).

$$TiF_4 + 4 (CH_3)_3 SiN_3 \xrightarrow{CH_3 CN} Ti(N_3)_4 + 4 (CH_3)_3 SiF$$
(1)

Removal of the volatile products (CH₃CN, (CH₃)₃SiF and excess (CH₃)₃SiN₃) at ambient temperature results in the isolation of $Ti(N_3)_4$ as an amorphous orange solid. All attempts to obtain single crystals by recrystallization or sublimation were unsuccessful.

As expected for a highly endothermic, covalent polyazide species, Ti(N₃)₄ is very shock sensitive and can explode violently when touched with a metal spatula or by rapid change in temperature (e.g. freezing with liquid nitrogen). Its identity was established by the observed material balance, and vibrational and NMR spectroscopy. The presence of covalent azido ligands^[10-15] is confirmed by the observed ¹⁴N NMR shifts of d = -134 ppm (N_{β}, $\Delta v_{1/2} = 26$ Hz), -195 ppm (N_{γ}, $\Delta v_{1/2} = 39$ Hz) and -255 ppm (N_{α}, extremely broad) in DMSO solution at 25°C.

The observed Raman and IR spectra of solid $Ti(N_3)_4$ are shown in Figure 2, and the observed frequencies and intensities are listed in the experimental section. The experimental vibrational spectra deviate significantly from those calculated for free $Ti(N_3)_4$ at the B3LYP level of theory,^[8, 16] and resemble those of higher-coordinated compounds with bent M-N-NN bonds. Therefore, the predicted^[8] tetrahedral structure with linear Ti-N-NN bonds could not be confirmed in this study. The discrepancy between the calculated and observed spectra arises from solid state effects. The Ti atoms in $Ti(N_3)_4$ are coordinatively unsaturated seeking higher coordination numbers by the formation of nitrogen bridges, and crystal-structure data will be

required for a reliable determination of the precise arrangement of the azido ligands in solid $Ti(N_3)_4$. The growing of single crystals for such a study will be difficult because the compound is hard to recrystallize and does not sublime without decomposition. The structure determination of free monomeric $Ti(N_3)_4$ and proof for its predicted tetrahedral structure with linear Ti-N-NN bonds will be even more difficult.

The reaction of $Ti(N_3)_4$ with one equivalent of ionic azides leads to the formation of the $[Ti(N_3)_5]^{-1}$ anion according to Equation (2).

$$Ti(N_3)_4 + [P(C_6H_5)_4]N_3 \longrightarrow [P(C_6H_5)_4][Ti(N_3)_5]$$
(2)

The $[P(C_6H_5)_4][Ti(N_3)_5]$ salt was isolated as an orange solid. It is less sensitive than $Ti(N_3)_4$ and does not explode upon freezing with liquid nitrogen. It was characterized by its ¹⁴N NMR spectrum, and by infrared and Raman spectroscopy. The observed Raman and IR spectra of $[P(C_6H_5)_4][Ti(N_3)_5]$ are shown in Figure 3. The experimental infrared and Raman frequencies and intensities are listed in the experimental section. Three well-resolved ¹⁴N NMR resonances were found in DMSO at 25°C. The spectrum shows a sharp signal at d = -133 ppm ($\Delta v_{b_2} = 28$ Hz) for the N_β atoms, a medium-sharp resonance at d = -194 ppm ($\Delta v_{b_2} = 40$ Hz) for the N_γ atoms, and a very broad signal at d = -263 ppm ($\Delta v_{b_2} = 160$ Hz) for the N_α atoms, in accord with our expectations for covalently bound azido groups. The calculated structure for the free gaseous anion is that of a trigonal bipyramid, and its calculated frequencies are listed in the Theoretical Methods section.

By reaction of $Ti(N_3)_4$ with two equivalents of ionic azide, the $[Ti(N_3)_6]^{2-}$ anion is formed according to Equation (3).

$$Ti(N_3)_4 + 2 [P(C_6H_5)_4]N_3 \longrightarrow [P(C_6H_5)_4]_2[Ti(N_3)_6]$$
(3)

 $[P(C_6H_5)_4]_2[Ti(N_3)_6]$ was isolated as an orange solid and is stable at room temperature. Because of the presence of two large counter-ions, it is much less sensitive and explosive than $Ti(N_3)_4$. It can even be heated to its melting point at 191°C without any signs of decomposition. The compound was characterized by its crystal structure,^[17] and vibrational and ¹⁴N NMR spectroscopy. Single crystals were obtained from a solution in CH₃CN.

 $[P(C_6H_5)_4]_2[Ti(N_3)_6]$ crystallizes in the triclinic space group $P\overline{1}$ and is the first structurally characterized binary titanium azide. Figure 4 depicts the structure of the $[Ti(N_3)_6]^{2^-}$ anion that is only slightly distorted from perfect \S symmetry. The structure consists of an asymmetric TiN₉ unit with three azido groups covalently bonded in a trigonal pyramidal fashion to the titanium. The remaining three azide groups are generated by symmetry (symmetry operation – *x*+2,-*y*+1,-*z*). The average Ti-N distance of 2.02 Å is in good agreement with the one reported for (C₅H₅)₂Ti(N₃)₂ (2.03(1) Å).^[6] In the $[P(C_6H_5)_4^+b_2[Ti(N_3)_6]^{2^-}$ salt, the anions are well separated by twice as many large counter-ions, and the closest Ti…N contacts between neighboring anions are 7.1 Å.

Further support for the presence of the $[Ti(N_3)_6]^2$ ion is provided by the NMR spectrum. In analogy with $Te(N_3)_4$ and $[Ti(N_3)_5]^2$, the ¹⁴N NMR spectra in DMSO show at d = -134 ppm $(N_{\beta}, \Delta v_{1/2} = 30 \text{ Hz})$, -199 ppm $(N_{\gamma}, \Delta v_{1/2} = 35 \text{ Hz})$ and -264 ppm $(N_{\alpha}, \Delta v_{1/2} = 165 \text{ Hz})$ resonances characteristic for covalent azides.

The observed Raman and IR spectra of $[P(C_6H_5)_4]_2[Ti(N_3)_6]$ are shown in Figure 5, and the observed frequencies and intensities are listed in the experimental section. Assignments of the observed spectra were made by comparison with those calculated at the B3LYP/SBKJC+(d) level of theory^[16] and are given in the experimental section.

Although the occurrence of linear Ti-N-NN bond angles could not be confirmed experimentally, we have confirmed by independent computations the correctness of the previous predictions^[8] for free gaseous Ti(N₃)₄. Furthermore, it was shown, both computationally and experimentally by our crystal structure determination, that in higher coordinated binary titanium azide species, such as $[Ti(N_3)_6]^{2^-}$, and in the main group tetra-azides $M(N_3)_4$ (M = Si, Ge, Sn) the M-N-NN bond angles are normal and strongly bent. It was also shown that the occurrence of linear M-N-NN bond angles should not be limited to group 4 tetra-azides, but might also occur in compounds, such as $Fe(N_3)_2$.^[18] Normal coordinate analyses were carried out for the binary Ti-azides of this study and showed that the modes due to the azido ligands were highly characteristic, while those of the TiN_n skeletal modes were strongly mixed and, therefore, are not reported in this paper. However, the TiN_n stretching mode clusters clearly exhibit the frequency decreases expected for increasing polarities of the Ti-N bonds with increasing negative formal charges. Thus, the range of the TiN_n stretching modes decreases from 472 - 371 cm⁻¹ in Ti(N₃)₄ to 448 - 355 cm⁻¹ in Ti(N₃)₅⁻ and 398 - 307 cm⁻¹ in [Ti(N₃)₆]²⁻.

In the previous theoretical paper on the linearity of the M-N-NN bond angles for M = Ti, Zr and Hf, the linearities of the M-N-NN angles was explained by M-N-N-N conjugation.^[8] In our opinion, the linearity of the M-N-NN angles is due to a nearly ideal overlap between the three valence electron pairs on the *a*-nitrogen atoms of the azide ligands and the lobes of the *d*orbitals on the central metal atom. Because the *d*-orbitals of the (+IV) central metal atoms are unoccupied, the three valence electron pairs of the *a*-nitrogens can donate electron density equally into three lobes of the empty *d*-orbitals resulting in the *a*-nitrogen acting as a tridentate donor ligand. The symmetry of the central axes of these *d*-orbitals is tetrahedral. Therefore, this tridentate type of overlap is possible only for the tetra-coordinated azides of group 4 transition

metals with d^0 configurations. For other coordination numbers or electron configurations, the *a*nitrogens act usually as monodentate donors with only one pair donating and two sterically active free valence electron pairs, thus resulting in strongly bent M-N-NN bond angles. An excellent example for the validity of our explanation is the known crystal structure of Zr(BH₄)₄ (see Figure 6) involving a (+IV) group 4 d^0 transition metal central atom and 4 trihapto BH₄ ligands.^[19]Another example for a special geometry, dictated by the empty d^0 -orbitals of a (+IV) group 4 transition metal is Ti(O₂ClO₂)₄ in which the perchlorato groups act as bidentate ligands.^[20]

Experimental Section

Caution! Covalent azides are potentially hazardous and can decompose explosively under various conditions! They should be handled only on a scale of less than 2 mmol with appropriate safety precautions (safety shields, safety glasses, face shields, leather gloves, protective clothing, such as leather suits, and ear plugs). Teflon containers should be used, whenever possible, to avoid hazardous fragmentation. Pure $Ti(N_3)_4$ has to be cooled or heated carefully and slowly. A sample at ambient temperature must not be cooled directly with liquid nitrogen. Doing so can result in violent explosions. **Ignoring safety precautions can lead to serious injuries.**

Materials and Apparatus: All reactions were carried out in Teflon-FEP ampules that were closed by stainless steel valves. Volatile materials were handled in a Pyrex glass vacuum line. All Teflon reaction vessels were passivated with ClF₃ prior to use. Nonvolatile materials were handled in the dry argon atmosphere of a glove box.

Raman spectra were recorded at -80 °C in the range 4000–80 cm⁻¹ on a Bruker Equinox 55 FT-RA spectrophotometer using a Nd-YAG laser at 1064 nm with power levels of less than 200 mW. Pyrex melting point tubes that were baked out at 300 °C for 48 h at 10 mTorr vacuum or Teflon-FEP tubes with stainless steel valves that were passivated with ClF₃ were used as sample containers. Infrared spectra were recorded in the range 4000-400 cm⁻¹ on a Midac, M Series, FT-IR spectrometer using KBr or AgCl pellets. The pellets were prepared inside the glove-box using an Econo press (Barnes Engineering Co.).

 14 N NMR spectra were recorded unlocked at 36.13 MHz on a Bruker AMX 500 spectrometer using solutions of the compounds in DMSO in sealed standard glass tubes. Neat CH₃NO₂ (0.00 ppm) was used as the external reference.

The starting materials TiF_4 and $[P(C_6H_5)_4]I$ (both from Aldrich) were used without further purification. $(CH_3)_3SiN_3$ (Aldrich) was purified by fractional condensation prior to use. Solvents were dried by standard methods and freshly distilled prior to use. $[P(C_6H_5)_4]N_3$ was prepared from $[P(C_6H_5)_4]I$ and AgN_3 .

Preparation of $Ti(N_3)_4$: A sample of TiF₄ (0.96 mmol) was loaded into a Teflon-FEP ampule, followed by the addition of 3 mL CH₃CN and (CH₃)₃SiN₃ (4.32 mmol) *in vacuo* at -196 °C. The mixture was allowed to warm to room temperature. Within minutes, the mixture turned yellow, and the color intensified while the reaction proceeded. After 1 hour, all volatile material was pumped off, leaving behind an orange solid (0.20 g, weight calculated for 0.96 mmol Ti(N₃)₄ = 0.21 g). The obtained orange solid was characterized by vibrational and NMR spectroscopy. IR (AgCl): $tt = 2141(vs)/2093(vs)/2051(vs) (v_{as}N_3), 1376(s, br)/1253(s) (v_sN_3),$ $684(s)/668(s)/665(s)/575(m)/561(m) (\delta N_3), 458(w) cm⁻¹ (vTiN_n). Raman (50mW, -80°C):$ $<math>tt = 2160(10.0)/2141(4.0)/2108(2.9)/2100(1.8)/2079(2.8) (v_{as}N_3), 1407(1.0)/1385(0.2)/1366(0.2)/$

1283(0.3)/1269(0.3) (v_8N_3), 671(0.3)/574(0.4) (δN_3), 472(9.6)/454(4.9)/391(2.2)/371(1.8) (vTiN_n), 310(1.7), 270 (1.7), 227 (0.8), 168 (2.1), 135(2.0) cm⁻¹.

Preparation of [PPh₄][Ti(N₃)₅]. A solution of Ti(N₃)₄ (0.5 mmol) in 3 mL CH₃CN was added to a mixture of PPh₄N₃ (0.5 mml) in 2 mL CH₃CN at -64 °C. The reaction mixture was warmed to ambient temperature where it formed a clear orange solution. After 2 hours, all volatiles were slowly removed in a dynamic vacuum at 20 °C, leaving behind an orange solid (0.450 g, weight calculated for 0.5 mmol [PPh₄][Ti(N₃)₅] = 0.443 g). IR of [Ti(N₃)₅]⁻ (KBr): tt=2100(vs)/2070(vs)/2058(vs) ($v_{as}N_3$), 1373(m)/1356(s)/1320(s) (v_sN_3), 621(m)/595(w)/ 591(w)/580(vw) (δN_3), 448(w)/440(vw) cm⁻¹ (vTiN_n). Raman of [Ti(N₃)₅]⁻ (50 mW, -80 °C): tt=2133(10.0)/2110(4.4)/2083(2.9)/2070(2.2) ($v_{as}N_3$), 1367(0.3)/1342(0.3)/1325(0.2)/1314(0.1) (v_sN_3), 623(0.2)/608(0.1)/597(0.1) (δN_3), 445(3.4)/438(1.0)/412(0.6)/398(2.2)/363(0.9)/355(0.7) (vTiN_n), 306(0.9), 245(1.3), 210(0.9), 174(1.6) cm⁻¹.

Preparation of [PPh₄]₂[Ti(N₃)₆]. A solution of Ti(N₃)₄ (0.5 mmol) in 3 mL CH₃CN was added to a mixture of PPh₄N₃ (1.0 mml) in 2 mL CH₃CN at -64 °C. The reaction mixture was warmed to ambient temperature and an orange precipitate was formed. After 8 hours, all volatiles were removed in a dynamic vacuum at 20°C, leaving behind an orange solid (0.768 g, weight calculated for 0.5 mmol [PPh₄]₂[Ti(N₃)₆] = 0.778 g). Single crystals were grown from a solution in CH₃CN by slow evaporation in a dynamic vacuum. IR of [Ti(N₃)₆]²⁻ (KBr): tt =2109(m)/2061(vs)/2041(vs) (v_{as}N₃), 1356(s)/1322(s) (v_sN₃), 622(m)/615(m)/595(w) cm⁻¹ (δN₃). Raman of [Ti(N₃)₆]²⁻ (50 mW, -90 °C) tt =2110(10.0)/2063(1.2)/2038(0.3) (v_{as}N₃), 1366(0.7)/1341(0.7)/1325(0.4) (v_sN₃), 631(0.2)/622(0.3)/608(0.3) (δN₃), 398(5.4)/316(1.1)/307(1.1) (vTiN_n), 246(1.3), 210 (1.8), 103(0.5), 95(0.5) cm⁻¹.

Theoretical Methods. The molecular structures and harmonic vibrational frequencies were calculated at the DFT level using the B3LYP hybrid functional,^[16a] which included the VWN5 correlation functional.^[16b] The SBKJC^[16c] effective core potential and the corresponding valence-only basis set was used for titanium. The all-electron 6-31G basis set,^[16d] augmented with a d polarization function $^{[16e]}$ and a diffuse s+p shell $^{[16f]}$ and denoted as 6-31+G(d), was used for nitrogen. All calculations were performed using the GAMESS^[16g] quantum chemistry program. Unscaled calculated frequencies (cm⁻¹) and (infrared, km/mol) and [Raman, Å⁴/amu] intensities for Ti(N₃)₄: 2293 (0.0) [794], 2251 (5737) [1022], 1497 (0.0) [4.5], 1475 (2005) [48], 573 (110) [0.3], 571 (0.0) [0.3], 569 (0.0) [0.0], 516 (933) [67], 373 (0.0) [103], 192 (17) [7.3], 167 (0.0) [14.0], 14.8 (0.0) [50], 14.0 (0.0) [71]. [Γ i(N₃)₅]⁻ (pseudo-trigonal bipyramid of C₁ symmetry): 2248 (72) [751], 2212 (3406) [42], 2206 (1441) [160], 2202 (1836) [77], 2200 (142) [278], 1445 (48) [28], 1433 (390) [4.3], 1415 (180) [16], 1404 (174) [17], 1402 (384) [6.9], 620 (50) [0.3], 618 (18) [0.8], 615 (57) [0.7], 602 (13) [1.2], 601 (12) [0.1], 599 (10) [0.7], 595 (12) [0.8], 594 (1.0) [0.4], 584 (19) [0.9], 577 (43) [0.4], 487 (325) [1.2], 475 (347) [0.7], 456 (534) [1.8], 401 (1.2) [77], 322 (0.3) [4.2], 263 (2.6) [8.6], 252 (5.3) [7.3], 231 (3.9) [7.2], 228 (0.8) [19], 200 (0.7) [8.1], 131 (2.8) [4.8], 103 (1.1) [11], 94 (0.2) [12], 91 (0.5) [7.1], 66 (0.1) [1.9], 54 (1.1) [4.6], 40 (0.1) [18], 36 (0.1) [22], 33 (0.6) [18], 30 (0.2) [19], 24 (1.1) [2.4], 19 (0.7) [8.1]. $[Ti(N_3)_6]^{2-}$: 2234 (1.4) [798], 2186 (3723) [9.5], 2185 (2778) [90], 2180 (2776) [74], 2168 (25) [180], 2167 (468) [134], 1438 (3.6) [70], 1429 (161) [18], 1428 (19) [46], 1425 (182) [23], 1424 (186) [24], 1424 (165) [17], 626 (4.1) [5.9], 622 (34) [1.2], 621 (39) [2.8], 620 (59) [3.9], 620 (0.0) [2.9], 616 (3.1) [3.9], 608 (5.2) [0.2], 605 (8.8) [0.0], 604 (3.9) [0.1], 599 (9.4) [0.1], 599 (9.9) [0.0], 598 (0.6) [0.4], 409 (675) [0.4], 404 (540) [0.9], 404 (601) [1.2], 364 (1.2) [78], 284 (0.2) [11], 280 (0.1) [8.4], 264 (3.0) [7.0], 261 (5.4) [0.2], 250 (3.5) [6.1], 235 (3.6) [8.7],

209 (0.1) [19], 205 (0.2) [7.4], 151 (1.2) [25], 150 (0.2) [15], 142 (0.4) [25], 78 (1.9) [2.2], 63 (0.1) [11], 62 (2.7) [0.4], 44 (2.2) [20], 40 (0.5) [28], 39 (0.0) [14], 33 (1.0) [28], 28 (2.0) [2.7], 23 (0.1) [13], 21 (1.2) [4.1], 16 (0. 1) [4.5], 15 (1.2) [6.8].

Received: , 2004

Keywords: Crystal structure, Titanium azides, Vibrational spectra, NMR spectra, Theoretical calculations

References

- [1] K. Dehnicke, J. Inorg. Nucl. Chem. 1965, 27, 809.
- [2] V. Gutmann, O. Leitmann, A. Scherhaufer, H. Czuba, *Monats. Chem.* 1967, 98, 188.
- [3] K. Dehnicke, N. Krueger, *Chem.-Zt.* **1982**, *106*, 187.
- [4] R. Choukroun, D. Gervais, J. Chem. Soc. Dalton Trans. 1980, 9, 1800.
- [5] R. S. P. Coutts, P. C. Wailes, Aust. J. Chem. 1971, 24, 1075.
- [6] E. R. de Gil, M. De Burguera, A. Valentina Rivera, P. Maxfield, *Acta Crystallogr. Sect. B* 1977, 33, 578.
- C. J. Carmalt, A. H. Cowley, R. D. Culp, R. A. Jones, Y.-M. Sun, B. Fitts, S. Whaley, H.
 W. Roesky, *Inorg. Chem.* 1997, 36, 3108.
- [8] L. Gagliardi, P. Pyykkö, Inorg. Chem. 2003, 42, 3074.
- [9] For example, a) As(N₃)₃ and Sb(N₃)₃: R. Haiges, A. Vij, J. A. Boatz, S. Schneider, T. Schroer, M. Gerken, K. O. Christe, *Chem. Eur. J.* 2004, *10*, 508; b) [Te(N₃)₆]²⁻: R.

Haiges, J. A. Boatz, A. Vij, M. Gerken, S. Schneider, T. Schroer, K. O. Christe, Angew. *Chem. Int. Ed.* **2003**, *42*, 5847; c) [Te(N₃)₆]²⁻: T. M. Klapötke, B. Krumm, P. Mayer, I. Schwab, Angew. Chem. Int. Ed. 2003, 42, 5843; d) [Si(N₃)₆]²⁻: A. C. Filippou, G. Schnakenburg, J. Am. Chem. Soc. 2002, 124, 12396; e) [Ge(N₃)₆]²⁻: A. C. Filippou, P. Portius, D. U. Neimann, K.-D. Wehrstedt, Angew. Chem. Int. Ed. 2000, 39, 4333; f) [Sn(N₃)₆]²⁻: A. C. Filippou, G. Schnakenburg, *Angew. Che. Int. Ed.* **2000**, *39*, 4333; g) [As(N₃)₆]⁻: T. M. Klapötke, H. Nöth, T. Schütt, M. Warchhold, Angew. Chem. Int. Ed. **2000**, 39, 2108; h) $[Pt(N_3)_4]^{2-}$: B. Neumüller, F. Schmock, S. Schlecht, K. Dehnicke, Z. Anorg. Allg. Chem. 2000, 626, 1792; i) [Cu(N₃)₄]²⁻: W. Hiller, K. Hösler, K. Dehnicke, Z. Anorg. Allg. Chem. **1989**, 574, 7; j) $[Pd(N_3)_4]^{2-}$: W. Beck, T. M. Klapötke, J. Knizek, H. Nöth, T. Schütt, *Eur. J. Inorg. Chem.* **1999**, 523; k) [Au(N₃)₄]²⁻: W. Beck, H. Nöth, *Chem. Ber.* **1984**, *117*, 419; 1) [Zn(N₃)₄]²⁻: G. F. Platzer, H. Krischner, Z. Kristallogr. 1975, 142, 363; A. C. Brunner, H. Krischner Z. Kristallogr. 1975, 142, 24; H. Krischner, O. Baumgartner, H. E. Maier, A. I. Saracoglu, Z. Kristallogr. 1983, 164, 89; F. A. Mautner, H. Krischner, *Monatsh. Chem.* **1990**, *121*, 91; m) $[Mn(N_3)_4]^2$: K. Steiner, W. Willing, U. Müller, K. Dehnicke, Z. Anorg. Allg. Chem. 1987, 555, 7.

- [10] T. M. Klapoetke, B. Krumm, P. Mayer, O. P. Ruscitti, Inorg. Chem. 2000, 39, 5426.
- [11] T. M. Klapoetke, B. Krumm, P. Mayer, H. Piotrowski, O. P. Ruscitti, A. Schiller, *Inorg. Chem.* 2002, 41, 1184.
- [12] W. Fraenk, T. M. Klapoetke in *Inorganic Chemistry Highlights* (Eds.: G. Meyer, D. Naumann, L. Wesemann) Wiley-VCH, Weinheim, Germany, 2002.
- [13] T. M. Klapoetke, B. Krumm, P. Mayer, H. Piotrowski, I. Schwab, M. Vogt, Eur. J. Inorg. Chem. 2002, 2701.

- [14] J. Mason in *Multinuclear NMR* (Ed.: J. Mason) Plenum Press, New York, **1987**.
- [15] S. Berger, S. Braun, H. O. Kalinowski, NMR Spectroscopy of the Non-Metallic Elements, John Wiley & Sons, Chichester, England, 1997.
- [16] a) A. D. Becke, J. Chem. Phys. 1993, 98, 5648; P. J. Stephens, F. J. Devlin, C. F. Chablowski, M. J. Frisch, J. Phys. Chem. 1994, 98, 11623; R. H. Hertwig, W. Koch, Chem. Phys. Lett. 1997, 268, 345). b) S. H. Vosko, L. Wilk, M. Nusair, Can. J. Phys. 1980, 58, 1200. c) W. J. Stevens, H. Basch, M. Krauss, J. Chem. Phys. 1984, 81, 6026; W. J. Stevens, M. Krauss, H. Basch, P. G. Jasien, Can. J. Chem. 1992, 70, 612. d) W. J. Hehre, R. Ditchfield, J. A. Pople, J.Chem.Phys. 1972, 56, 2257. e) P. C. Hariharan, J. A. Pople, Theoret. Chim. Acta 1973, 28, 213. f) T. Clark, J. Chandrasekhar, G. W. Spitznagel, P. von R. Schleyer, J. Comput. Chem. 1983, 4, 294. g) M. W. Schmidt, K. K. Baldridge, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. J. Su, T. L. Windus, M. Dupuis, J. A. Montgomery, J. Comput. Chem. 1993, 14, 1347.

[17] Crystal data for C48H40N18P2Ti: $M_r = 978.82$, triclinic, space group $P\overline{1}$, a = 10.106(11), b = 10.272(16), c = 12.348(13) Å, a = 88.029(18), $\beta = 75.077(15)$, ? = 70.298(18)°, V = 1164(2) Å³, F(000) = 506, ?_{calcd}. (Z = 1) = 1.396 g·cm⁻³, $\mu = 0.310$ mm⁻¹, approximate crystal dimensions 0.65 x 0.52 x 0.19 mm³, ? range = 1.71 to 27.67°, Mo_{Ka} (? = 0.71073 Å), T = 153 K, 7267 measured data (Bruker 3-circle, SMART APEX CCD with ?-axis fixed at 54.74°, using the SMART V 5.625 program, Bruker AXS: Madison, WI, 2001), of which 5057 ($R_{int} = 0.0164$) unique. Lorentz and polarization correction (SAINT V 6.22 program, Bruker AXS: Madison, WI, 2001), absorption correction (SADABS program, Bruker AXS: Madison, WI, 2001). Structure solution by direct methods (SHELXTL 5.10, Bruker AXS: Madison, WI, 2000), full-matrix least-squares refinement on F^2 , data to parameters ratio: 16.2 : 1, final *R* indices [I > 2s(I)] : RI = 0.0404, wR2 = 0.1050, RI = 0.0476, wR2 = 0.1099 (all data), GOF on $F^2 = 1.030$. Further crystallographic details can be obtained from the Cambridge Crystallographic

Data Centre (CCDC, 12 Union Road, Cambridge CB21EZ, UK (Fax: (+44) 1223-336-033; e-mail: <u>deposit@ccdc.cam.ac.uk</u>) on quoting the deposition no. CCDC 232521.

- [18] M. Teichert, J. Boatz, private communication.
- [19] P. H. Bird, M. R. Churchill, *Chem. Commun.* **1967**, 403.
- [20] M. M. Fourati, M. Chaabouni, C. Belin, M. Charbonnel, J. L. Pascal, J. Potier, *Inorg. Chem.* 1986, 25, 1386.

Figure 1. Predicted tetrahedral structure of free gaseous $Ti(N_3)_4$ with linear Ti-N-NN bonds.

•



14







Figure 3. IR and Raman spectra of $[P(C_6H_5)_4][Ti(N_3)_5]$. The bands belonging to the $[Ti(N_3)_5]^-$ anion are marked with asterisks.

Figure 4. ORTEP drawing of the dianionic part of the crystal structure of $[P(C_6H_5)_4]_2[Ti(N_3)_6]$. Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths [Å] and angles [°]: Ti-N1 2.002(2), Ti-N4 2.019(2), Ti-N7 2.048(2), N1-N2 1.173(2), N2-N3 1.138(2), N4-N5 1.200(2), N5-N6 1.146(3), N7-N8 1.197(2), N8-N9 1.140(3), N1-N2-N3 177.5(2), N4-N5-N6 177.6(2), N(7)-N(8)-N(9) 176.9(2), N1-Ti-N4 91.68(9), N1-Ti-N7 90.18(7), N4-Ti-N7 89.73(9), Ti-N1-N2 140.64(14), Ti-N4-N5 126.01(13), Ti-N7-N8 129.67(14)



Figure 5. IR and Raman spectra of $[P(C_6H_5)_4]_2[Ti(N_3)_6]$. The bands belonging to the $[Ti(N_3)_6]^{2-}$ anion are marked with asterisks.



18

Figure 6. Crystal structure of $Zr(BH_4)_4$ demonstrating the perfect geometrical overlap of a tridentate donor into the empty d^0 orbitals of a (+IV) group 4 transition metal atom.



Synopsis

R. Haiges*, J. A. Boatz, S. Schneider, T. Schroer, M. Yousufuddin, K. O. Christe*

Binary Titanium Azides. Preparation and Characterization of $Ti(N_3)_4$, $[P(C_6H_5)_4][Ti(N_3)_5]$ and $[P(C_6H_5)_4]_2[Ti(N_3)_6]$ Theory predicts that $Ti(N_3)_4$ should exhibit unprecedented linear Ti-N-NN bond angles. $Ti(N_3)_4$ and its mono- and di-anions, the first examples of binary titanium azides, were prepared and characterized. They do not possess linear Ti-N-NN angles because their coordination numbers exceed 4. The predicted linearity of the Ti-N-N bonds is explained by the *a*-N atoms of the azide groups acting as tridentate donors into the empty tetrahedral d^0 orbitals of a (+IV) group 4 metal atom.

