

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 25-08-2004		2. REPORT TYPE Final technical		3. DATES COVERED (From - To) Oct 2002-Sept 2003	
4. TITLE AND SUBTITLE Sedimentology of a Morphologically Complex Seafloor Environment, New Jersey Continental Shelf			5a. CONTRACT NUMBER N/A		
			5b. GRANT NUMBER N00014-03-1-0032		
			5c. PROGRAM ELEMENT NUMBER N/A		
			5d. PROJECT NUMBER N/A		
6. AUTHOR(S) Sommerfield, Christopher K.			5e. TASK NUMBER N/A		
			5f. WORK UNIT NUMBER N/A		
			8. PERFORMING ORGANIZATION REPORT NUMBER N/A		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Delaware College of Marine Studies 700 Pilottown Rd. Lewes, DE 19958			10. SPONSOR/MONITOR'S ACRONYM(S) ONR		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Ballston Centre Tower One 800 North Quincy Street Arlington, VA 22217-5660			11. SPONSOR/MONITOR'S REPORT NUMBER(S) N/A		
			12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release: distribution is unlimited		
13. SUPPLEMENTARY NOTES None					
14. ABSTRACT The overarching goal of this research was to advance our understanding of "geologic clutter", acoustic anomalies created by natural sub-seafloor features such as infilled paleochannels, using the continental shelf off New Jersey as a natural laboratory. As part of the ONR Geoclutter initiative, the objectives of this project were as follows: (1) to identify modern Holocene sedimentary processes responsible for the morphology and sedimentological heterogeneity of the seafloor; and (2) to determine the physical basis for seismic discontinuities and associated stratigraphic surfaces in the shallow subbottom. To meet these objectives, coordinated geophysical and geological studies were performed to characterize the sedimentology and stratigraphy at priority Geoclutter sites on the continental shelf. The geological data collected in this study will be used to design signal-processing algorithms that distinguish natural features and man-made targets present at (and just below) the seafloor.					
15. SUBJECT TERMS Continental shelf, marine geology, sedimentation					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON Dr. Christopher K. Sommerfield
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (Include area code) (302) 645-4255

20040915 145

Sedimentology of a Morphologically Complex Seafloor Environment, New Jersey Continental Shelf

Final Report to the Office of Naval Research

Submitted by:

Dr. Christopher K. Sommerfield
College of Marine Studies, University of Delaware
Lewes, DE 19958-1298
Phone: (302) 645-4255 fax: (302) 645-4007 email:cs@udel.edu
Grant #: N00014-03-1-0032

LONG-TERM GOAL

The overarching goal of this research is to advance our understanding of "geologic clutter", acoustic anomalies created by natural sub-seafloor features such as infilled paleochannels, using the continental shelf off New Jersey as a natural laboratory.

OBJECTIVES

As part of the ONR Geoclutter initiative, the objectives of this project are as follows: (1) to identify modern–Holocene sedimentary processes responsible for the morphology and sedimentological heterogeneity of the seafloor; and (2) to determine the physical basis for seismic discontinuities and associated stratigraphic surfaces in the shallow subbottom. To meet these objectives, coordinated geophysical and geological studies have been performed to characterize the sedimentology and stratigraphy at priority Geoclutter sites on the continental shelf.

APPROACH

The research approach involved geological groundtruthing of two sonar datasets previously generated for the Geoclutter study area: (1) multibeam acoustic backscatter derived from Simrad EM1000 (95 kHz) multibeam bathymetry (Goff et al., 1999); and (2) chirp seismic stratigraphy using a system developed by Florida Atlantic University (1.5–15 kHz pulse bandwidth). To correlate acoustic backscatter and seabed sediment type, the sedimentology of short cores collected aboard the RV *Cape Henlopen* in 2001 (cruise CH01–17) was characterized by the PI (UDel Task 1/2). To determine the physical basis for subbottom seismic reflectors at selected Geoclutter sites, AHC-800 drillcores were recovered aboard the RV *Knorr* in 2002 (cruise KN-168) by a team of scientists from DOSECC, University of Texas Institute of Geophysics (UTIG), Skidaway Institute of Oceanography (SKIO), University of Delaware (UDel), and Georgia State University (GSU). Specifically, the PI conducted aboard-ship measurements of sediment physical properties on the drillcores using a GeoTek multi-sensor core logger (UDel Task 2/2).

WORK COMPLETED

Task 1 has been completed, and portion of the data have been incorporated in a manuscript submitted for publication (Goff et al., 2003). Task 2 has also been completed; sediment physical properties data

for three drillcores (4–13 m length) have been analyzed and used to develop synthetic seismograms for the cored sections. Details of the drilling effort are provided in a cruise report authored by C. Alexander and J. Austin. Preliminary results of the ongoing drillcore studies, which will continue during FY03–04, will be presented at the Fall 2003 AGU meeting in a series of papers coauthored by the collaborating investigators (see Publications).

RESULTS

Seabed Studies. In the Geoclutter study area the seabed is a heterogenous mosaic of late Pleistocene fluvio-deltaic muds and medium-grained sands deposited subsequently during the Holocene. The Holocene sand sheet, continuous throughout the greater middle shelf, wedges-out at ~110 m leaving the Pleistocene muds exposed at the seafloor. These muds are over-consolidated (stiff), having been exposed throughout the Holocene. Acoustic backscatter in the area is negatively correlated with seabed porosity, and though this was expected, the broad range of porosities measured (0.35–0.55) was not. The spatial variation in porosity is largely due to the coexistence of sand and stiff mud; sand porosities ranged from 0.55 to 0.45, whereas the stiff muds were less porous at 0.35–0.45 (Goff et al., 2003). Another cause for spatial variability is internal sedimentary structures such as shell deposits and rip-up clasts, evidence of post-depositional erosion and re-sedimentation during the Holocene transgression (Duncan et al., 2000).

Correlation of seismic profiles and lithostratigraphy. Significant progress has been made in relating chirp seismic reflectors and sediment lithology via physical properties analysis of AHC-800 drillcores; seismic discontinuities within the channel fills have been convincingly correlated with sedimentary units at decimeter length scales. One important finding is that pronounced changes in acoustic impedance downsection are associated with depositional *and* erosional surfaces, contacts of strata with disparate sediment types and (or) consolidation states (Figure 1). The majority of these surfaces were created by shallow-marine processes as the channels backfilled during the Holocene transgression, the details of which are being investigated via C-14 geochronology (C. Alexander) and microfossil analysis (B. Christensen). Some of these surfaces span the channel cross-section, whereas others are discontinuous and of variable reflectivity. Intervals of high and extremely variable acoustic impedance are produced by thick beds of shell fragments, material originally transported within the channel environment. Other changes in impedance occur across surfaces separating strata of nearly identical sediment type, yet differing in consolidation state. Such boundaries were most likely produced by a hiatus or erosion in the paleoenvironment. In sum, these results reveal that the small-scale seismic reflectors (decimeters to several meters thick) are produced by highly heterogeneous sedimentary strata with physical characteristics related to the depositional and post-depositional processes.

IMPACT/APPLICATIONS

The goal of the Geoclutter Phase II was to characterize seafloor geology at a number of selected sites through analysis of multibeam and chirp sonar datasets, sediment sampling, and studies of sediment physical properties from drillcores. When modeled by acousticians, the geological data will be used to design signal-processing algorithms that distinguish natural features and man-made targets present at (and just below) the seafloor. In addition, new insights on the nature of infilled paleochannels on the continental shelf have resulted from this work, knowledge with potential applications to shallow-marine environments worldwide.

TRANSITIONS

No specific products have stemmed from this research to date.

RELATED PROJECTS

The geological/geophysical component of the ONR Geoclutter program is composed of several coordinated subprojects, each with a particular emphasis. UTIG is the lead group on multibeam backscatter groundtruthing (Goff), as well as the analysis of chirp stratigraphy (Austin, Fulthorpe, Goff, and Gulick). J. Austin (UTIG) and C. Alexander (SKIO) were the co-chief scientists on the drilling cruise, and D. Nielson (DOSECC) coordinated the drilling operation itself. The PI was responsible for operating the GeoTek logger aboard ship and subsequently for reducing the physical properties data. Detailed sedimentological and chronological studies of the drillcores by C. Alexander and B. Christensen (GSU) are underway.

REFERENCES

- Duncan, C.S., Goff, J.A., Austin, J.A., and Fulthorpe, C.S., 2000. Tracking the last sea-level cycle: seafloor morphology and shallow stratigraphy of the latest Quaternary New Jersey middle continental shelf. *Marine Geology*, 170: 395–421.
- Goff, J.A., Swift, D.J.P., Duncan, C.S., Mayer, L.A., and Hughes-Clarke, J., 1999. High-resolution swath sonar investigation of sand ridge, dune and ribbon morphology in the offshore environment of the New Jersey margin. *Marine Geology*, 161: 307–337.

PUBLICATIONS

- Alexander, C.R., Sommerfield, C.K., Austin, J.A., Christensen B., Fulthorpe, C.S., Goff, J.A., Gulick, S.P.S., Nordfjord, S., Nielson, D.L., and Schock, S. (2003). Sedimentology and age control of latest Quaternary New Jersey shelf deposits. *EOS, Transactions of the American Geophysical Union*.
- Austin, J., Goff, J., Gulick, S., Fulthorpe, C., Nordfjord, S., Wiederspahn, M., Sastrup, S., Schock, S., Wulf, J., Gjerding, Mayer, L., and Sommerfield, C. (2001). Assessing the "Geo" in GeoClutter: New Chirp sonar, sampling, and compressional wave velocity results from the New Jersey Shelf. *EOS, Transactions of the American Geophysical Union*, 82 (47), F655.
- Gulick, S.P.S., Fulthorpe, C.S., Goff, J.A., Austin, J.A., Nordfjord S, Sommerfield, C.K., Alexander, C.R., Christensen, B., Schock, S., Nielson, D.L. (2003). Mapping a pre-Last Glacial Maximum paleo-seafloor and shelf-slope sediment wedges beneath the New Jersey shelf. *EOS, Transactions of the American Geophysical Union*.
- Nordfjord, S., Goff, J.A., Austin, J.A., Fulthorpe, C.S., Gulick, S.P.S., Sommerfield, C.K., Alexander, C.R., Christensen, B., Schock, S., and Nielson, D.L. (2003). Geomorphologic comparisons of shallowly buried, dendritic drainage systems imaged on the outer New Jersey shelf with modern records of both fluvial and tidal influences. *EOS, Transactions of the American Geophysical Union*.
- Goff, J.A., Kraft, B.J., Mayer, L.S., Schock, S.G., Sommerfield, C.K., Olson, H.C., Gulick, S.P., and Nordfjord, S., 2003. Seabed characterization on the New Jersey middle and outer shelf: Correlability and spatial variability of seafloor sediment properties. Submitted manuscript.




Figure 1. Example of correlated seismic and sediment physical properties data for Geoclutter drillcore Site 2. (Top) Chirp profile showing a buried channel with internal seismic reflectors of variable amplitude. The 13-m section cored during cruise KN-168 is shown (black rectangle). (Bottom) Downcore physical properties and photograph for the core section corresponding to a prominent seismic discontinuity near the channel base (white rectangle). The discontinuity is created by a contrast in acoustic impedance (density x velocity) across a stratal surface formed by the contact of shell-gravel-sand (high impedance) and clayey silt (low impedance). Seismic profile courtesy of UTIG.