

Woods Hole Oceanographic Institution



A High-Resolution Bathymetry Map for the Marguerite Bay and Adjacent West Antarctic Peninsula Shelf for the Southern Ocean GLOBEC Program

by

S. T. Bolmer¹
R. C. Beardsley¹
C. Pudsey²
P. Morris²
P. Wiebe¹
E. Hofmann³
J. Anderson⁴
A. Maldonado⁵

¹Woods Hole Oceanographic Institution, Woods Hole, MA 02543, U.S.A.

²British Antarctic Survey, Cambridge, CB3 0ET, U.K.

³Old Dominion University, Norfolk, VA 23529, U.S.A.

⁴Rice University, Houston, TX 77251, U.S.A.

⁵Universidad de Granada Facultad de Ciencias, 18002, Granada, Spain

May 2004

Technical Report

Funding was provided by the Office of Naval Research under Contract No. N00014-99-1-0213.

Approved for public release; distribution unlimited.

WHOI-2004-02

**A High-Resolution Bathymetry Map for the Marguerite Bay
and Adjacent West Antarctic Peninsula Shelf
for the Southern Ocean GLOBEC Program**

by

S. T. Bolmer¹
R. C. Beardsley¹
C. Pudsey²
P. Morris²
P. Wiebe¹
E. Hofmann³
J. Anderson⁴
A. Maldonado⁵

¹Woods Hole Oceanographic Institution, Woods Hole, MA 02543, U.S.A.

²British Antarctic Survey, Cambridge, CB3 0ET, U.K.

³Old Dominion University, Norfolk, VA 23529, U.S.A.

⁴Rice University, Houston, TX 77251, U.S.A.

⁵Universidad de Granada Facultad de Ciencias, 18002, Granada, Spain

May 2004


Technical Report

Funding was provided by the National Science Foundation under Grant Number OPP-0234163.

Reproduction in whole or in part is permitted for any purpose of the United States Government. This report should be cited as Woods Hole Oceanog. Inst. Tech. Rept., WHOI-2004-02.

Approved for public release; distribution unlimited.

Approved for Distribution:



Nelson G. Hogg, Chair

Department of Physical Oceanography

SO GLOBEC Bathymetry
WHOI-2004-02

Abstract

One objective of the U.S. Southern Ocean Global Ocean Ecosystems Dynamics (SO GLOBEC) program is to gain a better understanding of the sea floor bathymetry in the program study area. Much of Marguerite Bay and the adjacent shelf west of the Antarctic Peninsula were poorly charted when the SO GLOBEC program started in 2000. Before the first SO GLOBEC cruise, an improved local area version (ETOPO8.2A) was created from the Smith and Sandwell (1997) topo_8.2.img 2-minute digital gridded bathymetry for the study area. The first SO GLOBEC mooring cruise on the *R/V Lawrence M. Gould* (March 2001) showed that the 2-minute spatial resolution of ETOPO8.2A did not resolve many of the canyons and abrupt changes in topography that characterize Marguerite Bay and the inner- to mid-shelf region. It also was not particularly accurate in the more uniform terrain regions. We then decided to collect as much multibeam bathymetry data as possible during the SO GLOBEC broad-scale survey cruises on the *R/VIB Nathaniel B. Palmer* and combine these data with all other available multibeam and trackline bathymetry data to construct a digital bathymetry database and map for the study area. The resulting database has high-resolution data over much of the shelf and parts of Marguerite Bay gridded at 2 seconds in latitude and 6 seconds in longitude spacing between 65° to 71° S and 65° to 78° W. This technical report describes the steps taken to assemble and construct this database and how to access the data via the Internet.

SO GLOBEC Bathymetry
WHOI-2004-02

Table of Contents

<i>Abstract</i>	<i>1</i>
<i>Table of Contents</i>	<i>2</i>
<i>Table of Figures</i>	<i>4</i>
2 Data Sources and Types	7
2.1 ETOPO8.2A	7
Figure 1. Smith & Sandwell topo_8.2.img data.	8
Figure 2. ETOPO8.2A.	9
Figure 3. Comparison of depth data from recent cruise tracks and old navigation charts.	10
2.2 GEBCO	11
Figure 4. GEBCO 2003 1-minute map for SO GLOBEC study area.	12
Figure 5. GEBCO 2003 depth contours.	13
Figure 6. GEBCO 2003 and SO GLOBEC contours.	14
2.3 SO GLOBEC	15
2.4 Other United States Antarctic Program Bathymetry Data	15
Figure 7. Tracks of SO GLOBEC cruises that collected digital bathymetry data.	16
Figure 8. Tracks of other (non-GLOBEC) USAP cruises that collected bathymetry.	17
Figure 9. USAP <i>R/SV Laurence M. Gould</i> JGOFS Bathymetry used in this compilation.	18
2.5 National Geophysical Data Center	19
2.6 British Antarctic Survey	19
2.7 Spanish Data	20
2.8 Lamont-Doherty Antarctic Multibeam Bathymetric Synthesis Database	20
2.9 Other Contributions	21
Figure 10. Tracks of all other cruises in NGDC that collected bathymetry data used in this study.	22
Figure 11. Map of BAS, Spanish, and Russian multibeam bathymetry data contributions.	23
3 Analysis Methods	24
3.1 Multibeam Data	25
3.1.1 RVIB Nathaniel B. Palmer SeaBeam	25
3.1.2 NGDC Multibeam	25
3.1.3 British Antarctic Survey Multibeam	25
3.1.3.1 ASCII XYZ multibeam data	25
3.1.3.2 RAW Multibeam data	26
3.2 Conventional echo sounding data	26
3.2.1 NGDC single point track line	26
3.2.2 R/V Laurence M. Gould along-track data	27
3.3 Coastline Data	27
3.4 DEM Topography	28
Figure 12. All the track coverage with DEM GTOPO30 and SCAR ADD data.	29

SO GLOBEC Bathymetry
WHOI-2004-02

4 Steps in Getting to a Final Grid	30
Figure 13. Final SO GLOBEC bathymetric map.	32
Figure 14. Final SO GLOBEC topographic map with elevation coloring.	33
Figure 15. Final SO GLOBEC data, contour interval 100 meters.	34
5 Data Distribution	35
6 Summary	36
7 Suggestions for improving the compilation	37
8 A Brief Time Line of the Compilation	39
9 Acknowledgements	41
10 References	42
TABLE 1:	43
Navigational Charts used to augment ETOPO2-v8.2	43
TABLE 2:	44
Cruises Used to Create the Marguerite Bay Region Bathymetry Chart. 02/01/04	44
TABLE 3:	46
LAND DATA used to Create the Marguerite Bay Region Bathymetry Chart.	46
TABLE 4:	47
Programs used in processing the data	47
APPENDIX 1: MB-System and GMT scripts	48
Script to grid the Nathaniel Palmer Sea Beam data	48
Script to grid the Other Multibeam Data	49
Script to grid the ASCII along track data	50
Script to merge Land, Gridded at sea data and the SCAR Coastline	51
Script to find widely divergent points	54
MatLab Scripts to edit JGOFS data	57
JGOFSglobal_start.m	57
Figure 16. Main window in JGOFSplot MATLAB GUI tool.	58
Figure 17. JGOFSplot MATLAB GUI tool zoom window.	59
getJGOFS.m	60
JGOFS2mat.m	61
JGOFSdisplay.m	68
JGOFSgros.m	77

SO GLOBEC Bathymetry
WHOI-2004-02

Table of Figures

<i>Figure 1. Smith & Sandwell topo_8.2.img data.</i>	8
<i>Figure 2. ETOPO8.2A.</i>	9
<i>Figure 3. Comparison of depth data from recent cruise tracks and old navigation charts.</i>	10
<i>Figure 4. GEBCO 2003 1-minute map for SO GLOBEC study area.</i>	12
<i>Figure 5. GEBCO 2003 depth contours.</i>	13
<i>Figure 6. GEBCO 2003 and SO GLOBEC contours.</i>	14
<i>Figure 7. Tracks of SO GLOBEC cruises that collected digital bathymetry data.</i>	16
<i>Figure 8. Tracks of other (non-GLOBEC) USAP cruises that collected bathymetry.</i>	17
<i>Figure 9. USAP R/SV Laurence M. Gould JGOFS Bathymetry used in this compilation.</i>	18
<i>Figure 10. Tracks of all other cruises in NGDC that collected bathymetry data used in this study.</i>	22
<i>Figure 11. Map of BAS, Spanish, and Russian multibeam bathymetry data contributions.</i>	23
<i>Figure 12. All the track coverage with DEM GTOPO30 and SCAR ADD data.</i>	29
<i>Figure 13. Final SO GLOBEC bathymetric map.</i>	32
<i>Figure 14. Final SO GLOBEC topographic map with elevation coloring.</i>	33
<i>Figure 15. Final SO GLOBEC data, contour interval 100 meters.</i>	34
<i>Figure 16. Main window in JGOFSplot MATLAB GUI tool.</i>	58
<i>Figure 17. JGOFSplot MATLAB GUI tool zoom window.</i>	59

1 Introduction

During the initial planning for the U.S. Southern Ocean (SO) GLOBEC field program, it became very clear that the program would need a better knowledge of the sea floor bathymetry in the program study area if the program was to achieve its' scientific objectives. Much of Marguerite Bay and the adjacent western Antarctic Peninsula (WAP) shelf were poorly charted, and the coverage of high-quality digital sounding data with GPS-quality navigation data was sparse. Examination of existing maps and along-track data in the U.S. National Geophysical Data Center (NGDC) showed the WAP shelf to be rugged. The area has a mean depth of roughly 500 m but with many canyons and depressions to depths of 800 m interspersed with shallow banks. Marguerite Bay has a more complex bathymetry with George VI Sound having a maximum depth of 1600 m, along with extensive shallows and island groups. Such topographic variability should strongly influence local currents, water properties, and regional circulation, plus biological processes, including animal distributions and behavior (e.g., seal dive depth). Thus, a good knowledge of the bottom topography is essential for both physical and biological sampling and experimental design, data interpretation and physical/biological modeling.

It was evident by the December 2000 SO GLOBEC planning meeting that the program needed to obtain as much new high-quality bathymetry data as possible during the field program if significant improvements to the bathymetry data in the study area were to be achieved. Thus, it was decided to collect high-quality swath bathymetry data during all *R/VIB Nathaniel B Palmer (NBP)* cruises in the SO GLOBEC study area during 2001-2002. We would merge these data with along-track data collected on the *R/V Laurence M. Gould (LMG)* and all other multibeam and centerline data that we could obtain. These data would be gridded into a bathymetry database that could be used in planning and scientific analysis by program investigators. Peter Wiebe, Chief Scientist of the *NBP* 2001 and 2002 austral "fall" and "winter" SO GLOBEC survey cruises, requested that the swath data be collected, and Bob Beardsley agreed to organize and oversee the post-cruise processing and construction of a composite digital bathymetry database for the program.

As a first step towards constructing a better bathymetry we used a digitizing table to determine the latitude and longitude of depth values on

SO GLOBEC Bathymetry
WHOI-2004-02

existing navigational charts and merged these data with the Smith and Sandwell (1997) topo_8.2.img 2-min digital gridded bathymetry to create the ETOPO8.2A data set. Comparisons with along-track data collected during the March 2001 *LMG* mooring cruise showed that the 2-min resolution of the ETOPO8.2A data set did not resolve many of the canyons and abrupt changes in topography which characterize Marguerite Bay and the inner- to mid-shelf region, nor was it particularly accurate. At several mooring sites, the pre-deployment site depth survey made with the *LMG* fathometer bore little resemblance with ETOPO8.2A. This strengthened our desire to collect as much new high-resolution digital bathymetry data as possible in the SO GLOBEC study area during the program field work.

The work of building a composite digital bathymetry database started in earnest in the second half of 2001. High-quality swath bathymetry data were collected on the austral "fall" and "winter" SO GLOBEC broad-scale surveys using the *NBP* SeaBeam system, and along-track digital bathymetry data from the *LMG* "fall" and "winter" SO GLOBEC process-oriented cruises. In addition, an extensive multibeam survey was conducted along the deep trough exiting from Marguerite Bay onto the shelf aboard the *RRS James Clark Ross* in 2001 and made available to us by the British Antarctic Survey (BAS). Center trackline and multibeam data already available at the U.S. Geophysical Data Center (NGDC) also were obtained and incorporated into the map. A first attempt to merge these new data sets with ETOPO8.2A exhibited so much variation with the actual data collected in the study region that we decided to exclude ETOPO8.2A data in future compilations.

Additional multibeam and center trackline data were collected during the 2002 SO GLOBEC "fall" and "winter" *NBP* broad-scale survey cruises and *LMG* process-oriented cruises and mooring cruise. Other bathymetric data were from British and Russian sources also were added to the data set. We stopped actively looking for additional data, following the last SO GLOBEC *LMG* mooring cruise in March 2003, and focused on assembling a final quality-controlled digital bathymetry data set for the SO GLOBEC study region. This new compilation could then be used by interested investigators to produce high-resolution gridded bathymetry maps. This report describes the steps taken to assemble and edit the individual data sets collected in this effort, and the final data set and how to access this data via the Internet. Figure 12 shows the coverage in the area we have assembled from cruise data.

2 Data Sources and Types

2.1 ETOPO8.2A

As a first step towards building an improved bathymetry database before any SO GLOBEC cruises in the area, Bob Beardsley and Mike Caruso obtained the Sandwell and Smith topo_8.2.img 2-min digital gridded bathymetry for the SO GLOBEC study area (Figure 1, see <http://www.ngdc.noaa.gov/mgg/bathymetry/predicted/explore.HTML> for more information about ETOPO2). This bathymetry data set is a merger of satellite-derived data and digital along-track and swath data that had been obtained from the NGDC (see <http://www.ngdc.noaa.gov/mgg/geodas/trackline.html>) as of July 15, 2000.

The positions of individual soundings on four published navigational charts (Table 1) in the study area were determined using an XY digitizing table. The resulting digital XYZ data were combined with the above topo_8.2.img data using the GMT program surface. The resulting digital bathymetry data ETOPO8.2A (Figure 2) showed more spatial structure around Adelaide Island and within Marguerite Bay as a result of the addition of the navigational chart data which are not part of the NGDC.

However, as mentioned above, the difference between ETOPO8.2A and depth data collected on the first *LMG* mooring cruise in March 2001 was so large (Figure 3) that this initial merged product was eventually discarded later that year. We decided not to include the digitized navigational chart data nor the satellite-derived data in ETOPO8.2A in our final data set presented here. The navigational chart data were collected in times when navigation and depth sounding were very crude relative to today's methods.

Smith & Sandwell ETOPO2 Version 8.2




Figure 1. Smith & Sandwell topo_8.2.img data. This is a 2-min gridded bathymetry for the SO GLOBEC study area. Note how the shelf break is interpreted in this data set. The Generic Mapping Tools (GMT) coastline is in green. The ADD edge of the ice is shown in blue.

*Sandwell & Smith Version 8.2
plus Digitized Charts*




Figure 2. ETOPO8.2A.

The increased bumpiness reflects the addition of the navigational chart data before re-gridding. The ADD coastline is in black and ADD edge of the ice is in blue.

Latest Gridded Track Data versus Digitized Chart Data




Figure 3. Comparison of depth data from recent cruise tracks and old navigation charts.

The digitized chart data are plotted as pluses (+) overlying track data. Note the large number of red pluses over areas of yellow and green showing the large differences from the two depth data sets. The black area is deeper than the coloring scheme used. This was left to illustrate the differences in the data sets.

2.2 GEBCO

In 2003, the General Bathymetric Chart of the Oceans (GEBCO) published under the auspices of the International Hydrographic Organization (IHO) and the United Nations' (UNESCO) Intergovernmental Oceanographic Commission (IOC) a 1-min bathymetric data set for the world ocean (see <http://www.ngdc.noaa.gov/mgg/gebco/gebco.html>). This data set was compiled from a large collection of spot soundings, swath data and satellite-derived data. The compiled data were then plotted and contoured by hand with the resulting contours digitized and gridded to form the final 1-min data set.

The GEBCO interpretation (Figure 4) does not fit well with the multibeam data we have collected in the SO GLOBEC study area. An over-plot of GEBCO contours on a shaded relief map from our data (Figure 5) shows general agreement in some areas and radical differences in others. This is emphasized by the plot of GEBCO contours on our contours (Figure 6), especially over the outer shelf and shelfbreak region. There are also notable differences in coastline and ice shelf edge positions. For this reason, we decided not to use the GEBCO 1-min data. (A discussion group email from Walter Smith about the strengths and weaknesses of the GEBCO 1-min data set can be found at: <http://op.gfz-potsdam.de/GMT-Help/Archive/msg09701.html>. This note helps explain our omission of the GEBCO bathymetry.)

GEBCO 2003




Figure 4. GEBCO 2003 1-minute map for SO GLOBEC study area.

GEBCO 2003 Contours over SO GLOBEC Data




Figure 5. GEBCO 2003 depth contours. The depth contours plotted in black over the shaded relief plot of the SO GLOBEC data set. The contour interval here is 500 meters.

GEBCO 2003 compared with SO GLOBEC Data




Figure 6. GEBCO 2003 and SO GLOBEC contours. The GEBCO 2003 and SO GLOBEC contours plotted in blue and red respectively. The ADD coastline is plotted in black and the GEBCO coastline in green. The contour interval here is 500 meters.

2.3 SO GLOBEC

SeaBeam swath bathymetry data were collected on the *R/VIB Nathaniel B. Palmer (NBP)* 2001 “fall” and “winter” and 2002 “fall” broad-scale survey cruises. For the *NBP* 2002 “winter” cruise there was no operating multibeam system aboard. Thus, only digital centerline echo sounder data were collected and recorded in JGOFS format with the other underway along-track data. The *NBP* multibeam data were edited at sea during each cruise and are now in good condition. The *NBP* JGOFS formatted bathymetry data needed extensive post-cruise editing since only limited editing was done at sea. The multibeam data had the sound velocity profiles used in the data processing adjusted at sea. The multibeam soundings data were consistent between each cruise and with the sound velocity profiles agree with data collected from CTD casts taken during the cruise.

On all seven SO GLOBEC *R/V Laurence M. Gould (LMG)* cruises, digital centerline echo sounder data were recorded in JGOFS format along with other underway along-track data. This JGOFS bathymetry data also needed extensive cleaning as no editing was done at sea. The tracks and swaths where the *NBP* and *LMG* collected bathymetry data during the SO GLOBEC field program are shown in Figure 7. The documentation of the sound velocity profile used in collecting the JGOFS data is very poor to non-existent. The good agreement in crossings with the *NBP* multibeam data showed that this data was useable in our compilation.

2.4 Other United States Antarctic Program Bathymetry Data

Multibeam and centerline bathymetry data collected on other U.S. Antarctic Program (USAP) *NBP* and *LMG* cruises in the study area have been obtained for inclusion in our compilation (Figure 8). John Anderson (Rice University) has contributed SeaBeam data from his 1999 *NBP* geophysics cruise (*NBP* 99-02) and selected segments of SeaBeam data from his 2002 *NBP* geophysics cruise (*NBP* 02-01) in Marguerite Bay. Both of these cruises cover areas that needed coverage and the swath data have been included in the compilation. Centerline data collected on many of the USAP *LMG* cruises (Figure 9) in the study area from 2000 to early 2003 also have been obtained and included into the compilation (Table 2). As with the other centerline data collected in JGOFS format, the *LMG*

SO GLOBEC Bathymetry
WHOI-2004-02

data have required extensive editing to produce accurate depth data (see Appendix 2).

SO GLOBEC Collected Data used in compilation




Figure 7. Tracks of SO GLOBEC cruises that collected digital bathymetry data.

Also shown in gray are tracks of other cruises that contributed data to the final data set.

Other USAP Data used in compilation




Figure 8. Tracks of other (non-GLOBEC) USAP cruises that collected bathymetry.

USAP Laurence M. Gould JGOFS Data Tracks




Figure 9. USAP R/SV Laurence M. Gould JGOFS Bathymetry used in this compilation.

2.5 National Geophysical Data Center

A search of the U.S. National Geophysical Data Center (NGDC) for bathymetric data in our region of interest identified the cruises listed in Table 2 (Figure 10). The majority of these cruises (43 of 46) only had data from 3.5 or 12 kHz centerline recording systems. The bathymetry data have been checked to remove any spurious depths. Since these data were generally not rigorously edited before submission to NGDC and may have been taken with a variety of sound velocities in the water used to calculate depth, a thorough editing and comparison with other data at track crossovers is essential. NGDC has a system of reporting these velocities, but often this seems to be not reported or forgotten when those who collected the data returned from sea.

In addition to the centerline data, we were fortunate to find three multibeam data sets at NGDC. These were of a mixed quality. The data from one Lamont-Doherty Earth Observatory *R/V Maurice Ewing* cruise has the overall crude bathymetry in the right range but the actual data points need further editing. These data are in areas that fills gaps in our coverage but this cruise data must be used with care. Two multi-beam cruises attributed to NOAA seem to have data that fits in well and looks fairly cleanly edited.

2.6 British Antarctic Survey

We were fortunate to make contact with Carol Pudsey and Peter Morris at the British Antarctic Survey (BAS) near the beginning of the SO GLOBEC field program. They have provided us with extensive multibeam data from their 2001 and 2002 *RRS James Clark Ross (JCR)* geophysics cruises in Marguerite Bay (Figure 11). These cruises mapped large parts of Marguerite Trough, the channel that leads from to the shelf break from the mouth of Marguerite Bay and George VI Sound. Since the Marguerite Trough is such a dominant topographic feature in the region, the high-quality multibeam data from these BAS surveys have been crucial additions to our data set.

During the summer of 2003, Carol and Peter provided us with raw multibeam data from several other *JCR* cruises in the SO GLOBEC study area. These data had not been edited beforehand and after editing by Tom

SO GLOBEC Bathymetry WHOI-2004-02

Bolmer have been added to our data set. These surveys have added many new and useful miles of new data to this study. In the winter of 2004 Rob Lartner sent us trackline data from the JCR that go along the shelf and the rise.

Peter also provided edited multibeam data collected on a 1991 geophysics cruise aboard the Russian ship *R/V Akademik Boris Petrov*. This cruise included several tracks along the shelf that parallel themselves and cross the SO GLOBEC tracks at right angles, thus providing very useful tie lines. Some of the Russian data were discarded in the deeper water areas off the shelf, where they did not agree with our other data there. The data on the shelf areas is not without its' problems. But, since these data cover areas that need coverage it has been left in. We have done some further editing of these data in shallower areas. The editing has been cumbersome since the data are only in an ASCII XYZ format and thus it is hard to remove points from easily.

Rob Larter at the BAS provided MGD77 formatted data for 2 tracks from the *JCR* cruise JR04 in 1993 that follow the shelf break on the shelf and out on the continental rise (Figure 11).

2.7 Spanish Data

Andrés Maldonado (Instituto Andaluz de Ciencias de la Tierra, Universidad de Granada, Spain) contributed the gridded multibeam bathymetry data set from the ANTPAC97-98 geophysics cruise on the *R/V Bio-Hesperides*. This data set improved the density of coverage at the shelf break in one area (Figure 11).

2.8 Lamont-Doherty Antarctic Multibeam Bathymetric Synthesis Database

The Lamont-Doherty Earth Observatory Antarctic Multibeam Bathymetric Synthesis Database web site (<http://data.ldeo.columbia.edu/antarctic/>) provided us with several USAP data sets that were not available elsewhere. These cruises have helped fill in some areas in the north east regions that had few cruises. As of this date this site has two large multibeam data sets that have not been

SO GLOBEC Bathymetry
WHOI-2004-02

released. We are using the underway JGOFS bathymetry for both of these, but we do have a partial set of the multibeam data released to us for one of the cruises. These two cruises should be available in early 2004 after the 2 years proprietary time for these data is up. At the time of this report one of the chief scientists for these cruises has put an additional 6 month hold on releasing the data from his cruise.

2.9 Other Contributions

Laurie Padman (Earth and Space Research, Seattle) contributed the bathymetry data used in his tidal models for the study region. In particular, his data set has a 1/4 degree longitude by 1/12 degree latitude grid with coverage under the Wilkins Ice Sheet and George VI Sound. While this is very coarse spacing, it does help constrain the land and other bathymetry in our study area. Laurie based his data base on an Oregon State University global tidal solution based on the ETOPO-5 data. The web-site describing this data is at: <http://www.ers.org/antarctic/barotropic.html#CAT> .

From Maslanyj (1987) we used 3 profiles across George IV Sound. These data were collected using seismic techniques. There are only three profiles that we can use in the area but they show us that George IV Fjord is a channel around Alexander Island. These profiles can be seen in Figure 12.

Other NGDC Data used in compilation




Figure 10. Tracks of all other cruises in NGDC that collected bathymetry data used in this study.

SO GLOBEC Bathymetry
WHOI-2004-02

BAS and Spanish Data used in compilation




Figure 11. Map of BAS, Spanish, and Russian multibeam bathymetry data contributions.

The red and pink like BAS data and the blue Spanish data were sent as ASCII XYZ data and those points are plotted here instead of a cruise track.

3 Analysis Methods

We have relied on publicly available programs to create the final bathymetry data set presented here. In particular, we have used the MB-System software written by David Caress (Monterey Bay Aquarium Research Institute) and Dale Chayes (Lamont-Doherty Earth Observatory). This software package was developed with NSF support and is described at <http://www.ldeo.columbia.edu/MB-System/MB-System.intro.html>. This software allows editing and converting the multibeam data into latitude-longitude gridded data that we can then manipulate and merge with other data.

We also have used the Generic Mapping Tools (GMT) written by Paul Wessel (School of Ocean and Earth Science and Technology, University of Hawaii) and Walter Smith (NOAA Laboratory for Satellite Altimetry) (see <http://gmt.soest.hawaii.edu>). These programs were used to merge and do the rest of the gridding and plotting of the data. See Appendix I for examples of the scripts we had not previously noticed.

In the process of sorting, editing, and managing the various data sets, we have also developed PERL and MatLab programs as needed. We also have used the sh and csh shells, the generic shell scripting inherent in UNIX, to manage the data and run the processing steps we used. All final processing was done on a Sun Ultra 60 UNIX workstation running Solaris 8.0 with nearly 150 GB hard disk drive storage. Brief descriptions of the processing done on various data sets are presented next.

We were fortunate to recently have the Interactive Visualization Software company's Fledermaus program available to us. This program was able to read in the gridded NETCDF data from GMT and plot the data in Three Dimensions (3D). By being able to rotate and fly through the 3D figure we were able to find large highs and lows that were previously missed. We then went back to the raw data and removed the extraneous points and ran through the whole process to make a gridded data set again. By using this 3D and editing process several times were able to remove the abnormalities. There are some examples of the Fledermaus figures on the data web site referred to below.

3.1 Multibeam Data

Multibeam systems considered here provide up to 150 and more depth points on a swath (transect) perpendicular to the ship's track. The width of the swath is typically 1.5 times the water depth, so this swath measurement approach is much more effective in mapping areas than the along-track single-point echo sounder. The raw multibeam data requires careful visual editing of individual ping data, a major labor-intensive effort. We have used the MB-System mbedit program for editing raw multibeam as needed, and the MB-System mbgrid program to convert the edited multibeam swath data into equally-spaced latitude, longitude, depth (xyz) values.

3.1.1 RVIB Nathaniel B. Palmer SeaBeam

The multibeam data collected on the *R/VIB Nathaniel B. Palmer* were edited by the scientific party on board the ship at the time of collection. The efforts of the scientific party were coordinated and quality controlled by the Raytheon Polar Services shipboard technicians. In addition to overseeing the basic swath editing, the technicians also checked the navigation data input and made corrections as needed. This thankless job was done on all cruises and has made this digital data base construction possible.

3.1.2 NGDC Multibeam

NGDC had three multibeam data sets collected in the study area. At this time, we have used these data without editing, although we recommend further editing of these data sets.

3.1.3 British Antarctic Survey Multibeam

3.1.3.1 ASCII XYZ multibeam data

The British Antarctic Survey has supplied both edited processed and raw multibeam data. The processed data were in ASCII xyz format. These data sets have been used without change. They do exhibit some spatial distortion due to the space scale used in the gridding. If possible, we will attempt to obtain the original edited multibeam files to include in the final database.

3.1.3.2 RAW Multibeam data

The BAS raw multibeam data had been collected with a Simrad 120 multibeam system that used a data storage format that could not be edited directly by MB-System. The raw data were first converted into a format compatible with MB-System, then edited with mbedit. A few sections needed sound speed adjustment. These data are now edited and included in the current version of the SO GLOBEC data base.

3.2 Conventional echo sounding data

3.2.1 NGDC single point track line

A large part of the digital bathymetry data in the study area was collected using more traditional echo sounders. This type of data provided the largest number of miles of data along track. This type of echo sounding only provides a single depth point under the ship, unlike the swath data from the multibeam mapping systems.

These data need quality control checking. The data were provided to NGDC from many sources, and while many sources provide processed data, some only provide raw (unprocessed and unchecked) data. There are no real quality control tags on the data provided to NGDC, so care must be made to check the crossings of the along-track data between individual cruises. Sometimes the data will show differences of 10's (to even 100's) m between the depths from two cruises at a crossing point. This difference must be resolved before utilizing the data from either cruise.

Although these along-track data cover a small area of the map relative to that area covered by the multibeam data, the along-track data are critical in providing coverage of the region and "tie lines" to the multibeam data. These tie lines can be used to constrain valleys and ridges and help confirm regional trends.

The NGDC along-track data were processed using the MB-System program mgrid to create a gridded data set to use with data from other sources.

3.2.2 *R/V Laurence M. Gould along-track data*

We have been fortunate to get access to the along-track single-point bathymetry data collected during 2000-2003 on the *R/V Laurence M. Gould* (Figure 9). The ship's data acquisition system records 1-min data from a suite of underway meteorological and water sampling systems that include the ship's digital echo sounder and GPS latitude and longitude. The data are stored in the Global Ocean Flux Study (JGOFS) ASCII format. The echo sounder uses its own bottom detection software and the data must be carefully edited to remove the many spurious depth values that occur, especially when the ship is working in the ice. Tom Bolmer wrote a MatLab program (see Appendix II) to identify and remove spurious data plus obvious peaks and valleys in the data that are unreasonable relative to what is known about the area or relative to other data (Figures 16 and 17).

The resulting edited JGOFS depth and position data were then processed with the MB-System program *mbgrid* to create a gridded data set that can be merged with data from other sources.

3.3 *Coastline Data*

Peter Morris provided the Scientific Committee on Antarctic Research (SCAR) 1:1 million Antarctic coastline data set (http://www.nerc-bas.ac.uk/public/magic/add_home.html). This is the current version of the coastline for the region. The GMT coastline subroutine in the GMT suite of programs uses data from old navigation charts of the area so that the GMT coastline is only useful for gross features. During cruises in the area, ships have "sailed" past the GMT/navigational chart coastlines onto "land". We have used the SCAR coastline data to constrain the areas with very little data and define the coastline in our gridded data (figure 12).

SO GLOBEC Bathymetry
WHOI-2004-02

3.4 DEM Topography

We have downloaded the United States Geological Survey global Digital Elevation Model (DEM) (GTOPO30) (<http://edcdaac.usgs.gov/topo30/topo30.html>) to use for the land area (Figure 13). This data set has a horizontal grid spacing of 30 arc seconds (approximately 1 kilometer), and covers most of the region except for some of the smaller islands in the northeast and southwest which are likely too small to be resolved with the 30-sec grid spacing. The DEM data have helped constrain the bathymetry by having the topography to influence the gridding of the bathymetry.

Data used in compilation




Figure 12. All the track coverage with DEM GTOPO30 and SCAR ADD data.

The DEM data is shown on land. The SCAR coastline and ice edges are shown. There is good agreement between the two data sets within the 30-sec spacing of the GTOPO30 data

4 Steps in Getting to a Final Grid

All of the data sources and types mentioned above were used to create a final data set and bathymetric chart. We first created a series of grids from some of the different data types and then merged them together with some sources getting preference. The steps we followed to create the final products are listed below. The scripts used to create these files are included in Appendix 1. Table 4 has a list of the programs used in the processing. The final version of the data set can be seen in Figures 13, 14 and 15.

1. The *NBP* multibeam data were turned into a grid using *mbgrid*. We have good faith in this data set and gave it preference over the other data sources. Some padding was performed around the data points to fill in areas with no data and create a fuller grid. These data were gridded using a 75 meter grid spacing. We created a NetCDF-formatted gridded data set (“*grd*” file) that can then be used by the GMT programs to merge and plot the data.
2. The remaining multibeam data from other sources were gridded using *mbgrid*. A smaller amount of padding was used in this data set. This created a second *grd* file.
3. These two grids were then merged using the GMT program *grdmath*. The program was asked to do a logical AND on the data. If there was data at a grid location in the *NBP* *grd* file, it was used. If not, then a value from the other multibeam *grd* file was used. If there was no data there, then a “NaN” (Not a Number) was assigned to that grid location. This created a file of high confidence and high quality data.
4. The NDGC and *LMG* JGOFS trackline data were turned into a grid using the same *mbgrid* program. A smaller padding number was used to fill in empty areas in this grid.
5. The merged multibeam *grd* file was merged with the NDGC and JGOFS trackline *grd* file using the same technique with GMT’s *grdmath* program as discussed above. The multibeam data was given precedence over the trackline data as above. This *grd* file represents all data collected by ships in the area in our data base.
6. The GTOPO30 data were reformatted into an ASCII XYZ file to represent only the land areas.
7. The combined ship data *grd* file was also reformatted into an ASCII XYZ file using the GMT *grd2xyz* program.

SO GLOBEC Bathymetry
WHOI-2004-02

8. The SCAR ADD coastline data were converted into an ASCII XYZ data file by adding a value of 0.0 for the water depth to represent the coast line.
9. All of the ASCII XYZ data files were concatenated together into a single file. These files are: a) the ASCII XYZ ship bathymetry data, b) the ASCII XYZ GTOPO30 land elevation data, c) the ASCII XYZ SCAR ADD coastline data, d) the ASCII XYZ file of bathymetric soundings in the King George VI Sound, and e) the ASCII XYZ data sent by Laurie Padman for use in areas of ice sheets.
10. This single XYZ file was then searched using the UNIX program egrep to remove those locations where there was no data. The NaN values were removed in this step, leaving only XYZ points with real data.
11. The GMT program blockmedian was then run to get a median position and value for every non-empty block in a grid "cell". Grid locations with data in them were created, thus generating a grid with either data or no data in equally spaced locations from the irregularly spaced input data.
12. The resultant file from the program blockmedian was then processed with the GMT program surface. Surface smoothly fills in empty grid locations with an interpolation sheet passing through the existing grid cells with data. The resulting grid can then be used to create maps of the study area. We used a grid spacing of 6 seconds longitude and 2 seconds of latitude. The program surface could not use meters as a grid spacing. These grid spacing values in surface best fit those output from the mbgrid program.
13. The grid above was input into the IVS Fledermaus program to find largely divergent highs and lows in the data set. The raw data were plotted in these regions to help find and remove the bad data points from the data set.
14. The data center line data were then regridded, and merged with the multibeam data again. Then steps 10 to 12 from above were repeated. The process of doing steps 13 and then redoing 10 to 12 was done several times until no more large diversions were noted in the data set with Fledermaus.

SO GLOBEC Bathymetry




Figure 13. Final SO GLOBEC bathymetric map.

SO GLOBEC Bathymetry




Figure 14. Final SO GLOBEC topographic map with elevation coloring.

SO GLOBEC Contours
from surface_16_3_all_topo30.surface




Figure 15. Final SO GLOBEC data, contour interval 100 meters.

5 Data Distribution

To facilitate the initial distribution of this data set to the SO GLOBEC participants, the web site http://www.whoi.edu/science/PO/so_globec/ was created to allow participants to download the data set. This web site has a link to an ftp site. The ftp site has a user name and password associated with it. For the time period that the data are proprietary. Since the SO GLOBEC participants use such a large range of computer operating systems and software, it was hard to find a single kind of data format that would be useful to all. As a result, we decided to serve on the ftp site the data set in the following formats:

1. GMT-created NetCDF binary files with 3 grid resolutions to allow users with smaller computers to get the data and be able to work with the files. These were a 6seconds longitude and 2 seconds latitude grid, a 15 seconds on each side grid and a 1 minute on each side grid.
2. ASCII XYZ files with 2 grid resolutions for the same reason. Only the 15 seconds and 1 minute grids were used here due to file size limitations.
3. Separate ASCII XY files for each 100-m depth contour.
4. MatLab MAT files of the gridded data with two different grid resolutions. We used the same size grids as in number 2 above.
5. A TIFF file for GIS users. (We have had trouble distributing data to users of GIS software packages. These programs seem unable to read the NetCDF-formatted files nor re-grid the ASCII XYZ data easily. This TIFF file is an attempt to get the GIS users data they can use.)
6. GIS users have a separate directory of files provided by Ari Friedlaender (Duke University). (The format of these files is not understood by us.) Other GIS users have had trouble using these files also. These files are still being served in the hope that they are helpful to someone.) At the time of publication these GIS files are not of the current gridded data set. We hope to have that updated soon with help from other SO GLOBEC participants.
7. There are two scene files for use in the IVS iView3D program. These files used with the free program iView3D program will enable the user to "fly" though the data in a 3D grid.

SO GLOBEC Bathymetry WHOI-2004-02

With the publication of this technical report, the web site http://www.whoi.edu/science/PO/so_globec/ will be updated and the digital bathymetry ftp site [ftp.whoi.edu](ftp://ftp.whoi.edu), which has the username of `so_globec`, will be opened for all users. Another link to these sites will be added to the U.S. Southern Ocean GLOBEC website at Old Dominion University (http://www.odu.edu/Research/globec_menu.html) to help dissemination of the existence and availability of this data set. The GLOBEC web site at W.H.O.I. for the distribution of all U.S. GLOBEC data (Georges Bank, NEP, and SO GLOBEC) also has a link to this data at: <http://globec.whoi.edu/jg/dir/globec/soglobec/>. The complete SO GLOBEC digital bathymetry data set will be archived at WHOI by Tom Bolmer (tbolmer@whoi.edu) and investigators seeking data products not present on the ftp site should contact Tom directly.

6 Summary

By combining bathymetric data from many sources and using various interpolation methods, a data set of bathymetry gridded nominally at 75-m spacing and bounded by 65° to 71° S and 65° to 78° W has been created. This represents a significant improvement over what was previously available in this area. The final product in the form of several gridded depth and depth contour data sets using different horizontal resolutions will now be available to all users (both the SO GLOBEC investigators and the rest of the public) at the ftp site: [ftp.whoi.edu](ftp://ftp.whoi.edu) (ftp://so_globec@ftp.whoi.edu), at the web site: http://www.whoi.edu/science/PO/so_globec/, and at the US GLOBEC web site at: <http://globec.whoi.edu/jg/dir/globec/soglobec/>.

7 Suggestions for improving the compilation

This compilation is only a snap shot of the data available at the time of publication of this report. There are many things that could be done to improve upon this bathymetry data set. We list some ideas below.

- 1) We are confident that we have found most of the science cruises in region that collected well-located digital bathymetric data. But we feel that we have not gotten all of those available.
- 2) Some of the data that we have could be improved upon by having access to the processed multibeam files instead of more coarsely gridded ASCII XYZ data.
- 3) There are several cruises included in this compilation for which only a limited subset of the bathymetry data was made available to us. Including the missing data from these cruises would improve the accuracy in some areas.
- 4) In the process of editing the single channel track data, we may have erred and edited out good data points. It is very difficult to edit and process this type of data without access to the actual echo sounding records for comparison. Several examples of undiscovered shoals were found in the echo sounding data on the SO GLOBEC RVIB N. B. Palmer cruises that were not shown in the edited single channel track data. These omissions should be investigated further.
- 5) Some of the data could benefit from further processing. We perhaps would be able to improve upon the map we have if we could process the entire data set with the new bathymetry processing tool now in development and testing at the University of New Hampshire.
- 6) The final gridded product could be improved by having a surfacing routine that smoothes the areas with no data in a better more realistic manner. The currently used GMT program is not producing smoothly flowing surfaces.

SO GLOBEC Bathymetry
WHOI-2004-02

- 7) There should be some way to incorporate larger data sets like the Smith & Sandwell 2-minute gravity-derived topography, the GEBCO 2003 1 minute grid or the SCAR BEDMAP data set into this SO GLOBEC compilation. At this moment, there is a large difference between our product and these. Since we have a larger data set in a smaller area than these compilations, our data should be able to guide these data sets in our region of interest. If we could use our data set (which is more complete and accurate in our study region) as a ground truth data set for one of these products, perhaps we could utilize the resulting other product to make a more complete bathymetry map for our region.
- 8) There are regions in the map area where additional data could make a large improvement. These areas can be seen easily by looking at Figure 12. There are large areas where there is very sparse data coverage. Several of the areas we think that need more survey work done are:
- a) The region off of Lazareth Bay. It would be good to see if the deep area near the bay continues offshore. Does this deep continue further onshore?
 - b) Both the Wilkins and George VI Ice Shelves need more data. Only Maslanyj's profiles in George VI Sound contain any real data in these areas.
These areas probably contribute to the general circulation in the region with water passing under the ice shelves. Knowing how much water could possibly move in these regions would help modelers in the general region.
 - c) The general southwestern part of the map needs more data collected. There are hints of features but the existing data are so sparse that we need more data to constrain these features. Does the deep in the southwestern area continue over the continental shelf to the deep sea?
 - d) There has been scientific interest in a more-detailed description of the bathymetry in the Crystal Sound area. There are questions as to whether the deeper areas are connected to the shelf. Deeper water mass characteristics have been noted in the area and it would be good to know if there is a channel for this water to get there.
 - e) The southeastern part of Marguerite Bay to the east of Marguerite Trough shows no cruise tracks in the area. The general interior of Marguerite Bay is only sparsely surveyed. Having a better idea of the bathymetry here would help a lot in understanding the overall Marguerite Bay system.

8 A Brief Time Line of the Compilation

Get Smith & Sandwell 2 minute global elevation data
check and get NGDC track lines data
digitize Navigational charts
Merge all of this into a new ETOPO82A gridded data set

First *R/V Laurence M. Gould* SO GLOBEC Cruise LMG0102 Data are noted to be VERY different from the ETOPO82A data

Begin first of three *RVIB Nathaniel B. Palmer* SO GLOBEC cruises collecting SeaBeam data. Data are processed at sea.

After the first two cruises we tried to merge all of the data assembled to date. These data are noted to not merge very well with the ETOPO8.2, the ETOPO82A or the digitized Navigation charts.

A decision is made to not use any of the previously assembled data. We will only collect our own data and search for other available data sets.

Carol Pudsey at British Antarctic Survey offers the data from a multibeam cruise on the James Clark Ross.

Do a search of NGDC and get all bathymetry data available. This includes a few cruises that collected multibeam. Get the GTOPO30 30 second spaced topography data. This fills in the land areas and helps constrain the data set.

Peter Morris sends the SCAR ADD coastline data. We tried to get this from the ADD web site but it is only distributed in GIS files. Could not read the GIS data formats so the ASCII files from Peter are a great help. This constrains the bathymetry data set to the water area only when we are using surface to fill in unsurveyed areas.

Created a tool to edit out bad data points from the *R/V Laurence M. Gould* and *RVIB Nathaniel B. Palmer* underweigh "JGOFs" meteorology data. These data included bathymetry from the either the 3.5 or 12 kHz echo sounders. These data were NOT edited for quality control at sea. This MatLab tool helped by being able to remove extraneous points, using the GUI tools in MatLab. Wildly different peaks and valleys were removed this way.

Peter Morris at the BAS puts me in touch with Andras Maldonado in Spain who offers multibeam from the Hesperides.

SO GLOBEC Bathymetry
WHOI-2004-02

Presented a poster at the Ocean Sciences AGU Hawaii February 2002 to show the process of the work.

After SO GLOBEC cruise III on the *RVIB Nathaniel B. Palmer* some data were also released by previous chief scientists on the *RVIB Nathaniel B. Palmer*. SO GLOBEC cruise III is the last SO GLOBEC to collect multibeam data. The SeaBeam system on the *RVIB Nathaniel B. Palmer* was replaced by a Simrad system, which was not operational at cruise time for SO GLOBEC IV.

Created a web site to show the AGU poster on line. This was added to as new things were done to the data. Before the 2002 December SO GLOBEC meeting the first set of preliminary data was put up on the web for SO GLOBEC participants to download and use.

BAS sent raw multibeam data from cruises in the area. These were data that had not been processed due to lack of interest at BAS. Tom mbeditted the data in the summer of 2003 and added it to the collected data.

Searched through the data and added points to areas where no data existed. Data from Laurie Padman were used to fill in areas like under the Wilken's Ice Shelf where we had no data. We needed to edit Laurie's data so that it did not occur on land or in areas we had real ship track data. We also added points to areas that Laurie's data did not cover but where we had coastline and land elevation data. This helped constrain some areas to give surfaces that fit the general data available. This was basically an adjustment to the data to force the computer surfacing and gridding programs towards more realistic results.

Acquire the GEBCO 2003 world 1 minute gridded data set. This turns out to be of no use. The grid was made by digitizing hand contoured data and then re-gridding the data. The data utilized were not as current a data set as we have. This produced a product that is very different from the real data we have assembled. It had been hoped that the data set could be a back drop to the data we have collected.

IVS Fledermaus program available to 3D viewing. This was used to highlight regions with previously unnoticed spurious highs and lows. These points were then removed and the data re-gridded in mbgrid. The rest of the data flow was then repeated to get a final gridded product.

Rob Lartner at BAS sent 2 single channel tracks of bathymetry data from the *RRS James Clark Ross* as the final parts of this report were being checked prior to publication.

9 Acknowledgements

We want to thank the many individuals who have helped collect and edit the multibeam and along-track bathymetry data included in this SO GLOBEC digital bathymetry data set. Without their effort, both at sea and back home, this project could not have happened. The resulting data set and our improved knowledge of the bathymetry in Marguerite Bay and adjacent shelf is a direct result of many individual decisions to collect high-quality data and share these data with others.

We also wish to thank Walter Smith for the helpful advice and comments he provided during this project.

This work has been supported by the U.S. National Science Foundation through grants OPP 99-10092 and OPP 02-34163 to Bob Beardsley and Tom Bolmer (WHOI), OPP 99-10307 to Peter Wiebe (WHOI), OPP 99-09956 to Eileen Hoffman (ODU), OPP 95-278776 to John Anderson (Rice), and the British Antarctic Survey (Natural Environment Research Council). This support from NSF and BAS is greatly appreciated.

10 References

- Caress, D. W., and D. N. Chayes, New software for processing sidescan data from sidescan-capable multibeam sonars, *Proceedings of the IEEE Oceans 95 Conference*, 997-1000, 1995.
- Caress, D. W., and D. N. Chayes, Improved processing of Hydrosweep DS multibeam data on the R/V Maurice Ewing, *Mar. Geophys. Res.*, 18, 631-650, 1996.
- GTOPO30, Global Topographic Data. Land Process Distributed Active Archive Center (LP DAAC), U. S. Geological Survey's EROS Data Center, <http://edcdaac.usgs.gov> (<http://edcdaac.usgs.gov/gtopo30/gtopo30.html>).
- IOC, IHO, and BODC (2003). Centenary Edition of the GEBCO Digital Atlas. Published on CD-ROM on behalf of the Intergovernmental Oceanographic Commission and the International Hydrographic Organization as part of the General Bathymetric Chart of the Oceans; British Oceanographic Data Centre, Liverpool.
- Maslanyj, M. P. (1987). Seismic bedrock depth measurements and the origin of George VI Sound, Antarctic Peninsula. *British Antarctic Survey Bulletin*, 75, 51-65.
- SCAR Working Group on Geodesy and Geographic Information, Antarctic Digital Database, Version 3.0, http://www.nerc-bas.ac.uk/public/magic/add_main.html, July 2000, Scientific Committee on Antarctic Research.
- Smith, W. H. F., and D.T. Sandwell (1997). Global Sea Floor Topography from Satellite Altimetry and Ship Depth Soundings. *Science*, 277, 1956-1962. http://topex.ucsd.edu/WWW_html/mar_topo.html
- Wessel, P., and Smith, W. H. (1991). Free Software helps Display and Map Data. *EOS Trans.*, 72, 441,445-446.
- Wessel, P., and Smith W. H. (1998). New, improved version of the Generic Mapping Tools released. *EOS Trans.*, 79, 579.

SO GLOBEC Bathymetry
WHOI-2004-02

TABLE 1:

Navigational Charts used to augment ETOPO2-v8.2

Map	Description
GB 3571	Lavoisier Island to Alexander Island U.K. Hydrographic Office (fm) 1:500 000 10/1/1998
GB 3577	Adelaide Island, South Western Approaches U.K. Hydrographic Office (fm) 1:75 000 3/1995
US 29142	Alexander Island to Square Bay, including Marguerite Bay National Imagery and Mapping Agency (m) 1:200 000 11/2/1996
US 29141	Square Bay to Matha Strait, including Adelaide Island Defense Mapping Agency (m) 1:200 000 8/6/1988

SO GLOBEC Bathymetry
WHOI-2004-02

TABLE 2:

*Cruises Used to Create the Marguerite Bay Region Bathymetry
Chart. 02/01/04*

<u>Cruise</u>	<u>Center Beam</u>	<u>Multi-Beam</u>
NSF NBP0103		X
NSF NBP0104		X
NSF NBP0201	JGOFS (the rest)	X (selected)
NSF NBP0202		X
NSF NBP0204	JGOFS	
B.A.S. JR59		X (XYZ file)
B.A.S. JR69		X (XYZ file)
B.A.S. JR71		X (XYZ file)
LDEO EW9101		X
NOAA RITS93A		X
NOAA RITS94B		X
RUSSIA Petrov 1991		X (XYZ file)
SPAIN Hesperides ANTPAC97-98		X (XYZ file)
NSF LMG0009	JGOFS	
NSF LMG0101	JGOFS	
NSF LMG0102	JGOFS	
NSF LMG0103	JGOFS	
NSF LMG0104	JGOFS	
NSF LMG0105	JGOFS	
NSF LMG0106	JGOFS (Discarded)	
NSF LMG0201	JGOFS	
NSF LMG0201A	JGOFS	
NSF LMG0202	JGOFS	
NSF LMG0203	JGOFS	
NSF LMG0205	JGOFS	
NSF LMG0301	JGOFS	
NSF LMG0302	JGOFS	

SO GLOBEC Bathymetry
WHOI-2004-02

<i>Cruise</i>	<i>Center Beam</i>	<i>Multi-Beam</i>
LDEO C1503	MGD77	
LDEO ELT10	MGD77	
LDEO ELT43	MGD77	
LDEO THA80	MGD77	
LDEO THB80	MGD77	
W.H.O.I. PD388L01	MGD77	
W.H.O.I. PD190L02	MGD77	
RICE UNIV. DF85	MGD77	
RICE UNIV. DF86	MGD77	
RUSSIA Ioffe 1992	MGD77	
S.I.O. DSDP35GC	MGD77	
U K RRS Bransfield 1985	MGD77	
U K HMS Endurance 1985	MGD77	
U K RRS Discovery 1887	MGD77	
U K HMS Endurance 1987	MGD77	
U K HMS Endurance 1988	MGD77	
U K RRS Charles Darwin 1989	MGD77	
U K HMS Endurance 1991	MGD77	
U K HMS Endurance 1992	MGD77	
U K HMS Polar Circle 1992 (139)	MGD77	
U K HMS Polar Circle 1992 (144)	MGD77	
U K HMS Endurance 1994	MGD77	
U K HMS Endurance 1996	MGD77	
U K HMS Endurance 1997	MGD77	
U K HMS Endurance 1999	MGD77	
U K RRS James Clark Ross JCR10L01	MGD77	
U K RRS James Clark Ross JCR10L02	MGD77	
U K RRS James Clark Ross JCR04	MGD77	
BRAZIL TF88-89	MGD77	
ARGENTINA IRI93-94	MGD77	
ARGENTINA AI9495	MGD77	
ARGENTINA BHPD9798	MGD77	
NSF NBP92-8	MGD77	
NSF NBP93-7	MGD77	

SO GLOBEC Bathymetry
WHOI-2004-02

<i>Cruise</i>	<i>Center Beam</i>	<i>Multi-Beam</i>
NSF <i>NBP93-8</i>	MGD77	
NSF <i>NBP94-2</i>	MGD77	
NSF <i>NBP95-7</i>	MGD77	
NSF <i>NBP99-2</i>		X
NSF <i>NBP99-3</i>		X
NSF <i>NBP99-5</i>		X
NSF <i>NBP99-6</i>	MGD77	
NSF <i>NBP00-1</i>	MGD77	
NSF <i>NBP00-4</i>	MGD77	
NSF <i>NBP01-5</i>	MGD77	
NSF <i>NBP01-7</i>	MGD77	
UNIV TOKYO UM66-B	MGD77	
ODP LEG 178JR	MGD77	

TABLE 3:

LAND DATA used to Create the Marguerite Bay Region Bathymetry Chart.

<i>Data is in the form of XYZ and in the best spacing available.</i>
GTOPO30
SCAR ADD coastline data
GEORGE VI Sound Soundings traverses (Maslanjy)

SO GLOBEC Bathymetry
WHOI-2004-02

TABLE 4:

Programs used in processing the data

MB-System		Caress, Spitzak, and Chayes
GMT	(Generic Mapping Tools)	Wessel and Smith
PERL	(Practical Extraction and Report Language)	Wall
MatLab		MathWorks
Fledermaus		Interactive Visualization Systems

APPENDIX 1: MB-System and GMT scripts

Script to grid the Nathaniel Palmer Sea Beam data

```
#!/bin/csh -f
#
# Shellscrip to grid the NBP SeaBeam data
#
# This shellscrip created by following command line:
# mbm_grid -A2 -C3 -E100/100 -F-5 -G3 -Inew_grid_1.list -N -Onew_grid_1
-R-78/-66/-70/-65 -V
#
# Define shell variables used in this scrip:
set REGION          = -78/-65/-71/-65
set INPUT_FILE      = all_NBP_multibeam2.list
set INPUT_FORMAT    = -5
set ROOT            = all_NBP_multibeam_C5_2
#
# Make datalist file
echo Making datalist file...
echo $INPUT_FILE $INPUT_FORMAT >! datalist$$
#
# Run mbgrid
echo Running mbgrid...
mbgrid -I$INPUT_FILE \
    -R$REGION \
    -O$ROOT \
    -A2 -N \
    -F1 \
    -E75/75"meters" \
    -C5 \
    -G3 \
    -V
echo All done!
```

SO GLOBEC Bathymetry
WHOI-2004-02

Script to grid the Other Multibeam Data

```
#!/bin/csh -f
#
# Shellsript to grid the other Multibeam data
#
# Define shell variables used in this script:
set REGION      = -78/-65/-71/-65
set INPUT_FILE  = other_mbeam.list
set INPUT_FORMAT = -5
set ROOT        = other_mbeam_C5
#
# Make datalist file
echo Making datalist file...
echo $INPUT_FILE $INPUT_FORMAT >! datalist$$
#
# Run mbgrid
echo Running mbgrid...
mbgrid -I$INPUT_FILE \
    -R$REGION \
    -O$ROOT \
    -A2 -N \
    -F1 \
    -E75/75"meters" \
    -C5 \
    -G3 \
    -V

echo All done!
grdmath all_NBP_multibeam_C5_2.grd other_mbeam_C5.grd AND =
multibeam.grd
```

SO GLOBEC Bathymetry
WHOI-2004-02

Script to grid the ASCII along track data

```
#!/bin/csh -f
#
# Shellsript to grid the NGDC and Other ASCII data files
#
# Define shell variables used in this script:
set REGION      = -78/-65/-71/-65
set INPUT_FILE  = all_ASCII.list
set INPUT_FORMAT = -5
set ROOT        = all_ASCII_75m_C3
#
# Make datalist file
echo Making datalist file...
echo $INPUT_FILE $INPUT_FORMAT >! datalist$$
#
# Run mbgrid
echo Running mbgrid...
mbgrid -I$INPUT_FILE \
    -R$REGION \
    -O$ROOT \
    -A2 -N \
    -F1 \
    -E75/75"meters" \
    -C3 \
    -G3 \
    -V

echo All done!

grdmath multibeam.grd all_ASCII_75m_C3.grd AND = grid_14_1.grd
```

SO GLOBEC Bathymetry
WHOI-2004-02

Script to merge Land, Gridded at sea data and the SCAR Coastline

```
#!/bin/sh

# This script gets all of the data into an ASCII format with no NaNs.
# It then grids all of the data and fits the areas with no data to a
# "sheet" that passes through all of the known points.
# It then creates a postscript file to plot the data with.

rm .gmtdefaults
gmtset PAPER_MEDIA tom
gmtset PAPER_MEDIA archE+
gmtset BASEMAP_AXES WeSn
gmtset DEGREE_FORMAT 3
gmtset BASEMAP_TYPE PLAIN
gmtset COLOR_NAN 255/255/255
gmtset COLOR_BACKGROUND 0/0/0
gmtset COLOR_FOREGROUND 255/255/255
gmtset LABEL_FONT_SIZE 12
gmtset HEADER_FONT_SIZE 12
gmtset ANOT_FONT_SIZE 10
gmtset LABEL_FONT_SIZE 12

BOX1="-78/-65/-71/-65"
BOX="-78.1/-63.9/-71.1/-64.9"
SHIFT="-X1.5 -Y2.0"
MAP_PROJ="m0.45"

rm surface_16_2_all.ps

grd2xyz -V -R$BOX grid_16_2_75m.grd | egrep -v NaN | \
awk '{ if ($3 < 0.0) {print;}}' > grid_16_2_all.xyz

cat grid_16_2_all.xyz \
  /fal10/mb/BAS/george6dep_neg.dat \
  /fal10/mb/BAS/SCAR_Dick_coast_forgrid.psx \
  /fal10/mb/data/padman/padman_depths_ROSS_neg.dat \
  /fal10/mb/data/padman/WILKNINSICE.padman_depths2_neg2.dat \
  /fal10/mb/data/padman/padman_depths_south_neg.dat \
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
/fal10/mb/data/padman/dummy_topo.pos \  
/fal10/mb/data/padman/gtopo30_south_pos.psx \  
/fal10/mb/grids/surface_12/topo30_pos.xyz \  
/fal7/mb/checks/dummy_islands_topo30_pos.xyz > \  
surface_16_2_all_topo30.xyz
```

```
rm grid_16_2_all.xyz
```

```
blockmedian surface_16_2_all_topo30.xyz -R$BOX \  
-I6c/2c -V > surface_16_2_all_topo30.block
```

```
rm surface_16_2_all_topo30.xyz
```

```
# -I6.25c/2.4c -T.25i -T0b \  
# -I6.25c/2.4c -T.4i -T0b \  
# -I6c/2c -T.4i -T0b \  
# -I6c/2c -T.05 \ # pretty good  
# -I6c/2c -T.00 \ # many above sealevel arfeas created  
# -I6c/2c -T.10 \ # still have false above se level areas  
surface surface_16_2_all_topo30.block -R$BOX \  
-I6c/2c -T.5 \  
-Gsurface_16_2_all_topo30.surface -V
```

```
rm surface_16_2_all_topo30.block
```

```
grdhisteq -V -N surface_16_2_all_topo30.surface \  
-Gsurface_16_2_all_topo30_hist.grd  
grdgradient -V -A0 -Nt surface_16_2_all_topo30_hist.grd \  
-Gsurface_16_2_all_topo30_hist_grad.grd
```

```
rm surface_16_2_all_topo30_hist.grd
```

```
makecpt -Crainbow -T-4500/-0/25 -V > grid_15_1.cpt
```

```
grdimage surface_16_2_all_topo30.surface \  
-Isurface_16_2_all_topo30_hist_grad.grd \  
-Ba1g1f2/a1g1f1:"surface_16_2_all Tension of .50":WeSn \  
-R$BOX1 -J$MAP_PROJ -Cgrid_15_1.cpt \  
$SHIFT -V -K -P > surface_16_2_all.ps
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
psscale -Cgrid_15_1.cpt -E -B500:"Topography (meters)": \  
-D2.9/-8/5.5/3h -V -O -K >> surface_16_2_all.ps
```

```
pstext -R$BOX1 -J$MAP_PROJ -N -V -K -O <<EOF>>  
surface_16_2_all.ps  
-75.15 -69.95 10 0.0 5 MC Charcot Island  
-73.25 -69.50 10 0.0 5 BL Rothschild Island  
-71.00 -69.00 10 30.0 5 MC Alexander Island  
-68.60 -67.20 10 70.0 5 MC Adelaide Island  
EOF
```

```
psxy -R$BOX1 -J$MAP_PROJ -M /fal10/mb/BAS/SCAR_ice.psyx \  
-W3/0/0/200 \  
-V -O -K >> surface_16_2_all.ps
```

```
psxy -R$BOX1 -J$MAP_PROJ -M /fal10/mb/BAS/SCAR_coast.psyx -W3 \  
-V -O -U/0/-0.5/"Compiled by Tom Bolmer" >> surface_16_2_all.ps
```

```
gs surface_16_2_all.ps  
#gs -dNOPAUSE -sDEVICE=jpeg -sOutputFile=surface_16_2_all.jpg  
surface_16_2_all.ps -dBATCH
```

SO GLOBEC Bathymetry
WHOI-2004-02

Script to find widely divergent points

```
#!/bin/sh

# STB 2/15/4
# This script will plot the gridded multibeam data.
# On top of that the along track data is plotted as +'s.
# A box is selected which is drawn around points which are
# listed to the screen. The points listed also show the line number in
# the data file so that the points can be found and edited.

rm .gmtdefaults
gmtset PAPER_MEDIA tom
gmtset BASEMAP_AXES WeSn
gmtset DEGREE_FORMAT 3
gmtset BASEMAP_TYPE PLAIN
gmtset COLOR_NAN 255/255/255
gmtset COLOR_BACKGROUND 0/0/0
gmtset COLOR_FOREGROUND 255/255/255
gmtset LABEL_FONT_SIZE 12
gmtset HEADER_FONT_SIZE 12
gmtset ANOT_FONT_SIZE 10

BOX="-75/-73/-68/-67:15"
MAP_PROJ="m3"
SHIFT="-X1.5 -Y3.0"

makecpt -Crainbow -T-3500/-100/25 > get_bad_points.cpt

grdimage /fal7/mb/grids/surface_16/grid_16_2_C10_75m_good_MB.grd \
  -Ba5mg5mf1m:."NGDC points":WeSn \
  -R$BOX -J$MAP_PROJ -Cget_bad_points.cpt \
  $SHIFT -V -K -P > get_bad_points.ps

# select a box to list the data from
LEFT1='-75.0'
RIGHT1='-73'
TOP1='-67:15'
BOTTOM1='-68'
LEFT=`echo $LEFT1 | awk '{split($1,new,"."); \
  printf("%d.%3d",new[1],((new[2]/60.0)*1000));}'`
RIGHT=`echo $RIGHT1 | awk '{split($1,new,"."); \
  printf("%d.%3d",new[1],((new[2]/60.0)*1000));}'`
TOP=`echo $TOP1 | awk '{split($1,new,"."); \
```


SO GLOBEC Bathymetry
WHOI-2004-02

```

printf("%d.%3d",new[1],((new[2]/60.0)*1000));}"
BOTTOM=`echo $BOTTOM1 | awk '{split($1,new,":"); \
printf("%d.%3d",new[1],((new[2]/60.0)*1000));}"
# list to the screen the points in the box selected
nawk 'BEGIN { count = 0; } { \
count=count+1; \
if (($1 != ">") && \
($1 < right) && \
($1 > left) && \
($2 > bottom) && \
($2 < top)) { \
printf ("%f %f %f %d %d %s\n", $1, $2, $3, count, $4, $5);}' \
right=$RIGHT left=$LEFT top=$TOP bottom=$BOTTOM \
all_tracks_list.psx

psxy -R$BOX -J$MAP_PROJ \
-W3/255/0/255 \
-V -O -K <<EOF>> get_bad_points.ps
$LEFT $TOP
$LEFT $BOTTOM
$RIGHT $BOTTOM
$RIGHT $TOP
$LEFT $TOP
EOF

# Plot data that is thought to be good but not in raw multibeam
format.
psxy -R$BOX -J$MAP_PROJ \
-M /fal10/mb/grids/surface_12/topo30_pos.xyz \
-Sx.05 \
-V -O -K >> get_bad_points.ps
psxy -R$BOX -J$MAP_PROJ -M \
/fal10/mb/grids/surface_12/rightgtopo30.xyz \
-Sx.05 -W4/0/0/255 \
-V -O -K >> get_bad_points.ps
psxy -R$BOX -J$MAP_PROJ -M dummy_islands_topo30_pos.xyz \
-Sx.05 -W3/255/0/0 -G255/0/0 \
-V -O -K >> get_bad_points.ps

psxy /fal10/mb/data/BAS/JR04_1.psx -R -Jm -Sx.1 -
Cget_bad_points.cpt \
-V -O -K >> get_bad_points.ps
psxy /fal10/mb/data/BAS/JR04_2.psx -R -Jm -Sx.1 -
Cget_bad_points.cpt \
-V -O -K >> get_bad_points.ps

psscale -Cget_bad_points.cpt -E -B500:"Topography (meters)": \

```

SO GLOBEC Bathymetry
WHOI-2004-02

```
-D2.9/-.8/5.5/.3h -V -O -K >> get_bad_points.ps
psxy -R$BOX -J$MAP_PROJ -M \
    /fal10/mb/BAS/SCAR_Dick_coast.psx \
    -W3/0/255/0 \
    -U/0/-0.5/"Compiled by Tom Bolmer" -V -O -K >> get_bad_points.ps

# plot the points being printed to the screen
psxy -R$BOX -J$MAP_PROJ all_tracks_list.psx \
    -Sx.05 -Cget_bad_points.cpt -M \
    -V -O >> get_bad_points.ps

gs get_bad_points.ps
```

APPENDIX 2: Matlab Script

MatLab Scripts to edit JGOFS data

JGOFSglobal_start.m

```
% global setup for start of run
% this should be cut and pasted into the matlab window
% to initialize the globals
%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%

global CHANNEL_1;
global CHANNEL_2;
global CHANNEL_3;

CHANNEL_1 = [];
CHANNEL_2 = [];CHANNEL_3 = [];

global xnew ynew znew file_name
global savefile_button
global safde_day file_name;
global win_len x ddd;
global dddd hhhh;

safde_day = 400.0;
win_len = 1;
```

SO GLOBEC Bathymetry
 WHOI-2004-02




Figure 16. Main window in JGOFSplot MATLAB GUI tool.

SO GLOBEC Bathymetry
 WHOI-2004-02




Figure 17. JGOFSplot MATLAB GUI tool zoom window.

SO GLOBEC Bathymetry
WHOI-2004-02

getJGOFS.m

```
function getJGOFS()

% set up parameters to call the JGOFS2mat m-file
% to read JGOFS data
% Tom Bolmer 09/02/03

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

global CHANNEL_1;
global CHANNEL_2;
global CHANNEL_3;

CHANNEL_1 = [];
CHANNEL_2 = [];CHANNEL_3 = [];

global xnew ynew znew file_name
global savefile_button
global safde_day file_name;
global win_len x

safde_day = 400.0;
win_len = 1;
file_name = input('enter filename to read ','s');

skip=0;
ntraces=1;
% Call m-file to read in data

JGOFS2mat(file_name);
```

SO GLOBEC Bathymetry
WHOI-2004-02

JGOFs2mat.m

```
function JGOFs2mat(input)
%
% Tom Bolmer 09/02/03
%
% For use at sea as a VERY crude testing tool
%
% a specialized function for reading the headers of SWD binary files
% To run 1 arguments is needed:
% input:  Raw binary data file name.

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

global CHANNEL_1;
global CHANNEL_2;
global CHANNEL_3;
CHANNEL_1=[];
CHANNEL_2=[];CHANNEL_3=[];
global ddd dddd hhhh dd hh;

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
[CHANNEL_2,CHANNEL_3,
CHANNEL_1,dddd,hhhh]=textread(input,'%f%f%f%s%s\n');

dd=char(dddd);
hh=char(hhhh);
l=length(dddd)
months=[ 0 31 59 90 120 161 192 223 254 284 315 345];
for i=1:l

    day(i)=sscanf(dd(i,1:2),'%f');
    mon(i)=sscanf(dd(i,4:5),'%f');

    hour(i)=sscanf(hh(i,1:2),'%f');
    min(i)=sscanf(hh(i,4:5),'%f');
    sec(i)=sscanf(hh(i,7:8),'%f');
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
h(i)=(hour(i) + ((min(i)+(sec(i)/60.0))/60.0));
ddd(i)= day(i) + (h(i)/24.0) + months(mon(i));

end

%
mat=1;
JGOFSPlot.m

%function JGOFSPlot(action)

% JGOFSPlot
% for JGOF bathymetry data viewing .
%
% STB 09/02/03
% extensively copied from a file by Jim Doutt of 15 Feb 1994

% *****

REVIS = '03/07/02 11:15';

global savefile_button NNV

nversio = version;
NNV = str2num(nversio(1:1));

fprintf('\n JGOFSPlot %s version 1 %s\n\n',nversio,REVIS);

SAMPLE_RATE = 400.0;
SAMPLE_RATE = 1.0;
FREQUENCY = 1.0 / SAMPLE_RATE;

global CHANNEL_1;
global CHANNEL_2;
global CHANNEL_3;

global x xnew ynew znew file_name;
global safde_day map4 DEPTH;
safde_day = SAMPLE_RATE ;
day = 1;
```


SO GLOBEC Bathymetry
WHOI-2004-02

```
% graphics initialization
clf reset;
set(gcf,'Units','inches','position',[6 5 7 10]);
fig =(gcf;
clf;

% Set up some dummy data for plot routines
t = [1 2];
y1 = [0 0];
y2 = 2*y1;

big=length (CHANNEL_1);
x=[1:big];
x=ddd;
smin = x(1:1);
dmax = x(big:big);

% x=x/400.0;
smin = 1;
dmax = big;

g = 'y';
start_day_value = smin;
end_day_value = dmax;
win_len = 400;
win_len = 1;

% Set up all pushbuttons
% buttons and display for start day slider

start_day_lab = uicontrol('Style','Text',...
'Units','normalized', 'Position',[.38 0.95 .06 .05],...
'String','start','backgroundcolor','white');

start_day=uicontrol('Style','Slider',...
'Units','normalized',...
'Position',[.52 .95 .20 .05],...
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
'Min',smin,'Max',dmax,'Value',smin,...
'CallBack',[...
'start_day_value = (get(start_day,"Value"));',...
'set(start_day_valu,"String"),',...
'num2str(get(start_day,"Val"))),'],'backgroundcolor','yellow');

start_day_min = uicontrol('Style','Text',...
'Units','normalized','Position',[.44 0.95 .08 .05],...
'String',num2str(smin),'backgroundcolor','magenta');

start_day_max = uicontrol('Style','Text',...
'Units','normalized','Position',[.72 0.95 .08 .05],...
'String',num2str(dmax),'backgroundcolor','magenta');

start_day_valu = uicontrol('Style','Text',...
'Units','normalized','Position',[.80 0.95 .12 .05],...
'String',num2str(get(start_day,'Value')),'backgroundcolor','cyan');

% buttons and display for end day slider

end_day_lab = uicontrol('Style','Text',...
'Units','normalized','Position',[.38 0.9 .06 .05],...
'String','end','backgroundcolor','white');

end_day=uicontrol('Style','Slider',...
'Units','normalized',...
'Position',[.52 .9 .20 .05],...
'Min',smin,'Max',dmax,'Value',dmax,...
'CallBack',[...
'end_day_value = (get(end_day,"Value"));',...
'set(end_day_valu,"String"),',...
'num2str(get(end_day,"Val"))),'],'backgroundcolor','yellow');

end_day_min = uicontrol('Style','Text',...
'Units','normalized','Position',[.44 0.9 .08 .05],...
'String',num2str(smin),'backgroundcolor','magenta');

end_day_max = uicontrol('Style','Text',...
'Units','normalized','Position',[.72 0.9 .08 .05],...
'String',num2str(dmax),'backgroundcolor','magenta');
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
end_day_valu = uicontrol('Style','Text',...
'Units','normalized','Position',[.80 0.9 .12 .05],...
'String',num2str(get(end_day,'Value')),'backgroundcolor','cyan');

% button for stating program
start_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.00 .95 .09 .05],'Interruptible','on','String','Start',...
'Callback','JGOFSDisplay;','backgroundcolor','green');
% buttons and display for next full window and 1/2 window

back_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.1 .95 .05 .05],'String','<--',...
'Callback',[g = "y"; , 'ii = ii - fix(win_len * safde_day);]);
next_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.15 .95 .05 .05],'String','-->',... 'Callback',[g = "y"; , 'ii = ii +
fix(win_len * safde_day);]);

back2_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.1 .9 .05 .05],'String','<.5',...
'Callback',[g = "y"; , 'ii = ii - fix(win_len * (safde_day / 2.0));]);
next2_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.15 .9 .05 .05],'String','.5>',... 'Callback',[g = "y"; , 'ii = ii +
fix(win_len * (safde_day / 2.0));]);
% buttons for printing and zooming

copy_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.2 .95 .08 .05],'String','Print',...
'Callback','g = "p";');
zoomn=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.2 .9 .08 .05],'String','zoom',...
'Callback','g = "z";');
% button for STOP

exit_button=uicontrol('Style','Pushbutton','Units','normalized',...
'Position',[.92 .95 .08 .05],'String','STOP',...
'Callback','g="n";','backgroundcolor','red');

edit_button=uicontrol('Style','Pushbutton','Units','normalized',...
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
'Position',[.92 .90 .08 .05],'String','EDIT',...
'Callback','g="e";','backgroundcolor','yellow');
save_edit_button=icontrol('Style','Pushbutton',...
'Units','normalized',...
'Position',[.89 .85 .11 .05],...
'String','KEEP IT',...
'Callback','g="k";','backgroundcolor','green','visible','off');
% buttons and display for window length in days

day_label=icontrol('Style','text','Units','normalized',...
'Position',[.29 .95 .08 .05],...
'String','win len','backgroundcolor','white');

day_val = uicontrol('Style','edit','Units','normalized',...
'Position',[.29 .9 .08 .05],...
'String',num2str(win_len),...
'callback',[win_len = str2num(get(day_val,"string"));],...
'value',win_len,'backgroundcolor','cyan');

% Set up areas for figures
beg_y = .10;
beg_y = .10;
del_y = .25;

xa1 = .12; x2 = .90; xtdel = .8; xfdel = .45; xofst = .02;   xa1 = .12; x2 = .90;
xtdel = .8; xfdel = .45; xofst = .02;
% Figure 2 - channel 2

ax4 = axes('Position', [xa1 beg_y+0.05 xtdel .45],'nextplot','replace',...
'visible','off',...
'xticklabel',';','yticklabel',';...',
'xaxislocation','bottom');
ax4_line=image(LONG(1,:),LAT(:,1),DEPTH,'CDataMapping','scaled');

set(ax4,'ydir','normal',...
'clim',[-4500 2000]);
ax3 = axes('Position', [xa1 beg_y+0.05 xtdel .45],'nextplot','add',...
'Visible','off');
ax3_line = plot(t,y1,'EraseMode','normal','color','green');
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
ax5 = axes('Position', [xa1 beg_y+0.05 xtdel .45], 'nextplot', 'add', ...
    'visible', 'off', 'box', 'on');
ax5_line=plot(CHANNEL_2, CHANNEL_3, 'k', ...
    cx.x(5).contours, cy.y(5).contours, 'b', ...
    cx.x(8).contours, cy.y(8).contours, 'b', ...
    cx.x(10).contours, cy.y(10).contours, 'g', ...
    cx.x(12).contours, cy.y(12).contours, 'g', ...
    cx.x(15).contours, cy.y(15).contours, 'r', ...
    cx.x(18).contours, cy.y(18).contours, 'c', ...
    cx.x(20).contours, cy.y(20).contours, 'c', ...
    cx.x(24).contours, cy.y(24).contours, 'c', ...
    cx.x(2).contours, cy.y(2).contours, 'y', ...
coastx, coasty, 'g', ... 'EraseMode', 'normal');

    ylabel('Latitude');
    xlabel('Longitude');

axb=axes('Position', [xa1 .05 xtdel .05], 'nextplot', 'add', ...
    'Visible', 'off'); axbar=colorbar('horiz');

set(axbar, 'Position', [xa1 .05 xtdel .05], 'visible', 'on', ...
    'climmode', 'manual', 'clim', [-4500 2000]);
theLim=[-4500 2000];
relabel(theLim, 'x');

    ax1 = axes('Position', [xa1 beg_y+del_y+del_y+.05 xtdel
    .20], 'nextplot', 'replace', 'Visible', 'off');
    ax1_line = plot(t, y1, 'EraseMode', 'normal', 'color', 'yellow');
ax2 = axes('Position', [xa1 beg_y+del_y+del_y+.05 xtdel
    .20], 'nextplot', 'replace', 'Visible', 'on'); ax2_line =
plot(t, y1, 'EraseMode', 'normal');
    TITL=title(file_name);

    ylabel('meters');
```

SO GLOBEC Bathymetry
WHOI-2004-02

JGOFSDisplay.m

```
% JGOFSDisplay()
% function to display JGOFSD Bathymetry data.
% Tom Bolmer 09/02/03
% adapted form the below versions
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%
% MELTPLOT.M
% Plots SAFDE electrical Noise data for the X and Y channels.
% file: meltplot.m
% by: Tom Bolmer
% date: 7/22/97
% for: Alan Chave
% Copied and modeled extensively from a file by Jim Doust Feb 1994
% copied form safdeplot.m
% Stb 7/22/97

%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%

global x CHANNEL_1 CHANNEL_2 CHANNEL_3 xnew ynew znew
file_name NNV ddd;
global cx cy coastx coasty map4;

start_day_value=x(1);
end_day_value=x(end);

dev_name = file_name;
x11 = start_day_value;

% Set up the max and min X values to be used in looping through
% the data.

iiend= length(CHANNEL_1) ;
clear start;
clear start1;
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
if isempty(x11)
    start= '1';
else
    start = x11;
end
set(start_button, 'Visible','off');
x12=end_day_value;

clear iend;
clear iend1;
iend = length(x);
if isempty(x12)
    iend= length(x) ;
else
    if iend > length(x);
        iend = length(x);
    end
end

text ('units','normalized','position',[.025 .9],'string','Channel 1');text
('units','normalized','position',[.025 -.35],'string','Navigation');
ii = fix(start);
% Loop through and do the plotting
while ii < iend
% Find the Max and Min for the data in this window so all% the boxes
will have the same axes.

start = fix(ii);
stop = (ii+fix(win_len * safde_day));
yymin = -1 * (max(-1*CHANNEL_1(start:stop)));
ymax = max(CHANNEL_1(start:stop)) ;

if (ymax < 0)
    ymax=ymax*.95;
else
    ymax=ymax*1.05;
end
if (yymin < 0.0)
    yymin = yymin * 1.05;
else
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
yymin = yymin * 0.95;
end
if (yymin == yymax)
    if (yymin < 0.0)
        yymin = yymin * 1.05;
    else
        yymin = yymin * 0.95;
    end
    if (yymin == 0)
        yymin=-.001
    end
end
end

xxmin = x(start);
xxmax = x(stop);

lonleft = (-1 * (max(-1 * CHANNEL_2(start:stop))))-.25;
lonright = max(CHANNEL_2(start:stop))+.25;
latbot = (-1 * (max(-1 * CHANNEL_3(start:stop))))-.25;
lattop = max(CHANNEL_3(start:stop))+.25;
hold on;

set(ax4,'XLim',[lonleft lonright],'YLim',[latbot lattop],...
    'visible','on','clim',[-4500 2000]);
set(ax3,'XLim',[lonleft lonright],'YLim',[latbot lattop],'nextplot','add');
set(ax3_line,'Xdata',CHANNEL_2(ii:(ii+fix(win_len * safde_day))),:),...
    'Ydata',CHANNEL_3(ii:(ii+fix(win_len * safde_day))),:),...
    'visible','on','linestyle','-','marker','+','color','red');set(ax5,'XLim',[lonleft
lonright],'YLim',[latbot lattop],'box','on');

if (CHANNEL_2(start) < CHANNEL_2(stop))
    direction='normal';
else
direction='reverse'; end
set(ax2,'XLim',[xxmin xxmax],'YLim',[yymin yymax],'visible','on');
set(ax2_line,'Xdata',x(start:stop),...
    'Ydata',CHANNEL_1(start:stop),...
    'linestyle','-','marker','none','color','green');
set(ax1,'XLim',[xxmin xxmax],'YLim',[yymin yymax],'Visible','off');
set(ax1_line,'Xdata',x(start:stop),...
```


SO GLOBEC Bathymetry
WHOI-2004-02

```
'Ydata',CHANNEL_1(start:stop),...  
'linestyle','-','marker','none','color','green');  
  
hold on;  
title_str=sprintf ("%s samples %10d to  
%10d',dev_name,fix(ii),fix(ii+fix(win_len * safde_day)));  
set (get(ax2,'title'),'string',title_str); hold off;  
% Reset the start slider value  
  
set (start_day,'Value',ii);  
set (start_day_valu,'String',num2str(ii));  
% Wait for an option to be selected  
  
g = 'stopit';  
while g == 'stopit'  
pause(1); end  
% Print the current window  
  
if (g == 'p')  
print -dps;  
g = 'stopit';  
while g == 'stopit'  
pause(1);  
end  
end  
  
if isempty(g)  
g = 'Y';  
end  
  
% ZOOM in on a selected area  
% Use ginput to find the array arguments  
  
while g == 'z'  
clear z1;  
clear z2;  
[x11,y]=ginput(1);  
if isempty(x11)  
x11 = ii;  
end
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
cc=find(x < x11);  
z1 = x11;  
z1 = cc(end);  
if (z1 <= 0)  
z1=1; end
```

```
[x12,y]=ginput(1);  
if isempty(x12)  
x12 = ii+700;  
end
```

```
cc=find(x < x12);  
z2 = x12;  
z2 = cc(end);  
% Convert to array arguments from seconds.
```

```
z1=fix(z1);  
z2=fix(z2);
```

```
% Find the Max and Min for the data in this window so all% the boxes  
will have the same axes.
```

```
start = fix(z1);  
stop = fix(z2);  
yymin = -1 * (max(-1*CHANNEL_1(start:stop)));  
yymin = (max(CHANNEL_1(start:stop)));  
if (yymin < 0)  
yymin=yymin*.95;  
else  
yymin=yymin*1.05; end
```

```
if (yymin < 0.0)  
yymin = yymin * 1.05;  
else  
yymin = yymin * 0.95;  
end
```

```
xxmin = x(start);  
xxmax = x(stop);
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
lonleft = (-1 * (max(-1 * CHANNEL_2(z1:z2))))-.1;
lonright = max(CHANNEL_2(z1:z2))+.1;
latbot = (-1 * (max(-1 * CHANNEL_3(z1:z2))))-.1;
lattop = max(CHANNEL_3(z1:z2))+.1;

set(ax4,'XLim',[lonleft lonright],'YLim',[latbot lattop],...
    'visible','on','clim',[-4500 2000]);
hold on;
set(ax3,'XLim',[lonleft lonright],'YLim',[latbot lattop]);
set(ax3_line,'Xdata',CHANNEL_2(z1:z2),...
    'Ydata',CHANNEL_3(z1:z2),...
    'visible','on','linestyle','-','marker','+', 'color','red'); set(ax5,'XLim',[lonleft
lonright],'YLim',[latbot lattop]);
set(ax1,'XLim',[xxmin xxmax],'YLim',[yymin ymax]);
set(ax1_line,'Xdata',x(z1:z2), 'Ydata',CHANNEL_1(z1:z2),...
'linestyle','none','marker','+', 'color',[1 .5 1]);
set(ax2,'XLim',[xxmin xxmax],'YLim',[yymin ymax],'visible','on');
set(ax2_line,'Xdata',x(z1:z2),'Ydata',CHANNEL_1(z1:z2),...
'linestyle','none','marker','+', 'color',[1 .5 1]);
title_str=sprintf ("%s samples %10d to %10d",dev_name,fix(z1),fix(z2));
set (get(ax2,'title'),'string',title_str);

% button for EDIT
% NOT used for SWD data
% Kept here for future use.

set(edit_button, 'Visible','on');

g = 'stopit';
while g == 'stopit'
    pause(1);
end

if (g == 'p')
    print -dps;
    g = 'stopit';
    while g == 'stopit'
        pause(1);
    end
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
    if (g == 'y')
        g = 'Y';
    end
end
if isempty(g)
    g = 'Y';
end
while g == 'e'
[xnew, ynew, but] = JGOFSGros(CHANNEL_1) ;

    xhold=xnew;
    yhold=ynew;

    xll=length(xnew);
    xuse = xll;
    xuse = 0;
    clear xnew ynew;

    for xlll=z1:z2,
        xuse = xuse+1;
        xnew(xuse)= x(xlll:xlll);
        if but(1) == 1
            ynew(xuse) = CHANNEL_1(xlll:xlll);
            znew(xuse) = CHANNEL_3(xlll:xlll);
        else
            ynew(xuse) = CHANNEL_2(xlll:xlll);
            znew(xuse) = CHANNEL_3(xlll:xlll);
        end
    end
end

for xllll=1:xll,
    for xlll=1:xuse,
        if (xhold(xllll) == xnew(xlll))
            if (isnan(yhold(xllll)) == 1)
                ynew(xlll) = NaN;
            end
        end
    end
end
end
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
title_str=sprintf ("%s samples %10d to %10d',dev_name,fix(z1),fix(z2));
title (title_str); hold off;
hold on;
if but(1) == 1

    set (ax1,'visible','on','XLim',[xxmin xxmax],'YLim',[yymin yymax]);
    set
(ax1_line,'visible','on','Xdata',xnew,'Ydata',CHANNEL_1(z1:z2),'marker','o',
color','red');
    hold on;
    set (ax2,'XLim',[xxmin xxmax],'YLim',[yymin yymax],...
'nextplot','add','visible','off');
    set (ax2_line,'Xdata',xnew,'Ydata',ynew,'linestyle','-','color','blue');

    else
    set(ax2_line,'Xdata',x(z1:z2),
'Ydata',CHANNEL_1(z1:z2),'marker','+','color','yellow');
    end
% button for SAVE EDIT
    set(save_edit_button, 'Visible','on');

    g = 'stopit';
    while g == 'stopit'
        pause(1);
    end
    if (g == 'p')
        print -dps;
        g = 'stopit';
        while g == 'stopit'
            pause(1);
        end
    if (g == 'y')
        g = 'Y';
    end
end

if g == 'k'
    xll=length(xnew);
    xstart=find(x == xnew(1));
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
for xlll = 1:xll,
    xuse = xstart-1+xlll;
    if but(1) == 1
        CHANNEL_1(xuse:xuse) = ynew(xlll);
        if (isnan(ynew(xlll)) == 1)
            CHANNEL_2(xuse:xuse) = NaN;
            CHANNEL_3(xuse:xuse) = NaN;
        end
    else
        CHANNEL_2(xuse:xuse) = ynew(xlll);
    end
end
set(save_edit_button, 'Visible','off');
g = 'stopit';
while g == 'stopit'
    pause(1);
end
end
if (g == 'p')
    print -dps;
    g = 'stopit';
    while g == 'stopit'
        pause(1);
    end
    if (g == 'y')
        g = 'Y';
    end
end
end
end
```

```
set(edit_button, 'Visible','off');
set(save_edit_button, 'Visible','off'); end
% END of EDIT Section
```

```
if (g == 'n')
    g = 'N';
    set(start_button, 'Visible','on');
end
if (g == 'N')
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
set(start_button, 'Visible','on');  
break  
end  
hold off;  
end
```

JGOFSGros.m

```
% JGOFSGros()  
% STB 09/02/03  
% script to remove small spikes from the data  
% called by JGOFSDisplay.m  
%%%%%%%%  
%          splines across spikes. You may want to decrease  
%          the value of nwid (experiment to see what is best)  
%          There are also other spline options. Linear may be  
%          sufficient. % originally written for Alan Chave
```

```
function [xnew,ynew,but] = JGOFSGros(e1)  
% script to remove small spikes from the data  
%%%%%%%%  
global xnew ynew x
```

```
x1=1;  
d1=1;  
clear d1;  
clear x1;  
clear xnew;  
clear ynew;  
[xs,ys,but]=ginput(2);  
  
cc=find(x < xs(1));  
xl=cc(end);  
cc=find(x < xs(2));  
xr=cc(end);  
nwid= 10;  
xhead = xl-nwid;  
xr-xl
```

SO GLOBEC Bathymetry
WHOI-2004-02

```
for i=1:nwid
  if but(1) == 1
    d1(i) = e1(xhead+i);
    d1(nwid+i) = e1(xr+i);
  else
    d1(i) = e2(xhead+i);
    d1(nwid+i) = e2(xr+i);
  end
  x1(i)= xhead+i;
  x1(nwid+i) = xr+i;
  xx(i)=x(xhead+i);
end;

for i= (x1+1):(xr-1),
  ynew(i - x1) = NaN;
xnew(i - x1) = x(i); end;

return;
%
```


REPORT DOCUMENTATION PAGE	1. REPORT NO. WHOI-2004-02	2.	3. Recipient's Accession No.
4. Title and Subtitle A High-Resolution Bathymetry Map for the Marguerite Bay and Adjacent West Antarctic Peninsula Shelf for the Southern Ocean GLOBEC Program		5. Report Date May 2004	
7. Author(s) S. T. Bolmer , R. C. Bearsdley, C. Pudsey, P. Morris, P. Wiebe, E. Hofmann, I. Anderson, A. Maldonado		8. Performing Organization Rept. No. WHOI-2004-02	
9. Performing Organization Name and Address Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543		10. Project/Task/Work Unit No.	
		11. Contract(C) or Grant(G) No. (C) OPP-0234163 (G)	
12. Sponsoring Organization Name and Address National Science Foundation		13. Type of Report & Period Covered Technical Report	
		14.	
15. Supplementary Notes This report should be cited as: Woods Hole Oceanog. Inst. Tech. Rept., WHOI-2004-02.			
16. Abstract (Limit: 200 words) One objective of the U.S. Southern Ocean Global Ocean Ecosystems Dynamics (SO GLOBEC) program is to gain a better understanding of the sea floor bathymetry in the program study area. Much of Marguerite Bay and the adjacent shelf west of the Antarctic Peninsula were poorly charted when the SO GLOBEC program started in 2000. Before the first SO GLOBEC cruise, an improved local area version (ETOPO8.2A) was created from the Smith and Sandwell (1997) topo_8.2.img 2-minute digital gridded bathymetry for the study area. The first SO GLOBEC mooring cruise on the <i>R/V Lawrence M. Gould</i> (March 2001) showed that the 2-minute spatial resolution of ETOPO8.2A did not resolve many of the canyons and abrupt changes in topography that characterize Marguerite Bay and the inner- to mid-shelf region. It also was not particularly accurate in the more uniform terrain regions. We then decided to collect as much multibeam bathymetry data as possible during the SO GLOBEC broad-scale survey cruises on the <i>R/VIB Nathaniel B. Palmer</i> and combine these data with all other available multibeam and trackline bathymetry data to construct a digital bathymetry database and map for the study area. The resulting database has high-resolution data over much of the shelf and parts of Marguerite Bay gridded at 2 seconds in latitude and 6 seconds in longitude spacing between 65° to 71° S and 65° to 78° W. This technical report describes the steps taken to assemble and construct this database and how to access the data via the Internet.			
17. Document Analysis a. Descriptors Bathymetry Marguerite Bay SO GLOBEC b. Identifiers/Open-Ended Terms c. COSATI Field/Group			
18. Availability Statement Approved for public release; distribution unlimited.		19. Security Class (This Report) UNCLASSIFIED	21. No. of Pages 76
		20. Security Class (This Page)	22. Price

DOCUMENT LIBRARY

Distribution List for Technical Report Exchange - July 1998

University of California, San Diego
SIO Library 0175C
9500 Gilman Drive
La Jolla, CA 92093-0175

Hancock Library of Biology & Oceanography
Alan Hancock Laboratory
University of Southern California
University Park
Los Angeles, CA 90089-0371

Gifts & Exchanges
Library
Bedford Institute of Oceanography
P.O. Box 1006
Dartmouth, NS, B2Y 4A2, CANADA

NOAA/EDIS Miami Library Center
4301 Rickenbacker Causeway
Miami, FL 33149

Research Library
U.S. Army Corps of Engineers
Waterways Experiment Station
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Marine Resources Information Center
Building E38-320
MIT
Cambridge, MA 02139

Library
Lamont-Doherty Geological Observatory
Columbia University
Palisades, NY 10964

Library
Serials Department
Oregon State University
Corvallis, OR 97331

Pell Marine Science Library
University of Rhode Island
Narragansett Bay Campus
Narragansett, RI 02882

Working Collection
Texas A&M University
Dept. of Oceanography
College Station, TX 77843

Fisheries-Oceanography Library
151 Oceanography Teaching Bldg.
University of Washington
Seattle, WA 98195

Library
R.S.M.A.S.
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149

Maury Oceanographic Library
Naval Oceanographic Office
Building 1003 South
1002 Balch Blvd.
Stennis Space Center, MS, 39522-5001

Library
Institute of Ocean Sciences
P.O. Box 6000
Sidney, B.C. V8L 4B2
CANADA

National Oceanographic Library
Southampton Oceanography Centre
European Way
Southampton SO14 3ZH
UK

The Librarian
CSIRO Marine Laboratories
G.P.O. Box 1538
Hobart, Tasmania
AUSTRALIA 7001

Library
Proudman Oceanographic Laboratory
Bidston Observatory
Birkenhead
Merseyside L43 7 RA
UNITED KINGDOM

IFREMER
Centre de Brest
Service Documentation - Publications
BP 70 29280 PLOUZANE
FRANCE