

# Potential Surfaces and Dynamics of the $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}^1\text{A}_1) \rightarrow \text{OH}(\text{X}^2\Pi) + \text{OH}(\text{X}^2\Pi)$ Reaction

M. Braunstein\*, R. Panfili, R. Shroll, and L. Bernstein

*Spectral Sciences, Inc.*

*Burlington, MA 01803*

\*Electronic mail address: [matt@spectral.com](mailto:matt@spectral.com)

## ABSTRACT

We present global potential energy surfaces for the three lowest triplet states in  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}^1\text{A}_1)$  collisions and present results of classical dynamics calculations on the  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}^1\text{A}_1) \rightarrow \text{OH}(\text{X}^2\Pi) + \text{OH}(\text{X}^2\Pi)$  reaction using these surfaces. The surfaces are spline-based fits of  $\sim 20,000$  fixed geometry *ab-initio* calculations at the CASSCF+MP2 level with a  $\text{O}(4s3p2d1f)/\text{H}(3s2p)$  one electron basis set. Computed rate constants compare well to measurements in the 1,000-2,500 K range using these surfaces. We also compute the total, ro-vibrationally resolved, and differential angular cross sections at fixed collision velocities from near threshold at  $\sim 4 \text{ km s}^{-1}$  ( $16.9 \text{ kcal mol}^{-1}$  collision energy) to  $11 \text{ km s}^{-1}$  ( $122.5 \text{ kcal mol}^{-1}$  collision energy), and we compare these computed cross sections to available space-based and laboratory data. A major finding of the present work is that above  $\sim 40 \text{ kcal mol}^{-1}$  collision energy ro-vibrationally excited  $\text{OH}(\text{X}^2\Pi)$  products are a significant and perhaps dominant contributor to the observed 1-5  $\mu$  spectral emission from  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}^1\text{A}_1)$  collisions. Another important result is that  $\text{OH}(\text{X}^2\Pi)$  products are formed in two distinct ro-vibrational distributions. The ‘active’ OH products are formed with the reagent O-atom, and their ro-vibrational distributions are extremely hot. The remaining ‘spectator’ OH is relatively ro-vibrationally cold. For the active OH, rotational energy is dominant at all collision velocities, but the opposite holds for the spectator OH. Summed over both OH products, below  $\sim 50 \text{ kcal mol}^{-1}$  collision energy, vibration dominates the OH internal energy, and above  $\sim 50 \text{ kcal mol}^{-1}$  rotation is greater than vibrational energy. As the collision energy increases, energy is diverted from vibration to mostly translational energy. We note that the present fitted surfaces can also be used to investigate direct collisional excitation of  $\text{H}_2\text{O}(\text{X}^1\text{A}_1)$  by  $\text{O}(^3\text{P})$  and also  $\text{OH}(\text{X}^2\Pi) + \text{OH}(\text{X}^2\Pi)$  collisions.

Report Documentation Page		Form Approved OMB No. 0704-0188
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.		
1. REPORT DATE <b>DEC 2004</b>	2. REPORT TYPE	3. DATES COVERED -
4. TITLE AND SUBTITLE <b>Potential Surfaces and Dynamics of the O(3P) + H2O(X1A1) -&gt; OH(X3Pi) + OH(X2PI) Reaction</b>		5a. CONTRACT NUMBER <b>F04611-03-C-0015</b>
		5b. GRANT NUMBER
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S) <b>M Braunstein; R Panfili; R Shroll; L Bernstein</b>		5d. PROJECT NUMBER <b>BMSB</b>
		5e. TASK NUMBER <b>R2FT</b>
		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Air Force Research Laboratory (AFMC),AFRL/PRSA,10 E. Saturn Blvd.,Edwards AFB,CA,93524-7680</b>		8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>		
13. SUPPLEMENTARY NOTES		

## 14. ABSTRACT

We present global potential energy surfaces for the three lowest triplet states in  $O(3P) + H_2O(X1A1)$  collisions and present results of classical dynamics calculations on the  $O(3P) + H_2O(X1A1)$  and  $OH(X2\&\#61520;) + OH(X2\&\#61520;)$  reaction using these surfaces. The surfaces are spline-based fits of ~20,000 fixed geometry ab-initio calculations at the CASSCF+MP2 level with a  $O(4s3p2d1f)/H(3s2p)$  one electron basis set. Computed rate constants compare well to measurements in the 1,000-2,500 K range using these surfaces. We also compute the total, ro-vibrationally resolved, and differential angular cross sections at fixed collision velocities from near threshold at ~4 km s<sup>-1</sup> (16.9 kcal mol<sup>-1</sup> collision energy) to 11 km s<sup>-1</sup> (122.5 kcal mol<sup>-1</sup> collision energy), and we compare these computed cross sections to available space-based and laboratory data. A major finding of the present work is that above ~40 kcal mol<sup>-1</sup> collision energy ro-vibrationally excited  $OH(X2\&\#61520;)$  products are a significant and perhaps dominant contributor to the observed 1-5  $\mu$ m spectral emission from  $O(3P) + H_2O(X1A1)$  collisions. Another important result is that  $OH(X2\&\#61520;)$  products are formed in two distinct ro-vibrational distributions. The 'active' OH products are formed with the reagent O-atom, and their ro-vibrational distributions are extremely hot. The remaining 'spectator' OH is relatively ro-vibrationally cold. For the active OH, rotational energy is dominant at all collision velocities, but the opposite holds for the spectator OH. Summed over both OH products, below ~50 kcal mol<sup>-1</sup> collision energy, vibration dominates the OH internal energy, and above ~50 kcal mol<sup>-1</sup> rotation is greater than vibrational energy. As the collision energy increases, energy is diverted from vibration to mostly translational energy. We note that the present fitted surfaces can also be used to investigate direct collisional excitation of  $H_2O(X1A1)$  by  $O(3P)$  and also  $OH(X2\&\#61520;) + OH(X2\&\#61520;)$  collisions.

## 15. SUBJECT TERMS

## 16. SECURITY CLASSIFICATION OF:


a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>
----------------------------------	------------------------------------	-------------------------------------

17. LIMITATION  
OF ABSTRACT18. NUMBER  
OF PAGES**45**19a. NAME OF  
RESPONSIBLE PERSON


## I. INTRODUCTION

Collisions of  $O(^3P)$  with  $H_2O(X, ^1A_1)$  give rise to spectral radiation observed from the long wave infrared ( $\sim 20\text{-}10\text{ }\mu$ )<sup>1-3</sup>, through the infrared ( $10\text{-}1\text{ }\mu$ )<sup>1-9</sup> and up to the vacuum ultraviolet ( $\sim 0.25\text{ }\mu$ )<sup>10-12</sup> depending on the relative collision energy. These emissions are important to characterize because they can be a significant background noise source for telescopes mounted on spacecraft in low-Earth-orbit or LEO. In LEO ( $\sim 300\text{ km}$  altitude and  $\sim 8\text{ km s}^{-1}$  relative atmospheric velocity),  $O(^3P)$ , which is the main component of the atmosphere at these altitudes, collides with  $H_2O(X, ^1A_1)$  from outgassing spacecraft surfaces or from spacecraft engine exhaust.<sup>4</sup>

The mechanisms that give rise to the observed radiation in  $O(^3P) + H_2O(X, ^1A_1)$  collisions remain elusive, however, because the rarefied conditions and large relative collision velocities necessary are difficult to achieve in the laboratory, and so experimental data remain sparse and difficult to interpret. Focusing on the  $1\text{-}5\text{ }\mu$  region, Bernstein et al.<sup>7</sup> have shown through analysis of space experiments and laboratory data that the observed radiation from  $O(^3P) + H_2O(X, ^1A_1)$  collisions may be consistent with forming ro-vibrationally excited OH through hydrogen abstraction,



and/or forming ro-vibrationally excited  $H_2O$  through direct collisional excitation,



depending on the collision velocity. Other studies<sup>6</sup> have attributed all such radiation from  $1\text{-}5\text{ }\mu$  to Reaction (2). It is particularly difficult to distinguish between (1) and (2) in the  $1\text{-}5\text{ }\mu$  spectral region because spectral observations remain the main source of data and  $OH(X, ^2\Pi)$  and  $H_2O(X, ^1A_1)$  have highly overlapping spectral distributions in this region. Furthermore, since spectral observations are only sensitive to internally excited products, the magnitude and energy dependence of these observations minimize contributions from products formed with little internal excitation, products which may in fact carry most of the total reaction cross section and dynamical information. In addition, theoretical examination of the reactions (1) and (2) has been


problematic because they involve computation of high-spin open shell wave functions which presents special difficulties for theory. There have been theoretical studies which used non-reactive  $O(^3P) + H_2O(X, ^1A_1)$  potential surfaces to examine the collisional excitation channel (Eq. 2).<sup>13-16</sup> Although these studies have had some success in explaining measurements in the 1-5  $\mu$  region, there remain major differences between these studies and measurements at certain collision velocities and significant differences between the theoretical results themselves.

Besides being important in LEO conditions, collisions of  $O(^3P) + H_2O(X, ^1A_1)$  are also important in  $H_2/O_2$  combustion chemistry.<sup>17</sup> The reverse of Reaction (1) is an important step in such chemistry. In fact, the surfaces involved in Reaction (1) are part of a family of triplet  $H_2O_2$  surfaces leading to  $H_2(X^1\Sigma_g^+) + O_2(X^3\Sigma_g^-)$  and  $H(^2S) + OOH(X^2A')$ , pathways critical in understanding such combustion chemistry. The potential surfaces involved in  $O(^3P) + H_2O(X, ^1A_1)$  collisions have been probed recently using transition state photoelectron spectroscopy,<sup>18,19</sup> which has given information about the barrier height and other transition state properties for Reaction (1). Instead of translational energy as in previous space and laboratory measurements, Ref. 20 used vibrational energy of deuterated  $H_2O$  reagents to overcome the large endothermicity of Reaction (1),



In these experiments, product internal state distributions were measured directly through laser-induced fluorescence, and therefore the usual ambiguities in interpreting the overlapping spectra of Reactions (1) and (2) were removed. These measurements yield rich dynamical information on the surfaces involved in Reaction (1).

For completeness, we note that collisions of  $O(^3P) + H_2O(X, ^1A_1)$  at very large relative collision velocities can produce electronically excited  $OH(A, ^2\Sigma^+)$ ,



which has been observed in space-based<sup>11</sup> and laboratory molecular beam measurements.<sup>10</sup> The electronically excited OH is apparently formed through high-lying conical intersections in the potential energy surfaces of the four-atom complex.<sup>12</sup> The surfaces involved in Reactions (1-2) are part of a large manifold of electronic states which give rise to the products of Reaction (4).

In this paper, we present global potential energy surfaces for the three lowest triplet adiabatic surfaces involved in  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  collisions and present results for classical dynamical calculations on these surfaces. The main purpose of the paper is to investigate the role of Reaction (1) in producing spectral radiation in the 1-5  $\mu$  region. By comparing to available measurements, we will assess the validity of these surfaces for further dynamical studies including Reaction (2) in order to form a complete picture of  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  collisions over a broad energy range. The present work is also of interest as a guide to the design and analysis of possible future hyperthermal O-atom +  $\text{H}_2\text{O}$  beam measurements. In these experiments, it should be possible to measure the OH products and their internal state distributions directly, rather than through spectral emission, over a range of energies near 64 kcal mol<sup>-1</sup> ( $\sim 8$  km s<sup>-1</sup> relative collision velocity).<sup>21</sup>

The surfaces we report for  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  are spline-based fits based on  $\sim 20,000$  fixed geometry *ab initio* calculations at the CASSCF+MP2 level with a  $\text{O}(4\text{s}3\text{p}2\text{d}1\text{f})/\text{H}(3\text{s}2\text{p})$  one electron basis set. The surfaces are used in quasi-classical trajectory (QCT) calculations to compute rate constants of reaction (1) as a function of temperature. The computed rate constants compare very well to available measurements. We also compute the total, ro-vibrationally resolved, and differential angular cross sections for reaction (1) with QCT on the fitted surfaces at fixed velocities from threshold at  $\sim 4$  km s<sup>-1</sup> to 11 km s<sup>-1</sup> relative collision velocity (122.5 kcal mol<sup>-1</sup>). We compare the computed cross sections to available space-based and laboratory data. We also compute cross sections for reaction (3) and compare these results to results of Ref. 20 using the fitted surfaces.

A major finding of the present work is that the ro-vibrationally excited  $\text{OH}(\text{X}^2\Pi)$  product from  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  collisions are a significant and perhaps dominant contributor to 1-5  $\mu$  spectral emission above  $\sim 40$  kcal mol<sup>-1</sup> collision energy. This finding is in contrast to most interpretations of the experimental data which attribute this radiation to collisional excitation of  $\text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  exclusively.<sup>6</sup> Another important result is that  $\text{OH}(\text{X}^2\Pi)$  products of reaction (1) are formed in two distinct ro-vibrational distributions. The ‘active’ OH products are formed with the reagent O-atom, and their ro-vibrational distributions are extremely hot. The remaining ‘spectator’ OH is relatively ro-vibrationally cold. Summed over both OH products, below  $\sim 50$  kcal mol<sup>-1</sup> collision energy, vibration dominates the OH internal energy. Above  $\sim 50$  kcal mol<sup>-1</sup>,

rotation is greater than vibrational energy, but the total product energy fraction is mostly in translation. We note that the present surfaces can also be used to investigate collisional excitation of  $\text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$ , reaction (2), or  $\text{OH}(\text{X}, ^2\Pi) + \text{OH}(\text{X}, ^2\Pi)$  collisions, the reverse of reaction (1), but we leave such work for a separate study.

The methods used for the potential fit, the analytic representation of the fit, the fitted surface properties for all three triplet surfaces, and the QCT dynamics methods used are discussed in Section 2. Section 3 presents the results of the QCT calculations and compares results to available experimental data. Section 4 discusses possible future work.

## II. METHODS

### A. Potential Surfaces

Collisions of  $\text{O}(^3\text{P})$  with  $\text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  give rise to three electronic states which adiabatically lead to  $\text{OH}(^2\Pi) + \text{OH}(^2\Pi)$ . These three lowest triplet adiabatic states are the focus of the present study, and we label them ‘state 1’, ‘state 2’, and ‘state 3’, where state 1 is the lowest energy state and state 3 is the highest. We fit these adiabatic states independently. The  $\Delta H$  of the  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(^1\text{A}_1) \rightarrow \text{OH}(^2\Pi) + \text{OH}(^2\Pi)$  reaction is  $+16.9 \text{ kcal mol}^{-1}$ . From rate constant measurements, the activation energy is  $\sim 17.0 \text{ kcal mol}^{-1}$  (Refs. 22,23) or a relative collision velocity of  $4.1 \text{ km s}^{-1}$ , which implies a very small energy barrier of  $\sim 0.1 \text{ kcal mol}^{-1}$ . We note that crossings of the three lowest triplet states with higher triplet states can lead to electronically excited  $\text{OH}(\text{A}^2\Sigma^+)$  with an endothermicity of  $+110.5 \text{ kcal mol}^{-1}$  (Ref. 12), but we do not consider such interactions here. Nor do we consider crossings with excited singlet oxygen,  $\text{O}(^1\text{D}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  states which can also lead to  $\text{OH}(^2\Pi) + \text{OH}(^2\Pi)$ .<sup>24</sup> We further note that by symmetry there is a fourth triplet leading adiabatically to  $\text{OH}(^2\Pi) + \text{OH}(^2\Pi)$ , but this state must originate from electronically excited triplets of  $\text{O} + \text{H}_2\text{O}$ . Furthermore,  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1)$  reaction channels leading to  $\text{H}_2(^1\Sigma_g^+) + \text{O}_2(^3\Sigma_g^-)$ ,  $\Delta H = -1.6 \text{ kcal mol}^{-1}$ , and  $\text{H}(^2\text{S}) + \text{OOH}(^2\text{A}')$ ,  $\Delta H = +51.0 \text{ kcal mol}^{-1}$ , are not addressed here. These channels go through high lying transition states ( $60\text{-}70 \text{ kcal mol}^{-1}$ )<sup>17</sup> and should be accessible at hyperthermal energies. Fully global surfaces for triplet  $\text{O} + \text{H}_2\text{O}$  should take these channels into account. A summary of the  $\text{O} + \text{H}_2\text{O}$  reagent, transition states based on the work of Ref. 17, and product states is shown in Figure 1. In the present study we

follow the three lowest triplet states of  $O(^3P) + H_2O(^1A_1)$  adiabatically to  $OH(^2\Pi) + OH(^2\Pi)$  products.

The approach taken to fit the three lowest triplet adiabatic surfaces follows closely the work of Wu et. al.<sup>25</sup> for the reaction  $H + H_2O \rightarrow OH + H_2$ , but with some modifications which we discuss below. For each triplet surface, we treat the full  $(3N-6) = 6$  dimensions by defining three ‘active’ internal coordinates which most affect motion along the reaction path (bond breaking and forming) and three remaining ‘spectator’ internal coordinates. The active coordinate potential is a 3-dimensional spline determined from a large set of *ab-initio* calculations spanning the range of active coordinate values. The spectator potential is determined for a fixed set of active coordinates at a geometry near the transition state. An approximation is made that the spectator coordinate potential does not vary much along the reaction pathway (changes in the active coordinates). A global analytic representation of a particular surface is defined as a sum of active and spectator parts. The global potential surface connects reagents to products smoothly and includes the proper symmetrization with respect to O-atom attack on either of the  $H_2O$  H-atoms. The active potential spline coefficients, spectator potential parameters and other parameters which determine the surface analytic representation are based on new *ab-initio* calculations of the  $O(^3P) + H_2O(^1A_1)$  potential at the CASSCF+MP2 level. The internal coordinates, analytic potential representation, *ab-initio* calculations, and properties of the fitted surfaces are described below.

#### A. 1. Internal Coordinates

Figure 2 shows the internal coordinates used. The active coordinates are the set  $\{x,y,z\}$  which most affect reaction going to OH products. The coordinate  $x$  is the OH bond of  $H_2O$  being broken,  $y$  is the OH bond being formed, and  $z$  is the angle formed from between the breaking and forming bond, atoms  $O_a-H_b-O_b$ . The spectator coordinates are the set  $\{u,\theta,\tau\}$ , where  $u$  is the OH bond length of  $H_2O$  which is not attacked by the incoming O,  $\theta$  is the  $H_a-O_a-H_b$  bond angle, and  $\tau$  is the ‘out-of-plane’ torsion angle defined as the angle between  $H_a$  and  $O_b$  when viewed along the  $O_a-H_b$  bond. The active coordinates have the range  $\{x: 0 \rightarrow \infty, y: 0 \rightarrow \infty, z: 0 \rightarrow 180^\circ\}$ , while the spectator coordinates have the range  $\{u: 0 \rightarrow \infty, \theta: 0 \rightarrow 360^\circ, \tau: 0 \rightarrow 180^\circ\}$ .



## A. 2. Analytic Representation of the Surfaces

We first define the “active+spectator” potential,  $V_{a+s}$ , as a sum of active,  $V_a$ , and spectator,  $V_s$ , parts,

$$V_{a+s}(x, y, z, u, \theta, \tau) = V_a(x, y, z)_{u_0, \theta_0, \tau} + V_s(u, \theta)_{x_0, y_0, z_0, \tau_0}, \quad (1)$$

The parameters,  $(x_0, y_0, z_0, u_0, \theta_0, \tau_0)$ , are particular values of the internal coordinates defined further below. The active potential is further broken down into a weighted sum of two parts that depend on  $\tau$ ,

$$V_a = W(\tau) \cdot V_{spline}(x, y, z)_{u_0, \theta_0, \tau=0} + (1 - W(\tau)) \cdot V_{spline}(x, y, z)_{u_0, \theta_0, \tau=180}. \quad (2)$$

The potential  $V_{spline}$  is defined as a 3D spline interpolation<sup>26</sup> on a grid of calculated *ab-initio* points in  $(x, y, z)$  described further below. Notice that the spline potential only depends on planar geometries of the O + H<sub>2</sub>O system and for purposes of the spline interpolation,  $z$  is considered to be continuous from 0 to 360°.  $W(\tau)$  is a weighting function that weights the  $\tau=0^\circ$  and  $\tau = 180^\circ$   $V_{spline}$  values and provides an interpolation of the active potential for non-planar geometries,  $\tau \neq 0^\circ, 180^\circ$  based on the planar  $V_{spline}$  values.  $W(\tau)$  is defined

$$W(\tau) = \sum_{m=0}^6 a_m \cos^m \tau, \quad (3)$$

where the parameters  $\{a_m\}$  are parameters determined from a fit of the *ab-initio* data described below. The spectator potential,  $V_s$ , is a sum of two terms,

$$V_s = V_u(u)_{x_0, y_0, z_0, \tau_0, \theta_0} + V_\theta(\theta(x))_{y_0, z_0, \tau_0}. \quad (4)$$

Here  $V_u$  is a Morse potential given by,

$$V_u(u) = D_e \left[ 1 - e^{-b(u-u_o)} \right]^2, \quad (5)$$

with fit parameters  $\{D_e, u_o, b\}$ . The other part of the spectator potential,  $V_\theta$ , is given by,

$$V_\theta(\theta(x)) = \left( A \cdot (\cos \theta - \cos \theta_1)^4 + B \cdot (\cos \theta - \cos \theta_1)^3 \right) \cdot e^{-\alpha(x-x_1)} \quad (6)$$

where

$$\theta_1 = C + D \cdot e^{-\alpha'(x-x_1)} \quad (7)$$

with fit parameters  $\{A, B, C, D, \alpha, \alpha', x_1\}$ . The functional form of Eq. (6) is slightly different from the study of Wu et al.<sup>25</sup> to better match the *ab-initio* O + H<sub>2</sub>O surfaces.

Since functional forms for the H<sub>2</sub>O reagents<sup>27</sup> and OH products<sup>28</sup> have already been developed which match experimental data very well, we incorporate these known potentials in the present global fit. We do this in a similar fashion to Wu et al.<sup>25</sup> by employing switching functions that turn on and off the reagent, product, and  $V_{a+s}$  potentials in appropriate regions. We choose the following forms for these switching functions:

$$S_r = \frac{1}{2}[1 + \tanh(r_0 \cdot (y - r_1))] \quad (8a)$$

$$S_q = \frac{1}{2}[1 + \tanh((y - x))] \quad (8b)$$

$$S_p = \frac{1}{2}[1 + \tanh(p_0 \cdot (x - p_1))] \quad (8c)$$

with fit parameters  $\{r_0, r_1, p_0, p_1\}$ . Defining  $V_r$  to be the isolated H<sub>2</sub>O potential, and  $V_p$  as the potential of two isolated OH products, the global potential can then be expressed as,

$$V_{unsym} = S_r \cdot S_q \cdot V_r + (1 - S_r \cdot S_q - S_p \cdot (1 - S_q)) \cdot V_{a+s} + S_p \cdot (1 - S_q) \cdot V_p \quad (9)$$

An undesired side-effect of separating active and internal coordinate potentials is that the potential,  $V_{unsym}$ , treats the hydrogen atoms on the water reagent in an unsymmetrical fashion. This means that the potential  $V_{unsym}$  is different depending on whether the incoming oxygen attacks H<sub>b</sub> rather than H<sub>a</sub>, which is unphysical. To compensate, we follow Wu et al.<sup>25</sup> and introduce the switching function  $S_s$ , which makes attack on either hydrogen equivalent.

$$S_s = \frac{1}{2}[1 - \tanh(s_0 \cdot (R_{O_bH_a} - R_{O_bH_b}))], \quad (10)$$

where  $\{s_0\}$  is a fit parameter. The full global symmetrized potential is then the weighted sum of two unsymmetrized parts,

$$V_{sym} = S_s \cdot V_{unsym}(H_a - O_a - H_b + O_b) + (1 - S_s) \cdot V_{unsym}(H_b - O_a - H_a + O_b) \quad (11)$$

where  $V_{unsym}(H_a - O_a - H_b + O_b)$  is the potential of the atoms in their original configuration, and  $V_{unsym}(H_b - O_a - H_a + O_b)$  is the potential with the H atoms switched. This is essentially an averaging of two geometric configurations.

Table I gives the values of all the parameters for the potential for all three electronic states and how they were derived. Many of the values were obtained from *ab-initio* calculations described in the next section.

TABLE I. Surface fit parameters. All values are a.u.

Fit Parameters	Values (a.u.)	How Determined
“Transition State” $\{x_0, y_0, z_0, u_0, \theta_0, \tau_0\}$	(2.3501, 2.0558, 141.06°, 1.8287, 104.55°, 0.0)	Ref. 17
$W(\tau)$ , $\{a_0, \dots, a_6\}$ state1 state2 state3	( 0.672, 0.649, -0.311, -0.347, 0.373, 0.198, -0.234) (1.47, -0.296, -0.387, 0.511, -0.0724, 0.263, -0.496) (3.63, 0.480, -5.74, -0.614, 5.15, 0.627, -2.55)	Fit of <i>ab-initio</i> data
$V_u \{D_e, b, u_0\}$ state1 state2 state3	(0.1628, 1.2200, 1.8500) (0.1757, 1.2644, 1.8287) (0.1574, 1.3000, 1.8286)	Fit of <i>ab-initio</i> data
$V_\theta \{A, B, C, D, \alpha, \alpha', x_1\}$ state1 state2 state3	(0.7400, 0.065, 99.0, 110., 1.9, 1.6, 2.11) (0.221, 0.0164, 98.7, 3.3, 1.0525, 0.8, 2.232) (0.137, -0.0416, 103.5, 16., 2.307, 0.1, 2.35)	Fit of <i>ab-initio</i> data
switching functions $\{r_0, r_1, p_0, p_1, s_0\}$	(4.0, 6.0, 3.0, 4.0, 3.0)	Ref. 25
OH product $\{D_e, b, u_0\}$	(0.1698, 1.21, 1.85)	Ref. 28
H <sub>2</sub> O reagent	many	Ref. 27
$V_{spline}$ coefficients	many	Fit of <i>ab-initio</i> data

### A. 3. Electronic Structure Calculations

Most of our *ab-initio* calculations were based on a 10 electron 8 orbital (10e8o) CASSCF + MP2 (Complete-Active-Space Self-Consistent-Field + Second Order Perturbation Theory) treatment with a (4s3p2d1f) O-atom basis and 3s2p H-atom basis for a total of 90 basis functions. The (10e8o) CASSCF includes all excitations of eight electrons within the space of an initial configuration of two bonding, two lone-pair, two singly occupied, and two un-occupied anti-bonding orbitals. (The 1s and 2s oxygen orbitals are not included in the CASSCF space.) This totals 1512 configuration state functions. We state-average the three lowest electronic states and perform all these calculations with no symmetry. This choice of CASSCF space ensures that the three electronic states treated will be degenerate on the reagent and product sides of the reaction as they should be, and that the CASSCF space includes bonding and anti-bonding orbitals so that OH bonds can break and form in an even handed way along the reaction path. A summary of the electronic energies and energy differences of the reagents, products, and at the lowest transition state geometry (as determined from the study of Ref. 17) are given in Table II. All calculations were done with the GAMESS electronic structure package.<sup>29</sup>

TABLE II. Computed CASSCF-MP2 Energies.  $T_e$  is defined to be the calculated energy difference with respect to  $O + H_2O(R_e)$ , Energy-Energy( $O+H_2O(R_e)$ ).

Geometry	Energy (a.u.)	$T_e$ (kcal mol <sup>-1</sup> )
$O + H_2O(R_e)$	-151.289431	0.
$OH(R_e) + OH(R_e)$	-151.264587	15.59
Transition State (Following Ref. 17, Table I)		
state 1	-151.266063	14.66
state 2	-151.250827	24.22
state 3	-151.227117	39.10

The experimental  $T_e$  leading to OH + OH is 14.7 kcal mol<sup>-1</sup> which is 0.9 kcal mol<sup>-1</sup> below the present calculated *ab initio* results. Because we use switching functions to smoothly connect to the reagent and product parts of the potential, we have defined the  $T_e$  for the present fitted potential to be the experimentally measured  $T_e$  of 14.7 kcal mol<sup>-1</sup>. Computing transition state geometries at the CASSCF-MP2 level is problematic, especially with state averaging. So we have adopted the transition state geometry reported in Ref. 17 which uses a basis set extrapolation technique within a QCISD(T) treatment. Ref. 17 obtained a  $T_e$  for the first transition state of state 1 of +1.61 kcal mol<sup>-1</sup> above the OH + OH limit, while the present surface would be nearly equal in energy to the OH + OH limit for the lowest triplet. Deyerl et al.<sup>19</sup> obtain a lowest triplet transition state very similar in geometry to Karkach et al. within a QCISD(T)/6-311++G(3df,2p) treatment which is +0.10 kcal mol<sup>-1</sup> above the OH + OH limit (electronic energy -151.315271). As discussed further below, the second and third lowest triplets have potential surfaces quite similar to the lowest triplet, except that the barrier heights are larger by ~10 and ~15 kcal mol<sup>-1</sup> respectively. The second and third states should therefore play a role at hyperthermal energies.

To obtain the spline fit parameters, GAMESS CASSCF+MP2 calculations were done on a 34x34x14 grid of x,y, and z points for a total of 16,184 points. For these calculations ( $\theta=104.5^\circ$ ,  $u=1.8085$  a.u.,  $\tau=0^\circ$ ). The values of  $u$  and  $\theta$  are slightly different from the transition state geometry adopted here, but these differences will have minimal impact on the fit. The x and y coordinate grid spanned the range 1.3228 a.u. to 7.559 a.u. with spacings of 0.1890 a.u. in each direction. The z-coordinate values extend from 50° to 310° with a spacing of 20°. We fit the computed energies to a 3D cubic spline (with tension) after Ref. 26. For geometries lying outside the x- and y-value grid points, we map the results to the nearest extreme grid value. For z-values outside the calculated range, we perform a linear interpolation between the nearest extreme grid values. Also, for  $z \in [50^\circ, 70^\circ]$  we found it necessary to use a multi-dimensional linear (rather spline fit) to avoid un-physical oscillations in the interpolated potential. We note the cubic spline interpolation is simultaneously performed for all three active coordinates, so the z coordinate is treated on equal footing with the x and y coordinates. We also note that for subsequent dynamics calculations, splines are especially efficient for evaluations of the potential

as all interpolations are performed locally while the value of the function and its first derivative remain continuous from one grid point to the next.

To obtain the  $W(\tau)$  fit parameters for each state, we performed CASSCF+MP2 calculations as described above fixing all internal coordinates except  $\tau$  to the transition state geometry. We then varied  $\tau$  in steps varying from 5 to 20 degrees from 0 to 180. Similarly, to obtain the  $V_u$  fit we performed CASSCF+MP2 calculations fixing all internal coordinates except  $u$  to the transition state geometry and varied  $u$  over 61 points in the range 1.22 to 9.44 a.u. For the  $V_\theta$  fit we performed a series of CASSCF + MP2 calculations fixing all internal coordinates except  $\theta$  and  $x$  at their transition state geometries. We varied  $\theta$  and  $x$  on a two-dimensional grid, varying  $\theta$  every 5 degrees from 0 to 360 degrees and varying  $x$  from 1.16 to 8.019 a.u. over 24 unevenly spaced steps. Since  $V_{spline}$  already contains a dependence on  $x$ , we fit the residual dependence,  $V_{ab-initio} - V_{spline}$  in order to isolate the contribution of varying the spectator coordinate  $\theta$ .

#### A. 4. Properties of Potential Surfaces

The potential surface fits for the three lowest triplet states of the O + H<sub>2</sub>O system are shown in Figure 3 for planar geometries. In figure 3, the spectator coordinates,  $\{u, \theta, \tau\}$ , are fixed at their transition state values as given in Table I to isolate the  $V_a$  portion of the potential, and the  $z$ -coordinate is fixed at 80<sup>0</sup>, 140<sup>0</sup> and 240<sup>0</sup>. The active coordinate  $\{x\}$  varies along the abscissa the active coordinate  $\{y\}$  varies along the ordinate. Large values of  $y$  correspond to the separated O + H<sub>2</sub>O reagents and large values of  $x$  correspond to separated OH + OH products. The transition state barrier for this fitted potential occurs near  $z=140^0$  for all three surfaces and its energies are ~15, ~24, and ~39 kcal mol<sup>-1</sup>, respectively with respect to O + H<sub>2</sub>O(R<sub>e</sub>). The transition state saddle point for the lowest state occurs near  $x \approx 2.4$  and  $y \approx 2.1$  a.u., so that the OH bond that is breaking along  $x$  is larger than the OH bond that is forming along  $y$ .

Figure 4 shows the fitted potential  $V_a$  for non-planar geometries for all three fitted surfaces. In Figure 4, all coordinates are fixed at their transition state values as shown in Table I, and the torsion angle  $\tau$  is varied from 0 to 180<sup>0</sup> to isolate the effect of out-of-plane motion. For the lowest electronic state there is a preference for 0<sup>0</sup> (cis) geometries over 180<sup>0</sup> (gauche) by ~7 kcal mol<sup>-1</sup>. There appears to be an avoided crossing near  $\tau=130^0$  between the two lowest electronic

states. For the third electronic state there is an energy maximum near  $\tau=90^\circ$  and the 0 and  $180^\circ$  geometries have about the same energy. The barriers for out-of-plane motion are generally small.

Figure 5 shows the  $V_\theta$  component of the spectator potential, which is sensitive to the reagent  $\text{H}_2\text{O}$  internal bond angle, for all three fitted surfaces. For these plots, all internal coordinates except  $\theta$  are fixed at their transition state values as given in Table I. For all three surfaces there is a barrier at linear  $\text{H}_2\text{O}$  geometries, which is  $\sim 30 \text{ kcal mol}^{-1}$  for the lowest electronic state. There are local minima at  $\theta \approx 104.5^\circ$  and  $\theta \approx 255.5^\circ$  which correspond to cis and trans geometries of the combined  $\text{O} + \text{H}_2\text{O}$  system where the reagent is close to its isolated equilibrium bond angle.  $V_\theta$  is constructed to be zero here as the  $V_a$  part of the potential already treats these geometries. At small and large values of  $\theta$  correspond to ‘small angle’ geometries of the reagent  $\text{H}_2\text{O}$ , and so are dominated by the nuclear-nuclear repulsions of the  $\text{H}_2\text{O}$  hydrogens. Figure 5 is for fixed a value of  $x$ . As  $x$  increases and products are formed,  $V_\theta$  retains the same shape but decreases in magnitude for all three states as expected.

Figure 6 shows the full potential for all three fitted states,  $V_{\text{sym}}$  for planar geometries. In Figure 6, the coordinates  $\{x,u,\theta,\tau\}$  have been fixed at their transition state values as shown in Table I, and  $\{y,z\}$  have been allowed to vary. This view shows the potential for a fixed  $\text{H}_2\text{O}$  reagent at varying angles of approach of the O-atom. The chief feature of these surfaces is a well near angles of approach corresponding to cis geometries of the  $\text{O} + \text{H}_2\text{O}$  system, in regions centered near  $z \approx 140^\circ$  for the lowest electronic state. For the higher electronic states the well is not so deep and  $z \approx 180^\circ$ . These  $z$  angles are the preferred angles of approach of the O-atoms for chemical reaction. As the reagent OH bond lengthens ( $x$  increases) and the product OH bond forms ( $y$ -decreases) the shallow wells in Figure 6 deepen. Generally, other angles of approach are repulsive, except at large distances where there are shallow long-range type interactions of the approaching Oxygen and  $\text{H}_2\text{O}$  hydrogens.<sup>30</sup> Some of this long-range attraction corresponds to geometries affected by our separation of active and spectator coordinates and the way the present potential is symmetrized, and so we do not expect these regions to be as accurate as the transition state geometry regions which are fit by splines alone.

## II. B. Quasi-classical trajectory (QCT) calculations

We use the VENUS classical trajectory code<sup>31</sup> with an added interface modification in order to use our fitted potential surfaces and derivatives to generate rate constants and cross sections. For all calculations we use a fixed time step of  $\delta t = 1.e-16$  seconds and a maximum impact parameter  $b_{\max}$  ranging from 5.7 a.u. to 7.56 a.u. To generate rate constants at fixed temperatures, we perform 1.e5 to 5.e5 trajectories per electronic state per temperature. To generate cross sections at fixed translational energies, we fix the reagent water in its ground ro-vibrational state and perform 1.e5 to 5.e5 trajectories per each electronic state per energy. To obtain total rate constants or cross sections we add the trajectory contributions from each electronic state together and multiply by 1/3, the electronic degeneracy factor. For final state-resolved quantities the OH(v,j) states are binned in the usual way as outlined in Ref. 31. Final H<sub>2</sub>O products are binned according to the amount of internal energy in rotation and vibration. We note that our results do not adjust for zero-point energy of the reagents, which will sometimes tend to make such classical dynamical results more ‘reactive’ than the potential actually allows.

### III. RESULTS AND DISCUSSION

#### A. Rate constants at fixed temperatures for $O(^3P) + H_2O(X, ^1A_1) \rightarrow OH(X, ^2\Pi) + OH(X, ^2\Pi)$

Figure 7 shows the computed QCT rate constant,  $k_{rxn}$ , for the reaction,  $O(^3P) + H_2O(X, ^1A_1) \rightarrow OH(X, ^2\Pi) + OH(X, ^2\Pi)$ , as a function of temperature together with data from two measurements.<sup>22,23</sup> Calculations were performed from 1,000 K to 3,500 K at intervals of 500 K. There is generally good agreement between the measurements.<sup>22,23</sup> (especially Ref. 23), Generally, the present results are within experimental uncertainties (about a factor 2). Usually, neglect of a zero point energy correction as in the present QCT calculations would tend to over-estimate the rate constant, but there may be a compensating error in the neglect of a tunneling correction. We do not report results below 1,000 K, because of the difficulty of obtaining acceptable QCT statistics. The separate electronic state contributions to the QCT computed rate constant are also shown. The dominant contribution is from the lowest energy electronic state, state 1, as expected. The second lowest electronic state also makes an appreciable contribution at all temperatures studied and at the highest temperatures has a contribution within a factor of two of state 1.



## B. Cross sections at fixed collision energies for $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1) \rightarrow \text{OH}(\text{X}, ^2\Pi) + \text{OH}(\text{X}, ^2\Pi)$

### 1. Total and vibrationally resolved cross sections

Figure 8 shows the total calculated QCT ‘OH cross section’,  $\sigma_{OH}$ , for the reaction,  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1) \rightarrow \text{OH}(\text{X}, ^2\Pi) + \text{OH}(\text{X}, ^2\Pi)$ , as a function of collision energy along with component contributions from the three lowest triplet electronic states. Calculations were performed at the following collision energies: 16.2, 18.7, 20.5, 25.3, 30.6, 36.4, 49.6, 64.8, 82.0, 101.2, and 122.5 kcal mol<sup>-1</sup> which correspond to relative collision velocities of 4.0, 4.3, 4.5, 5.0, 5.5, 6.0, 7.0, 8.0, 9.0, 10.0, and 11 km s<sup>-1</sup>. We define the OH cross section,  $\sigma_{OH}$ , in the following way,

$$\sigma_{OH} = 2\sigma_{rxn} = \pi b_{\max}^2 \frac{2N_r}{N_{tot}}, \quad (12)$$

where  $\sigma_{rxn}$  is the reaction cross section,  $N_r$  is the number of reactive QCT trajectories and  $N_{tot}$  is the total number of trajectories. The cross section,  $\sigma_{OH}$ , is more convenient for comparisons to most of the available measurements which often report absolute cross sections by detecting and adding together signals from both OH products formed in the reaction. The OH cross section is relatively small near threshold and increases with energy. By 30 kcal mol<sup>-1</sup> collision energy, the cross section is relatively constant, although it continues to increase with collision energy. This implies that future measurements need to start below 5 km s<sup>-1</sup> to capture the rapid increase of the excitation function near threshold. As the energy increases past the energy barriers of state 2 and state 3, they begin to contribute significantly to the cross section. At the highest collision energies studied, the contribution from state 3 becomes comparable to that of state 2, where they are about 2.5 times smaller than state 1.

Figure 9a shows the QCT vibrationally resolved OH cross sections,  $\sigma_{OH}(v)$ , for the reaction,  $\text{O}(^3\text{P}) + \text{H}_2\text{O}(\text{X}, ^1\text{A}_1) \rightarrow \text{OH}(\text{X}, ^2\Pi) + \text{OH}(\text{X}, ^2\Pi)$ , at 25.3, 49.6, 64.8, 82.0, and 122.5 kcal mol<sup>-1</sup> collision energy or 5.0, 7.0, 8.0, 9.0, and 11.0 km s<sup>-1</sup> collision velocity, respectively, where we have summed the contributions from both OH products in a particular vibrational state (Eq. 12). At 5, 7, and 8 km s<sup>-1</sup> collision velocity, the distributions are well fit by thermal populations with temperatures of 1,900 K, 2,700 K, and 3,100 K respectively. At 9 and 11 km s<sup>-1</sup> collision

velocity we can fit the lower OH(v) states both with thermal temperatures of  $\sim 3,500$  K. The highest OH(v) populations for these higher velocities begin to show a different type of distribution. However these cross sections are very low, and proper fitting of these population requires better statistics. Figure 9b and 9c show the component ‘active’ and ‘spectator’ vibrationally resolved OH cross sections. The active OH products are from the incoming reagent atomic oxygen, and the spectator OH products are made from the oxygen in the reagent H<sub>2</sub>O, so that the total OH cross section (Fig. 9a) for a given vibrational level is the sum of active and spectator OH cross sections for that level. Although measurements of the active and spectator OH cross sections would probably only be available through detailed molecular beam isotopic studies, the computed active and spectator OH cross sections offer insight into the dynamics of the collision process. For example, it is clear that the active and spectator vibrationally resolved OH cross sections are quite distinct. The spectator cross sections are significantly colder than the active OH populations. The spectator OH’s mainly populate OH(v’=0,1) levels, with the v’=1 relative populations significantly less than that of the active OH(v’=1) populations. The spectator OH(v’>1) populations are almost negligible. This means that the total OH(v’>0) populations are due almost exclusively to the active OH, and the total OH(v=0) populations are mainly due to the spectator OH. The OH bond being formed in the reaction takes up most of the vibrational energy in the products. Joint active/spectator distributions would reveal more detailed information, but would require substantially better statistics.

The vibrationally resolved results of Figure 9a can be compared with data. Fixed energy spaced-based and laboratory measurements for  $O(^3P) + H_2O(X, ^1A_1) \rightarrow OH(X, ^2\Pi) + OH(X, ^2\Pi)$  focus on the detection of infrared radiation from vibrationally excited total OH (active + spectator, Fig. 9a) products summed over the populated vibrational states. Since the available spaced-based and laboratory measurements do not measure vibrationally resolved cross sections directly, it is necessary to briefly review the nature of these experiments to make appropriate comparisons with the present results.

As outlined in Ref. 7, space-based measurements (such as those of Refs. 5, 6 and 9) are characterized by a steady-state outgassing or exhausting source of H<sub>2</sub>O with a known or assumed rate,  $N_{H_2O}$  (number of H<sub>2</sub>O per second), interacting with atomic oxygen in the atmosphere. The signal is usually observed through a somewhat opaque atmosphere integrated over a wavelength

band and characterized by a transmittance,  $T_{\Delta\lambda}$ . The observed steady state intensity,  $I_{\Delta\lambda}^{space}$  (photons/second), can be written,

$$I_{\Delta\lambda}^{space} \approx \left[ \frac{\sigma^*}{\sigma_{tot}} \right] N_{H_2O} T_{\Delta\lambda}, \quad (13)$$

where  $\sigma_{tot}$  is the total (reactive and non-reactive) collision cross section for  $O + H_2O$  and  $\sigma^*$  is the photon excitation cross section which includes all contributions due to photon cascades from multi-quantum excited states. For detection in the 2-5  $\mu$  spectral region where most of the relevant optical measurements take place for this system, the intensity will be sensitive to single vibrational quantum changes, so we write  $\sigma^*$  as,

$$\sigma^* = \sum_{species} \sum_{v=1} v \sigma_v^{species}, \quad (14)$$

where  $v$  is the vibrational quantum number,  $\sigma_v^{species}$  is the cross section to a particular vibrational state of either collisionally excited  $H_2O$  or reaction product  $OH$  radiating in the detection band. It is important emphasize that observations usually include contributions from both species, and so the first sum extends over  $OH$  and  $H_2O$ . The second sum extends over all possible excited vibrational levels of a given species. Space-based measurements usually assume or otherwise employ models to estimate values for  $N_{H_2O}$ ,  $T_{\Delta\lambda}$  and  $\sigma_{tot}$ , and from the observed intensity,  $I_{\Delta\lambda}^{space}$ , report a photon excitation cross section,  $\sigma^*$ . Therefore, uncertainties in the estimated values of  $N_{H_2O}$ ,  $T_{\Delta\lambda}$  and  $\sigma_{tot}$  are reflected in the reported  $\sigma^*$ .

On the other hand, ground-based laboratory measurements involving optical detection typically use pulsed beam sources. Signals,  $I_{\Delta\lambda}^{lab}$ , from such measurements are related to the photon excitation cross section in the following way,

$$I_{\Delta\lambda}^{lab} \propto \sum_{species} \sum_{v=1} A_v^{species} \sigma_v^{species}, \quad (15)$$

where  $A_v^{species}$  are Einstein coefficients (photons/second) for single photon radiative decay. Because contributions from separate species and from individual vibrational levels usually cannot be distinguished, lab measurements typically report a single value for the right hand side

double sum of Eq. (15) at fixed O + H<sub>2</sub>O collision velocities, and for convenience we denote this quantity as  $A\sigma$  (length<sup>2</sup>/second). We note that these laboratory measurements are often difficult to carry out and calibrate absolutely with high precision due to uncertainties in reactant concentrations and the inability of optical systems to capture all collision produced photons within a limited field of view.

Table III gives two spaced-based results for the photon excitation cross section (Eq. 14), two laboratory results for the product cross sections times Einstein A (Eq. 15), and the corresponding quantities derived from the present calculated vibrationally resolved total OH cross section (Fig 9a). These quantities follow the analysis of Ref. 7 and assume  $A_v^{OH} = \nu \cdot 17 \text{ sec}^{-1}$ . It is estimated that the experimental results are uncertain by a factor of  $\sim 2$  to  $\sim 3$ . At  $4.3 \text{ km s}^{-1}$ , the laboratory results of Ref. 1 are about 12 times larger than the presently computed contribution from excited OH. This collision energy is too low to excite OH populations. Therefore, the experimental signal is nearly all H<sub>2</sub>O, collisionally excited in the  $\nu_1$  and  $\nu_3$  stretching modes. At  $8 \text{ km s}^{-1}$ , well above the reaction energy barrier to make OH, the measured space-based<sup>6</sup> results are about a factor of 2 smaller and laboratory results<sup>4</sup> about equal to the presently computed OH contribution to the overall signal. At  $11 \text{ km s}^{-1}$  there is good agreement between the space based combined signal and presently computed contribution from OH. The present results therefore imply that at and above  $8 \text{ km s}^{-1}$  collision velocity, the OH contribution is at least comparable and perhaps larger than the H<sub>2</sub>O contribution to the observed infrared signals. We note that the H<sub>2</sub>O collisional excitation cross sections have been computed in previous studies. Although these studies used the same non-reactive potential surface, the quantum scattering results of Ref. 14 differ by orders of magnitude from the classical scattering results of Ref. 13 even at high velocities, so it is difficult to assess the possible relative contributions from collisionally excited water. We note however, that at  $8 \text{ km s}^{-1}$  collision velocity there is fairly good agreement between the quantum results for collisional excitation of H<sub>2</sub>O and the measurements of Ref. 4. With the presently computed contribution from OH, the theoretically derived combined H<sub>2</sub>O/OH would still be within the experimental error bounds.

TABLE III. Excitation cross sections derived from spaced-based, laboratory, and presently calculated results for  $O(^3P) + H_2O(X^1A_1)$  collisions to produce radiation from 2-5  $\mu$ .

Relative Collision velocity (km s <sup>-1</sup> )	$\sum_{v=1} v \sigma_v^{OH}$ (Theory: a.u.)	$\sum_{species} \sum_{v=1} v \sigma_v^{species}$ (Exp: a.u.)	$\sum_{v=1} A_v^{OH} \sigma_v^{OH}$ (Theory: a.u./second)	$\sum_{species} \sum_{v=1} A_v^{species} \sigma_v^{species}$ (Exp. a.u./second)
4.3			0.09 (Ref. 1)	1.07
~8.	1.28	0.54 (Ref. 6)	21.75 (Ref. 4)	18.20
11	4.41	3.57 (Ref. 9)		

We note that the vibrational temperature of 3,100 K obtained from the present calculations at 8.0 km s<sup>-1</sup> is quite different from what has been conjectured by Bernstein et al.<sup>7</sup> in their fit of observed spectra of Ref. 4 obtained in hyperthermal O-atom beam experiments at similar collision energies. Bernstein et al.<sup>7</sup> derive a vibrational temperature of 7,000 K, which was mostly driven by the observed large OH overtone signals. The present results cannot reproduce the apparently large overtone populations observed in Ref. 4.

## 2. Rotationally resolved cross sections

Figure 10 shows the rotationally resolved total OH cross at 5.0, 7.0, 8.0, 9.0, and 11.0 km s<sup>-1</sup> relative collision velocity for  $O(^3P) + H_2O(X,^1A_1) \rightarrow OH(X,^2\Pi) + OH(X,^2\Pi)$ , where the OH(v')

populations have been summed over. These cross sections show a bi-model distribution, a narrow low OH( $j' < 10$ ) peak and a wide almost flat OH( $j'$ ) extending to very high OH( $j'$ ) which eventually falls off as the energetic limit is reached. As shown in Fig. 11 for 8 km s<sup>-1</sup> collision velocity, the two underlying rotational distributions come from the active and spectator OH's. The active OH is much more rotationally excited than the spectator OH, so that the active OH receives the vast majority of the internal energy imparted to products in the collision. The superimposed oscillatory structure for low active OH( $j$ ) populations appears to be real and not caused by statistical uncertainty.

We have fit the separate active and spectator OH( $j'$ ) rotational populations to a distribution derived from surprisal theory,<sup>32</sup>

$$P(vj) \propto \exp(\theta_r E_{vj}) (2j+1) \sum_{ml} (2l+1) (E - E_{vj} - E_{ml})^{1/2}, \quad (16)$$

where ( $vj$ ) are the vibrational and rotational quantum numbers, respectively, for OH product being fit, ( $ml$ ) are the vibrational and rotational quantum numbers, respectively, for the other OH product,  $E$  is the total energy available to be distributed to products in the reaction,  $\theta_r$  is a surprisal parameter, the sum of ( $ml$ ) extends over all the available energy states, and we have set  $\nu=0$ . The results are shown in Fig. 11 for separate fits to the active and spectator OH's, where  $\theta_r = 0.0$  (statistical) for the active OH and  $\theta_r = -0.41$  for the spectator OH. We note that the peak OH( $j$ ) products correspond to rotational temperatures of 11,400 K and 1,100 K for the active and spectator OH's respectively. The analysis of Ref. 7 of the data in Ref. 4 derives a rotational temperature for total OH of 1,500 K. This temperature should be dominated by the active vibrationally excited OH for which we calculate a much higher rotational temperature.

### 3. Product energy disposal

Figure 12a shows the total OH and Figure 12b shows the active and spectator OH energy disposal shown as energy fractions of translation, vibration, and rotation, as a function of the collision energy. We define the total energy,  $E = E_{\text{coll}} + E_{\text{z.p.}}(\text{H}_2\text{O}) - 14.7 \text{ kcal mol}^{-1}$ , and the energy fractions as  $f_v = \langle E_v \rangle / E$ ,  $f_r = \langle E_r \rangle / E$ , and  $f_T = 1 - \langle f_v \rangle - \langle f_r \rangle$ . We note that the average energies,  $\langle E_v \rangle$  and  $\langle E_r \rangle$ , were obtained before histogram binning and are summed over all

trajectories with the appropriate weighting of electronic state contributions. For the total OH energy disposal at energies below  $\sim 35$  kcal mol<sup>-1</sup> the vibrational and translation fractions are about the same and are appreciably larger than the rotational fraction. However, above 50 kcal mol<sup>-1</sup> the rotational fraction begins to dominate over the vibrational fraction as the vibrational energy becomes increasingly diverted to translational energy with increasing collision energy. Figure 12b shows the breakdown of the energy fractions into active and spectator components. It is quite clear that the active OH has the majority of product internal energy and this tendency increases dramatically above  $\sim 40$  kcal mol<sup>-1</sup> collision energy. The largest changes with collision energy occur with the active and spectator vibrational energy fractions which decrease dramatically above 25 kcal mol<sup>-1</sup> collision energy. As the collision energy increases, the vibrational energy fraction becomes diverted into rotation and translation, especially at the highest collision velocities.

#### 4. Differential angular cross sections

The differential angular cross section summed over all OH states is shown in Fig. 13 at 5.0, 7.0, 8.0, 9.0, and 11.0 km s<sup>-1</sup> collision velocity. At low collision velocities, the distribution is backward peaked, but as the collision velocity increases the angular distribution becomes more and more forward peaked, until at 11 km s<sup>-1</sup> products are mostly forward scattered. There appears to be a change in mechanism from backward to forward peaked with collision velocity. This mirrors the changes in energy deposition from vibration to rotation and translation with collision energy, and indicates different reaction mechanisms at low and high energies. The changeover occurs between 5 and 7 km s<sup>-1</sup>.

#### C. Comparisons to measurements of $O(^3P) + HOD(X,4,OH) \rightarrow OH(X,^2\Pi) + OD(X,^2\Pi)$

The same potential surfaces have been used to examine the measurements of Ref. 20 of the reaction  $O(^3P) + HOD(X,4,OH) \rightarrow OH(X,^2\Pi) + OD(X,^2\Pi)$ . In these measurements, atomic oxygen formed from photolysis of NO<sub>2</sub> collides at collision energies of 1.69 kcal mol<sup>-1</sup> with HOD in which the OH bond has been excited by 4 quanta. The necessary energy to create products is obtained mostly from excitation of the HOD reagent so that there is sufficient available energy (24.3 kcal mol<sup>-1</sup>) to excite up to OD( $v=3$ ) and OH( $v=2$ ) products. Product

internal energy distributions were probed with laser induced fluorescence. Table IV shows the experimental and theoretical relative populations of OD(v) and OH(v). The theoretical results only use electronic state 1. We note that classical trajectory products can be assigned to quantum bins that can exceed the total energy available to products, a problem inherent in conventional histogram binning in QCT. As described in Ref. 20, most of the initial vibrational energy in the OH bond of HOD is retained in the OH product, and the OD functions like a spectator in the reaction. The theoretical results agree well with the measured OD populations, but there is only qualitative agreement with measurements for the OH(v) populations. The theoretical results are largest for OH(v=2) but there is appreciable population in v=0 and v=1.

TABLE IV. Relative populations of OD(v) and OH(v) for the measurements of Ref. 20.

	OD(Ref. 20)	OD(theory)	OH(Ref. 20)	OH(theory)
v=0	0.93±0.03	0.81	-----	0.17
v=1	0.07±0.03	0.16	0.00±0.04	0.34
v=2	----	0.03	1.0±0.04	0.49

Figure 14 shows the relative populations of OD(v=0,j) and OH(v=2,j) states, for the measurements of Ref. 20 and the present theoretical results. There is fairly good agreement for OD(v=0,j) populations, but there is considerable disagreement for the OH(v=2,j) populations. Due to the histogram binning method in the presently implemented QCT, the theoretical results predict OH(j>10) states which are not energetically allowed and this is extremely problematic in the present case where the vibrational bins are quite wide. This problem can be addressed with the use of gaussian binning, something we plan to carry out in future work. In this case, Gaussian binning could have a dramatic impact on the rotational distributions. We also show in Fig. 14 the relative populations for OD(v=0,j) and OH(v=2,j) for a rotational temperature of 700 K which is fairly close to the temperature implied by the relative collision energy. The 700 K temperature populations are a fairly good fit to the data, and this could suggest that collisional relaxation plays a major role in determining these populations.



#### IV. FUTURE WORK

The present calculated results for reaction (1) are well suited for planning and analyses of hyperthermal O-atom measurements<sup>21</sup> which should be possible in the near future. If final state resolution is possible, such measurements would be an ideal way to assess the accuracy of the present surfaces and dynamics results and explore the rich dynamics of this system. The present surfaces can also be used to investigate direct collisional excitation of  $\text{H}_2\text{O}(\text{X}^1\text{A}_1)$  by  $\text{O}(\text{P})$ , reaction (2). For further analysis of the laboratory and space-based spectral data, direct collisional excitation results will be critical. This is probably best carried out with quantum scattering approaches, as resolution of final quantum states for polyatomic systems, although done previously,<sup>13</sup> presents special problems with classical approaches. It is also possible to model  $\text{OH}(\text{X}^2\Pi) + \text{OH}(\text{X}^2\Pi)$  collisions, the reverse of reaction (1) with the present surfaces, although the present surfaces only involve three of the four electronic states on the  $\text{OH} + \text{OH}$  side. Since  $\text{OH}(\text{X}^2\Pi) + \text{OH}(\text{X}^2\Pi)$  collisions to make  $\text{H}_2\text{O}$  are exothermic, measurements of the reverse of (1) are more experimentally accessible. We should mention also that the present work does not consider singlet surfaces nor electronic coupling which could be important in comparing to detailed measurements. Finally, we note that the present surfaces form the lower part of a large electronic state manifold leading to electronically excited  $\text{OH}(\text{A},^2\Sigma^+)$ .<sup>12</sup> The present approaches and surfaces together with information from Ref. 12 could be used as the basis to construct global surfaces with couplings leading to electronically excited OH observed in several measurements.<sup>10,11</sup>

#### ACKNOWLEDGEMENTS

We acknowledge support through a Small Business Innovative Research (SBIR) award from the Missile Defense Agency (MDA) Contract No. F04611-03-C-0015, and technical oversight from Dr. Marty Venner of the Air Force Research Laboratory. We also acknowledge support from the DoD through contract F19628-00-C-0006. This work has benefited greatly from discussions with George Schatz, Bisu Maiti, Timothy Minton, James Duff, and Steve Adler-Golden. We also acknowledge William Hase for making the VENUS trajectory code available to us.

## REFERENCES

1. M. G. Dunn, G. T. Skinner, and C. E. Treanor, AIAA Journal **13**, 803 (1975).
2. C. E. Kolb and J. B. Elgin, J. Chem. Phys. **66**, 119 (1977).
3. J. W. Duff, S. C. Richtsmeier, L. S. Bernstein, H. H. K. Burke, and M. Pietrzyk, J. Space. Rock. **30**, 755 (1993).
4. G. E. Caledonia, "Infrared Radiation Produced in Ambient/Spacecraft-Emitted Gas Interactions Under LEO Conditions", AIAA paper 00-0104 (2000).; and as reported in Ref. 7.
5. D. K. Zhou, W. R. Pendleton, Jr. G. E. Bingham, D. C. Thompson, W. J. Raitt, and R. M. Nadile, J. Geo. Res. **99**, 19,585 (1994).
6. R. E. Meyerott, G. R. Swenson, E. L. Schweitzer, and D. G. Koch, J. Geo. Res. **99**, 17,559 (1994).
7. L. S. Bernstein, J. B. Elgin, C. P. Pike, D. J. Knecht, E. E. Murad, T. F. Zehnpfennig, G. E. Galicia, and A. T. Stair, Jr., J. Geo. Res. **101**, 383 (1996).
8. D. P. Murtagh, E. J. Llewellyn, and P. J. Espy, Geo. Res. Lett, **24**, 85 (1997).
9. D. L. A. Rall, I. L. Kofsky, C. P. Pike, D. J. Knecht, and T. P. Zehnpfennig, Journal of Spacecraft and Rockets, **33**, 393 (1996).
10. O. J. Orient, A. Chutijan, and E. Murad, Phys. Rev. Lett, **65**, 2359 (1990).
11. L. Bernstein, Y. Chiu, J. A. Gardner, A. L. Broadfoot, M. Lester, M. Tsiouris, R. Dressler, and E. Murad, J. Phys. Chem. **107**, 10695 (2003).
12. S. Matsik and D. R. Yarkony, J. Chem. Phys. **117**, 3733 (2002).
13. M. J. Redmon, G. C. Schatz, and B. C. Garrett, J. Chem. Phys. **84**, 764 (1986).
14. B.R. Johnson, J. Chem. Phys. **84** 176 (1986).
15. R. J. Bartlett, and M. J. Redmon, Report No. AFRL-TR-81-27.

16. R. J. Redmon, R. J. Bartlett, B. C. Garrett, G. D. Purvis III, P. M. Saatzler, G. C. Schatz, and I. Shavitt, *Potential Energy Surfaces and Dynamics Calculations*, edited by D. G. Truhlar (Plenum, New York, 1981), pp. 771-803.
17. S. P. Karkach and V. I. Osherov, *J. Chem. Phys.* **110**, 11918 (1999).
18. D. W. Arnold, C. Xu, D. M. Neumark, *J. Chem. Phys.* **102**, 6088 (1995).
19. H.-J. Deyerl, T. G. Clements, A. K. Luong, and R. E. Continetti, *J. Chem. Phys.* **115**, 6931 (2001).
20. J. M. Pfeiffer, E. Woods, III, R. B. Metz, and F. Fleming Crim, *J. Chem. Phys.* **113**, 7982 (2000).
21. D. J. Garton, T. K. Minton, D. Troya, R. Pascual, and G. C. Schatz, *J. Phys. Chem. A* **107**, 4583 (2003).
22. W. Tsang and R. F. Hampson, *J. Phys. Chem. Ref. Data* **15**, 1087 (1986).
23. A. Lifschitz and J. V. Michael, *Symp. Int. Combust. Proc.* **23**, 59 (1991).
24. R. Sayos, C. Oliva, and M. Gonzalez, *J. Chem. Phys.* **115**, 8828 (2001).
25. G. Wu, G. C. Schatz, G. Lendvay, D.-C. Fang, and L. B. Harding, *J. Chem. Phys.* **113**, 3150 (2000).
26. D. McCune, Princeton Spline and Hermite Cubic Interpolation Routines, Princeton Plasma Physics Laboratory, <http://w3.pppl.gov/NTCC/PSPLINE/> (2000).
27. H. Partridge and D. W. Schwenke, *J. Chem. Phys.* **106**, 4618 (1997).
28. K. P. Huber and G. Herzberg, "Constants of Diatomic Molecules", Van Nostrand, 1979.
29. M. W. Schmidt, K. K. Baldridge, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. J. Su, T. L. Windus, M. Dupuis, J. A. Montgomery, *J. Comput. Chem.* **14**, 1347 (1993).
30. L. Harding, *J. Phys. Chem.* **95**, 8653 (1991).
31. W. L. Hase, R. J. Duchovic, X. Hu, A. Komornicki, K. F. Lim, D.-H. Lu, G. H. Peslherbe, K. N. Swamy, S. R. V. Linde, A. Varandas, H. Wang, and R. J. Wolfe, *Quantum Chem. Program Exchange* **16**, 671 (1996).

32. R. D. Levine and R. B. Bernstein, "Molecular Reaction Dynamics and Chemical Reactivity", chapter 5, Oxford University Press, (1987).

## FIGURE CAPTIONS

Figure 1. Energy level diagram of low-lying triplets (solid lines) and singlets (dashed lines) showing reagents, transition states and products for  $O + H_2O$  collisions. The transition state geometries and energy levels are for triplets based on the work of Ref. 17. The energy and corresponding relative collision velocities are shown in parentheses with respect to the  $O(^3P) + H_2O(X, ^1A_1)$  asymptote at zero energy. The energies of the fundamental vibrational modes of  $H_2O$  are shown for reference. In the present study, we follow the three lowest triplet states of adiabatically  $O(^3P) + H_2O(X, ^1A_1)$  to  $OH(^2\Pi) + OH(^2\Pi)$ .

Figure 2. Internal coordinates used in the description of the  $O(^3P) + H_2O(X, ^1A_1)$  potential energy surfaces. Active coordinates (dashed lines) are the set  $\{x, y, z\}$  and spectator coordinates (solid lines) are the set  $\{u, \theta, \tau\}$ . (a) illustrates the  $\{x, y, z, u, \theta\}$  'in-plane' coordinates and (b) illustrates the 'out-of-plane'  $\tau$  coordinate looking along the  $O_a-H_b$  bond.

Figure 3. Fitted potential energy surfaces of the three lowest triplet states of the  $O(^3P) + H_2O(X, ^1A_1)$  system. On each frame, the abscissa is the x-coordinate and the ordinate is the y-coordinate. Each column shows a different electronic state. Each row shows a different fixed z-coordinate ( $80^\circ, 140^\circ$  and  $240^\circ$ ). The spectator coordinates  $\{u, \theta, \tau\}$  are fixed at their transition state values as given in Table I. Contour levels are at  $\{1, 5, 10, 15, 20, 25, 30, 40, 50\}$  kcal mol $^{-1}$ . Distances are in a.u.

Figure 4. The  $\tau$  dependence of the active potential,  $V_a$  for the three lowest triplet states of the  $O(^3P) + H_2O(X, ^1A_1)$  system. Here it is defined with respect to the energy of  $V_a$  for state 1 at the transition state given in Table I.

Figure 5. The  $V_\theta$  portion of the spectator potential,  $V_s$ , as a function of  $\theta$  for the three lowest triplet states of the  $O(^3P) + H_2O(X, ^1A_1)$  system: —, state 1; - - -, state 2; ----, state 3.

Figure 6. Full potential,  $V_{sym}$ , for planar geometries of the  $O(^3P) + H_2O(X, ^1A_1)$  system where the reagent  $H_2O$  bond lengths and angles are fixed at the transition state geometry, as shown in Table I for the three lowest triplet adiabatic states. The position of the reagent O-atom is varied along the abscissa and ordinate coordinates. Distances are in a.u. Contour values are at  $\{1, 5, 10, 15, 20, 30, 40, 50\} + 16.8$  kcal mol $^{-1}$  (which is the value of the potential for the O-atom far away for state 1).

Figure 7. Rate constants as a function of temperature for the reaction,  $O(^3P) + H_2O(X, ^1A_1) \rightarrow OH(X, ^2\Pi) + OH(X, ^2\Pi)$ .  $\square$ , experimental results of Ref. 22;  $\times$ , experimental results of Ref. 23;  $\bigcirc$ , total calculated results including contributions from all three states;  $\diamond$ , contribution from state 1;  $\nabla$ , contribution from state 2;  $\triangle$ , contribution from state 3.

Figure 8. Total calculated OH cross section as a function of collision energy for the reaction  $O(^3P) + H_2O(X, ^1A_1) \rightarrow OH(X, ^2\Pi) + OH(X, ^2\Pi)$  according to Eq. 12:  $\bigcirc$ , total calculated results

including contributions from all three states;  $\diamond$ , contribution from state 1;  $\nabla$ , contribution from state 2;  $\triangle$ , contribution from state 3. Relative collision velocity is shown on the top scale.

Figure 9. (a) Calculated vibrationally resolved cross sections as a function of collision energy for the reaction  $O + H_2O \rightarrow OH(v) + OH(v)$  according to Eq. 12. For a given  $v$ , contributions from both  $OH(v)$  products are summed. (a),  $\bigcirc$ , results at  $5 \text{ km s}^{-1}$  collision velocity;  $\diamond$ , results at  $7 \text{ km s}^{-1}$  collision velocity;  $\triangle$ , results at  $8 \text{ km s}^{-1}$  collision velocity;  $\nabla$ , results at  $9 \text{ km s}^{-1}$  collision velocity;  $\square$ , results at  $11 \text{ km s}^{-1}$  collision velocity. (b), same as (a) except results only include  $OH(v)$  products formed from the incoming O-atom reagent (active OH). (c), same as (a) except results only include  $OH(v)$  products formed from reagent  $H_2O$  O-atom (spectator OH).

Figure 10. Calculated rotationally resolved cross sections as a function of collision energy for the reaction  $O + H_2O \rightarrow OH(j) + OH(j)$  according to Eq. 12. For a given OH rotational level  $j$ , contributions from both active and spectator OH products are summed over all  $OH(v)$  vibrational levels.  $\bigcirc$ , results at  $5 \text{ km s}^{-1}$  collision velocity;  $\diamond$ , results at  $7 \text{ km s}^{-1}$  collision velocity;  $\triangle$ , results at  $8 \text{ km s}^{-1}$  collision velocity;  $\nabla$ , results at  $9 \text{ km s}^{-1}$  collision velocity;  $\square$ , results at  $11 \text{ km s}^{-1}$  collision velocity.

Figure 11. Calculated relative populations at  $8 \text{ km s}^{-1}$  relative collision velocity for the reaction  $O + H_2O \rightarrow OH_{\text{active}}(j) + OH_{\text{spectator}}(j)$ .  $\text{---}\square\text{---}$ , presents results for  $OH_{\text{active}}(j)$ ;  $\text{--}\square\text{--}$  - surprisal theory, Eq. 16, with  $\theta_r=0.0$ ;  $\text{---}\bigcirc\text{---}$ , presents results for  $OH_{\text{spectator}}(j)$ ;  $\text{--}\bigcirc\text{--}$  - surprisal theory, Eq. 16, with  $\theta_r=-0.41$ .

Figure 12. Calculated product energy fraction as a function of collision energy for the reaction  $O(^3P) + H_2O(X, ^1A_1) \rightarrow OH(X, ^2\Pi) + OH(X, ^2\Pi)$ . (a) total summed over OH active and spectator contributions,  $\text{---}\triangle\text{---}$ , translational energy fraction,  $f_T$ ;  $\text{---}\diamond\text{---}$ , rotational energy fraction,  $f_r$ ;

—○—, vibrational energy fraction,  $f_v$ . (b) active OH and spectator OH contributions to the internal energy fraction.

Figure 13. Calculated differential angular cross sections as a function of collision velocity. ○, results at 5 km s<sup>-1</sup> collision velocity; ◇, results at 7 km s<sup>-1</sup> collision velocity; △, results at 8 km s<sup>-1</sup> collision velocity; ▽, results at 9 km s<sup>-1</sup> collision velocity; □, results at 11 km s<sup>-1</sup> collision velocity. To be consistent with previous results, we follow the convention for Eq. 12 for the total cross section.

Figure 14. Rotational relative populations for the reaction  $O(^3P) + HOD(4vOH) \rightarrow OD(v,j) + OH(v,j)$ . (a) calculated results for OD( $v=0$ ) product, - - □ - -, experimental results of Ref. 20; —○—, present calculated results; --△--, population of OD( $v=0,j$ ) at 700 K. (b) same as (a) except for OH( $v=2,j$ ).

Fig. 1






Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9






Fig. 9




Fig. 9




Fig. 10




Fig. 11




Fig. 12




Fig. 13




Fig. 14

