

DTIC COPY

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

Introduction to violent Sun-Earth connection events of October–November 2003

N. Gopalswamy,¹ L. Barbieri,¹ E. W. Cliver,² G. Lu,³ S. P. Plunkett,⁴ and R. M. Skoug⁵

Received 14 June 2005; revised 16 June 2005; accepted 20 June 2005; published 30 September 2005.

[1] The solar-terrestrial events of late October and early November 2003, popularly referred to as the Halloween storms, represent the best observed cases of extreme space weather activity observed to date and have generated research covering multiple aspects of solar eruptions and their space weather effects. In the following article, which serves as an abstract for this collective research, we present highlights taken from 61 of the 74 papers from the *Journal of Geophysical Research*, *Geophysical Research Letters*, and *Space Weather* which are linked under this special issue. (An overview of the 13 associated papers published in *Geophysics Research Letters* is given in the work of Gopalswamy et al. (2005a)).

Citation: Gopalswamy, N., L. Barbieri, E. W. Cliver, G. Lu, S. P. Plunkett, and R. M. Skoug (2005), Introduction to violent Sun-Earth connection events of October–November 2003, *J. Geophys. Res.*, 110, A09S00, doi:10.1029/2005JA011268.

1. Introduction

[2] The violent solar eruptions of October–November 2003 are one of the best observed outbreaks of intense solar activity to date. These events, referred to as the Halloween storms, are extreme events in terms of both their source properties at the Sun and their heliospheric consequences. The plasma, particle, and electromagnetic consequences of these events were detected at several locations in the heliosphere thanks to the distributed network of spacecraft. Disturbances associated with two of the October–November 2003 eruptions arrived at Earth in less than a day. Historically, only 13 such “fast transit” events, including the Carrington event of 1 September 1859, have been observed. Remarkably, the two fast transit events in October 2003 occurred on consecutive days, following a delay of over 30 years from the previous such event on 4 August 1972. Several aspects of the Halloween storms, including active region size and potential energy, flare occurrence rate and peak intensity, CME speed and energy, shock occurrence rate, SEP occurrence rate and peak intensity, and the geomagnetic storm intensity, displayed extreme behavior [Gopalswamy et al., 2005b].

[3] As expected, this outbreak of strong solar activity resulted in a broad spectrum of space weather impacts. About 59% of the reporting spacecraft and about 18% of the onboard instrument groups were affected by these storms; electronic upsets, housekeeping and science noise, proton

degradation to solar arrays, changes to orbit dynamics, high levels of accumulated radiation, and proton heating were observed. Most Earth-orbiting spacecraft were put into safe mode to protect from the particle radiation. Major societal impacts also occurred: ~50,000 people in southern Sweden (Malmö) experienced a blackout when the oil in a transformer heated up by 10 degrees; surge currents were observed in Swedish pipelines; and several occurrences were noted of degradation and outage of GPS systems. Teams climbing Mount Everest experienced interference on high-frequency radio communication paths.

[4] The solar energetic particle event on 28 October resulted in significant ozone depletion between 40 and 90 km from the ground. A tenfold enhancement in the ionospheric total electron content over the US mainland occurred during 30–31 October. Extraordinary density enhancements in both the magnetosphere and ionosphere coinciding with intervals of southward IMF and high-speed solar wind were observed.

[5] Effects of the eruptions were observed progressively later beyond Earth to the farthest reaches of the heliosphere [Lario et al., 2005; McKibben et al., 2005; Crider et al., 2005; Intriligator et al., 2005]. At Mars, the MARIE instrument on board the Mars Odyssey mission was completely damaged by particle radiation. The disturbances continued to the orbits of Jupiter and Saturn as detected by Ulysses and Cassini, respectively. Wind, Ulysses, and Cassini radio instruments observed a radio burst resulting from colliding CMEs on 4 November from widely different vantage points. Finally, the disturbances reached Voyager 2 after about 180 days, piled up together as a single merged interaction region (MIR), which led a large depression in cosmic ray intensity, lasting more than 70 days.

[6] In summary, the Halloween 2003 events serve as a useful benchmark of the extreme solar activity and its terrestrial and heliospheric effects [Gopalswamy et al., 2005b; Cliver and Svalgaard, 2004; Penna and Quillen, 2005; Malandraki et al., 2005; Ebihara et al., 2005; Dmitriev

¹Laboratory for Extraterrestrial Physics, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

²Space Vehicles Directorate, Air Force Research Laboratory, Hanscom Air Force Base, Massachusetts, USA.

³High Altitude Observatory, National Center for Atmospheric Research, Boulder, Colorado, USA.

⁴Naval Research Laboratory, Washington, D.C., USA.

⁵Los Alamos National Laboratory, Los Alamos, New Mexico, USA.

et al., 2005a, 2005b; *Belov et al.*, 2005; *Hu et al.*, 2005; *Nose et al.*, 2005; *Rosenqvist et al.*, 2005; *Miroshnichenko et al.*, 2005]. The following provides a synopsis of results obtained by analyzing data acquired during this interval. At this early stage in the data analysis, the emphasis is on the severity of the disturbances and their impacts. Nonetheless, the dynamic range provided by such disturbances has yielded, and continues to yield, insight to their physics.

2. Overview

[7] *Gopalswamy et al.* [2005b] summarize the properties of all the CMEs during this period in comparison with those of all the CMEs observed during SOHO's mission life until the end of 2003. They find that an unusually large fraction of fast and wide CMEs and halo CMEs occurred during this period. They report the observation of at least 16 shocks near the Sun using radio data, while eight of them were intercepted by spacecraft along the Sun-Earth line [see also *Terasawa et al.*, 2005]. The CMEs impacting the magnetosphere resulted in intense geomagnetic storms, some of them among the largest ones of solar cycle 23. Very intense SEP events, including three ground level enhancements (GLEs) occurred in association with the CMEs. *Gopalswamy et al.* find that the extreme CME kinetic energy in the Halloween eruptions is consistent with the largest energy extractable from the huge associated active regions. A plot summarizing solar, interplanetary, and geomagnetic conditions from 19 October to 21 November is given in Figure 1. Note the large numbers of flares, CMEs, and SEP events in the top three plots. The lull in flare activity during 6–11 November is because the three main active regions rotated off to the backside of the Sun. The activity returned when one of the active regions (AR 484) returned as AR 501. The number of CMEs is possibly an underestimate because the SOHO detectors were temporarily saturated by the SEPs during 28–30 October. The bottom two plots show extreme solar wind speeds and superintense geomagnetic storms. Detailed conditions in the corona overlying the eruptive regions can be found in the work of *Butala et al.* [2005] and *Grechnev et al.* [2005].

3. Solar Sources

[8] *Woods et al.* [2004] report on total solar irradiance (TSI) measurements in the UV and EUV spectral regimes during this active period. They find that the TSI drops by an unprecedented 0.34% due to the presence of large sunspots on the solar disk. They also report the first definitive detection of a flare in TSI on 28 October.

[9] Using riometer measurements at 20.1 MHz, *Brodrick et al.* [2005] reconstructed the X-ray flare on 4 November which was found to be saturated in the GOES-12 data. The authors suggested that an approximated energy flux of 3.8 mW/m^2 (X38) flare seems to be a more suitable value than the X28 flare estimated from the GOES data. This was the largest soft X-ray flare yet recorded.

4. Disturbance Propagation

[10] *Reiner et al.* [2005] report on a combined analysis of radio and white-light observations of a CME on 2

November using SMEI and Wind/WAVES data. They used these observations to constrain the parameters of a simple kinematic model of CME propagation and to derive the radial speed profile for this CME from the Sun to 1 AU. Their method may provide a framework for more accurate predictions of the arrival of CMEs at 1 AU and thus improved forecasts of space weather events. *Tokumaru et al.* [2005] used interplanetary scintillation measurements to establish unambiguous associations between interplanetary shocks and solar events in the period from 21 October to 8 November. Together, these papers illustrate the importance of tracking disturbances continuously from the Sun to 1 AU in order to establish the link between solar events and in situ measurements of the solar wind near the Earth.

[11] *Jackson et al.* [2005] use a kinematic solar wind density model to perform a three-dimensional (3-D) reconstruction of the 28 October CME from SMEI observations. (For a CME reconstruction for this event based on cosmic ray observations, see *Kuwabara et al.* [2004].) *Jackson et al.* [2005] also derive an estimate for the total mass of this CME in the inner heliosphere. This is the first 3-D reconstruction of a CME from SMEI white light data.

[12] *Wu et al.* [2005] describe the use of a 1.5-D MHD model to study the evolution and interaction of a series of shocks associated with the events from 28 October to 2 November. Their results show the importance of including shock interactions when considering the geomagnetic impacts of successive solar events. *Dryer et al.* [2004] evaluate the application of their "fearless forecasts" to the epoch from 19 October to 20 November. During this period, a total of 19 solar flares were accompanied by metric type II radio bursts, the triggering event for a forecast. The authors compare forecasts of the time of interplanetary shock arrival at Earth obtained by four different (analytic/heuristic, MHD, or kinematic) models. Best results are obtained for the Hakamada-Akasofu-Fry kinematic model with a success rate of 74% (defined as the ratio of hits (forecast arrival time within ± 15 hours of observed time) plus correct nulls divided by the total number of forecasts).

5. Solar Energetic Particles (SEPs)

[13] *Mewaldt et al.* [2005] determine that high-energy particle fluence recorded during the late October to early November 2003 period constituted 20% of that observed from 1997–2003. The authors estimate that the energy in the energetic particles in each of the major events during this 2 week interval ranged from ~ 1 to 25% of the kinetic energy in the associated coronal mass ejections. For each event, they construct energy spectra for H, He, and O over the range from ~ 0.1 to >100 MeV, and for electrons from 40 keV to 8 MeV. Both the ion and electron spectra can be fitted with double power laws.

[14] *Cohen et al.* [2005] combine SIS and ULEIS data from ACE to construct heavy ion spectra over more than 3 decades of energy for the five large events of October–November 2003. Despite considerable event-to-event variation, two interesting trends are observed: (1) the ratios of abundances at SIS (12–60 MeV) to ULEIS (0.64–0.91 MeV) energies increased in all cases with ionic charge to mass ratio (decreased with nuclear charge); and (2) fluence spectra of O, Ne, Mg, Si, Ca, and Fe within each




Figure 1. An overview plot showing (from top to bottom) the GOES X-ray flares, CME height-time plots, the SEP flux (>10 MeV protons), the 1 AU solar wind speed from ACE, and the *Dst* index. The nominal quiet condition is marked by the horizontal dashed lines. The solid lines in the CME height-time plots represent halo CMEs.

event could be organized remarkably well by assuming that the positions of spectral breaks for the different elements were governed by their diffusion coefficients. The latter result finds support in the study by *Mewaldt et al.* [2005]. *Cohen et al.* [2005] argue that knowledge of the charge states of heavy ions, and their variation with energy, is critical for obtaining further insights into the abundance variations observed in large SEP events.

[15] The current state of understanding of the acceleration and release of SEPs in conjunction with solar eruptions can be seen in a comparison of analyses by *Klassen et al.* [2005] and *Simnett* [2005] of the electron event associated with the X17 flare on 28 October. These authors independently identify phases of SEP injection during this event and suggest acceleration mechanisms for the various phases. From an analysis of radio and electron data, *Klassen et al.* deduce three phases of particle injection: (1) acceleration of ~ 30 keV electrons associated with an intense type III radio burst; (2) a delayed impulsive injection of <300 keV electrons and GeV protons; and (3) a further delayed injection of electrons with a hard spectrum at energies above ~ 100 keV. While the first of these components is attributed to the flare impulsive phase, the origin of the second and third components could lie either in acceleration in a coronal shock or in a reconnection related process in the wake of the CME. In the electron data, *Simnett* identifies a precursor, a main pulse, and a delayed prolonged component. *Simnett* attributes the main pulse to a fast CME and the delayed component to the flare. In order to explain the isotropic distribution of the delayed electrons at 1 AU, *Simnett* postulates that flare electrons are trapped within the CME magnetic structure, from which they leak out over time to fill the inner heliosphere, and are subsequently backscattered from a boundary somewhere in the heliosphere beyond 1 AU.

6. Magnetospheric Impacts

[16] *Looper et al.* [2005; see also *Lopez et al.*, 2004] describe the profound impact that major ICMEs can have on Earth's inner radiation belt. On 29 October 2003, SAMPEX observed that the usual belt of >20 MeV protons around $L = 2$ almost completely disappeared, to be replaced over the next several months by a belt of >10 MeV electrons that diffused from higher altitudes. Such inner belt disturbances are rare; the only comparable event was the first recognized disturbance of this type, observed by CRRESS in March 1991.

7. Ionospheric and Thermospheric Responses

[17] From an analysis of DMSP ion drift measurements, *Hairston et al.* [2005] concluded that the polar cap electric potential drop was saturated during the 29–31 October superstorm, with the saturation limit at about 260 kV. The ionosphere was severely disturbed during the storms. A highly elevated F2 layer was observed by an ionosonde in Kazakhstan, where $hmF2$ (the height of the F layer peak electron density) was raised as high as 700 km, along with a 60% decrease of $foF2$ (the critical frequency of the F layer peak electron density) [*Gordienko et al.*, 2005]. In addition, the unusual formation of the E, E2, and F1 layers at night as well as the sporadic E layer was also detected. *Sahai et al.*

[2005] showed that the spread F features formed over Brazil and wave-like disturbances in the F region height and electron density in both the Brazilian and east Asian longitudinal sectors were observed.

[18] A dramatically decreased plasma density was reported in the southern midlatitude and high-latitude regions following the storm commencement on 29 October [*Yizengaw et al.*, 2005]. The plasma depletion was accompanied by a deep oxygen dayglow depletion observed by IMAGE/FUV, and the region remained depleted for more than 24 hours until 31 October when the second storm began. The depletion of plasma density extended up to ~ 800 km as measured by DMSP. *Lin et al.* [2005] showed an expanded equatorial ionization anomaly (EIA) up to 40° latitude during the 29–30 October storm interval, and they attributed it to the strong upward $\mathbf{E} \times \mathbf{B}$ drift that produces a strong plasma fountain effect. Suppression of the EIA during the storm recovery phase was also found to be associated with the downward drift. A negative storm effect was observed in the Southern Hemisphere, which was corroborated by a reduction in O/N2 ratio in the TIMED/GUVI observations. During the 20–21 November storm, a phenomenon known as a tongue of ionization (TOI) was formed when a continuous stream of cold and dense plasmas is being transported from middle latitudes into the polar region [*Foster et al.*, 2005]. The TIMED/GUVI measured a severe depleted zone of the O/N₂ column density which extended from high latitudes to near the equator at the peak of the storm [*Meier et al.*, 2005].

[19] The storm also caused significant disturbances in the thermosphere. Enhanced meridional and zonal neutral winds of 400 m/s were observed over Scandinavia [*Thuillier et al.*, 2005]. The CHAMP satellite measured a dramatic increase in neutral mass density by 200–300% in the thermosphere at an altitude of ~ 410 km [*Liu and Lühr*, 2005; *Sutton et al.*, 2005]. The CHAMP measurements displayed a significant hemispheric asymmetry in the neutral density variations, with the Northern Hemisphere showing a greater density increase than the Southern Hemisphere.

[20] The solar forcing was felt on Mars. The Mars Global Surveyor Magnetometer/Electron Reflector experiment detected strong magnetic field oscillations at and below the oxygen gyrofrequency, an indication that ions of planetary origin are interacting with the solar wind plasmas. *Espley et al.* [2005] speculated that such an interaction may result in a significant atmospheric loss during the passage of large solar storms at Mars.

8. Impact of SEPs on the Earth's Atmosphere

[21] The October–November 2003 solar proton events were ranked as the fourth largest period of SEPs over the past 40 years [*Jackman et al.*, 2005]. The highly energetic protons penetrate into the mesosphere and stratosphere where they produce excitations, ionizations, dissociations, and dissociative ionizations. A strong depletion of ozone by 50–70% was observed in the mesosphere and stratosphere in the northern polar cap and a smaller (40%) reduction in the southern lower mesosphere [*Jackman et al.*, 2005; *Rohen et al.*, 2005; *López-Puertas et al.*, 2005a]. The ozone depletion was attributed to the enhanced production of HO_x

(H, OH, HO₂) and NO_x (NO and NO₂) by energetic solar protons. Model simulations carried out by Verronen *et al.* [2005] showed that an order of magnitude enhancement in HO_x and NO_x in the D region could cause a 20–95% reduction in ozone at 40–85 km. The HALOE (Halogen Occultation Experiment) on board the UARS (Upper Atmospheric Research Satellite) detected an increase of NO_x by more than 20 ppbv over the southern polar region [Jackman *et al.*, 2005], and the MIPAS (Michelson Interferometer for Passive Atmospheric Sounding) instrument on ENVISAT measured an elevated NO_x density of 20–70 ppbv in the Northern Hemisphere and 10–35 ppbv in the Southern Hemisphere in the altitude range of 40–60 km [López-Puertas *et al.*, 2005a]. Enhancement of other ozone-destruction compounds was also measured by MIPAS/ENVISAT, including a 0.2–0.4 ppbv increase of ClO, a more than 0.3 ppbv increase of HOCl, a 2 ppbv increase of HNO₃, a 0.5–1.2 ppbv increase of N₂O₅, and an increased ClONO₂ of 0.4 ppbv [von Clarmann *et al.* 2005; López-Puertas *et al.*, 2005b]. Gardner *et al.* [2005] showed that auroral activity during the storm may also lead to an increased production of N(4S) and N(2D), resulting in enhanced chemical formation of NO in the thermosphere and enhanced 5.3 um emissions such as measured by ENVISAT/MIPAS.

9. Space Weather Forecasting and Its Application

[22] During the Halloween storms spacecraft in all orbits, LEO, MEO, GEO, HEO, as well as interplanetary missions, were affected by the hostile radiation environment. Barbieri and Mahmot [2004] focused on benchmarking the mission effects for this period of atypical severe space weather. Approximately 59% of reporting spacecraft and ~18% of their instrument groups experienced some effect from solar activity. The benchmark shows that even in one of the most severe space weather events of recent years the effects on and costs to the spacecraft and missions were relatively modest: existing design practices and operations strategies mitigated effects.

[23] Oler [2004] provides a study of the prediction performance of five space weather forecast centers for the five strongest interplanetary coronal mass ejections (ICMEs) during this period. The evaluation is particularly intended for the Northeast Power Coordinating Council (NPCC), realizing that accurate time-of-arrival predictions and rapid responses to the upstream detection of strong ICMEs are of paramount importance to such critical infrastructures. Results indicate that the average time-of-arrival error for all forecast centers was 9.26 hours, which is consistent with the guidance errors associated with the leading shock propagation prediction models; overall, the strongest ICME impact events of 29 and 30 October were the most poorly predicted.

[24] In addressing the risk to aircrew and passengers at aircraft altitude from observations made during flights on 29–30 October, Getley [2004] presents data from a very rare occurrence capturing a large unpredictable event with monitoring equipment rarely used on board an aircraft, as well as on an aircraft at a significant latitude and altitude at the time of an event. The author concludes that, while major solar particle events are rare, the increase in equiv-

alent dose rate was ~37%. Thus solar events can significantly affect the total absorbed dose on longer flights.

10. Conclusion

[25] We have presented an overview of key findings on the size/impact of the Halloween storms of 2003 as presented in AGU journals. This overview is representative, not comprehensive. All of the papers collected electronically in this special series are listed in the reference section. Despite the substantial amount of work that has been completed, the cited references represent only a first installment of observations, analyses, and models of a series of events that provide the definitive example of an outburst of extreme space weather activity.

[26] **Acknowledgments.** Some of these results were presented at the Fall 2003 and Spring 2004 meetings of the American Geophysical Union in special sessions on the October–November 2003 events. We thank S. Yashiro for help with Figure 1. The special sections team acknowledges the efforts by A. Richmond, L. Lanzerotti, and M. Moldwin in making the trijournal special section happen. The effort of NG was supported by NASA/LWS program.

[27] Shadia Rifai Habbal thanks the reviewers for their assistance in evaluating this paper.

References

- Barbieri, L. P., and R. E. Mahmot (2004), October–November 2003's space weather and operations lessons learned, *Space Weather*, 2, S09002, doi:10.1029/2004SW000064.
- Belov, A., *et al.* (2005), Magnetospheric effects in cosmic rays during the unique magnetic storm on November 2003, *J. Geophys. Res.*, 110, A09S20, doi:10.1029/2005JA011067.
- Brodrick, D., *et al.* (2005), X-ray magnitude of the November 4 2003 solar flare inferred from the ionospheric attenuation of the galactic radio background, *J. Geophys. Res.*, 110, A09S36, doi:10.1029/2004JA010960.
- Butala, M. D., *et al.* (2005), Three-dimensional estimates of the coronal electron density at times of extreme solar activity, *J. Geophys. Res.*, 110, A09S09, doi:10.1029/2004JA010938.
- Cliver, E. W., and L. Svalgaard (2004), The 1859 solar terrestrial disturbance and the current limits of extreme space weather activity, *Solar Phys.*, 224, 407.
- Cohen, C. M. S., *et al.* (2005), Heavy ion abundances and spectra from the large SEP of October–November 2003, *J. Geophys. Res.*, 110, A09S16, doi:10.1029/2005JA011004.
- Crider, D. H., *et al.* (2005), Mars Global Surveyor observations of the Halloween 2003 solar super-storm's encounter with Mars, *J. Geophys. Res.*, 110, A09S21, doi:10.1029/2004JA010881.
- Dmitriev, A., *et al.* (2005a), Indirect estimation of the solar wind conditions in 29–31 October 2003, *J. Geophys. Res.*, 110, A09S02, doi:10.1029/2004JA010806.
- Dmitriev, A., J.-K. Chao, M. Thomsen, and A. Suvorova (2005b), Geosynchronous magnetopause crossings on October 29–31, 2003, *J. Geophys. Res.*, 110, A08209, doi:10.1029/2004JA010582.
- Dryer, M., Z. Smith, C. D. Fry, W. Sun, C. S. Deehr, and S.-I. Akasofu (2004), Real-time shock arrival predictions during the "Halloween 2003 epoch," *Space Weather*, 2, S09001, doi:10.1029/2004SW000087.
- Ebihara, Y., *et al.* (2005), Ring current and the magnetosphere-ionosphere coupling during the super storm of 20 November 2003, *J. Geophys. Res.*, 110, A09S22, doi:10.1029/2004JA010924.
- Espley, J. R., *et al.* (2005), Low frequency plasma oscillations at Mars during the October 2003 solar storm, *J. Geophys. Res.*, 110, A09S33, doi:10.1029/2004JA010935.
- Foster, J. C., *et al.* (2005), Multi-radar observations of the polar tongue of ionization, *J. Geophys. Res.*, 110, A09S31, doi:10.1029/2004JA010928.
- Gardner, J. L., *et al.* (2005), Rotational and spin-orbit distributions of NO observed by MIPAS/ENVISAT during the solar storm of October/November 2003, *J. Geophys. Res.*, 110, A09S34, doi:10.1029/2004JA010937.
- Getley, I. L. (2004), Observation of solar particle event on board a commercial flight from Los Angeles to New York on 29 October 2003, *Space Weather*, 2, S05002, doi:10.1029/2003SW000058.
- Gopalswamy, N., L. Barbieri, G. Lu, S. P. Plunkett, and R. M. Skoug (2005a), Introduction to the special section: Violent Sun-Earth connection

- events of October–November 2003, *Geophys. Res. Lett.*, **32**(3), L03S01, doi:10.1029/2005GL022348.
- Gopalswamy, N., et al. (2005b), Coronal mass ejections and other extreme characteristics of the 2003 October–November solar eruptions, *J. Geophys. Res.*, **110**, A09S15, doi:10.1029/2004JA010958.
- Gordienko, G. I., et al. (2005), Ionospheric disturbances over Alma-Ata during the October–November 2003 magnetic storms, *J. Geophys. Res.*, **110**, A09S35, doi:10.1029/2004JA010945.
- Grechnev, V. V., et al. (2005), CORONAS-F/SPRITE EUV observations of October/November 2003 solar eruptive events in combination with SOHO/EIT data, *J. Geophys. Res.*, **110**, A09S07, doi:10.1029/2004JA010931.
- Hairston, M. R., et al. (2005), Saturation of the ionospheric polar cap potential during the October–November 2003 superstorms, *J. Geophys. Res.*, **110**, A09S26, doi:10.1029/2004JA010864.
- Hu, Q., et al. (2005), On the magnetic topology of October/November 2003 events, *J. Geophys. Res.*, **110**, A09S03, doi:10.1029/2004JA010886.
- Intriligator, D. S., et al. (2005), From the sun to the outer heliosphere: Modeling and analyses of the interplanetary propagation of the October/November (Halloween) 2003 solar events, *J. Geophys. Res.*, doi:10.1029/2004JA010939, in press.
- Jackman, C. H., et al. (2005), Neutral atmospheric influences of the solar proton events in October–November 2003, *J. Geophys. Res.*, **110**, A09S27, doi:10.1029/2004JA010888.
- Jackson, B. V., et al. (2005), Preliminary 3D analysis of the heliospheric response to the 28 October 2003 CME using SMEI white-light observations, *J. Geophys. Res.*, doi:10.1029/2004JA010942, in press.
- Klassen, A., et al. (2005), Solar energetic electrons related to the October 28, 2003 flare, *J. Geophys. Res.*, **110**, A09S04, doi:10.1029/2004JA010910.
- Kuwabara, T., et al. (2004), Geometry of an interplanetary CME on October 29, 2003 deduced from cosmic rays, *Geophys. Res. Lett.*, **31**, L19803, doi:10.1029/2004GL020803.
- Lario, D., et al. (2005), Heliospheric energetic particle observations during the October–November 2003 events, *J. Geophys. Res.*, **110**, A09S11, doi:10.1029/2004JA010940.
- Lin, C. H., et al. (2005), Large scale variations of the low latitude ionosphere during the October–November, 2003 superstorm: Observational results, *J. Geophys. Res.*, **110**, A09S28, doi:10.1029/2004JA010900.
- Liu, H., and H. Lühr (2005), Strong disturbance of the upper thermospheric density due to magnetic storms: CHAMP observations, *J. Geophys. Res.*, **110**, A09S29, doi:10.1029/2004JA010908.
- Looper, M. D., J. B. Blake, and R. A. Mewaldt (2005), Response of the inner radiation belt to the violent Sun–Earth connection events of October–November 2003, *Geophys. Res. Lett.*, **32**, L03S06, doi:10.1029/2004GL021502.
- Lopez, R. E., et al. (2004), Sun unleashes Halloween storm, *Eos Trans. AGU*, **85**(11), 105–108.
- López-Puertas, M., et al. (2005a), Observation of NO_x enhancement and ozone depletion in the northern and southern hemispheres after the October–November 2003 solar proton events, *J. Geophys. Res.*, doi:10.1029/2005JA011050, in press.
- López-Puertas, M., et al. (2005b), HNO₃, N₂O₅ and ClONO₂ enhancements after the October–November 2003 solar proton events, *J. Geophys. Res.*, **110**, A09S20, doi:10.1029/2005JA011067.
- Malandraki, O. E., et al. (2005), October/November 2003 ICMEs: ACE/EPAM solar energetic particle observations, *J. Geophys. Res.*, **110**, A09S06, doi:10.1029/2004JA010926.
- McKibben, R. B., et al. (2005), Energetic particle observations from the Ulysses COSPIN instruments obtained during the October–November 2003 events, *J. Geophys. Res.*, **110**, A09S19, doi:10.1029/2005JA011049.
- Meier, R. R., et al. (2005), First look at the November 20, 2003 superstorm with TIMED/GUVI: Comparisons with a thermospheric global circulation model, *J. Geophys. Res.*, **110**, A09S41, doi:10.1029/2004JA010990.
- Mewaldt, R. A., et al. (2005), Proton, helium, and electron spectra during the large solar particle events of October, November, *J. Geophys. Res.*, **110**, A09S18, doi:10.1029/2005JA011038.
- Miroshnichenko, L. I., et al. (2005), Relativistic nucleon and electron production in the 2003 October 28 solar event, *J. Geophys. Res.*, **110**, A09S08, doi:10.1029/2004JA010936.
- Nose, M., et al. (2005), Overwhelming O⁺ contribution to the plasma sheet energy density during the October 2003 superstorm: Geotail/EPIC and IMAGE/LENA observations, *J. Geophys. Res.*, **110**, A09S24, doi:10.1029/2004JA010930.
- Oler, C. (2004), Prediction performance of space weather forecast centers following the extreme events of October and November 2003, *Space Weather*, **2**, S08001, doi:10.1029/2004SW000076.
- Penna, R. F., and A. C. Quillen (2005), The decay of interplanetary coronal mass ejections and Forbush decrease recovery times, *J. Geophys. Res.*, **110**, A09S05, doi:10.1029/2004JA010912.
- Reiner, M. J., et al. (2005), CME kinematics deduced from white-light (SMEI) and radio (Wind/WAVES) observations, *J. Geophys. Res.*, **110**, A09S14, doi:10.1029/2004JA010943.
- Rohen, G., et al. (2005), Ozone depletion during the solar proton events of Oct./Nov. 2003 as seen by SCIAMACHY, *J. Geophys. Res.*, **110**, A09S39, doi:10.1029/2004JA010984.
- Rosenqvist, L., et al. (2005), Extreme solar-terrestrial events of October 2003: High latitude and Cluster observations of the large geomagnetic disturbance on October 30, *J. Geophys. Res.*, **110**, A09S23, doi:10.1029/2004JA010927.
- Sahai, Y., et al. (2005), Effects of the major geomagnetic storms of October 2003 on the equatorial and low-latitude *F* region in two longitudinal sectors, *J. Geophys. Res.*, doi:10.1029/2004JA010999, in press.
- Simnett, G. M. (2005), Near relativistic electron emission following the 28 October, 2003 X17 flare, *J. Geophys. Res.*, **110**, A09S01, doi:10.1029/2004JA010789.
- Sutton, E. K., et al. (2005), Global thermospheric neutral density and wind response to the severe 2003 geomagnetic storms from CHAMP accelerometer data, *J. Geophys. Res.*, **110**, A09S40, doi:10.1029/2004JA010985.
- Terasawa, T., et al. (2005), Determination of shock parameters for the very fast interplanetary shock on 29 October 2003, *J. Geophys. Res.*, **110**, A09S12, doi:10.1029/2004JA010941.
- Thuillier, G., et al. (2005), Dynamics in the polar thermosphere after the coronal mass ejection of 28 October 2003 observed with the EPIS interferometer at Svalbard, *J. Geophys. Res.*, **110**, A09S87, doi:10.1029/2004JA010966.
- Tokumaru, M., M. Kojima, K. Fujiki, M. Yamashita, and D. Baba (2005), Interplanetary consequences caused by the extremely intense solar activity during October–November 2003, *J. Geophys. Res.*, **110**, A01109, doi:10.1029/2004JA010656.
- Verronen, P. T., et al. (2005), Diurnal variation of ozone depletion during the October–November 2003 solar proton events, *J. Geophys. Res.*, **110**, A09S32, doi:10.1029/2004JA010932.
- von Clarmann, T., et al. (2005), Experimental evidence of perturbed odd hydrogen and chlorine chemistry after the October 2003 solar proton events, *J. Geophys. Res.*, **110**, A09S45, doi:10.1029/2005JA011053.
- Woods, T. N., F. G. Eparvier, J. Fontenla, J. Harder, G. Kopp, W. E. McClintock, G. Rottman, B. Smiley, and M. Snow (2004), Solar irradiance variability during the October 2003 solar storm period, *Geophys. Res. Lett.*, **31**, L10802, doi:10.1029/2004GL019571.
- Wu, C.-C., et al. (2005), Flare-generated shock evolution and geomagnetic storms during the “Halloween 2003 epoch”: 29 October to 2 November 2003, *J. Geophys. Res.*, **110**, A09S17, doi:10.1029/2005JA011011.
- Yizengaw, E., et al. (2005), The southern hemisphere ionosphere and plasmasphere response to the interplanetary shock event of 29–31 October 2003, *J. Geophys. Res.*, **110**, A09S30, doi:10.1029/2004JA010920.
- L. Barbieri, Laboratory for Extraterrestrial Physics, NASA Goddard Space Flight Center, Code 444, Greenbelt, MD 20771, USA.
- E. W. Cliver, Space Vehicles Directorate, Air Force Research Laboratory, 29 Randolph Road, Hanscom Air Force Base, MA 01731, USA.
- N. Gopalswamy, Laboratory for Extraterrestrial Physics, NASA Goddard Space Flight Center, Code 695, Greenbelt, MD 20771, USA. (gopals@fugee.gsfc.nasa.gov)
- G. Lu, High Altitude Observatory, National Center for Atmospheric Research, Boulder, CO 80301, USA.
- S. P. Plunkett, Naval Research Laboratory, Code 7660, Washington, DC 20375, USA.
- R. M. Skoug, Los Alamos National Laboratory, Group NIS-1, MS D466, Los Alamos, NM 87545, USA.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 17-01-2006		REPRINT			
4. TITLE AND SUBTITLE Introduction to violent Sun-Earth connection events of October-November 2003				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 61102F	
				5d. PROJECT NUMBER 2311	
6. AUTHOR(S) Gopalswamy, N.*, L. Barbieri*, E.W. Cliver, G. Lu**, S.P. Plunkett#, and R.M. Skoug##				5e. TASK NUMBER RD	
				5f. WORK UNIT NUMBER A1	
				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-VS-HA-TR-2006-1011	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory/VSBXS 29 Randolph Road Hanscom AFB MA 01731-3010					
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S) AFRL/VSBXS	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited. *NASA Goddard Space Flt Ctr, Greenbelt, MD, **National Ctr for Atmos Res, Boulder, CO, #NRL, Washington, DC, ##Los Alamos National Lab, Los Alamos, NM					
13. SUPPLEMENTARY NOTES REPRINTED FROM: JOURNAL OF GEOPHYSICAL RESEARCH, Vol 110, A09S00, doi: 10.1029/2005JA011268, 2005.					
14. ABSTRACT [1] The solar-terrestrial events of late October and early November 2003, popularly referred to as the Halloween storms, represent the best observed cases of extreme space weather activity observed to date and have generated research covering multiple aspects of solar eruptions and their space weather effects. In the following article, which serves as an abstract for this collective research, we present highlights taken from 61 of the 74 papers from the <i>Journal of Geophysical Research</i> , <i>Geophysical Research Letters</i> , and <i>Space Weather</i> which are linked under this special issue. (An overview of the 13 associated papers published in <i>Geophysics Research Letters</i> is given in the work of Gopalswamy et al. (2005a)).					
15. SUBJECT TERMS Sun Corona Corona mass ejections Particle emissions					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON S.. W. Kahler
a. REPORT UNCLAS	UNCLAS	c. THIS PAGE UNCLAS			19b. TELEPHONE NUMBER (include area code) 781-377-9665