

## Orthopaedic Field Experience at a Level II Navy Surgical Facility during Operation Iraqi Freedom

**A. Taneja, M.D.**

CDR, MC, USN

Orthopaedic Surgeon

Naval Ambulatory Care Center Groton

1 Wahoo Drive

Box 600

Groton, CT 06349-5600

[ataneja@us.med.navy.mil](mailto:ataneja@us.med.navy.mil)

**C. Elsner, M.D.**

CDR, MC, USNR

Orthopaedic Surgeon

Naval Medical Center San Diego

### **ABSTRACT**

*A review of the orthopaedic surgical experience at a level II Navy field hospital during Operation Iraqi Freedom was undertaken. A retrospective data review was performed to evaluate include demographics of casualties treated; mechanism of injury; and procedures performed.*

*Results confirm that a majority of the injuries sustained on the battlefield will be to the musculoskeletal system. Both battle and non-battle injuries were administered to. The surgical environment remains austere at the level of the echelon II field hospital, requiring tolerance and improvisation. Management of the wounds remains unchanged, i.e. debridement and stabilization, and may be referred to in present day terms as damage control orthopaedics. Our results differ from the etiology of the injuries seen from other reports of recent campaigns. The majority of the battle injuries attended to at this facility were due to missile wounds as opposed to blast injuries as seen in other recent campaigns. Despite this the majority of wounds remained orthopaedic. The Orthopaedic surgeon is a vital asset to our fighting force in the field.*

### **1.0 OBJECTIVES**

Review of the field orthopaedic surgical experience at a Navy level II facility assigned to the US Marine Corps during Operation Iraqi Freedom.

### **1.1 METHODS**

A retrospective review was performed of the orthopaedic procedures performed at Bravo Surgical Company covering the period of 3 April 2003 to 4 May 2003 (28 operational days). Data was retrieved from operating room and personal logs. The data reviewed included all surgical cases performed in the Bravo Surgical Co.

*Paper presented at the RTO HFM Symposium on "Combat Casualty Care in Ground Based Tactical Situations: Trauma Technology and Emergency Medical Procedures", held in St. Pete Beach, USA, 16-18 August 2004, and published in RTO-MP-HFM-109.*

# Report Documentation Page

*Form Approved  
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>01 SEP 2004</b>	2. REPORT TYPE <b>N/A</b>	3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Orthopaedic Field Experience at a Level II Navy Surgical Facility during Operation Iraqi Freedom</b>		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Ambulatory Care Center Groton 1 Wahoo Drive Box 600 Groton, CT 06349-5600</b>		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>			
13. SUPPLEMENTARY NOTES <b>See also ADM001795, Combat Casualty Care in Ground-Based Tactical Situations: Trauma Technology and Emergency Medical Procedures (Soins aux blessés au combat dans des situations tactiques : technologies des traumatismes et procédures médicales d'urgence), The original document contains color images.</b>			
14. ABSTRACT			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	<b>UU</b>
			18. NUMBER OF PAGES <b>14</b>
			19a. NAME OF RESPONSIBLE PERSON

OR and the orthopaedic procedures performed at Forward Resuscitative Surgical System 5 (FRSS 5) while it was co-located with Bravo Surgical Co (while the FRSS and Surgical Company were co-located all patients were staged through the Surgical Company's resuscitation area, but varied as to which operating room they were taken to).

The data was reviewed to evaluate: (1) casualties attended to by an orthopaedic surgeon; (2) make-up of casualties i.e. Iraqi vs. U.S. Forces; (3) procedures performed; (4) cause of injury i.e. battlefield or non-battle injuries. Only those patients with which direct contact was made and who subsequently received intervention by the orthopaedic surgeon are presented. Patients managed via verbal consultation with non-orthopaedic colleagues are not presented.

The environment was a Navy level II surgical facility (i.e. field hospital) in support of the US Marine Corps during Operation Iraqi Freedom. Tents were used for resuscitations areas, operating rooms (Fig. 1a), and holding wards. The floors were the desert sand and included unexpected visitors (Figs. 1b and 1c). The mission was for the Surgical Company to be in position so that life and limb saving procedures could be performed prior to the patients being transported to more definitive care. Equipment available included digital x-rays, military individually packed external fixation devices (Fig. 2), dressing material, and rudimentary surgical equipment (transportability being the limiting factor). The Surgical Company has a 72-hour holding capacity by doctrine and as such all attempts were made to transfer patients to higher levels of care within that constraint.

### **1.1.1 RESULTS**

Bravo Surgical Co. had 66 casualties treated in the operating room. 41/66 (62%) of these casualties were attended to by an orthopaedic surgeon (Fig. 3), a general surgeon attended to the remaining casualties. Staffing of Bravo Surgical Co included 2 general surgeons, 3 ob/gyns, and 1 orthopaedic surgeon. Staffing of FRSS 5 included 1 general surgeon and 1 orthopaedic surgeon.

An orthopaedic surgeon treated a total of 53 casualties from 3 April 2003 to 4 May 2003 (Table 1); on 3 April and from 8 April to 13 April Bravo Surgical Company and FRSS 5 were co-located and both authors attended to procedures performed on those days. 27/53 (51%) of the patients and 41 % of the procedures reviewed were attended to by both authors (AT and CE), the remaining casualties reviewed were attended to by AT.

41 casualties were treated in the Bravo Surgical Co. OR, 7 casualties in the FRSS 5 OR, and 5 casualties were treated at Bravo Surgical Co outside the formal operating room (Fig. 4). 26/53 (49%) of the casualties treated were Iraqi; 27/53 (51%) were U.S. Forces (Fig. 5). 39/53 (74%) of the casualties sustained battlefield injuries: 27 from missile wounds, and 12 from blast injuries (Fig. 7). 14/53 (26%) of the casualties sustained non-battle injuries: 6 MVA/MCA, and 8 occupational or recreational injuries (Fig. 8). No patient was operated on more than once, as all casualties (Iraqi and US), were transferred to higher echelon of care within 48 hours.

124 total procedures were performed on the 53 casualties. The procedures included: fasciotomy /compartment release (61); I & D and stabilization of open wounds / fractures (24); amputations (15); ex-fix (9); arthrotomy (6); closed reduction and stabilization (6); radial artery repair (1); sagittal band repair (1); removal of hardware (1) (Fig. 9). All wounds were left open and dressed with a wet to dry dressing. 79/124 (63%) procedures on 35/53 (66%) of the casualties were performed in the first 11 days (9 operational days). Further delineation, by body region, of fasciotomies / compartment releases, amputations, and external fixators can be seen in Figs. 10, 11, and 12 respectively.

External fixators were limited in supply and were used most often to manage unstable injuries that were not otherwise amenable to splinting (Figs. 12a and 12b). Severity of wounds and their management are represented in Figs. 13a through 15c.

With some severe wounds improvisation was necessary for optimal management (Figs. 15a, 15b and 15c). In this case dental acrylic was used, as it was incidentally discovered that the material used by our dental colleagues for capping teeth was similar in character to PMMA used in orthopaedic practice.

### 1.1.1.1 CONCLUSIONS

Shortfalls in this review include lack of follow-up as all pts were transferred from our facility in short order and received further care at the next echelon of care. Another deficiency is that data on all orthopaedic injuries that were treated at the Surgical Company are not presented, as many of the non-operative orthopaedic injuries were treated by non-orthopaedic colleagues, i.e. FP's, Dentists, and PA's via verbal consultation since the tempo of operations was too great to allow for direct intervention by the surgeon.

A consequence of any military campaign will be taking of casualties. Recent campaigns have shown that 60-70% of battlefield injuries will be to the musculoskeletal system<sup>1-5</sup>. The majority of penetrating injuries sustained in recent wars are reported to have been the result of blast injuries.<sup>1,3,6</sup> Our experience does confirm that the majority of injuries sustained on the battlefield are to the musculoskeletal system. The cause, however, of the majority of wounds was a result of missile rather than blast injuries as seen previously. This was probably due to the way this campaign was waged, with an early and heavy ground assault. The orthopaedic surgeon is an essential and vital asset in the field setting. The majority of the surgical expertise that will be needed in the field for future campaigns will be from an orthopaedist.

The type of surgery performed could be called damage control orthopaedics, in keeping with the principles used to manage femur fractures with temporary external fixators prior to definitive osteosynthesis in the multiply injured civilian trauma patient.<sup>7-11</sup> The same principles are applied in the field (i.e. temporary stabilization with delayed definitive fracture fixation). The limiting factor in the field not only being the severity of the injuries, but also of the environment in which the treatment is rendered. The flow of casualties, as expected, followed the pace of the battles being waged and our experience showed the high intensity of the battles being waged in a short duration of time. Despite advances in our specialty, modern day field orthopaedics remains a practice in an extremely austere environment with only rudimentary and limited equipment requires adherence to the principles of battlefield wound management<sup>2, 12</sup> and necessitates tolerance, improvisation and innovation on the part of the surgeon.



**Fig. 1a** Operating Room



Fig. 1b Operating Room



Fig. 1c Operating Room




Fig. 2 Individually packed military external fixator




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9






Fig. 10





Fig. 11



**Fig. 12a**



**Fig. 12b** fasciotomy and external fixator bridging elbow for GSW to distal humerus



**Fig. 13a** GSW to Hand



**Fig. 13b** wound debrided & wrist disarticulation performed



**Fig. 14a** High Energy GSW to knee with 2<sup>o</sup> injuries to proximal tibia from bone fragments. Wounds grossly contaminated with dirt, rocks, etc.



**Fig. 14b** debridement and stabilization prior to transfer completed



**Fig. 15a** 4x6cm segmental defect radius and ulna s/p high energy GSW



Fig. 15b placement of intercalary spacer using dental acrylic



Fig. 15c dental acrylic

## References

1. Burkle FM Jr, Newland C, Meister SJ, Blood CG. Emergency medicine in the Persian Gulf War—Part3: Battlefield casualties. *Ann Emerg Med.* 1994;23:755-60.
2. Ryan JM, Cooper GJ, Haywood IR, Milner SM. Field Surgery on a future conventional battlefield: strategy and wound management. *Ann R Coll Surg Engl.* 1991;73:13-20.
3. Jackson DS, Batty CG, Ryan JM, McGregor WS. The Falklands war: Army field surgical experience. *Ann R Coll Surg Engl.* 1983;65:281-5
4. Mehran R, Connelly P, Boucher P, Berthiaume E, Cote M. Modern war surgery: the experience of Bosnia. 1: Deployment. *Can J Surg.* 1995;38:266-74.
5. Mehran R, Connelly P, Boucher P, Cote M. Modern war surgery: the experience of Bosnia. 2: The clinical experience. *Can J Surg* 1995;38:338-46.
6. Covey DC. Blast and fragment injuries of the musculoskeletal system. *J. Bone Joint Surg Am.* 2000;84:1221-1234.
7. Nowotarski PJ, Turen CH, Brumback RJ, Scarboro JM. Conversion of external fixation to intramedullary nailing for fractures of the shaft of the femur in multiply injured patients. *J Bone Joint Surg Am* 2000;82:781-8.
8. Scalea TM, Boswell SA, Scott JD, Mitchell KA, Kramer ME, Pollak AN. External fixation as a bridge to intramedullary nailing for patients with multiple injuries and with femur fractures: Damage control orthopedics. *J Trauma* 2000;48:613-623.
9. Pape HC, Hildebrand F, Pertschy S, Zelle B, Garapati R, Grimme K, Krettek C. Changes in the management of femoral shaft fractures in polytrauma patients: from early total care to damage control orthopedic surgery. *J Trauma* 2002;53:452-462.
10. Przkora R, Bosch U, Zelle B, Panzica M, Garapati R, Krettek C, Pape HC. Damage control orthopedic: a case report. *J Trauma* 2002;53:765-769
11. Pape HC, Giannoudis P, Krettek C. The timing of fracture treatment in polytrauma patients: relevance of damage control orthopedic surgery. *Am J Surg* 2002;183:622-629.
12. Weitz-Marshall AD, Bosse MJ. Timing of closure of open fractures. *J Am Acad Orthop Surg* 2002;10:379-384.

