

REPORT DOCUMENTATION PAGE

AFRL-SR-AR-TR-05-

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the data, reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

0514

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE	3. REPORT TYPE AND DATES COVERED Final	
4. TITLE AND SUBTITLE Frontiers in Applied and Computational Mathematics '05'			5. FUNDING NUMBERS FA9550-05-1-0373	
6. AUTHOR(S) Dr. Miura				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) New Jersey Institute of technology			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR- Dr. Nachman			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release, distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) FACM 2005 Conference from March 13-15, 2005				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

SESSION SPEAKERS (61)							
		First	Last	Talk Date	Talk Time	Session	add1
1	Dr.	Larry	F. Abbott	Friday, May 13	2:15 p.m.	Plenary II	Brandeis Univers
2	Dr.	Jonathan	Bell	Saturday, May 14	11:30 a.m.	Mathematical Fluid Dynamics III (M3)	University of Man Baltimore County
3	Dr.	Andrew	Belmonte	Friday, May 13	5:15 p.m.	Mathematical Fluid Dynamics II (M2)	Department of M:
4	Dr.	Richard	Bertram	Friday, May 13	4:15 p.m.	Mathematical Biology II (M2)	Florida State Uni
5	Dr.	Lora	Billings	Saturday, May 14	4:00 p.m.	Mathematical Biology IV (M4)	Montclair State U
6	Dr.	Victoria	Booth	Friday, May 13	5:45 p.m.	Mathematical Biology II (M2)	University of Mict
7	Dr.	Jared	Bronski	Saturday, May 14	11:30 a.m.	Nonlinear Waves (M3) Statistics in	Department of M:
8	Dr.	Javier	Cabrera	Friday, May 13	10:30 a.m.	Biology/Genetics and Education (M1)	Department of St:
9	Dr.	Russell	Caffisch	Saturday, May 14	11:00 a.m.	Mathematical Fluid Dynamics III (M3)	Department of M:
10	Dr.	Fioralba	Cakoni	Friday, May 13	5:15 p.m.	Computational Electromagnetics (M2)	Department of M: Sciences
11	Dr.	Nilanjan	Chatterjee	Friday, May 13	11:30 a.m.	Statistics in Biology/Genetics and Education (M1)	Senior Principal I
12	Dr.	Stephen	Childress	Saturday, May 14	3:30 p.m.	Mathematical Biology IV (M4)	Courant Institute Mathematical Sci
13	Dr.	Vittorio	Cristini	Friday, May 13	12 Noon	Mathematical Biology I (M1)	Department of Bi Engineering
14	Dr.	Pamela	Cook	Saturday, May 14	4:30 p.m.	Mathematical Fluid Dynamics IV (M4)	Department of M: Sciences
15	Dr.	Dipak	Dey	Friday, May 13	11:00 a.m.	Statistics in Biology/Genetics and Education (M1)	Head and Profes: Statistics
16	Dr.	Avner	Friedman	Saturday, May 14	5:15 p.m.	Keynote	Mathematical Bio Institute
17	Dr.	Anna	Georgieva	Saturday, May 14	12 Noon	Mathematical Fluid Dynamics III (M3)	Novartis Pharmar 436/1252A
18	Dr.	Martin	Golubitsky	Saturday, May 14	9:00 a.m.	Plenary III	University of Hou
19	Dr.	Richard	Haberman	Saturday, May 14	11:00 a.m.	Nonlinear Waves (M3) Statistics in	Department of M:
20	Dr.	Shelby W.	Haberman	Friday, May 13	12 Noon	Biology/Genetics and Education (M1)	Director
21	Dr.	Jackson	Hall	Saturday, May 14	3:30 p.m.	Biostatistics and Genomics (M4)	Professor of Bios
22	Dr.	John	Harris	Sunday, May 15	12:00 p.m.	Waves Computational	Emeritus Division of Applie
23	Dr.	Jan	Hesthaven	Friday, May 13	4:15 p.m.	Electromagnetics (M2)	Mathematics
24	Dr.	Philip	Holmes	Friday, May 13	11:00 a.m.	Mathematical Biology I (M1)	Princeton Univer
25	Dr.	Frank Anette	Hoppensteadt	Saturday, May 14	4:30 p.m.	Mathematical Biology IV (M4)	Courant Institute Mathematical Sci
26	Dr.	(Peko)	Hosoi	Friday, May 13	4:45 p.m.	Mathematical Fluid Dynamics II (M2)	Department of M: Engineering
27	Dr.	Leo	P. Kadanoff	Sunday, May 15	10:30 a.m.	Plenary VI	The James Franc
28	Dr.	Ashwani	K. Kapila	Sunday, May 15	12:00 p.m.	Mathematical Fluid Dynamics V (M5)	Department of M: Sciences

20060404033

add1	add2	add3	add4	add5	add6	add7
Brandeis University	MS 013 Brandeis University	Waltham, MA 02454-9110				
University of Maryland, Baltimore County	Department of Mathematics and Statistics	1000 Hilltop Circle	Baltimore, MD 21250			
Department of Mathematics	Pennsylvania State University	University Park, PA 16802				
Florida State University	Department of Mathematics (or Institute of Molecular Biophysics)	Tallahassee, FL 32306				
Montclair State University	Dept. of Mathematical Sciences	Upper Montclair, NJ 07043				
University of Michigan	Department of Mathematics	Ann Arbor, MI 48109-1109				
Department of Mathematics	University of Illinois	1409 W. Green Street	Urbana, IL 61801			
Department of Statistics	471 Hill Center, Busch Campus	Rutgers University	Piscataway, NJ 08854			
Department of Mathematics	University of California Los Angeles	405 Hilgard Avenue	Los Angeles, CA 90095-1555			
Department of Mathematical Sciences	University of Delaware	Newark, DE 19716-2553				
Senior Principal Investigator	Biostatistics Branch	Division of Cancer Epidemiology and Genetics	National Cancer Institute, NIH, DHHS	6120 Executive Blvd.	Rockville, MD 20852	
Courant Institute of Mathematical Sciences	251 Mercer Street	New York, NY 10012				
Department of Biomedical Engineering	University of California at Irvine	Irvine, CA 92697				
Department of Mathematical Sciences	University of Delaware	Newark, DE 19716				
Head and Professor of Statistics	Department of Statistics	CLAS Building, U-4120	University of Connecticut	215 Glenbrook Road	Storrs, CT 06269	
Mathematical Biosciences Institute	Ohio State University	231 W. 18th Avenue	Columbus, OH 43210			
Novartis Pharmaceuticals 436/1252A	1 Health Plaza	East Hanover NJ 07936				
University of Houston	Department of Mathematics	Houston, TX 77204-3008				
Department of Mathematics	Southern Methodist University	Dallas, TX 75275-0156				
Director	Center for Statistical Theory and Practice	Educational Testing Service	Rosedale Road	Princeton, NJ 08541		
Professor of Biostatistics	Department of Biostatistics and Computational Biology	University of Rochester Medical Center	601 Elmwood Avenue, Box 630	Rochester, NY 14642		
Emeritus	Department of Theoretical and Applied Mechanics	University of Illinois	Ctr. QEFP Catalysis Bldg., Rm. 327	Northwestern University	2137 N. Sheridan Rd.	Evanston, IL 60208- 3020
Division of Applied Mathematics	Brown University	182 George Street, Box F	Providence, RI 02912			
Princeton University	Department of Mechanical and Aerospace Engineering	Princeton, NJ 08544				
Courant Institute of Mathematical Sciences	New York University	70 Washington Square, 1232	New York, NY 10012			
Department of Mechanical Engineering	MIT	77 Massachusetts Avenue	Cambridge, MA 02139			
The James Franck Institute	Office 109	The University of Chicago	5640 South Ellis Avenue	Chicago, IL 60637		
Department of Mathematical Sciences	Rensselaer Polytechnic Institute	Troy, NY 12180				

FACM '05 Conference Attendees May 13-15, 20

29	Dr.	William		Kath	Sunday, May 15	9:00 a.m.	Plenary V	Northwestern University	McCormick School
30	Dr.	Joseph	B.	Keller	Saturday, May 14	2:00 p.m.	Plenary IV	Department of Mathematics	Stanford University
31	Dr.	Thomas	B.	Kepler	Friday, May 13	11:30 a.m.	Mathematical Biology I (M1)	Duke University	Center for Biostatistics Bioinformatics
32	Dr.	J. Nathan		Kutz	Saturday, May 14	12 Noon	Nonlinear Waves (M3)	University of Washington	Department of Applied Mathematics
33	Dr.	Herbert		Levine	Friday, May 13	10:30 a.m.	Mathematical Biology I (M1)	University of California at San Diego	Department of Physics
34	Dr.	Bernard	J.	Matkowsky	Sunday, May 15	11:30 a.m.	Mathematical Fluid Dynamics V (M5)	John Evans Chair in Applied Mathematics	Northwestern University
35	Dr.	Masayasu		Mimura	Saturday, May 14	10:30 a.m.	Mathematical Biology III (M3)	Department of Mathematics	Meiji Institute for Math Sciences (MIMS)
36	Dr.	André		Nachbin	Friday, May 13	4:15 p.m.	Mathematical Fluid Dynamics II (M2)	Department of Pure and Applied Mathematics	IMPA
37	Dr.	Arje		Nachman	Friday, May 13	3:15 p.m.	Funding Agency	Air Force Office of Scientific Research	875 North Randolph R
38	Dr.	Monika		Nitsche	Saturday, May 14	11:30 a.m.	Mathematical Fluid Dynamics III (M3)	Department of Mathematics and Statistics	University of New Mex
39	Dr.	Andrew		Norris	Sunday, May 15	12:30 p.m.	Waves	Chairman	Department of Mechar Aerospace Engineering
40	Dr.	Hillary		Ockendon	Saturday, May 14	4:00 p.m.	Mathematical Fluid Dynamics IV (M4)	OCIAM	Mathematical Institute
41	Dr.	John		Ockendon	Friday, May 13	9:00 a.m.	Plenary I	OCIAM	Mathematical Institute
42	Dr.	Johan		Paulsson	Sunday, May 15	12:00 p.m.	Mathematical Biology V (M5)	Cambridge Computational Biology Institute	Centre for Mathematic Sciences
43	Dr.	John		Pearson	Saturday, May 14	11:00 a.m.	Mathematical Biology III (M3)	Los Alamos National Laboratory	P.O. Box 1663
44	Dr.	Fernando		Reitich	Friday, May 13	4:45 p.m.	Computational Electromagnetics (M2)	University of Minnesota	127 Vincent Hall
45	Dr.	John		Rinzel	Friday, May 13	5:15 p.m.	Mathematical Biology II (M2)	Courant Institute of Mathematical Sciences	New York University
46	Dr.	Gareth	J.	Russell	Sunday, May 15	12:30 p.m.	Mathematical Biology V (M5)	Columbia University	Department of Ecology and Environmental Biol
47	Dr.	William		Sallas	Saturday, May 14	4:30 p.m.	Biostatistics and Genomics (M4)	Novartis Pharmaceuticals Corporation	Clinical Development C
48	Dr.	Clyde		Scandrett	Sunday, May 15	11:30 a.m.	Waves	Chairman	Mathematics Department MA/Sd
49	Dr.	Arnd		Scheel	Saturday, May 14	10:30 a.m.	Nonlinear Waves (M3)	University of Minnesota	School of Mathematics
50	Dr.	Pranab	K.	Sen	Saturday, May 14	4:00 p.m.	Biostatistics and Genomics (M4)	Department of Biostatistics	McGavran-Greenberg I CB#7420
51	Dr.	Michael		Shelley	Friday, May 13	5:45 p.m.	Mathematical Fluid Dynamics II (M2)	Courant Institute of Mathematical Sciences	New York University
52	Dr.	Arthur		Sherman	Friday, May 13	4:45 p.m.	Mathematical Biology II (M2)	National Institutes of Health	N.I.H.-N.I.D.D.K.-M.R.E
53	Dr.	Kate		Stebe	Friday, May 13	10:30 a.m.	Mathematical Fluid Dynamics I (M1)	Department of Chemical and Biomolecular Engineering	The Johns Hopkins Uni
54	Dr.	Paul	H.	Steen	Friday, May 13	12 Noon	Mathematical Fluid Dynamics I (M1)	School of Chemical and Biomolecular Engineering	Cornell University
55	Dr.	D. Scott		Stewart	Sunday, May 15	12:30 p.m.	Mathematical Fluid Dynamics V (M5)	Department of Theoretical and Applied Mechanics	University of Illinois at C Champaign
56	Dr.	Howard		Stone	Friday, May 13	11:30 a.m.	Mathematical Fluid Dynamics I (M1)	Department of Engineering and Applied Sciences	Harvard University
57	Dr.	Jean-Marc		Vanden-Broeck	Saturday, May 14	12 Noon	Mathematical Fluid Dynamics III (M3)	School of Mathematics	University of East Angli
58	Dr.	Miguel	R.	Visbal	Friday, May 13	5:45 p.m.	Computational Electromagnetics (M2)	Computational Services Branch	Air Vehicles Directorate
59	Dr.	Z. Jane		Wang	Friday, May 13	11:00 a.m.	Mathematical Fluid Dynamics I (M1)	Dept. of Theoretical and Applied Mechanics	Cornell University

Attendees May 13-15, 2005

2

12/6/2005

g	McCormick School	Dept of Eng Sciences & Appl Math	2145 Sheridan Road	Evanston, IL 60208-3125		
atics	Stanford University	450 Serra Mall Bldg. 380	Stanford, CA 94305-2125			
	Center for Biostatistics and Bioinformatics	Durham, NC 27708				
on	Department of Applied Mathematics	Box 352420	Seattle, WA 98195-2420			
at San	Department of Physics	La Jolla, CA 92093				
ppled	Northwestern University	2145 Sheridan Road	Evanston, IL 60208-3125			
atics	Meiji Institute for Mathematical Sciences (MIMS)	School of Science and Technology	Meiji University	1-1-1 Higashimita	Yamaku, Kawasaki 214-8571	Japan
id	IMPA	Estrada Dona Castorina 110	Rio de Janeiro 22460-320	Brazil		
ntific	875 North Randolph Road	Ste 325, Room 3112	Arlington, VA 22203			
atics	University of New Mexico	Humanities 415	Albuquerque, NM 87131-1141			
	Department of Mechanical and Aerospace Engineering	Rutgers University	98 Brett Road	Piscataway, NJ 08854		
	Mathematical Institute	24-29 St. Giles'	Oxford OX1 3LB	United Kingdom		
	Mathematical Institute	24-29 St. Giles'	Oxford OX1 3LB	United Kingdom		
nal	Centre for Mathematical Sciences	Wilberforce Road	Cambridge, CB3 OWA	United Kingdom		
	P.O. Box 1663	Theoretical Biology and Biophysics	T-10 K710	Los Alamos, NM 87545		
a	127 Vincent Hall	206 Church St., S.E.	Minneapolis, MN 55455			
s	New York University	New York, NY 10012				
	Department of Ecology, Evolution and Environmental Biology	MC 5557	1200 Amsterdam Avenue	New York, NY 10027		
als	Clinical Development Operations	One Health Plaza	East Hanover, NJ 07936-1080			
	Mathematics Department, Code MA/Sd	Naval Postgraduate School	Monterey, CA 93943			
a	School of Mathematics	206 Church Street S.E.	Minneapolis, MN 55455			
stics	McGavran-Greenberg Hall CB#7420	Chapel Hill, NC 27599-7420				
s	New York University	251 Mercer Street	New York, NY 10012			
health	N.I.H.-N.I.D.D.K.-M.R.B.	Building 12A, Room 4007	12 South Dr. MSC 5621	Bethesda, MD 20892-5621		
al and	The Johns Hopkins University	221 Maryland Hall	3400 North Charles Street	Baltimore, MD 21218		
ing	Cornell University	120 Olin Hall	Ithaca, NY 14853-5201			
ical and	University of Illinois at Urbana-Champaign	216 Talbot Laboratory	104 S. Wright Street	Urbana, IL 61801		
ring	Harvard University	308 Pierce Hall	Cambridge, MA 02139-4307			
s	University of East Anglia	Norwich, England NR4 7TJ	United Kingdom			
s	Air Vehicles Directorate	Air Force Research Laboratory	WPAFB, OH 45433			
id	Cornell University	323 Thurston Hall	Ithaca, NY 14853-1503			

FACM '05 Conference Attendees May 13-15, 200

60	Dr.	Henry		Warchall	Friday, May 13	3:15 p.m.	Funding Agency	Program Director, Applied Mathematics	National Science Found
61	Dr.	Wendy		Zhang	Saturday, May 14	3:30 p.m.	Mathematical Fluid Dynamics IV (M4)	James Franck Institute	University of Chicago
POSTER PRESENTERS (121)									
1	Dr.	Silas		Alben	Harvard University		Division of Engineering and Applied Sciences	29 Oxford Street	Cambridge, MA 02138
2	Ms.	Christina		Ambrosio	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
3	Dr.	Roman		Andrushkiw	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
4	Dr.	Eliana		Antoniou	Assistant Prof.		William Paterson University	300 Pompton Road	Wayne, NJ 07470
5	Dr.	Felix	J.	Apfaltrer	Courant Institute of Mathematical Sciences		New York University	251 Mercer St.	New York, NY 10012
6	Dr.	Paul		Atzberger	Rensselaer Polytechnic Institute		Mathematical Sciences	8th Street	Troy, NY 12180
7	Dr.	Karim		Azer	Merck & Co., Inc.		Applied Computer Science & Mathematics	RY84-202, 126 E. Lincoln Avenue	Rahway, NJ 07065
8	Mr.	Dipankar		Bandyopadhyay	University of Georgia		266 Statistics Bldg	Athens, GA 30602	
9	Ms.	Anisha		Banerjee	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
10	Mr.	Sibabrata		Banerjee	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
11	Dr.	Lyudmyla		Barannyk	University of Michigan		Department of Mathematics	4855 East Hall	Ann Arbor, MI 48109-1109
12	Mr.	Michael		Bateman	University of Kansas		Department of Mathematics	1801 Kentucky #2	Lawrence, KS 66044
13	Dr.	John		Bechtold	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
14	Dr.	Manish		Bhattacharjee	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
15	Dr.	Denis		Blackmore	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
16	Dr.	Mark		Blyth	University of East Anglia		Mathematics Department	Norwich, England	NR4 7TJ, UK
17	Dr.	Michael		Booty	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
18	Dr.	Amitabha		Bose	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
19	Dr.	Bruce		Bukiet	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
20	Mr.	Jean		Cadet	SUNY at Stony Brook		Applied Mathematics and Statistics	Stony Brook, NY 11794-3600	
21	Dr.	Jerry		Chen	Case Western Reserve University		Department of Mathematics	10900 Euclid Avenue	Cleveland, OH 44106-7058
22	Mr.	Yiming		Cheng	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
23	Mr.	Shawn		Chester	NJIT		Mechanical Engineering Department	University Heights	Newark, NJ 07102
24	Dr.	Wooyoung		Choi	University of Michigan		Dept. of Naval Arch. & Marine Eng.	2600 Draper Rd.	Ann Arbor, MI 48109
25	Mr.	Sohae		Chung	SUNY at Stony Brook		Department of Applied Mathematics and Statistics	Stony Brook, NY 11794	
26	Dr.	Robert		Clewley	Cornell University		Department of Mathematics	Ithaca, NY 14853	
27	Dr.	Dean	T.	Crommelin	Courant Institute of Mathematical Sciences		New York University	251 Mercer St.	New York, NY 10012
28	Dr.	R.E.	Le	Deville	Courant Institute		New York University	251 Mercer St.	New York, NY 10012
29	Dr.	Sunil		Dhar	NJIT		Mathematical Sciences	University Heights	Newark, NJ 07102
30	Dr.	Ramana		Dodla	New York University		Center for Neural Science	4 Washington Place	New York, NY 10003
31	Dr.	Brent		Doiron	New York University		Center for Neural Science	4 Washington Place	New York, NY 10003
32	Mr.	Graham		Donovan	Northwestern University		Engineering Science and Applied Mathematics		
33	Dr.	Jonathan	D.	Drover	University of Pittsburgh		Mathematics Department	504 Cabot Way	Pittsburgh, PA 15203
34	Mr.	Srabasti		Dutta	SUNY at Stony Brook		Applied Mathematics and Statistics	Stony Brook, NY 11794-3600	

FACM '05 Conference Attendees May 13-15, 2005

35	Dr.	Alan		Elcrat	Wichita State University	Mathematics Department	1840 N. Fairmount	Wichita, KS
36	Dr.	Christopher	E.	Elmer	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
37	Mr.	Samiran		Ghosh	University of Connecticut	Department of Statistics	215 Glenbrook Rd. U-4120	Storrs, CT 06269
38	Dr.	Daniel		Goldman	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
39	Dr.	Jorge		Golowasch	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
40	Dr.	Roy		Goodman	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
41	Dr.	Arnaud		Goulet	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
42	Mr.	Muhammad		Hameed	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
43	Dr.	Alex		Haro	Universitat de Barcelona	Dept. de Matematica Aplicada I Anlisi	Gran Via 585	08007 Barcelona, Spain
44	Dr.	Cristel		Hohenegger	Georgia Institute of Technology	School of Mathematics	686 Cherry Street	Atlanta, GA 30332-0160
45	Dr.	David		Hornthrop	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
46	Dr.	Maureen		Howley	NJIT	Chemical Eng. Dept.	University Heights	Newark, NJ 07102
47	Dr.	Aridaman		Jain	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
48	Dr.	Shidong		Jiang	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
49	Dr.	Ning		Ju	Oklahoma State University	Department of Mathematics	401 Mathematical Sciences	Stillwater, OK 74078-1058
50	Mr.	Yeona		Kang	SUNY at Stony Brook	Applied Mathematics and Statistics	Stony Brook, NY 11794	
51	Dr.	Said		Kas-Danouche	Universidad de Oriente	Departamento de Matematicas	Cumana, Sucre 6101	Venezuela
52	Dr.	Aslan		Kasimov	University of Illinois at Urbana-Champaign	Department of Computer Science	216 Talbot Lab, 104 S. Wright St.	Urbana, IL 61801
53	Mr.	Navodit		Kaushik	University of Southern Mississippi	Department of Computer Science	Hattiesburg, MS 39406-0001	
54	Mr.	Adnan	A.	Khan	Rensselaer Polytechnic Institute	Mathematical Sciences	8th Street	Troy, NY 12180
55	Dr.	Hafiz	M.	Khan	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
56	Dr.	Christopher	E.	Khedouri	Center for Drug Evaluation and Research (CDER)	Food and Drug Administration (FDA)	9201 Corporate Blvd., Room S311	Rockville, MD 20850
57	Mr.	Nickolas		Kintos	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
58	Dr.	Lou		Kondic	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
59	Dr.	Gregory	A.	Kriegsmann	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
60	Dr.	Stephen		Kunec	Boston University	Department of Mathematics	Boston, MA 02215	
61	Ms.	Soumi		Lahiri	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
62	Mr.	Sanyogita		Lakhera	University of Southern Mississippi	Department of Mathematics	118 College Drive 5045	Hattiesburg, MS 39406-0001
63	Mr.	Joe		Latulippe	Montana State University	Mathematics Department	2-214 Wilson	Bozeman, MT 59717-2400
64	Dr.	Eric		Lauga	Harvard University	Division of Engineering and Applied Sciences	Cambridge, MA 02138	
65	Dr.	Kevin	K.	Lin	Courant Institute	New York University	251 Mercer St.	New York, NY 10012
66	Mr.	Xing		Liu	University of Maryland	Baltimore County	9 Merrill Road, Apt. A	Baltimore, MD 21228
67	Mr.	Xinfeng		Liu	SUNY at Stony Brook	Applied Mathematics and Statistics	Stony Brook, NY 11794	
68	Dr.	Dawn		Lott	Delaware State University	Mathematics and Biotechnology	1200 N. DuPont Hwy	Dover, DE 19901
69	Mr.	Tianshi		Lu	SUNY at Stony Brook	Department of Applied Math and Statistics	Stony Brook, NY 11794	
70	Dr.	Jonathan		Luke	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
71	Dr.	Valery		Lukyanov	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
72	Dr.	Marc	Q.	Ma	NJIT	Computer Science Department	University Heights	Newark, NJ 07102
73	Dr.	Victor		Matveev	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102

FACM '05 Conference Attendees May 13-15,

74	Dr.	Roberto	Mauri	University of Pisa	Department of Chemical Engineering	Pisa, Italy		
75	Dr.	Zoi-Heleni	Michalopoulou	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
76	Mr.	Yuriy	Mileyko	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
77	Dr.	Petronije	Milojevic	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
78	Dr.	Victor	Miroshnikov	College of Mount Saint Vincent	Math & Computer Science Department	6301 Riverdale Avenue	Riverdale, NY 10471	
79	Mr.	Milind	Misra	NJIT	Department of Chemistry and Environmental Science	University Heights	Newark, NJ 07102	
80	Dr.	Robert	M. Miura	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
81	Dr.	Brian	E. Moore	McGill University	Mathematics and Statistics	805 Sherbrooke Street W.	Montreal, Quebec H3A 2K6 Canada	
82	Dr.	Richard	Moore	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
83	Dr.	Pascal	Moyal	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
84	Dr.	Cyrril	Muratov	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
85	Dr.	Duane	Nykamp	University of Minnesota	Mathematics Department	206 Church St., SE	Minneapolis, MN 55455	
86	Dr.	Ozgur	Ozen	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
87	Dr.	Demetrios T.	Papageorgiou	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
88	Dr.	John	A. Pelesko	University of Delaware	Department of Mathematical Sciences	Ewing Hall Room 406	Newark, DE 19716-2553	
89	Dr.	Mark	Pernarowski	Montana State University	Mathematical Sciences	Bozeman, MT 59717		
90	Dr.	Peter	Petropoulos	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
91	Mr.	Valentin	Polishchuk	SUNY at Stony Brook	Applied Mathematics and Statistics	Stony Brook, NY 11794-3600		
92	Mr.	Joshua	L. Proctor	University of Washington	Department of Applied Mathematics	Box 352420	Seattle, WA 98195-2420	
93	Dr.	Christophe	Raymond	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
94	Dr.	Gregory T.	Reeves	Princeton University	Chemical Eng. Dept.	Princeton, NJ		
95	Mr.	Max	Roman	NJIT	Mechanical Engineering Department	University Heights	Newark, NJ 07102	
96	Mr.	Satrajit	Roychoudhury	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
97	Dr.	Roman	Samulyak	Brookhaven National Laboratory	Computational Science Center	Upton, NY 11973		
98	Dr.	Tobias	Schaefer	CUNY	Mathematics Department	2800 Victory Boulevard	Staten Island, NY 10314	
99	Dr.	Eric	Shea-Brown	Courant Institute	New York University	251 Mercer St.	New York, NY 10012	
100	Dr.	Asya	Shpiro	New York University	Center for Neural Science	4 Washington Place	New York, NY 10003	
101	Dr.	Michael	Siegel	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
102	Dr.	Linda	Smolka	Bucknell University	Department of Mathematics	Lewisburg, PA 17837		
103	Mr.	Seongho	Song	University of Connecticut	Department of Statistics	215 Glenbrook Rd. U-4120	Storrs, CT 06269-4120	
104	Dr.	David	Stickler	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
105	Dr.	Louis	Tao	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
106	Dr.	Burt	Tilley	Franklin W. Olin College of Engineering	Department of Mathematics	1000 Olin Way	Needham, MA 02492-1200	
107	Ms.	Natalia	Toporikova	Department of Biological Science	Florida State University	Tallahassee, FL 32306		
108	Dr.	A. David	Trubatch	United States Military Academy	Department of Mathematical Sciences	West Point, NY 10996		
109	Mr.	Dmitri	Tseluiko	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
110	Mr.	A. Kerem	Uguz	University of Florida	Chemical Eng. Dept.	Gainesville, FL 32611		
111	Mr.	Suleyman	Ulusoy	Georgia Institute of Technology	School of Mathematics	Atlanta, GA 30332-0160		
112	Dr.	Vianey	Villamizar	Brigham Young University	Mathematics Department	Provo, UT 84602		
113	Dr.	X. Sheldon	Wang	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
114	Dr.	Wonsuk	Yoo	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	

2

71						
H3A 2K6						
5455						
?						
?						
6-2553						
2						
95-2420						
2						
12						
12						
Y 10314						
0012						
0003						
02						
9-4120						
02						
02						
12492-1200						
102						
102						
102						

115	Mr.	Yun	Yoo	Drexel University	Department of Mathematics	Korman Center 252, 3141 Chestnut Street	Philadelphia, PA 19104	
116	Dr.	Yuan-Nan	Young	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
117	Mr.	Kai	Zhang	NJIT	Computer Science Department	University Heights	Newark, NJ 07102	
118	Ms.	Yili	Zhang	Rutgers University	Biological Sciences Department	Newark, NJ 07102		
119	Dr.	Yongmin	Zhang	SUNY at Stony Brook	Applied Mathematics and Statistics	Stony Brook, NY 11794		
120	Ms.	Lin	Zhou	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
121	Mr.	Ivan	Zorych	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	

POSTER CO-AUTHORS WHO ATTENDED (10)

1	Dr.	Nadine	Aubry	NJIT	Mechanical Engineering Department	University Heights	Newark, NJ 07102	w/Gouillet
2	Mr.	Amit	Banerjee	NJIT	Mechanical Engineering Department	University Heights	Newark, NJ 07102	w/Misra
3	Dr.	Jyoti	Champanerkar	William Paterson University	300 Pompton Road	Wayne, NJ 07470		w/Blackmore
4	Dr.	Tae-Chang	Jo	New Mexico Tech	Mathematics Department	801 Leroy Place	Socorro, NM 87801-4796	w/Choi
5	Dr.	Farzan	Nadim	NJIT	Mathematical Sciences Department	University Heights	Newark, NJ 07102	w/Golowasch
6	Dr.	Anthony	Rosato	NJIT	Mechanical Engineering Department	University Heights	Newark, NJ 07102	w/Chester
7	Dr.	Patricia	Soteropoulos	Public Health Research Institute	Center for Applied Genomics	225 Warren St.	Newark, NJ 07103	w/Ma
8	Mr.	Kentaro	Sugino	NJIT	Computer Science Department/Comp. Biology	University Heights	Newark, NJ 07102	w/Ma
9	Mr.	Tongsheng	Wang	Public Health Research Institute	Center for Applied Genomics	225 Warren St.	Newark, NJ 07103	w/Ma
10	Mr.	Yu	Wang	NJIT	Computer Science Department	University Heights	Newark, NJ 07102	w/Ma

NON-PRESENTING ATTENDEES (60)

1	Ms.	Shuchi	Agrawal	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
2	Dr.	Daljit	S. Ahluwalia	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
3	Dr.	Robert	A. Altenkirch	NJIT	President	University Heights	Newark, NJ 07102	
4	Dr.	Egbert	Ammicht	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
5	Mr.	Sultan	Babar	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
6	Dr.	Alex	Barnett	Courant Institute	New York University	251 Mercer Street	New York, NY 10012	
7	Ms.	Briti	Biswas	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
8	Dr.	Gareth	Block	Lawrence Livermore National Laboratory	Earth Science Division	7000 East Avenue, L-206	Livermore, CA 94550	
9	Dr.	Ruben	Moreno Bote	Science Center for Neuronal	New York University	4 Washington Place	New York, NY 10003	
10	Ms.	Lakshmi	Chandrasekaran	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
11	Dr.	Cameron	Connell	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
12	Dr.	Amy	L. Davidow	New Jersey Medical School	Dept. of Preventive Medicine & Community Health	Newark, NJ 07102		
13	Dr.	Fadi	Deek	NJIT	Dean of CSLA	University Heights	Newark, NJ 07102	
14	Dr.	Charles	R. Dees, Jr.	NJIT	Office of University Advancement	University Heights	Newark, NJ 07102	
15	Dr.	Rafael	De La Llave	University of Texas at Austin	Department of Mathematics	1 Univ. Station C1200	Austin, TX 78712	
16	Mr.	Leonardo	Espin	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102	
17	Dr.	Fariba	Fahroo	Air Force Office of Scientific Research	875 North Randolph Road	Ste 325, Room 3112	Arlington, VA 22203	
18	Dr.	Urmi	Ghosh-Dastidar	NY City College of Technology	City University of New York	Rm N726, 300 Jay Street	Brooklyn, NY 11201-2983	
19	Dr.	Juan	Gomez	1009 Louisa St, 2nd floor	Elizabeth, NJ 07201			

FACM '05 Conference Attendees May 13-15, 20

20	Mrs.	Padma	Gulati	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
21	Mr.	Joon	Ha	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
22	Ms.	Rashi	Jain	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
23	Dr.	Kenneth	Johnson	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
24	Dr.	Theodore	Johnson	NJIT	Human Resources Department	University Heights	Newark, NJ 07102
25	Mr.	Yogesh	Joshi	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
26	Mr.	Kei	Kaneko	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
27	Ms.	Manmeet	Kaur	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
28	Dr.	Paul	LaVergne	Vaughn College of Aeronautics and Technology	86-01 23rd Avenue	Flushing, NY 11369	
29	Mr.	Hyunsun	Lee	SUNY at Stony Brook	Applied Mathematics and Statistics	Stony Brook, NY 11794- 3600	
30	Mr.	Pil Hwa	Lee	Courant Institute of Mathematical Sciences	251 Mercer Street	New York, NY 10012	
31	Dr.	Dorothy	Levy	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
32	Mr.	Seth	Levy	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
33	Mr.	Ming	Lu	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
34	Mr.	Kamyar	Malakuti	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
35	Dr.	Colin	McKinstry	Lucent Technologies	101 Crawford Corner Rd.	Holmdel, NJ 07733	
36	Mr.	Yoichiro	Mori	Courant Institute	New York University	251 Mercer Street	New York, NY 10012
37	Mr.	Nebojsa	Murisc	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
38	Dr.	Priscilla	Nelson	NJIT	Provost	University Heights	Newark, NJ 07102
39	Ms.	Myongkeun	Oh	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
40	Mr.	Kunj	Patel	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
41	Dr.	Manuel	Perez	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
42	Mr.	Jonathan	Porus	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
43	Mr.	Filippo	Posta	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
44	Ms.	Karen	Roach	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
45	Dr.	Joel	Rogers	Polytechnic University	Mathematics Department	Brooklyn, NY 11201	
46	Ms.	Susan	Sutton	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
47	Mr.	Tsezar	Seman	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
48	Dr.	Thomas	Spencer	Walden University	Applied Management and Decision Sciences	103 Bergen Avenue	Princeton, NJ 08540
49	Ms.	Holly	Stern	NJIT	Chief Legal Counsel	University Heights	Newark, NJ 07102
50	Mr.	Alok	Tiwari	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102
51	Dr.	Christopher	Tong	Merck & Co., Inc.	Biometrics Research RY33- 300	P.O. Box 2000	Rahway, NJ 07065
52	Dr.	David	Ullmann	NJIT	Computing Services Division	University Heights	Newark, NJ 07102
53	Dr.	Ignatios	Vakalis	Capital University	Math/CS Department	1 College and Main	Columbus, OH 43209
54	Dr.	William	Van Buskirk	NJIT	Biomedical Engineering Department	University Heights	Newark, NJ 07102
55	Ms.	Paula	Vasquez	University of Delaware	Department of Mathematical Sciences	Newark, DE 19716	
56	Dr.	Jorge	Vñals	McGill University	Physics Department	3600 University Street	Montreal, Canada QC H3A 2T8
57	Dr.	Stuart	Walker	Northrop Grumman Corporation	600 Hicks Road	Rolling Meadows, IL 60008	
58	Mr.	Chunsheng	Yang	NJIT	Mathematical Sciences Computer Sciences	University Heights	Newark, NJ 07102
59	Mr.	Ijer	Yeh	NJIT	Department	University Heights	Newark, NJ 07102
60	Mr.	Yu	Zhang	NJIT	Mathematical Sciences	University Heights	Newark, NJ 07102

ance Attendees May 13-15, 2005

2

12/6/2005

I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I0012						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
I02						
8540						
I02						
I02						
065						
I02						
43209						
I02						
da QC H3A						
I02						
I02						
I02						

KEYNOTE SPEAKER

AVNER FRIEDMAN

Mathematical Biosciences Institute, Ohio State University, 231 W. 18th Avenue, Columbus, OH 43210

Mathematical Biology – A Newcomer to Applied Mathematics

Applied mathematics is traditionally concerned with mathematical models that arise in the physical sciences and in engineering. Such models are well established in areas of classical physics and chemistry, but are not as mature perhaps in materials science, in areas such as polymers and complex fluids. A new challenge in applied mathematics arises from the biosciences, where tremendous amounts of experimental data are currently being generated. In order to draw knowledge from these data, mathematical models need to be developed. The difficulty in developing and analyzing such models arise from the complexity of the biological processes, which flow from the molecular level to the organ level. How useful at this time are the mathematical sciences to the biosciences? In this talk I shall give you some examples from my experiences at the Mathematical Biosciences Institute.

PLENARY SPEAKERS

LARRY F. ABBOTT

Volen Center for Complex Systems, Brandeis University, Waltham, MA 02454-9110

Scale-Invariant Adaptation

Neuronal responses to constant stimuli adapt so that, for example, firing rates that are high when a stimulus first appears decrease over time. Such responses are often divided into transient and sustained components, and the transition between them is modeled as an exponential decay. More careful examination, however, often reveals that adaptation is better described by a power-law function of time rather than an exponential. Because such dynamics are scale-invariant, the division of the response into transient and sustained components is ambiguous, and extracted "time constants" of adaptation depend on the duration of the experiment rather than reflecting underlying features of the dynamics. Scale-invariance has the advantage that the time scale of the adaptation matches the temporal nature of the stimuli being encountered rather than being rigidly fixed by the system. I will discuss ways of modeling and thinking about scale-invariant adaptation and examine its functional implications for sensory processing.

MARTIN GOLUBITSKY

Department of Mathematics, University of Houston, Houston, TX 77204-3008

Coupled Cell Systems: Theory and Examples

A coupled cell system is a collection of interacting dynamical systems. Coupled cell models assume that the output from each cell is important and that signals from two or more cells can be compared so that patterns of synchrony can emerge. We ask: How much of the qualitative dynamics observed in coupled cells is the result of network architecture and how much depends on the specific equations?

The ideas will be illustrated through a series of examples and three theorems. The first theorem classifies spatio-temporal symmetries of periodic solutions; the second gives necessary and sufficient conditions for synchrony in terms of network architecture; and the third shows that synchronous dynamics may itself be viewed as a coupled cell system through a quotient construction. We also show how nongeneric bifurcations with nongeneric results arise in bifurcations in coupled systems.

LEO P. KADANOFF

The James Franck Institute, The University of Chicago, 5640 South Ellis Avenue, Chicago, IL 60637

Effective Scientific Simulations

I discuss a group of computer simulations, mostly ones related to astrophysical situations. My basic question relates to the effectiveness of simulations in the generation of new scientific knowledge. Examples under consideration include studies of solar neutrino production, the early history of the universe, sonoluminescence, "cold fusion", and mechanisms for mixing in novae, supernovae, and other turbulent flows.

I point to some very successful pieces of work, and to other pieces which are quite the opposite. I argue that scientific simulations can be used either as tools for discovering new knowledge or alternatively as tools for constructing arguments about scientific or technical issues. Supported in part by the Flash Center at the University of Chicago.

WILLIAM KATH

Engineering Sciences and Applied Mathematics, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208-3125

Simulating Rare Events in Lightwave Systems with Importance Sampling

Lightwave communication systems transmit information at extremely high rates. Transmission errors are handled at slower electronic speeds, however, and thus, systems must be designed to have extremely small error rates, typically one error per 10^9 or more bits. Since overall system performance is determined by extremely rare events, the accurate modeling of such systems presents a severe mathematical and computational challenge. In this talk, recent work aimed at overcoming this difficulty will be described. In particular, the application of importance sampling (one member of a general family of variance reduction techniques) to the numerical simulation of transmission impairments induced by amplified spontaneous emission noise in soliton-based optical transmission systems will be discussed. The method allows numerical simulations to be concentrated on the noise realizations that are most likely to result in transmission errors, leading to speedups of several orders of magnitude over standard Monte Carlo methods.

JOSEPH B. KELLER

Department of Mathematics, Stanford University, 450 Serra Mall Bldg. 380, Stanford, CA 94305-2125

Sound Source Location

The problem of how an animal determines the location of a source of sound will be formulated as an inverse problem. To solve it, the ears of the animal produce the Fourier transforms of the acoustic pressure signals in the two ears. At a higher level in the brain, the difference between the phases in the two ears is formed at each frequency. Similarly the difference between the sound pressure levels at each frequency is formed. It will be explained how these interaural differences are processed to determine the direction of the source.

JOHN OCKENDON

OCIAM, Mathematical Institute, 24-29 St. Giles', Oxford OX1 3LB, United Kingdom

Industry-Driven Challenges in Applied Mathematics

This talk will describe some industry-driven problems that pose interesting open questions in mathematical modelling and analysis. The topics will include:

- the dynamics of a shaped charge jet as it penetrates a target. For metal targets, it is the modelling of the elastic-plastic deformation that is crucial to the determination of the penetration depth.
- the effect of small dust particles on the flow of a cold plasma. There is an interesting analogy between the simplest continuum model for the ion velocity and the Euler equations of inviscid fluid dynamics.
- high-frequency asymptotics in the aircraft industry. One problem is that of how to incorporate localised diffraction effects into ray theory algorithms; another is that of the acoustic field generated by aircraft accelerating through sonic speed which still poses some interesting focusing problems.

INVITED TALKS

JONATHAN BELL

Department of Mathematics and Statistics, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250

Waves of Excitation in Neural Field Models

The neocortex is a distinctly layered, densely packed, neural structure. Thus, to consider large-scale patterns and dynamic activity, it has been reasonable to model the layers as continuum fields of excitable cells. With the increasing use of multiple electrode recordings and fMRI imaging studies, traveling waves of activity have become more important objects of study. I will discuss some recent work in existence, shape, and stability of waves of excitation in simplified neural field models. Here the neural connections are locally excitatory, distally inhibitory, and there is a single threshold for cells to achieve firing. I will then discuss ongoing work dealing with two-level firing rates, and self-organizing neural fields with neural plasticity.

ANDREW BELMONTE

W.G. Pritchard Laboratories, Department of Mathematics, Pennsylvania State University, University Park, PA 16802

Impact of a Solid into a Viscoelastic Micellar Fluid

We present an experimental study of the impact of a solid sphere on the free surface of a viscoelastic worm-like micellar fluid. Spheres of various sizes and densities are dropped from different heights above the fluid surface, below which transient oscillations are observed. It is well known that entry of a solid into a Newtonian fluid is characterized by the Froude number; our measurements of sphere penetration scale with the ratio of kinetic energy to the elastic modulus of the fluid. The cavity formed by the sphere also undergoes transitions from a smooth to fractured surface texture, dependent on both the Deborah number and the ratio of gravitational force to elasticity. Analogies between this system and impact in granular materials will also be discussed.

~~ANDREA BERTOZZI~~

~~Department of Mathematics, University of California at Los Angeles, 405 Hilgard Avenue, Los Angeles, CA 90095-1555~~

~~***Shocks in Driven Liquid Films***~~

~~Driven contact line problems in thin liquid films are an active area of research. The mathematical theory of shock waves has recently been shown to play an important role in our understanding of basic properties of the contact line motion. I will present the theory for two recently studied experimental systems: (1) Thermally driven films counterbalanced by gravity are described by a scalar conservation with a non-convex flux. Such systems are known to produce "undercompressive shocks" in which characteristics emerge from the shock on one side. (2) A related problem is that of particle laden flow driven by gravity. The differential settling rate of the particles with respect to the fluid results in the formation of double shock fronts which are solutions of a system of two conservation laws for the motion of the species. Comparison between theory and experiment will be discussed, along with open mathematical problems directly related to the experiments.~~

RICHARD BERTRAM

Department of Mathematics and Institute of Molecular Biophysics, Florida State University, Tallahassee, FL 32306

Modeling Network Interactions Between the Hypothalamus and Pituitary

The pituitary is the primary hormone-releasing gland. It contains a number of different cell types, including lactotrophs, gonadotrophs, thyrotrophs, melanotrophs, somatotrophs, and corticotrophs. These cells secrete different hormones, and each hormone has numerous cell targets in the brain and elsewhere in the body. The timing of hormone release is crucial to its function, and is controlled by the hypothalamus. This brain region is composed of nuclei that influence the pituitary cells in a variety of ways. As one might expect for a highly regulated system, there is two-way feedback between the pituitary and the hypothalamus. We will discuss our current effort to understand the network interactions involved in the regulated secretion of the hormone prolactin from pituitary lactotrophs. During the first ten days of pregnancy in rats, this hormone is secreted in a rhythmic fashion, consisting of two pulses per day. These pulses have functional significance, since pregnancy is aborted if the prolactin secretory pattern is suppressed. The model we have developed consists of coupled nonlinear delay differential equations. Model variables represent the gross activity level of neurons in two nuclei of the hypothalamus, a population of interneurons, and pituitary lactotrophs. This model is currently being used to design experiments in collaborating laboratories.

LORA BILLINGS

Department of Mathematical Sciences, Montclair State University, Upper Montclair, NJ 07043

Chaotic Desynchronization of Multi-Strain Diseases

As we become more sophisticated in our resources to fight disease, pathogens become more resilient in their means to survive. Multi-strain viruses now exhibit antibody-dependent enhancement (ADE), in which infection with a single serotype is asymptomatic, but infection with a second serotype leads to serious illness accompanied by greater infectivity. This talk will present a multi-strain model with ADE based on the dynamics of dengue in Thailand. We study the general dynamics and bifurcation structure of the system. For sufficiently small ADE, the number of infectives of each serotype synchronizes, with outbreaks occurring in phase. However, when the ADE increases past a threshold, the system becomes chaotic, and infectives of each serotype desynchronize. A thorough understanding of the dynamics is the best way to prescribe appropriate defensive measures without unintended consequences.

VICTORIA BOOTH

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-1109

Mean Theta Phase of Model CA1 Pyramidal Cell Firing Changes with Location of Synaptic Stimuli

In recordings from rat hippocampal CA1 place cells during learning and subsequent sleep episodes, Poe et al. (2000) observed changes in place cell activity during reactivation of cell firing in REM sleep. Specifically, the mean phase of place cell firing in relation to the theta rhythm membrane oscillation reversed during REM reactivation over the course of several days as the animal became familiar with the environment. Cells with place fields in an initially novel environment switched from firing near the theta rhythm peaks to firing near the theta troughs during REM while maintaining their theta peak activity during waking exploration. In a computational modeling study, we investigated neural mechanisms underlying this experience-dependent theta phase reversal by incorporating the differential in the phase of the theta rhythm drive at the two excitatory afferent pathways to CA1 and testing model CA1 cell responses to inputs at these two paths. We will discuss simulation results that show that firing in response to proximal (Schaffer collateral) synaptic

input occurs preferentially near the peaks of the theta membrane oscillation while firing in response to distal dendritic (temporo-ammonic) synaptic input occurs near the theta troughs. These results support the hypothesis that the gradual shift to theta trough firing would result from a growing potentiation of synapses in the direct, temporo-ammonic pathway while initially-potentiated Schaffer collateral synapses are depotentiated.

JARED BRONSKI

Department of Mathematics, University of Illinois, 1409 W. Green Street, Urbana, IL 61801

The Periodic Modulational Instability

We consider the stability of standing wave solutions to the nonlinear Schrodinger (NLS) equation, $i\psi_t = \psi_{xx} + V(x)\psi \pm |\psi|^2 \psi$, where the potential $V(x)$ has period 1 and the standing wave solution is either periodic ($\psi(x+1,t) = \psi(x,t)$) or antiperiodic ($\psi(x+1,t) = -\psi(x,t)$). We study the Floquet spectrum of the non-self-adjoint operator governing stability. We use the Hamiltonian structure of the problem, together with some symmetries, to derive a sufficient condition for the existence of a modulational instability. In particular, we show that the stability of plane waves of the integrable NLS equation ($V=0$) is unusual, in that one typically has a modulational instability in the defocusing as well as the focusing cases. This is joint work with Zoi Rapti.

JAVIER CABRERA

Department of Statistics, Rutgers University, 471 Hill Center, Busch Campus, Piscataway, NJ 08854

Statistical Analysis of Data from Comparative DNA Microarray Experiments

DNA microarray technology has enabled researchers in functional genomics to monitor gene expression profiles for thousands of genes at a time.

Most microarray experiments are comparative in nature. For example, suppose that we want to know which genes are expressed differently in two or more types of cells (e.g., a cancer cell and a normal cell). DNA microarray experiments will measure the expression levels of large sets of genes across the different types of cells. By comparing these expression levels, we will know which genes are differentially expressed in the two types of cells.

While these data are often analyzed via a series of t tests, it has been observed that, if the sample size per group is small (as it often is), the dependence between the t test statistic and the pooled standard error estimate leads to an excessively high false positive rate for low variance genes and an excessively high false negative rate for high variance genes. Idiosyncratic features of the data exacerbate the problem: the data are longer tailed than a Gaussian, the variability of a gene depends on its expression level, and the genes are co-dependent in clumps. We propose a model, which posits minimal distributional assumptions, and a conditional t suite of tests, which produces a critical envelope instead of a critical value, to analyze such data. The model is also generalized to statistical methods for multiple comparisons. This research was done jointly with Dhammika Amaratunga, J&J PRD.

RUSSELL CAFLISCH

Department of Mathematics, University of California Los Angeles, 405 Hilgard Avenue, Los Angeles, CA 90095-1555

Singularities in Incompressible Fluid Dynamics

Singularities can occur in incompressible fluid flow due to amplification of vorticity along a fluid interface and possibly in the interior of the flow. This talk will discuss results on singularity formation for a variety of different fluid flows and the physical significance of these results.

FIORALBA CAKONI

Department of Mathematical Sciences, University of Delaware, Newark, DE 19716-2553

Electromagnetic Inverse Scattering for a Buried Object

We are concerned with the problem of determining the shape of an obstacle embedded in a known inhomogeneous background from a knowledge of the scattered electric and magnetic fields measured on a given surface.

Our contribution is to present a method for solving this problem that

1. avoids the need for any a priori knowledge of the physical properties of the scattering object.
2. avoids the need to compute the Green's function for the background media.

The method is a combination of the linear sampling method with the gap reciprocity functional. Like the linear sampling method, our method is based on the characterization of the boundary of the scattering object by the solution of a linear integral equation that is set up independently of the physical properties of the object. But, since we are making use of both the (measured) electric and magnetic fields as our data, we can avoid the need to know the Green's function of the background media.

Numerical examples will be presented showing the performance of our method. This research is joint work with Houssein Haddar and M'Barek Fares from INRIA, Paris.

NILANJAN CHATTERJEE

Biostatistics Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, DHHS, 6120 Executive Blvd., Rockville, MD 20852

Semiparametric Maximum Likelihood Estimation Exploiting Gene-Environment Independence in Case-Control Studies

We consider the problem of maximum-likelihood estimation in case-control studies of gene-environment associations with disease when genetic and environmental exposures can be assumed to be independent in the underlying population. Traditional logistic regression analysis may not be efficient in this setting. We study the semiparametric maximum likelihood estimators of logistic regression parameters that exploit the gene-environment independence assumption and leave the distribution of the environmental exposures to be nonparametric. We use a profile-likelihood technique to derive a simple algorithm for obtaining the estimator and study the asymptotic theory. The results are extended to situations where genetic and environmental factors are independent, conditional on some other factors. Simulation studies investigate small sample properties. The method is illustrated using data from a case-control study designed to investigate the interplay of BRCA1/2 mutations and oral contraceptive use in the aetiology of ovarian cancer.

STEPHEN CHILDRESS

Courant Institute of Mathematical Sciences, 251 Mercer Street, New York, NY 10012

Flapping Flight as a Bifurcation in Frequency Reynolds Number

We describe observations of swimming by a swimming mollusc which suggest that the small wings used by it in "forward flight" become effective for forward motion only above a critical Reynolds number based upon flapping frequency. Models for this bifurcation are developed. We also describe experiments and computations carried out in the Courant Applied Mathematics Laboratory, involving a freely flapping blade, which elaborate the nature of the bifurcation. A new experiment utilizing a small passive flapper in an oscillating airflow suggests a similar bifurcation to hovering flight.

VITTORIO CRISTINI

Department of Biomedical Engineering, University of California at Irvine, Irvine, CA 92697

Two-Dimensional Chemotherapy Simulations Demonstrate Fundamental Transport and Tumor Response Limitations Involving Nanoparticles

Zheng, Wise, and Cristini, *Bulletin of Mathematical Biology* (2005) developed a multiscale, two-dimensional tumor simulator with the capability of showing tumoral lesion progression through the stages of diffusion-limited dormancy, neo-vascularization (angiogenesis) and consequent rapid growth and tissue invasion. In this paper, we extend their simulator to describe delivery of chemotherapeutic drugs to a highly perfused tumoral lesion and the tumor cells' response to the therapy. We perform 2-D simulations based on a rigorous parameter estimation that demonstrate fundamental convective and diffusive transport limitations in delivering anticancer drug into tumors, whether this delivery is via free drug administration (e.g., intravenous drip), or via 100 nm nanoparticles injected into the bloodstream, and releasing the drug that then diffuses into the tumoral tissue, or via smaller 1-10 nm nanoparticles that are capable of diffusing directly and targeting the individual tumor cell. Even in a best-case scenario involving: constant ("smart") drug release from the nanoparticles; a homogenous tumor of one cell type, which is drug-sensitive and does not develop resistance; targeted nanoparticle delivery, with resulting low host tissue toxicity; and for model parameters calibrated to ensure sufficient drug or nanoparticle blood concentration to rapidly kill all cells in vitro; our analysis shows that fundamental transport limitations are severe and that drug levels inside the tumor are far less than in vitro, leaving large parts of the tumor with inadequate drug concentration. A comparison of cell death rates predicted by our simulations reveals that the in vivo rate of tumor shrinkage is several orders of magnitude less than in vitro for equal chemotherapeutic carrier concentrations in the blood serum and in vitro, and after some shrinkage the tumor may achieve a new mass equilibrium far above detectable levels. We also demonstrate that adjuvant anti-angiogenic therapy "normalizing" the vasculature may ameliorate transport limitations, although leading to unwanted tumor fragmentation. Finally, our results suggest that small nanoparticles equipped with active transport mechanisms (e.g., chemotaxis) would overcome the predicted limitations and result in improved tumor response. This study is published in Sinek, Frieboes, Zheng, and Cristini, *Biomedical Microdevices* (2004).

L. PAMELA COOK

Department of Mathematical Sciences, University of Delaware, Newark, DE 19716

Flows of Worm-Like Micellar ("Living Polymer") Solutions: Modeling and Predictions

Modeling of, and the resultant predictions and comparison with experiment for, flow of worm-like micellar solutions is presented. Worm-like micelles are self-assembling very long cylindrical structures composed of amphiphilic surfactant molecules. The resultant flexible structures can entangle and thus behave much like polymers in solution, but in addition they break and reform; thus, they are referred to as "living polymers". The microstructure affects the macroscopic flow properties and vice versa. Both a "bead-spring" model (valid for dilute solutions) and a network model (valid for concentrated solutions) are discussed. Through a mesoscale bead-spring description a coupled stress/density model for the flow of micellar solutions is presented. The coupled nonlinear partial differential equation macroscale model comes from a mesoscale derivation which systematically includes finite extent of the bead-spring and slippage/tumbling of the bead-spring. The non-affine motion is related to micellar break-up. Numerical results for cylindrical Taylor-Couette flow are presented. The stress/strain-rate curve exhibits a plateau as observed in experiments, and as well shear banding is predicted both in velocity gradients and in the alignment/orientation of the bead-springs. Future work involving extensional flows and network models is discussed. Joint work with Gareth McKinley (MIT) and Lou Rossi and Paula Vasquez (University of Delaware). This work was supported by an NSF-DMS grant.

DIPAK DEY

Department of Statistics, University of Connecticut, 215 Glenbrook Road, Storrs, CT 06269

Classification of Shapes using Complex Elliptical Shape Distribution

In this presentation, we discuss classification procedures in a shape analysis context. We derive discriminant rules in shape space, while considering shape distributions as members of the complex elliptical family of shape distributions for the data. In particular, we consider the complex Watson distribution as a special case and develop MAP (Maximum A Posteriori Estimates) of parameters involved and calculate misclassification probabilities using Monte Carlo methods. Our proposed methodologies are exemplified through an example, where we are interested in classifying patients into the normal or schizophrenic groups, based on shapes created by MRI (Magnetic Resonance Image) of their brains. The methods provide us with a new way of diagnosis of this medical condition while controlling the error of misclassification. This is joint work with Athanasios Micheas of the University of Missouri, Columbia.

ANNA GEORGIEVA

Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, NJ 07936

Mechanistic Systems Biology Modeling Applied to the Pre-Clinical Cardiac Safety Assessment of a Pharmaceutical Compound: From Channels to Cells to Tissue

This presentation will focus on an integrative, mechanistic systems modeling approach to cardiac electrophysiology. Via an integrated suite of models from channel to cellular to ultimately, tissue levels, we used specific compound data from limited pre-clinical cardiac electrophysiological studies, such as channel assay as well as one action potential assay, to: (i) estimate missing compound data at the channel level through the application of reverse-engineering techniques; (ii) integrate measured and estimated channel values into action potentials at the cellular level; and (iii) further integrate such cellular responses to predict transmural ventricular responses, which in effect, represent ECG-like features at the tissue level, such as QT prolongation and transmural dispersion of repolarization.

The combination of experimental data and computer-based predictions of compound-induced changes at every level of cardiac biology organization (channel, cell, tissue) provides valuable, quantifiable information to aid in the pre-clinical cardiac safety assessment of new compounds. Joint work with G. Helmlinger, D. Bottino (Novartis Pharmaceuticals, USA), B. Dumotier, M. Traeberdt (Novartis Pharmaceuticals, CH), S. Lett (The BioAnalytics Group LLC, USA), C. Penland (Predix Pharmaceuticals, USA), and A. Stamps (Dept. of Chemical Engineering, University of South Carolina).

RICHARD HABERMAN

Department of Mathematics, Southern Methodist University, Dallas, TX 75275-0156

Vector Soliton Collision Dynamics in Nonlinear Optical Fibers

We consider the interactions of two identical, orthogonally polarized vector solitons in a nonlinear optical fiber with two polarization directions, described by a coupled pair of nonlinear Schrödinger equations. We study a low-dimensional model system of Hamiltonian ordinary differential equations (ODEs) derived by Ueda and Kath and also studied by Tan and Yang. We derive a further simplified model, which has similar dynamics but is more amenable to analysis. Sufficiently fast solitons move past each other without much interaction, but below a critical velocity, the solitons may be captured. In certain bands of initial conditions, the solitons are initially captured, but separate after passing each other twice, a phenomenon known as the two-bounce or two-pass resonance. We derive an analytic formula for the critical velocity. Using matched asymptotic expansions for separatrix crossing, we determine the location of these resonance windows. Numerical simulations of the ODE models show they compare quite well with our asymptotic theory. This is joint work with Roy Goodman (NJIT).

SHELBY HABERMAN

Center for Statistical Theory and Practice, Educational Testing Service, Rosedale Road, Princeton, NJ 08541

Identifiability of Parameters in Item Response Models with Unconstrained Ability Distributions

If a parametric model for the ability distribution is not assumed, then the customary two-parameter and three-parameter logistic models for item response analysis present identifiability problems not encountered with the Rasch model. These problems impose substantial restrictions on possible models for ability distributions. Attempts to circumvent these problems by use of latent-class versions of two-parameter and three-parameter logistic models are strongly limited by problems of near singularity of the information matrix, even for a modest number of latent classes. Thus efforts to use ability distributions that are not normal must be made with great caution.

W. JACKSON HALL

Department of Biostatistics and Computational Biology, University of Rochester Medical Center, 601 Elmwood Avenue, Box 630, Rochester, NY 14642

Secondary Inference after a Sequential Clinical Trial

Consider the following prototype situation: A two-arm randomized, staggered entry, clinical trial, designed to test hypotheses about a treatment effect, has just been completed, utilizing a planned sequential stopping rule. The primary analysis may have been a survival analysis, with treatment arm as the sole risk factor, or one based on the difference between mean responses after a prescribed observation period. No doubt, the trial statistician knew how to take the stopping rule into account in carrying out primary inferences, both hypothesis testing and estimation. But now secondary questions are raised: (i) Do the data provide evidence of a main effect for gender? (ii) Do the data provide evidence of a treatment-by-gender interaction? (iii) How big are any such effects?

Standard analyses that ignore the stopping rule are biased. We review a very simple example to clarify the issues.

We then describe the sequential primary inference problem as a stopping problem for a Brownian motion that approximates the score process for the trial on an information time scale. The secondary inference may then be based on the conditional distribution of another stopped Brownian motion (efficient score) given the primary process with modified stopping boundaries. The formal solution is summarized.

We exemplify with an analysis of a 2-strata clinical trial of implanted cardioverter defibrillators-- the problem that stimulated the research.

JOHN HARRIS

Department of Theoretical and Applied Mechanics, University of Illinois, Urbana-Champaign, IL; Ctr. QEFP Catalysis Bldg., Northwestern University, 2137 N. Sheridan Rd., Evanston, IL 60208

Coupled Elastic Waveguide Modes

The talk begins by describing an example of elastic-waveguide mode coupling at low frequencies: the curvature of an elastic ring couples the longitudinal and flexural propagating modes. Then a more complicated problem containing coupling is analyzed in detail; this is the main subject of the talk: A layer of homogeneous, isotropic, elastic material overlays a faster substrate. At the interface, a long inclusion, whose shear wavespeed is less than that of the layer and whose thickness varies, is introduced. It is imagined that the lowest surface-wave mode of the structure is incident to the growing inclusion. Numerical calculations show that the growth of the slow inclusion draws this lowest mode into an interval where it couples to the second mode, thus exciting it. This process is repeated when the second mode is drawn into

an interval where it couples to the third. Outside of these localized intervals, the modes propagate independently of one another (that is, adiabatically). The problem is formulated and solved within a framework of coupled local modes. An indication of a future asymptotic analysis of the wave fields in the coupling interval closes the talk.

JAN HESTHAVEN

Division of Applied Mathematics, Brown University, 182 George Street, Box F, Providence, RI 02912

Solving the Time-Domain Maxwell's Equations with Uncertainties

While the modeling of EM processes for known geometries and materials has advanced dramatically over the last decade, these advances are often lost when comparing with actual measurements. This can, to a large extent, be attributed to insufficient knowledge of the geometries, materials, dynamics, illumination, etc. of the actual experiment. To achieve a better agreement, one must include this uncertainty into the computational models.

In the past, statistical methods, e.g., Monte-Carlo-type methods, have been the dominant tool for such studies. However, this approach is expensive due to the slow convergence in the number of samples, and it is not feasible to use such techniques for large scale problems with uncertainty, in particular, when higher order statistical moments are needed.

In this talk, we introduce the use of homogeneous chaos expansions for the direct modeling of uncertainty in EM processes, modeled by the linear time-domain Maxwell's equations. This approach, which is probabilistic rather than statistical in nature, effectively transforms the stochastic problem into a sequence of deterministic equations, which we subsequently solve efficiently using a high-order accurate discontinuous Galerkin method.

We consider examples involving uncertainties in illumination/source, material properties, and geometric shapes to show the potential for obtaining results 50-100 times faster than computed to similar accuracy using Monte Carlo methods. Joint work with Laura Lurati (Division of Applied Mathematics, Brown University) and Cedric Chauviere (Laboratoire de Mathématiques Appliquées, Université Blaise Pascal, 63177 Aubière Cedex, France).

PHILIP HOLMES

Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, NJ 08544

A Central Pattern Generator for Insect Locomotion

We model a central pattern generator (CPG) for insect locomotion. We reduce ion-channel, Hodgkin-Huxley (HH)-type models of bursting neurons to single-variable phase oscillators and assemble a circuit of six representative interneurons that each feed a fast and a slow motoneuron. Our phase description retains sufficient detail to allow investigation and prediction of biophysical parameter changes, and it encompasses stepping frequency, duty cycle, and motoneuron output variations observed in cockroaches. The model's modular form allows dynamical analyses of individual components and the addition of other components, so we expect it to be more generally useful.

The methods involve computation of phase response curves, averaging, and elementary ideas from symmetric networks. Joint work with Raffaele Ghigliazza.

FRANK HOPPENSTEADT

Courant Institute of Mathematical Sciences, New York University, 70 Washington Square, New York, NY 10012

Dynamics of Coalitions in Aggregates and Networks of Oscillators

Consider a large array of harmonic oscillators whose center frequencies are arranged in increasing order, $w_1 < w_2 < w_3 < \dots < w_N$. These oscillators need not be arranged in any order in the three dimensional space where they reside; this addressing is in terms only of their center frequencies. We first consider an aggregate where the center frequencies are perturbed by a common external signal:

$$d^2x_j/dt^2 + (w_j^2 + p(t)) x_j = 0,$$

$j=1,2,3,\dots,N$. Here the common parametric forcing is $p(t) = \epsilon \cos \mu t$, which has frequency μ and amplitude ϵ (note that μ and ϵ do not depend on j). Such equations are well-studied, and it is known how to determine the number and identity of oscillators that are entrained at a common frequency. We refer to such a group of synchronized oscillators as being a *coalition*. There is no physical connection between these oscillators, but if the center frequencies are sufficiently dense between w_1 and w_N , the diagram of response frequencies across the array for a given value of μ resembles a staircase having treads located at rational number multiples of μ . The treads describe coalitions. Moreover, as μ is changed, the staircase changes. Thus, the coalitions move through the population of oscillators, as determined by relationships between μ and the center frequencies. There is a rich structure of coalitions of various sizes (i.e., the lengths of treads) that is indexed by (ϵ, μ) in this simple network. We use observations about the linear system to reveal coalitions in aggregates of nonlinear oscillators of the form

$$d^2x_j/dt^2 + \alpha dx_j/dt + (w_j^2 + p(t)) \cos x_j = 0$$

Such equations are well studied in physics, engineering, and mathematical neuroscience. We demonstrate how, within a coalition, information can be carried in the timing of oscillations (i.e., in phase deviations). Finally, we consider networks where

$$p(t) = \epsilon \cos \mu t + C_j(\sin x(t))$$

where $C_j(\sin x(t))$ describes connections to site j from all other oscillators. We use known methodologies from mathematical neuroscience to construct connections C that embed desired patterns of phase deviations, and we show that these patterns can be made to appear or disappear by changing the single parameter μ . Supported in part by MARCO grant 2003-NT-1107.

ANETTE (PEKO) HOSOI

Department of Mechanical Engineering, MIT, 77 Massachusetts Avenue, Cambridge, MA 02139

From Sliding Paper to Crawling Snails: Novel Applications of Thin Films

Liquid thin films have long been studied in the context of industrial, biological, and geophysical applications from spin coating in microcircuit fabrication to the liquid lining in the lung. In general, typical length-scales in these systems are set by surface tension. However, when the film is bounded by a flexible membrane, elasticity takes on the role of surface tension. We discuss some of the consequences of substituting elastic effects for surface tension in the context of several commonly observed phenomena, such as sliding paper and floppy drives, and in more exotic applications such as microfluidic switches and robotic snails.

ASHWANI K. KAPILA

Department of Mathematical Sciences, Rensselaer Polytechnic Institute, Troy, NY 12180

Detonations in Heterogeneous Explosives: Model and Computational Results

A heterogeneous explosive is morphologically complex. Upon application of an igniting stimulus the microstructure responds heterogeneously in two salient ways. First, the mechanical response is nonuniform in terms of local deformation and the associated deposition of energy. Second, the mechanical nonuniformity gives rise to thermal, and hence kinetic nonuniformity, so that ignition occurs at certain preferential sites, or hot spots, where the local temperature is significantly higher than the bulk temperature. In due course, the hot spots grow and merge, and reaction spreads from the hot spots to the bulk. As a rule, the scale of the device employing the explosive is substantially larger (of the order of tens of centimeters) than the micro scales (of the order of tens of microns) at which energy is released. Thus, predicting macro-scale device behavior by direct computation at the micro scale would be prohibitively expensive. One gets around this difficulty by posing continuum models that contain approximate submodels for the micro-scale input. Either implicitly or explicitly, these approximations represent an average behavior, in effect homogenizing the fine-scale description if such a description were available. This talk will describe one such model, its predictions, and its strengths and weaknesses.

THOMAS B. KEPLER

Duke University, Center for Biostatistics and Bioinformatics, Durham, NC 27708

Inducible Reorganization of the Immune System

The vertebrate immune system consists of a great diversity of motile cells whose activities become coordinated during infection. This orchestration is mediated by signaling molecules either secreted (cytokines) or engaged by direct cell-cell contact. Pathogenic microorganisms (and other stimuli) induce internal changes in the responding immune cells which, in turn, lead to spatial reorganization of these cells in a process arguably akin to a phase transition.

The models we are developing to explore these phenomena represent the cells of the immune system as individual "agents" with non-trivial internal states. The motion of these agents in a three-dimensional continuum is described by a continuous-time stochastic process, as are their internal dynamics. Soluble factors, such as cytokines, are represented as fields obeying reaction-diffusion equations on the continuum. Both the internal states of the agents and their motions are responsive to the state of the cytokine fields, which, in turn, are influenced by the agents, which act as time-dependent sources and sinks. I will present these models and illustrate them with examples of the inflammation-mediated spatial reorganization of the immune system.

J. NATHAN KUTZ

Department of Applied Mathematics, University of Washington, Seattle, WA 98195-2420

Computing Spectra of Linear Operators Using Hill's Method

In order to establish the stability of an equilibrium solution \mathbf{U} of an infinite-dimensional dynamical system $\dot{\mathbf{u}} = \mathbf{X}(\mathbf{u})$, one is interested in the spectrum of the linear operator $\mathbf{L}[\mathbf{U}]$ obtained by linearizing the dynamical system around \mathbf{U} . We use a spectrally accurate method for the computation of such spectra. In essence, the method is due to Hill, who used it in the study of the equation that now bears his name. The method is particularly well suited to the case of periodic \mathbf{U} , but is not restricted to it. By incorporating the fundamentals of Floquet theory, an almost uniform approximation to the entire spectrum is obtained, as opposed to an approximation of a few selected elements. The numerical component of the method is limited to (i) choosing the size of the matrices to be used, and (ii) an eigenvalue solver, such as the QR algorithm. Compared to often-used finite-difference approaches, the method is orders of magnitude faster for comparable accuracy.

HERBERT LEVINE

Department of Physics, University of California at San Diego, La Jolla, CA 92093

Fluctuation Effects in Intracellular Calcium Signaling

Most biophysical models of the nonlinear spatially-extended dynamics of intracellular calcium assume that one can neglect stochastic fluctuations. Recent experiments have, however, directly implicated noise due to the opening and closing of finite numbers of channels in a variety of dynamical behaviors. This talk focuses on some of the mathematical issues which arise when one adds fluctuations to non-equilibrium pattern-forming systems.

BERNARD J. MATKOWSKY

Engineering Sciences and Applied Mathematics, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208-3125

Flames as Discontinuity Surfaces in Gasdynamic Flows

Viewed on a hydrodynamic scale, flames in experiments are often thin so that they may be described as gasdynamic discontinuities separating the dense, cold fresh mixture from the light, hot burned products. The original model of a flame as a gasdynamic discontinuity was due to Darrieus and to Landau. In addition to the fluid dynamical equations, the model consists of a flame speed relation describing the evolution of the discontinuity surface, and jump conditions across the surface which relate the fluid variables on the two sides of the surface. The Darrieus-Landau model predicts, in contrast to observations, that a uniformly propagating planar flame is absolutely unstable and that the strength of the instability grows with increasing perturbation wave number so that there is no high wave number cutoff of the instability. The model was modified by Markstein to exhibit a high wave number cutoff if a phenomenological constant in the model has an appropriate sign. Both models are postulated, rather than derived from first principles, and both ignore the flame structure, which depends on chemical kinetics and transport processes within the flame. At present, there are two models which have been derived, rather than postulated, and which are valid in two nonoverlapping regions of parameter space. Sivashinsky derived a generalization of the Darrieus-Landau model which is valid for Lewis numbers (ratio of thermal diffusivity to mass diffusivity of the deficient reaction component) bounded away from unity. Matalon and Matkowsky derived a model valid for Lewis numbers close to unity. Each model has its own advantages and disadvantages. Under appropriate conditions, the Matalon-Matkowsky model exhibits a high wave number cutoff of the Darrieus-Landau instability. However, since the Lewis numbers considered lie too close to unity, the Matalon-Matkowsky model does not capture the pulsating instability. The Sivashinsky model does capture the pulsating insta-

bility, but does not exhibit the high wave number cutoff of this instability. Here, we derive a model consisting of a new flame speed relation and new jump conditions, which is valid for arbitrary Lewis numbers. It captures both monotonic and pulsating instabilities, and exhibits a high wave number cutoff for each. The flame speed relation includes the effect of short wave lengths, not previously considered, which leads to stabilizing transverse surface diffusion terms.

MASAYASU MIMURA

Department of Mathematics, Meiji Institute for Mathematical Sciences (MIMS), School of Science and Technology, Meiji University, 1-1-1 Higashimita, Tamaku, Kawasaki 214-8571, Japan

Self-Organized Patterns in Bacterial Colonies

It is observed by Budrene and Berg [1], [2] that chemotactic bacteria form regularized but complex patterns. They emphasized that such patterns are generated in a self-organized way due to the balance among substrate consumption, cell proliferation, excretion of attractant, and chemotactic motility. My talk is to understand theoretically how these four elements can possibly generate such patterns by using a mesoscopic continuous model.

[1] E. O. Budrene and H.C. Berg: Nature, 349 (1991) 630-633.

[2] E. O. Budrene and H.C. Berg: Nature, 376 (1995) 49-53.

ANDRÉ NACHBIN

Department of Pure and Applied Mathematics, IMPA, Estrada Dona Castorina 110, Rio de Janeiro 22460-320, Brazil

New Nonlinear Water Wave Models over Highly Variable Topographies

Recently, we formulated a weakly nonlinear, weakly dispersive terrain-following Boussinesq system (SIAP 2003) in order to study solitary waves over highly variable (random) topographies. The modeling allows for multiply-valued topography profiles. The reduced model is obtained through the standard power series expansion for the velocity potential along the free surface. In this talk, we will give a brief overview of this earlier study and present a new, fully dispersive, Boussinesq system (Phys.Rev.Lett., 2004) that generalizes the terrain-following system mentioned above. The full linear (hyperbolic tangent) dispersion relation is entirely retained by constructing a Dirichlet-to-Neumann (DtN) map along the top boundary of the highly corrugated strip representing the channel. Moreover, an efficient FFT-based method naturally arises from the DtN boundary integral formulation.

MONIKA NITSCHKE

Department of Mathematics and Statistics, University of New Mexico, Humanities 415, Albuquerque, NM 87131-1141

Regularizations of Vortex Sheet Motion

The vortex sheet is a mathematical model for a shear layer in which the layer is approximated by a surface. Generally, vortex sheets develop singularities in finite time. To approximate the fluid past this time, the motion is regularized, and the sheet is defined as the limit of zero regularization. However, very little is known about this limit, not even whether it is unique or depends on the regularization. I will discuss several regularization mechanisms, including physical ones such as fluid viscosity, and purely numerical ones such as the vortex blob and the Euler-alpha methods. I will show results for a model problem and discuss some of the unanswered questions of interest.

ANDREW NORRIS

Department of Mechanical and Aerospace Engineering, Rutgers University, 98 Brett Road, Piscataway, NJ 08854

Crack Front Waves using Matched Asymptotic Expansions

A steadily propagating crack with a straight edge is not stable under perturbation. This surprising result was found ten years ago, but is still not fully understood despite many studies using analytical, computational, and experimental techniques. The problem is exacerbated by the difficulty involved in the analysis of even the simplest case of mode I crack propagation. This talk provides a simple and physically appealing method for analyzing the crack front stability based on matched asymptotics. The inner region of the crack with the wavy edge is considered separately from the outer problem of the steadily propagating straight crack. By matching the solutions, the stability criterion is found. The simplicity of this method allows us to consider more complicated crack scenarios, including a crack on a bimaterial interface, and a crack with a cohesive zone. The talk will emphasize the power of using matched asymptotics to analyze stability of propagating material fronts, of which the crack is but one example.

HILLARY OCKENDON

OCIAM, Mathematical Institute, 24-29 St. Giles', Oxford OX1 3LB, United Kingdom

Turbulent Flow in Long Thin Channels

Turbulent compressible flow in long thin channels arises in many practical situations from a train in a long tunnel to a pressure transducer to the effect of a rock fall in a mine. This talk will describe the Fanno model for such flows and show how it can be applied in some of these examples.

JOHAN PAULSSON

Cambridge Computational Biology Institute, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge, CB3 0WA, United Kingdom

Suppressing Fluctuations in Living Cells

Life in single cells is dictated by chance: Reactions that involve small numbers of molecules generate spontaneous fluctuations that enslave all dependent processes. Such noise can be suppressed by negative feedback control where a molecule directly or indirectly represses its own synthesis. I will use a reinterpretation of the Fluctuation-Dissipation Theorem to demonstrate generic trade-offs where suppressing one type of fluctuations automatically amplifies other types. Strongly nonlinear strategies for breaking the trade-offs will also be discussed.

JOHN PEARSON

Los Alamos National Laboratory, P.O. Box 1663, Los Alamos, NM 87545

Equivalence and Identification of Markov Models for Ion Channel Kinetics

Deducing plausible reaction schemes from single channel current traces is time consuming and difficult. The goal is to find the simplest possible scheme, but there are many ways to connect even a small number of states (more than 2.5 million schemes with four open and four closed states). Many of those schemes make identical predictions, and even an exhaustive search over model space does not address the problem of how to represent the many equivalent schemes that may result.

We have found a canonical form that can express all reaction schemes for binary channels. This form has the minimal

number of rate constants unlike other canonical forms such as the well-established "uncoupled" scheme. For many models proposed in the literature, our form has dramatically fewer links than the uncoupled form. The new form, which is based on the number of independent open-closed transitions, leads naturally to new strategies for searching for the simplest model.

FERNANDO REITICH

University of Minnesota, 127 Vincent Hall, 206 Church St., S.E., Minneapolis, MN 55455

A New High-Order High-Frequency Integral-Equation Method for the Solution of Wave Scattering Problems

The effort and interest in the design of improved algorithms for computational electromagnetics and acoustics applications has consistently grown over the last twenty years as these simulations have become relevant in an increasing number of fields and have been facilitated by remarkable developments in computing resources. Still, current state-of-the-art algorithms are limited by the competing demands of accuracy, which typically requires an increasing number of degrees of freedom to resolve on the scale of a wavelength, and efficiency, which favors coarse discretizations. In this talk, we will present a new strategy for the solution of the integral equations of electromagnetic and acoustic scattering that successfully deals with these requirements by avoiding the need to discretize on the scale of the wavelength at high-frequencies, while retaining error-controllability and high-order convergence characteristics. The approach is based on the derivation of an appropriate ansatz for the phase of the (unknown) currents, on explicit treatment of shadow boundaries, and on localized high-order integration around critical points. Joint work with O. Bruno and C. Geuzaine.

JOHN RINZEL

Courant Institute of Mathematical Sciences, New York University, New York, NY 10012

Fast-Decaying Inhibition Can Facilitate Spiking

In classical postinhibitory rebound (PIR), a neuron can generate a spike when it is suddenly released from a long-lasting inhibitory input. During the inhibition, the membrane becomes hyperpolarized and some of the negative feedback (e.g., K^+ -conductance) that is present at rest is reduced – thereby making the cell hyperexcitable. We find that even very brief inhibition $g_{inh}(t)$ can induce a spike or facilitate the response to a subthreshold excitatory input $g_{ex}(t)$ -- enough to cause a spike, if g_{inh} decays fast enough and occurs in a favorable time window preceding g_{ex} . Such pairings can occur by chance during the presentation of random trains of g_{inh} and g_{ex} , and can form a significant fraction of spikes seen in simulated spontaneous states. Such effects occur in various spike-generating models, including the standard HH model. Phase-plane analysis of 2-variable models shows that if PIR occurs, then so does the extreme behavior of spike generation for the limiting brief case of a delta-function pulse of hyperpolarizing current. Such neuron models have two thresholds, for instantaneous jumps in membrane potential, in both the depolarizing and hyperpolarizing direction. Joint work with R. Dodla (Center for Neural Science, New York University).

GARETH J. RUSSELL

Columbia University, New York, NY, and NJIT, Newark, NJ 07102

Modeling Epidemics Based on Uncertain Timing Data

We adapt a likelihood model for the spread of a contagious process on a network of susceptible nodes so that it incorporates uncertainty in the timing of the node-level events (initial infection, the onset of contagiousness, etc.) that comprise the epidemic. The timing uncertainty is modeled as a uniform distribution of probability between known limits

(although other distributions are possible in principle), over which basic transmission functions are integrated. This general model is made specific to a particular case by incorporating functions that describe the nature of transmission process between nodes (e.g., nearest-neighbor only, rate as a function of Euclidean distance, etc.). Each specific model may be fit to data by maximizing its overall likelihood as a function of those parameters that describe the transmission process. Because the output of the model is a likelihood, it can be used to choose between alternative models of the transmission process. Increasing timing uncertainty reduces our ability to distinguish between models, and we present two case studies of the spread of tree pathogens which provide contrasting examples of how much can be learned from efforts to track the spread of the disease. Joint work with Jacqueline W. T. Lu (City of New York, Department of Parks and Recreation).

WILLIAM M. SALLAS

Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, NJ 07936

Deconvolution and Regularization: Connection to Linear Least Squares and Application to Estimating Insulin Secretion in Patients with Type 2 Diabetes

Deconvolution is applied to estimating insulin secretion in patients with type 2 diabetes in a clinical study with a meal challenge to assess β -cell function. C-peptide, which is cosecreted with insulin in an equimolar ratio, is preferred to insulin as a basis for estimating insulin secretion. Mathematically, we need to solve for the insulin secretion rate r given the impulse response h , and plasma C-peptide c , where

$$c(t) = \int_{-\infty}^t h(t-s)r(s)ds.$$

The impulse response is the pharmacokinetic model for an intravenously administered unit bolus dose of C-peptide when the endogenous C-peptide production has been suppressed, i.e.,

$$h(t-s) = (1/V)[fe^{-\alpha(t-s)} + (1-f)e^{-\beta(t-s)}]$$

where V , f , α , and β are a subject's parameters, which are generally unknown. However, from clinical studies, these parameters have been estimated in terms of a subject's disease status, body surface area, age, and gender (van Cauter et al. 1992). Thus, h becomes known. Three approaches to solving for $r(s)$ are reviewed: 1) solutions of linear systems of equations assuming either a piecewise constant function or a piecewise continuous linear function $r(s)$, 2) linear least squares solutions assuming smoothness criteria on the insulin secretion rates, and 3) a model that estimates insulin secretion rates accounting for changes in plasma glucose (Mari et al. 2002). Joint work with Huadong Tang (Wyeth Pharmaceuticals, Pearl River, New York) and Andrea Mari (C.N.R. Institute of Biomedical Engineering, Padova, Italy).

CLYDE SCANDRETT

Mathematics Department, Code MA/Sd, Naval Postgraduate School, Monterey, CA 93943

Cancellation Techniques in Scattering from Fluid Loaded Plates

Reduction and elimination of scattered acoustic signals by cancellation techniques involving piezoelectric materials is considered. Following formulation of the problem and a brief description of underlying principles, results on a few canonical problems will be given. These include radiation and scattering from heavy, fluid-loaded plates with either singly or periodically attached piezoelectric elements. Analytical treatments employing invariant embedding methods, Floquet theory, and asymptotics, will be compared with finite difference/boundary integral techniques.

ARND SCHEEL

Institute for Mathematics and Its Applications and School of Mathematics, University of Minnesota, 206 Church Street S.E., Minneapolis, MN 55455

Following Coherent Structures

Spiral waves, line defects, sources, and sinks, are examples of localized structures embedded in a background of waves with characteristic wavenumbers. Although these structures often appear to be very robust, the persistence under variations of system parameters turns out to be a delicate question due to essential spectrum caused by the background waves. We show how methods from dynamical systems can be used to prove robustness and bifurcation theorems, hence enabling a path-following type exploration of large-scale nonequilibrium systems. We also present examples where our approach exhibits structural barriers to pathfollowing: coherent structures can suddenly disappear.

PRANAB K. SEN

Departments of Biostatistics and Statistics & Operations Research, University of North Carolina at Chapel Hill, NC 27599-7420

The Curse of Dimensionality in Genomics: Beyond the Likelihood Paradigm

Genomic sequences involve an enormously large number of positions or sites (loci), each one having a purely qualitative categorical response (4 categories A, C, G, T for DNA nucleotides and some 20 amino acids in RNA codons). Thus, there is a very high dimensional categorical data model with the number of positions (K) often much larger than the number of sequences (n), so that $K \gg n$. The prospect of incorporating the asymptotics for the likelihood function and all its ramifications (including the empirical likelihood) depends not only on managing the complex categorical data models arising in this context but also on low sample size high-dimensional perspectives. As of now, there has not been any great methodologic advances in this respect. Statistical learning or knowledge discovery and data mining tools are useful from the computational point of view, but they may lack, to a certain extent, statistical methodologic support.

Although, it is often assumed that the positions have independent responses, in reality, inter-site stochastic dependence is very much perceptible. Hamming distance-based methodology, therefore, has been advocated recently. They allow a dimension reduction taking into account the inter-site stochastic dependence to a certain extent. Pinheiro et al. (2000, 2005) have considered some MANOVA procedures based on variants of the Hamming distance. There has been some further research in this direction which incorporate second-order decomposability of Hoeffding's (1948) U-statistics, and achieving asymptotic normality of the pseudo-U-statistics arising in this context. Applications to real data have also been considered. Joint work with Hildete P. Pinheiro and Aluisio S. Pinheiro (University of Campinas, Sao Paulo, Brazil).

MICHAEL SHELLY

Courant Institute of Mathematical Sciences, New York University, New York, NY 10012

Swimming Worms, Swimming Sheets

Motivated by experiments on "swimming" by active materials and the locomotion of nematodes, I will discuss different models and simulations of dynamic flexible bodies interacting with fluids. In two very different instances, we study how locomotion is generated by the propagation of large amplitude waves through a body, and how locomotion is affected by changes in the fluidic medium.

ARTHUR SHERMAN

L.B.M., N.I.D.D.K., National Institutes of Health, Building 12A, 12 South Dr. MSC 5621, Bethesda, MD 20892-5621

Metabolic and Electrical Oscillations in Insulin-Secreting Pancreatic Beta-Cells

The hormone insulin is the primary regulator of glucose concentrations in the blood and is important for regulation of consumption and storage of energy from carbohydrates, protein, and fat in general. Insulin concentrations in the blood oscillate with a period of about 5 minutes, and such oscillations have been shown to be disturbed in patients with diabetes. The whole-body oscillations are believed to be driven by oscillations in calcium in the beta-cells of the pancreatic islets of Langerhans, but there has been contention over whether the oscillations are metabolic or electrical in origin. We will discuss a beta-cell model with interacting metabolic and electrical oscillators. Both oscillators serve as readouts of the metabolic state of the cells, but are semi-independent in the sense that either can occur without the other. The two oscillators can also phase-lock in interesting ways. We will illustrate a wide variety of phenomena that can be explained by the combined model, but not readily with only one of the two sub-systems, including the glucose dose response curve and compound oscillations (bursts of bursts).

KATE STEBE

Department of Chemical and Biomolecular Engineering, The Johns Hopkins University, 221 Maryland Hall, 3400 North Charles Street, Baltimore, MD 21218

Surfactant Effects on Drop Detachment

Surfactant effects on the processes leading to drop detachment are studied for the case of a viscous drop detaching in a viscous fluid. In the absence of surfactants, neck pinch-off occurs at the primary neck and displays self-similar dynamics that are governed locally by the balance of capillary and viscous stresses in creeping flow. For viscous drops surrounded by viscous liquids, this regime has been predicted by scaling arguments, verified in experiment, and studied using boundary integral simulations. We study these effects numerically in the presence of surfactants in either the insoluble or adsorption-desorption limits.

In the absence of surfactants, the rate of surface dilatation is most negative just above the primary neck. When surfactants are present, they accumulate there. Since surface tension response is highly nonlinear at concentrations comparable to the maximum packing, regions of high surfactant packing have low local surface tensions and hence low capillary stresses. Necking dynamics are significantly altered. Drops break at the primary neck at low surfactant coverage, at the secondary neck at moderate coverages, or fail to neck at elevated coverages, suggesting a transition from dripping to jetting modes. These transitions are highly dependent on the surfactant sorption dynamics; phase diagrams for drop detachment regimes as a function of surfactant transport dynamics are developed. Joint work with Fang Jin and Nivedita Gupta (Department of Chemical & Biomolecular Engineering, Johns Hopkins University).

PAUL H. STEEN

School of Chemical and Biomolecular Engineering, Center for Applied Mathematics, Cornell University, Ithaca, NY 14853-5201

Dynamics and Stability of Capillary Surfaces: Designing Droplet Switches

A “capillary surface” is a liquid/liquid or liquid/gas interface whose shape is determined by surface tension. For typical liquids (e.g., water) against gas, capillary surfaces occur on the millimeter-scale and smaller where shape deformation due to gravity is unimportant (or on larger scales for soap-film surfaces). Capillary surfaces can be combined to make a switch – a system with multiple stable states. The simplest capillary switch consists of two droplets (or bubbles), coupled by common pressure.

The static response and stability (thermodynamics) of a capillary switch is given by its energy landscape. We use a theorem attributed to Poincaré to predict energy landscapes. Transition from one stable state to another (kinetics) involves the flow of fluid and is necessarily dynamical. Trajectories of the corresponding dynamical system can be computed directly. Or, alternatively, basins-of-attraction can be mapped (robustness). In this case, the fate of certain finite-amplitude disturbances is pre-determined by identifying trapping regions -- regions of disturbances that must always return to (or can never return to) a specified stable base state. We illustrate the prediction of thermodynamics, kinetics, and robustness of capillary switches using bifurcation theory, boundary-integral computations, and energy inequalities, respectively. Predictions are compared to observation of soap-film and droplet-scale realizations in the laboratory.

D. SCOTT STEWART

Mechanical and Industrial Engineering, University of Illinois, 216 Talbot Laboratory, 104 S. Wright Street, Urbana, IL 61801

Verification of Detonation Shock Dynamics by Numerical Simulation in Complex Geometries and in Multi-Material Systems

An extensive body of theoretical work has been carried out to derive front evolution equations that can describe the motion of detonation shocks in both condensed and gaseous explosives. The basic mathematical model for the explosive is the compressible Euler equations for pre-mixed materials, with a single one-step reaction from reactants to products. The explosive is represented by a pressure, volume, reactant progress variable equation of state (EOS), with a reaction rate law with similar arguments. The asymptotic theory of detonation shock dynamics (DSD) is a hydrodynamic flow theory that corrects a planar detonation to account for changes due to the shock curvature and unsteadiness. The asymptotics are based on the assumption that the radius of curvature of the shock is much larger than the reaction zone that supports the shock, and that changes to the shock shape are slow when measured on a time scale of a particle passage time through the reaction zone. Both are realistic for many systems of physical interest. A higher order theory includes unsteady effects such as shock acceleration and other higher derivatives in the surface.

Different forms of the detonation shock evolution equation can be obtained, and their validity and suitability for quantitative prediction of the flow states and the lead shock dynamics depends on the initial state of the flow and how close that state is to the assumption used in the asymptotic description. Of course, the theory for the discrepancy between the asymptotic predictions and the realization of the full system (i.e., a direct numerical simulation, properly resolved or alternatively the realization of the physical experiments themselves) is absent. To make these assessments, one relies on simulation and experiment.

In this talk, the current state of what is known about how well the asymptotic front theory predicts the detonation shock evolution of full systems, will be discussed. This will include a review of the different types of evolution equations obtained, such as the relation between detonation normal velocity and total shock curvature, and inclusion of higher order terms that can even lead to cellular detonation dynamics and chaos. The computed systems will include those with converging and diverging geometries, and complex multi-material interactions of shock with embedded inerts. Finally, if time permits, we will include applications of computed results to the design of novel miniaturized explosive systems currently be developed at University of Illinois.

This work is supported by: U.S. Department of Energy, Los Alamos National Laboratory, U.S. Air Force Research Laboratory, Eglin AFB, Munitions Directorate, and U.S. Air Force Office of Scientific Research, Physical Mathematics.

HOWARD STONE

Department of Engineering and Applied Sciences, Harvard University, 308 Pierce Hall, Cambridge, MA 02139-4307

The Reciprocal Theorem as a Way to Get Something for (Almost) Nothing: (I) The Mobility of Membrane-Trapped Particles and (II) The Normal Force in Sliding Lubrication with Soft Materials

The Stokes equations, which are the linearized version of the Navier-Stokes equations, describe a large number of flow problems in biophysics and the engineering and physics of thin films. In this talk, we describe two variants of these problems, which have a common solution feature, even though the physical contexts are very different. First, we consider the translation mobility of an ellipsoidal particle, which is trapped in a membrane or surfactant-coated interface, and protrudes into the surrounding fluid. The theoretical predictions for the mobility are compared with experimental measurements. Second, we consider the influence of substrate or particle deformability on the sliding lubrication of a sphere near a wall; deformation is responsible for the existence of a force normal to the wall. In both problems, neither of which had been treated successfully before, the Reciprocal Theorem, which is a variant of Green's Theorem, is applied to obtain the final results while bypassing much of the (tedious) calculations associated with a direct approach to these problems.

DE WITT SUMNERS

Department of Mathematics, Florida State University, Tallahassee, FL 32306

DNA Knots Reveal Chiral Packing of DNA in Phage Capsids

Bacteriophages are viruses that infect bacteria. They pack their double-stranded DNA genomes to near-crystalline density in viral capsids and achieve one of the highest levels of DNA condensation found in nature. Despite numerous studies, some essential properties of the packaging geometry of the DNA inside the phage capsid are still unknown. Although viral DNA is linear double-stranded with sticky ends, the linear viral DNA quickly becomes cyclic when removed from the capsid, and for some viral DNA the observed knot probability is an astounding 95%. This talk will discuss comparison of the observed viral knot spectrum with the simulated knot spectrum, concluding that the packing geometry of the DNA inside the capsid is non-random and writhe-directed.

JEAN-MARC VANDEN-BROECK

School of Mathematics, University of East Anglia, Norwich, England NR4 7TJ, United Kingdom

Nonlinear Capillary Waves in Electrified Fluid Sheets

Nonlinear capillary waves propagating on fluid sheets are calculated in the presence of uniform electric fields acting in directions parallel or perpendicular to the undisturbed configuration. The fluids are taken to be inviscid, incompressible, and irrotational. Both conducting and non-conducting fluids are considered. Waves of arbitrary amplitude and wavelengths are calculated and the effects of the electric fields are studied. Fully nonlinear solutions are obtained by boundary integral equation methods. In addition, long-wave nonlinear waves are also constructed using asymptotic methods. As time permits, recent new results for Stokes flows will also be presented. Joint work with Demetrios Papageorgiou (NJIT).

MIGUEL R. VISBAL

Computational Services Branch, Air Vehicles Directorate, Air Force Research Laboratory, WPAFB OH 45433

High-Order Finite-Difference Schemes for Time-Domain Computational Electromagnetics and Acoustics

This presentation will review recent progress achieved in the application of high-order finite difference schemes to the solution of multi-physics conservation laws on curvilinear geometries. We focus primarily on 4th- and 6th-order compact schemes coupled with up to 10th-order low-pass spatial Pade-type filters. Unlike standard numerical damping, the high-order filter operator provides selective dissipation of only the high-wavenumber content of the solution, which is already corrupted by the dispersion errors of the baseline discretization. These spatial algorithms are combined with explicit and implicit time integration methods to examine wave propagation in electromagnetics and acoustics. It is shown that without the incorporation of the filter, application of the high-order compact scheme to non-smooth collocated meshes results in spurious oscillations which inhibit their applicability. Inclusion of a discriminating low-pass high-order filter restores the advantages of the high-order approach even in the presence of localized large grid discontinuities. The filter operator in combination with highly stretched meshes also provides an alternative robust treatment of far field radiation conditions. Proper evaluation of coordinate transformation metrics is adopted to enforce the geometric conservation law on highly distorted and dynamically deforming curvilinear meshes. Finally, incorporation of high-order interpolation techniques and one-sided filter operators enable the extension of the approach to non-coincident overlapped meshes typically encountered in complex geometries.

Z. JANE WANG

Department of Theoretical and Applied Mechanics, 23 Thurston Hall, Cornell University, Ithaca, NY 14853

Fore- and Hind-Wing Interactions in Dragonfly Flight

Dragonflies are one of the most maneuverable insects. A distinctive feature of dragonflies is their use of two pairs of wings instead of one pair. This reflects their ancient origin. As such, understanding the coupling between their fore- and hind-wings might shed light on the evolution of flight based on four wings to that based on two. In this talk, I will describe the tracking of 3D wing motion of a tethered dragonfly, our computational method for simulating multiple wings, and our findings of the role of wing interactions and its effects on the forces and efficiency in hovering dragonfly flight. Joint work with David Russell.

WENDY ZHANG

James Franck Institute, University of Chicago, 5640 S. Ellis Avenue, Chicago, IL 60637

Viscous Entrainment: Singular and Nearly Singular Liquid Spouts

A small air bubble rising in syrup remains spherical. A larger air bubble deforms, developing an increasingly tapered trailing end. For an even larger air bubble, the rising movement is so severe in deforming the bubble that a thin tendril of air is deposited behind the rising bubble. Inspired by such familiar examples of viscous entrainment and recent experiments, we analyze the entrainment dynamics in a simple model problem where a long-wavelength model describes the essential dynamics. We show that both continuous and weakly discontinuous entrainment transitions are possible when the interface shape on the largest length-scales is constrained so that the base of the entrained tendril approaches a conical shape. Finally, we show that two kinds of critical transition exist in the full problem because the scale-invariant dynamics supports a saddle-node bifurcation. After the bifurcation, scale-invariant solutions which can link onto physical large-scale conditions do not exist.

CONTRIBUTED PAPERS

Silas Alben

Harvard University, Division of Engineering and Applied Sciences, 29 Oxford Street, Cambridge, MA 02138

The Fin Ray as a Motion Transducer

Our work considers the mechanics of swimming fish. In collaboration with the Lauder lab at Harvard, we are studying the structure of fish fin rays. Approximately half of all fish species utilize the same basic structure to transduce fin ray shape and motion from a given input force. We present a simple coupled elastica model which uses only geometry and a single elastic constant to obtain the scalings of forces and displacements. We will also present fin ray structures which are optimal for force output.

Christina Ambrosio

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

The Control of Frequency of an Excitable Network Simultaneously Subjected to Multiple Oscillatory Inputs

Biological networks responsible for the generation of rhythmic motor output often involve a pair or pairs of reciprocally inhibitory neurons that burst in alternation. The triggering of such bursts often requires multiple rhythmic inputs, thus, making the network a conditional oscillator. Here, we study a set of reciprocally inhibitory neurons lying in the stomatogastric ganglion (STG) of the stomatogastric nervous system of the crab *Cancer borealis*. This pair of neurons is responsible for generating the gastric mill rhythm which is involved in the digestive process of the animal. The aim of this work is to mathematically explain the role of two oscillatory inputs, a fast pyloric input and a slow modulatory input, that act in coordination to set the frequency of the gastric mill rhythm. In particular, we mathematically explain experimental results in which it is shown that the pyloric input increases the gastric mill frequency when the modulatory input is tonic but does not increase the frequency when the modulatory input is rhythmic. Reduction techniques are used to reduce the modeling set of equations to a study on lower dimensional slow manifolds and the fast transitions between them. This enables us to geometrically show through phase plane analysis, the exact role that each of the inputs plays in determining the frequency of the gastric mill rhythm and it reveals the importance of the timing between such inputs. We conclude that for a small delay between the timing of the pyloric and modulatory inputs, the gastric mill rhythm frequency is only determined by the dynamics of the modulatory input as is seen experimentally in the biological network. For a larger delay, however, the inputs work together to speed the frequency. Joint work with Amitabha Bose (NJIT) and Farzan Nadim (NJIT and Rutgers University).

Roman Andrushkiw

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

On the Spectral Theory of Operator Valued Functions

We study a class of operator-valued functions that are nonlinearly dependent on the spectral parameter and involve operators that may be unbounded and non-symmetric. Spectral properties of this class of functions are investigated in a Hilbert space setting, and sufficient conditions for the existence of the eigenvalues are derived. An iterative method for approximating the eigenvalues is developed, based on a variational characterization of the spectral problem, and some examples are given to illustrate the theory.

Eliana Antoniou

William Paterson University, Mathematics Department, 300 Pompton Road, Wayne, NJ 07470

A Theoretical Simulation of Hematopoietic Stem Cells During Oxygen Fluctuations: Prediction Of Bone Marrow Responses During Hemorrhagic Shock

Assembly of the human genome shows that the biological system of human cannot be explained by ~30,000-40,000 genes. This study is based on the premise that studies on bone marrow (BM) stem cells (lymphohematopoietic stem cell, LHSC) by proteomics, computational biology and genomics could be aided by mathematical models. We use a mathematical model of acute blood loss, which is associated with a sharp decrease in pO_2 , to predict the responses of LHSC in an area distant from the main sinus. The equations were modified to present the responses of two population of LHSC (Proliferating, P; Non-proliferating, N). Hematopoietic responses by the most primitive LHSC were simulated for otherwise healthy individuals who have been subjected to various degrees of blood loss, as represented by 3%, 5% and 20% O_2 . The model is robust and could predict for hematopoietic activity in the area close to the endosteum during low pO_2 when there is acute blood loss. Steady state hematopoiesis at physiological oxygen level (80% O_2) could not be simulated with the equations. Functional assays tested the model with a modified long

term culture initiating cell assay. In the presence of 1%, 3-5% and 20% O₂, highly purified LHSC showed significant increases in cell proliferation when compared to 80% O₂. Thus, the functional studies show that the theoretical model is robust and could be used to gain insights into the biology of LHSC during different degrees of blood loss. The utility of such a model in surgical trauma is discussed.

Felix J. Apfaltrer

Courant Institute of Mathematical Sciences, New York University, 251 Mercer St., New York, NY 10012

Population Density Methods in Two Spatial Dimensions and Application to Neural Networks with Realistic Synaptic Kinetics

We explore the extension to two "spatial" dimensions of a computationally efficient method of simulating networks of neurons. The method is applied to integrate-and-fire neurons with realistic synaptic kinetics. In this method neurons are grouped into populations of similar biophysical properties, and for each population a probability density function (PDF) is constructed. This PDF represents the distribution of neurons over state-space. The evolution equation of the probability density functions is a partial differential-integral equation. To begin with, we model neurons with only excitatory synaptic input.

In the case where the unitary postsynaptic events are fast enough to be considered instantaneous (Nykamp and Tranchina, 2000), the PDF is one-dimensional, as the state of a neuron is completely determined by its transmembrane voltage. When synaptic kinetics are not assumed to be fast on the time-scale of the resting membrane time constant, and when the unitary postsynaptic conductance event has a single exponential time course, the state-space and the PDF are 2-dimensional; the state of the neuron is determined by its random membrane voltage and random excitatory postsynaptic conductance.

The population firing rate is given by the integral of the flux of probability per unit conductance across the threshold voltage over all possible excitatory postsynaptic conductance values.

We formulate a pair of coupled partial differential-integral equations, one for the neurons in their non-refractory state and the second one for the neurons in the refractory pool. The higher dimensionality causes an increase in computation time, which we tackle numerically by using an operator-splitting method to solve our partial-differential integral equations. We compare our two-dimensional results to Monte-Carlo simulations for simple neural networks and check for their speed and accuracy in such instances. We then extend the population density method by adding inhibitory synaptic input. We consider a method for reduction from three to two dimensions and we apply it to small sample neural networks, as well as to a model orientation selectivity network for 1 hypercolumn of the visual cortex.

Limitations of the method are discussed and possible improvements and directions for future study are suggested.

Paul Atzberger

Rensselaer Polytechnic Institute, Mathematical Sciences, 8th Street, Troy, NY 12180

A Stochastic Immersed Boundary Method for Biological Fluid Dynamics at Small-Length Scales

In modeling many biological systems it is important to take into account the interaction of a fluid with deformable structures. At the length scale of individual cells and organelles thermal fluctuations of the aqueous environment become significant. To model and simulate the mechanics of such systems consistently taking into account these features presents a variety of theoretical and numerical challenges. For macroscopic systems the immersed boundary method has found wide use as an efficient numerical method for simulation of deformable structures in a fluid. In this work we show how the framework can be extended to small length scales to include thermal fluctuations. We then develop an efficient stochastic numerical method. Applications of the method, including simulation of diffusing particles, polymer knots, and self-avoiding membranes, are also presented.

Karim Azer

Merck & Co., Inc., Applied Computer Science & Mathematics, RY84-202, 126 E. Lincoln Avenue, Rahway, NJ 07065 and Courant Institute of Mathematical Sciences

A One Dimensional Model with Friction of Blood Flow in Arteries: Analysis, Numerical Solution, Simulation and Validation

In this poster, we present a one-dimensional model for computing blood flow in arteries, without assuming an a priori shape for the velocity profile across an artery. Combining the equations for conservation of mass and momentum with the Womersley model in an iterative way gives us a second order method. We are able to use this method to get the velocity profile at any particular location along an artery, which is very valuable in many applications. Moreover, we are able to get a more accurate representation of the flow and pressure waves in the arterial system since the friction term is based on the more realistic velocity profile that we compute. We also present flow simulations using both structured trees and pure resistance models for the small arteries, and the resulting flow and pressure waves under various friction models.

Joint work with Charles Peskin (Courant Institute of Mathematical Sciences).

Dipankar Bandyopadhyay

University of Georgia, Department of Statistics, 266 Statistics Bldg, Athens, GA 30602

A Novel Approach to Testing Equality of Survival Distributions when the Group Memberships are Censored

This paper introduces a novel nonparametric approach for testing the equality of two or more survival distributions when the group membership information is not available for the right censored individuals. Although such data structures arise in practice very often, this problem has received less than satisfactory treatment in the nonparametric testing literature. Currently there is no nonparametric test for this hypothesis in its full generality in the presence of right censored data. We propose a novel approach leading to a division of the samples that enables one to use any class of tests for comparing survival curves based on independent samples. We study a class of weighted log-rank tests obtained this way through extensive simulation. An asymptotic linear representation of our test statistic is obtained and two resampling alternatives were proposed which might be easier to use in practice. The testing methodology is illustrated using two real data sets. This is joint work with Prof. Somnath Datta, Dept of Statistics, University of Georgia.

Keywords: log-rank tests; competing risk model; independence; equality of survival curves.

Anisha Banerjee

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Modeling Steady-State and Time-Dependent Oxygen Delivery in Working Muscle

Advisor: Dr. Daniel Goldman

Oxygen delivery and distribution in muscle is studied using Krogh-type oxygen transport models. A series of modifications to the classic steady-state Krogh tissue cylinder model is used to study oxygen transport from individual capillaries in exercising skeletal muscle. Physiologically important features studied include intravascular resistance, myoglobin facilitation of oxygen diffusion, mitochondrial clustering near capillaries and Michaelis-Menten oxygen consumption kinetics. When consumption does not depend on oxygen concentration, intravascular resistance to oxygen diffusion is shown to significantly lower the partial pressure of oxygen in the tissue, suggesting a decrease in consumption rate. Mitochondrial clustering is shown to decrease the partial pressure of oxygen in most of the tissues, giving shallow tissue oxygen gradients as has been observed experimentally. Myoglobin facilitation is found to play only a minor role in steady state oxygen transport. By including oxygen-dependent consumption, the model makes it possible to study hypoxia as seen in working muscle. This also permits more accurate calculation of tissue oxygen distributions and the total oxygen consumption rate (oxygen extraction). The steady-state model is generalized to allow study of time dependent oxygen transport. In particular, time-dependent tissue oxygen distributions and oxygen consumption are calculated for coordinated changes in muscle activity and blood flow.

Sibabrata Banerjee

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

A Comparison Study of Models for the Human Sex Ratio

The human sex ratio is always very close to unity. If it moves too far away from unity, we face the risk of extinction and thus it is a matter of great interest in population statistics. Overdispersion models are often used to fit the human sex ratio. Branching processes are also very powerful tools for the modeling human sex ratio. This study would involve fitting several models using classical and modern techniques and a comparison would be made between these methods.

Keywords: overdispersion, non-linear regression, quasi-likelihood (multiplicative) generalization of binomial model, double exponential family.

Lyudmyla L. Barannyk

University of Michigan, Department of Mathematics, 4855 East Hall, Ann Arbor, MI 48109-1109

Numerical Simulations of Kelvin-Helmholtz Instability in Slightly Stratified Fluid in the Channel

A system of density-stratified incompressible inviscid fluids in an infinite horizontal channel is considered. The approach is based on a boundary integral formulation in which the fluid interface is modeled as a free vortex sheet and the channel walls as bound vortex sheets. The density variations are assumed to be small (Boussinesq regime) and the motion of the interface between fluids is studied numerically using the vortex blob method. The goal is to simulate the flow in the inclined channel and compare the numerical results with the experimental results obtained by Thorpe [J. Fluid Mech. 46 (1971) 299-319]. Joint work with Robert Krasny.

Michael D. Bateman

University of Kansas, Department of Mathematics, Lawrence, KS 66044

Traveling Wave Solutions to a Coupled System of Spatially Discrete Nagumo Equations

We consider a coupled system of discrete Nagumo equations and derive traveling wave solutions to this system using McKean's caricature of the cubic. A certain form of this system is used to model ephaptic coupling between pairs of nerve axons; we present the derivation of this particular system, but consider a slightly more general version. We study the difference $g(c) = a_1 - a_2$ between the detuning parameters a_1 that is required to make both waves move at the same speed c . Of particular interest is the effect of a coupling parameter α and an "alignment" parameter A on the function g . The introduction of a time delay β is used to show that the effects of perturbation away from $A = 0$ are minimal. Numerical investigation indicates that for fixed A , there exists a β value that results in $g = 0$; and for large enough wave speeds, multiple such β values exist. Also, numerical results indicate that the perturbation of α away from zero will yield additional solutions with positive wave speed when $A = \frac{1}{2}$. We rely on both numerical and analytical results to demonstrate our claims. Joint work with Erik S. Van Vleck (University of Kansas).

John K. Bechtold

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Nonlinear Oscillations in Diffusion Flames

Recent experiments of diffusion flames have identified fascinating nonlinear behavior, especially near the extinction limit. In this work, we carry out a systematic analysis of pulsating instabilities in diffusion flames. We consider the simple geometric configuration of a planar diffusion flame situated in a channel at the interface between a fuel being supplied from below, and oxidant diffusing in from a stream above. Lewis numbers are assumed greater than unity in order to focus on pulsating instabilities. We employ the asymptotic theory of Cheatham and Matalon, and carry out a bifurcation analysis to derive a nonlinear evolution for the amplitude of the perturbation. Our analysis predicts three possible flame responses. The planar flame may be stable, such that perturbations decay to zero. Second, the amplitude of a perturbation can eventually become unbounded in finite time, indicating flame quenching. And finally, our amplitude equation possesses time-periodic (limit cycle) solutions, although we find that this regime cannot be realized for parameter values typical of combustion systems. The various possible burning regimes are mapped out in parameter space, and our results are consistent with available experimental and numerical data. Joint work with H. Y. Wang and C. K. Law (Princeton University).

Manish C. Bhattacharjee

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Strong Versions of the DFR Property and Applications

We explore two nonparametric classes of life distributions nested within the class of decreasing failure rate (DFR) distributions, motivated by the observation that the standard parametric families of such distributions possess properties which are stronger than the decreasing failure rate property. Our results characterize such distributions via several representation theorems. Associated consequences such as closure properties and reliability bounds are considered, along with examples of how such strongly DFR distributions arise naturally in many applied contexts.

Denis Blackmore

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Periodic and Quasiperiodic Motion of Point Vortices

We study the dynamics of any finite number of point vortices moving in an ideal fluid in both the plane and half-plane. If there are n point vortices, the governing dynamical equations are Hamiltonian and have n -degrees of freedom. These equations are completely integrable for n small, but not integrable for larger numbers of point vortices. Among the results obtained is the following: If the vortices all have strengths of the same sign, there is a set of initial configurations of positive measure for which the Hamiltonian equations of motion have invariant KAM tori containing quasiperiodic orbits, as well as periodic solutions of arbitrarily large periods. This result is proved using a modified version of the KAM theorem, and a new version of the Poincare-Birkhoff fixed point theorem that we developed to study small perturbation of completely integrable Hamiltonian systems. Joint work with Jyoti Champanerkar (William Paterson University).

Mark G. Blyth

University of East Anglia, Norwich, Norfolk, England, NR4 7TJ, UK

Stability of Two-Layer Channel Flow in the Presence of Surfactants

The effect of an insoluble surfactant on the stability of two-layer viscous flow in an inclined channel confined by two parallel walls is considered both under conditions of Stokes flow and at arbitrary Reynolds numbers. At zero Reynolds number, a long-wave model is developed leading to a nonlinear evolution equation for the interfacial position. Numerical boundary-integral simulations for waves of arbitrary length agree with linear stability predictions at small amplitude, and predict that waves may grow and saturate into steep profiles or begin to overturn and break. The results are then extended to include inertia. A generalized form of the Orr-Sommerfeld equation is derived describing the growth of small disturbances at arbitrary Reynolds number and solved numerically. For general Reynolds numbers and arbitrary wavenumbers, the surfactant is found to either provoke instability or significantly lower the rate of decay of infinitesimal

perturbations, while inertial effects widen the range of unstable wavenumbers. The nonlinear evolution is computed numerically using a finite-difference method. The results are consistent with the linear theory and show that the interfacial waves steepen and eventually overturn under the influence of the shear flow.

Michael R. Booty

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Scattering from a Thin Inhomogeneous Cylinder in a Waveguide

A thin cylindrical target is placed with almost arbitrary orientation in a rectangular TE₁₀ waveguide and is subjected to an imposed electromagnetic field. The scattered far-field is expressed in terms of the scattered field inside the target and the far-field expansion of the dyadic Green's function for the waveguide. To capture features of interest in microwave heating applications, we allow the target material's electrical properties to be arbitrary functions of position along the target's axis. Reflection and transmission coefficients are derived together with an expression for the rate of deposition of electromagnetic energy within the target. Joint work with Gregory A. Kriegsmann (NJIT).

Amitabha Bose

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Localized Activity Patterns in Excitatory Neuronal Networks

The existence of localized activity patterns, or bumps, has been investigated in a variety of spatially distributed neuronal network models that contain both excitatory and inhibitory coupling between cells. Here we show that a neuronal network with purely excitatory synaptic coupling can exhibit localized activity. Bump formation ensues from an initial transient synchrony of a localized group of cells, followed by the emergence of desynchronized activity within the group. Transient synchrony is shown to promote recruitment of cells into the bump, while desynchrony is shown to be good for curtailing recruitment and sustaining oscillations of those cells already within the bump. These arguments are based on the geometric structure of the phase space in which solutions of the model equations evolve. We explain why bump formation and bump size are very sensitive to initial conditions and changes in parameters in this type of purely excitatory network, and we examine how short-term synaptic depression influences the characteristics of bump formation. Joint work with Jonathan Rubin.

Bruce Bukiet

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Mathematical Modeling of Postural Stability

Understanding postural stability (human balance) is important because millions of Americans experience dizziness and balance problems in their lifetimes. The quality of many people's lives could be improved through development of high quality diagnostic methods that could reliably determine who is in danger of falling. With this in mind, we have developed mathematical models of postural stability, including 2 and 4 link models. We have also created an index that can be used as a diagnostic tool to quantify postural stability. We discuss our model, the properties of our Postural Stability Index and applications of our work in the diagnosis and evaluation of treatment options. Joint work with Hans Chaudhry, Zhiming Ji, Thomas Findley, Karen Quigley, and Miriam Maney.

Jean Cadet

SUNY at Stony Brook, Department of Applied Mathematics and Statistics, Stony Brook, NY 11794-3600

Robustness in Drosophila Melanogaster

Driever and Nusslein-Volhard showed that the bicoid gene protein determines position in the Drosophila embryo in a concentration-dependent manner. We combine our knowledge of this result with the gene circuit model which is based on 4 steps: formulate a theoretical model, acquire data, optimize using simulated annealing and learn new biology. Finally, we propose a manner to characterize statistically biological robustness despite changes in the bicoid dosage.

Jerry J. Chen

Case Western Reserve University, Department of Mathematics, 10900 Euclid Avenue, Cleveland, OH 44106

Bifurcation and Chaos in Discrete Lotka-Volterra Equations

A discrete Lotka-Volterra (L-V) population model for a pair of competing species is analyzed. It is known that the one-dimensional logistic map exhibits a variety of dynamical features such as period-doubling bifurcation and chaotic cobweb. It is shown that the L-V map demonstrates its own brands of period-doubling bifurcation and a two-dimensional twisted horseshoe. With initial values outside the Cantor-like set, the weaker species ultimately becomes extinct, which agrees with the continuous version of the Lotka-Volterra equations. A controlling scheme that diminishes the intrinsic growth rate of the dominant species is provided to prevent the probable extinction of the weaker species. Some numerical results are given to portray its effectiveness.

Yiming Cheng

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Analysis of Equivalent Distorted Ratchet Potentials

Separation technology is important in biotechnology for certain processes such as separation of microscopic particles undergoing Brownian motion. These particles can be separated by subjecting them periodically to an asymmetrical spatially periodic electric field, sometimes referred to as a Brownian ratchet. It has been found numerically that two different potentials under special equivalent distortions (Chen et al., 1999; Yan et al., 2001) can produce equal fluxes of particles. Although a simple physical explanation for this can be given, there is no analytical proof of this equivalency. Here, we pose the mathematical problem and discuss the difficulty in carrying out the analytical proof.

Shawn Chester

NJ Institute of Technology, Granular Science Laboratory, Mechanical Engineering Department, University Heights, Newark, NJ 07102

Discrete Element Simulations of Floor Pressure in a Granular Material in a Cylindrical Vessel

Advisor: Anthony Rosato

At the end of the 19th century, H. A. Janssen discovered that the bottom floor pressure in a cylindrical container of granular material asymptotes exponentially to a value less than the weight of the material, i.e., the pressure becomes independent of the fill height of the column. Applications are prevalent in the handling, processing and transport of industrial bulk materials. Janssen's theoretical prediction is based on a constitutive assumption that the vertical (axial) stress in the material is proportional to the horizontal (radial) stress. We model this problem using particle level (discrete element) simulations in a cylindrical vessel in which the floor is slowly moved down. The objective of our investigation is to reproduce the theoretical prediction, and to make quantitative comparisons with experiments in the literature. A key parameter in the analysis is the selection of an appropriate friction coefficient between the particles and the cylinder walls, a topic which has seen much discussion in the literature. Results obtained thus far have shown a floor pressure that is smaller than the static value (due to the material weight), in qualitative agreement with the theory. However, the rate at which the graph of pressure versus fill height asymptotes to its constant value does not agree with the theory. We are now exploring a wide range of friction coefficients to understand how this parameter affects our simulated results. Work is also underway to test the major assumption in the model concerning the proportionality between the vertical and horizontal stresses.

Wooyoung Choi

University of Michigan, Dept. of Naval Arch. & Marine Eng., 2600 Draper Rd., Ann Arbor, MI 48109

Nonlinear Surface Waves in a Linear Shear Current

Nonlinear surface gravity waves interacting with a linear shear current are studied for both infinitely deep and shallow water. For deep water, a closed system of the exact evolution equations is obtained using the conformal mapping technique and is solved numerically via a pseudo-spectral method to study the evolution of slowly modulated periodic surface waves. For shallow water, we consider an asymptotic model for long waves and discuss its solitary wave solutions including a solution with infinite slope.

Wooyoung Choi

University of Michigan, Dept. of Naval Arch. & Marine Eng., 2600 Draper Rd., Ann Arbor, MI 48109

Propagation of Large Amplitude Internal Waves Over Variable Bottom Topography

The evolution of large amplitude internal waves is investigated using a strongly nonlinear long wave model for a system of two layers of different densities. A local stability analysis is presented to show that the solitary wave solution of this 'inviscid' model suffers from a Kelvin-Helmholtz type instability due to a velocity discontinuity across the interface between two layers. A numerical filter is used to eliminate the short-wave instability (that is absent in real observations) and its effects on long-term numerical simulations are discussed. Joint work with Taechang Jo (New Mexico Institute of Mining and Technology).

Sohae Chung

SUNY at Stony Brook, Department of Applied Mathematics and Statistics, Stony Brook, NY 11794

Mathematical Erosion as a Measure for Osteoporosis: Quantifying Topological Change

The evaluation of topological changes (hole formation, breakage) in the trabecular bone region provides insight into how trabecular bone protects itself against the effects of osteoporotic erosion and may provide essentials for fracture risk assessment at an early time point that are not reflected by density measurements (eg, bone mineral density). Based on 3D digital images obtained using micro-CT, a mathematical erosion model has been developed that enables the analysis of the relationship between virtual bone loss generated by mathematical erosion and the bone's structural alteration during progressive bone loss. The results demonstrate that erosion induced stresses are relieved through the preferential production of holes over breaks in the trabeculae. Trabecular bone appears constructed so that such preferential relief is

able to continue down to extremely eroded forms of the structure. Further, the topological analysis indicates that trabecular structure is better protected against isotropic rather than unidirectional erosive action. Joint work with W. Brent Lindquist and Yi-Xian Qin (Stony Brook University), and John Pinezich (Advanced Acoustic Concepts, Inc., Hauppauge, NY 11788).

Robert Clewley

Cornell University, Department of Mathematics and Center for Applied Mathematics, Ithaca, NY 14853

Dynamical Models of Finger Biomechanics and Neuromuscular Control

We analyze the periodic motor patterns of index finger control of a trackball, during a psychophysics task in which the subject has to match a constant angular velocity. We explore minimal hybrid (DAE) kinematic models of the system, and use parameter estimation to investigate what control strategies provide maximally-robust achievement of the target velocity. This work is also intended to develop the computational tools necessary to study a related question, in insect locomotion: What are the relative roles of 'pre-flex' and CNS feedback control strategies in cockroach locomotion? The cockroaches are mounted on a Kugel (a freely rotating styrofoam ball) at Robert Full's Berkeley laboratory, and so this resembles the finger manipulation problem under certain assumptions. Joint work with John Guckenheimer, Francisco Valero-Cuevas, Dan Brown, and Erik Sherwood.

Dean T. Crommelin

Courant Institute of Mathematical Sciences, New York University, 251 Mercer St., New York, NY 10012

Extracting Low Order Stochastic Models from Data

We present a numerical technique to derive optimal stochastic models (Markov chain or SDEs) for the description of the evolution of a few interesting collective variables in large-sized dynamical systems. The technique is based on constructing the stochastic model whose eigenfunctions and eigenvalues are the closest (in some appropriate norm) to the one gathered from the observations. The technique is validated on an example arising from climate dynamics, namely by extracting a stochastic model for the evolution of the first few principal orthogonal modes observed in a general circulation model. Joint work with E. Vanden-Eijnden.

R.E. Lee Deville

Courant Institute, New York University, 251 Mercer St., New York, NY 10012

Self-Induced Stochastic Resonance for Excitable Systems

Excitable systems arise in a wide variety of areas, including climate dynamics and lasers, and are especially common in biology. One can think of them schematically as dynamical systems possessing a rest state, an excited state, and a recovery state. Recently, there has been much interest in coherent response of these systems to small amplitude stochastic forcing. In particular, the well-known mechanism of coherence resonance has been successful in explaining some of these phenomena. Here we discuss an alternate mechanism to coherence resonance which takes place in different parameter regimes and gives rise to qualitatively different behavior. Moreover, in a suitable limit, we recover deterministic behavior which is far from the unforced dynamics. We will treat the cases of stochastically forcing excitable ODE and excitable media, i.e., the spatially extended PDE. This is joint work with E. Vanden-Eijnden (Courant Institute) and C. Muratov (NJIT).

Sunil K. Dhar

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

On the Characterization of a Bivariate Geometric Distribution

In this paper it is shown that a bivariate random variable has constant failure rate and mixture geometric marginals if and only if it has the loss of memory property and the discrete Freund distribution. This characterization is achieved by extending a key lemma in this area. The mixture geometric can be collapsed to geometric marginals, thus validating the characterization as a generalization of the past work. Joint work with Srinivasan Balaji (George Washington University).

Sunil K. Dhar

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Improved Methods for Establishing Noninferiority in Clinical Trials

We consider a randomized, active-control clinical trial setting in which the objective is to test for noninferiority of the experimental treatment compared to control. An approach for defining a noninferiority margin is based on the concept of preserving a certain fraction of the active control effect. Noninferiority is established if the ratio of the lower limit of the two-sided 95% confidence interval for the treatment difference to the estimated mean of the active control is greater than a prespecified fraction. The nominal significance level, depending on values of location and scale parameters, may not be maintained by this confidence-interval approach to testing noninferiority. We examine bootstrap methods to derive a more accurate lower limit of the confidence interval for known functions of the treatment means. This, in turn, improves the power

of the test which will be demonstrated through simulation. Other methods of establishing noninferiority or equivalence will also be explored. Joint work with Michael Chen and Farid Kianifard (Biometrics, U.S. Clinical Development and Medical Affairs, Novartis Pharmaceuticals, East Hanover, NJ 07936).

Ramana Dodla

New York University, Center for Neural Science, 4 Washington Place, New York, NY 10003

Dynamics of Weakly Coupled Kuramoto Oscillators

Phase coupled Kuramoto oscillators have become classic examples of phaselocking among non-identical oscillators. While there are many studies on the limiting case of the number of oscillators becoming infinity, the dynamics of finitely large number of oscillators is still a largely unexplored and fascinating area of research. In this poster we present some preliminary results on the finitely large number of globally coupled phase oscillators when the coupling strength is weak. Under special frequency distributions, the oscillators could show signatures of a completely integrable system. For example, a perturbation decays in time, but can be recovered with a periodicity that is characteristic of the number of coupled oscillators.

Brent Doiron

New York University, Center for Neural Science, 4 Washington Place, New York, NY 10003

Coding with Inter-Spike Intervals and Dendrites

Many neurons use precise spike timing to reliably encode dynamic stimuli. We present experimental and computational results from sensory pyramidal neurons showing that the time interval between two precise spikes can also encode relevant sensory information. Electrosensory pyramidal neurons produce high frequency two spike bursts in response to upstrokes of dynamic stimuli. Using pattern classifier techniques we show that the inter-spike interval duration of a burst can accurately encode the scale of stimuli upstrokes. Electrosensory neurons have well characterized nonlinear dendritic processes that support backpropagation of action potentials along the dendritic arbor. We introduce a simple modified integrate-and-fire neuron that accounts for how nonlinear dendritic effects shape spike train statistics in response to dynamic stimuli. Using this model we study how active dendrites are critical for this inter-spike interval coding scheme. Specifically, the amplitude of the dendritic spike sets the range over stimuli space that the inter-spike intervals can accurately span. Interestingly, the amplitude of the dendritic spike that maximizes the information in the model's interval code is the value that best matches experimental observations of dendritic spikes in real electrosensory neurons.

Graham Donovan

Northwestern University, Engineering Science and Applied Mathematics, Evanston, IL 60208

Non-Gaussian Optical Field Statistics in a Long-Haul Soliton-Based DPSK Transmission System

We demonstrate with importance-sampled Monte-Carlo simulations that the tails of the optical field's probability distribution at the end of a long-haul soliton-based differential phase-shift keyed (DPSK) transmission system are strongly non-Gaussian.

Jonathan D. Drover

University of Pittsburgh, Mathematics Department, 504 Cabot Way, Pittsburgh, PA 15203

Nonlinear Coupling Near a Degenerate Hopf Bifurcation

A nonlinearly coupled network of oscillators near a Hopf bifurcation is analyzed. The equations are derived from the normal form of a subcritical Hopf bifurcation where the unstable branch of periodics 'turns around', creating a parameter dependent region of bistability. Solutions such as waves and localized regions of excitation, which may be analogous to patterns of activity used to model working memory, are found to exist numerically. A mechanism where waves do not exist because of large phase gradients is explored.

Srabasti Dutta

SUNY at Stony Brook, Applied Mathematics and Statistics, Stony Brook, NY 11794-3600

LES Simulations of Turbulent Combustion in a Type Ia Supernovae

We propose a 2D axisymmetric model of a type Ia supernova explosion, based on a front tracking sharp flame model. The calculation is free from adjustable turbulent transport parameters, and in this sense it is in the spirit of Large Eddy Simulation (LES) turbulence simulations. Since the mixing is dominated by the largest eddies, we resolve these and not the smaller ones. We believe this method results in a tolerable error, which, in any case understates the success of the explosion. We report successful explosions. Both the 2D and the LES nature of the model greatly simplify parameter identification. The 2D model allows multiple simulations and an exploration of unknown parameters, while the LES model removes parameters from the simulation. Joint work with James Glimm and Yongmin Zhang (SUNY at Stony Brook).

Alan Elcrat

Wichita State University, Mathematics Department, 1840 N. Fairmount, Wichita, KS

Free Surface Waves in Equilibrium with a Vortex

Finite amplitude solitary waves of uniform depth which interact with a stationary point vortex are considered. Waves both with and without a submerged triangular obstacle are computed. The method of solution is collocation of Bernoulli's equation at a finite number of points on the free surface coupled with equations for the equilibrium of the point vortex. The stream function and vortex location are found by computing a conformal map from the flow domain to an infinite strip using Driscoll's SCTOOLBOX. The solutions are parameterized with respect to Froude number, vortex circulation, and, when an obstacle is present, obstacle height. When no obstacle is present there are two families of solutions. In one the amplitude increases with increasing circulation, while in the other it decreases. Beyond a certain critical Froude number the maximum amplitude wave has a sharp crest with an angle of 120 degrees. Similar behavior is observed for flow past an obstacle except that below a certain Froude number there is no solution at all.

Christopher E. Elmer

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Functional Differential Equations of Mixed Type

We illustrate techniques for deriving and for computing solutions to functional differential equations of mixed type. Also included are examples of applicable models.

Samiran Ghosh

University of Connecticut, Department of Statistics, 215 Glenbrook Rd. U-4120, Storrs, CT 06269

Statistical Approach to Metabonomic Analysis of Rat Urine Following Surgical Trauma

Motivation: Acute trauma is often associated with the progressive deterioration of multiple organ systems in humans and is the leading cause of death in trauma care units. Previous studies have suggested that multiple organ failure is likely related to uncontrolled systemic inflammation; however, causal mechanisms remain unknown. Current methods of assessing trauma patient status and predicting outcome are based on a variety of anatomical and/or physiological scoring models. While useful, these are labor intensive and do not allow for real-time analysis of patient status. In this study, we develop a metabonomic based approach using a rodent model of acute trauma in order to determine whether statistically significant differences exist between the quantitative and qualitative profile of urinary metabolites of control rats and rats that have experienced surgical trauma. This approach incorporates statistical, analytical and computational tools in order to identify metabolites that are unique to trauma and may be used to predict trauma outcome.

Results: We used electrospray ionization/time-of-flight (ES/TOF) mass spectrometry to analyze the urine metabonomes of eleven male rats 48 hours following surgical implantation of a biotelemetry transmitter and vessel cannulation as a model of surgical trauma. Statistical analysis showed significant differences between trauma and control urinary metabonomes. These included metabolites that were detected ES/TOF both positive and negative ionization modes. Principle component analysis was used to categorize metabonomes into either control or trauma with greater than 80% accuracy. We also show, using Bayesian methods that we could classify subjects as being traumatic or control with a given credible interval. These results suggest that metabonomics may prove useful for quantifying and identifying biomarkers of trauma status as well as trauma outcome in humans. Joint work with Dennis Hill, David F. Grant, and Dipak K. Dey (University of Connecticut).

Daniel Goldman

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Calculations of Tissue Oxygen Delivery and Utilization During the Time Course of Microvascular Injury in a 6-Hour Acute Sepsis Model

Sepsis produces severe disturbances to the microcirculation, including loss of functional capillary density (fCD) and increased blood flow heterogeneity. We have previously presented a mathematical model of microvascular oxygen delivery in rat skeletal muscle during sepsis (Goldman et al., AJP Heart, 2004), based on measurements of fCD, hemodynamics and blood oxygen saturation in a 24-hour cecal ligation and perforation (CLP) model. Here we utilize recent measurements of these same transport parameters, as well as capillary geometry, in a single microvascular network at two, three and four hours after surgery in an acute 6-hour CLP model. This data was obtained to study changes in microvascular blood flow during the time course of sepsis, but also allows us to study progressive changes in oxygen delivery and utilization. We will present results showing how changes in functional microvascular geometry and blood flow result in changes

in oxygen transport, as the effects of sepsis on the microcirculation become more severe over time. Joint work with Graham Fraser and Christopher Ellis (University of Western Ontario, London, Ontario, Canada).

Jorge Golowasch

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Gap-Junction Conductance Determines an Optimal Coupling Diameter in Passive Fibers

Neurons communicate via chemical synapses but also via direct electrical connections called gap junctions. These are molecular channels that directly link cells for electrical and biochemical signal transfer. Gap junctions, although discovered almost 50 years ago, are only now being studied in more detail at the molecular level. At the computational level gap junctions are thought to have a synchronizing role of the electric behavior of coupled cells. However, even some very basic questions have not been addressed. We are interested in understanding the behaviors that may arise from the combination of current flow along cables and their coupling gap junctions. Current flow through gap junctions depends on their conductance as well as the electrical properties of the coupled cells. We show that an optimal current transfer between two gap-junctionally coupled cables occurs as a function of cable diameter as well as a function of the gap junctional conductance. This non-monotonic behavior does not depend on any active properties. The optimal current transfer is a local phenomenon that depends on the diameters of only the coupled processes. We predict that in biological networks coupled processes may have optimal diameters that maximize current transfer between neurons. Supported by NIH grants MH-60605 (F.N.) and MH-64711 (J.G.). Joint work with Farzan Nadim (NJIT).

Roy Goodman

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Heads Up! A Capstone Applied Math Course in the Mechanics of Coin Flipping

In a recent preprint, Diaconis, Holmes, and Montgomery have shown that under general conditions, a coin tossed in the air that starts heads up spends more than half of its time in the air heads up, and thus is inherently biased to land that way. This effect does not diminish as the coin is thrown higher or with more vigorous rotation. This result formed the basis of my capstone course for graduating seniors in applied mathematics at NJIT. This bias arises because the coin precesses as it tumbles, and is a straightforward, though novel, application of results due to Euler in about 1750! We reproduce experiments in which flipping coins are filmed using a high speed video camera. We use Matlab's image processing toolbox to analyze the motion and quantify the bias. We also analyze at "Feynman's plate" and the motion of gyroscope under the influence of gravity using the same techniques. The experiments are incredibly simple, yet analyzing them in the framework of Euler's equation requires a great deal of knowledge and computational work. Joint work with the students of Math 451H-04.

Arnaud Goulet

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Control of Mixing

In this work, we address the control of Lagrangian chaos or chaotic advection in a two-dimensional incompressible fluid flow. Due to the fact that the motion of passive tracers (particles) can be described by a Hamiltonian function, we apply a control technique recently developed for Hamiltonian systems. The control is performed by introducing a suitable small term in the original Hamiltonian which is derived explicitly. The control term is in the form of a secondary flow which slightly deforms the streamlines of the original flow and reduces the chaoticity of the particle trajectories.

The technique is implemented on two different fluid flow systems consisting of Rossby waves and Rayleigh-Bénard convection which can both exhibit chaotic advection for certain parameter values. For each system, we derive the control term and demonstrate its effect on chaotic advection through numerical simulations of the time dependent material lines as well as Poincaré sections. Joint work with Nadine Aubry (NJIT).

Muhammad Hameed

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Influence of Surfactant on Necking and Pinch-Off in Two Fluid Jets

The effect of surfactant on the breakup of a periodic fluid jet immersed in another viscous fluid at low Reynolds number is studied both numerically and experimentally. Evolution equations for the jet interface and surfactant concentration are derived using long wavelength approximations. These one dimensional partial differential equations are solved numerically for given initial interface and surfactant concentration. It is found that the presence of surfactant at the interface retards pinch-off and results in the formation of a thin and long filament. To check the predictions of our model, we performed experiments both for clean interface and as well as in presence of surfactants. The experimental results support the prediction of theoretical model that the presence of surfactant slows down the pinch-off process.

Alex Haro

Universitat de Barcelona, Dept. de Matemàtica Aplicada I Anàlisi, Gran Via 585, 08007 Barcelona, Spain

Invariant Manifolds in Quasiperiodic Systems: Theory, Computation and Applications

The parameterization method is useful to prove the existence of invariant manifolds in quasi-periodic systems, and it also provides effective algorithms to compute these manifolds. We apply the method to two paradigmatic examples: the quasiperiodic Hénon map and the quasiperiodic standard map. Joint work with Rafael de la Llave.

Cristel Hohenegger

Georgia Institute of Technology, School of Mathematics, 686 Cherry Street, Atlanta, GA 30332

Statistical Reconstruction of Velocity Profiles for Nano-PIV

Velocities and Brownian effects at nano-scales near microchannel walls have been measured by evanescent-wave illumination techniques [R. Sadr et al., J. Fluid Mech. 506, 357-367 (2004)]. Assuming mobility of spherical particles is dominated by hydrodynamic interaction between the particle and wall, and that fluid velocity is directed in one in-plane direction, the out-of-plane dependence of mobility and velocity are clearly coupled. We investigate such systems computationally, using a Milstein algorithm that is both weak- and strong-order 1. We demonstrate that a maximum likelihood algorithm can reconstruct the out-of-plane velocity profile given known mobility dependence and ideal particle identification. We further test this reconstruction for measurements obtained by cross-correlation techniques applied to windowed simulation data. Application to physical data is proposed via analytical results about the influence of Brownian motion in this setting on the correlation peak, combined with simulation results to help identify nearly-optimal parameters. Joint work with P.J. Mucha (School of Mathematics, Georgia Institute of Technology).

David Horntrop

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Mesoscopic Simulation of Domain Coarsening in Surface Processes

Self-organization of components of two phase mixtures through a diffusive mechanism is known as Ostwald ripening and is an example of a multiscale phenomenon which is well-suited to study using mesoscopic models. Mesoscopic models are stochastic partial differential equations which are directly derived from the underlying microscale behavior. In order to computationally study these models, new spectral schemes for stochastic partial differential equations are introduced and validated using exactly solvable benchmark problems. These schemes are then applied to the mesoscopic model and simulation results are compared with various theoretical results such as the Lifshitz-Slyozov growth law.

Maureen Howley

NJ Institute of Technology, Chemical Engineering Department, University Heights, Newark, NJ 07102

A Comparison of One-Dimensional Traveling Waves in Inverse and Normal Fluidized Beds

The state of uniform fluidization is usually unstable to small disturbances, and this can lead to the formation of vertically traveling voidage waves. In inverse fluidization, when particle density is less than fluid density ($\rho_p < \rho_f$), particles fluidize in the direction of gravity when the drag force exerted by the fluid overcomes buoyancy. Inverse fluidization thus provides a unique parameter space, which augments the study of instability behavior in normal fluidization when $\rho_f < \rho_p$. Using continuum equations of continuity and motion, we compared the linear stability of normal and inverse bed modes to examine the effect of the Froude number Fr and fluid to solid density ratio ($\delta = \rho_f / \rho_p$). Making use of numerical bifurcation analysis and continuation, periodic solutions in the form of one-dimensional traveling waves (1D-TWs) were computed. Based on wave growth rates and bifurcation structure, we identified the Fr as an important parameter for predicting instability strength. However, δ affects instability onset, or the point at which the base state is rendered unstable. In the case studies we examined, traveling waves were shown to propagate in the direction of fluidization, and asymmetrical, high amplitude 1D-TW profiles suggest fully developed bubble-like structures are oriented in the direction of fluidization.

Aridaman K. Jain

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Small-Sample Non-Parametric Tests for the Effectiveness of Liposuction Surgery for Breast Hypertrophy

Previous studies on the effectiveness of treatments for breast hypertrophy have been based on the use of normal approximations for the averages of the responses from a large number of patients treated at several plastic surgeon practices. In those studies, patients were asked to fill out questionnaires containing many questions on measures of health/pain, on a 3-point or a 5-point scale, both before and after the surgery. We describe the results from a prospective research study of breast-reduction, which is unique in two aspects: (i) it is the first such study focused exclusively on African-American women, (ii) it investigates the liposuction breast reduction technique. Due to the newness of this surgical procedure, data are available only for a limited

number of patients. Since the use of the normal approximation for a small sample number of patients may be questionable, we decided to use non-parametric tests that do not make any assumptions about the shape of the probability distribution for the self-reported responses on 3-point or 5-point scales. After the surgery, the pain was reduced and the general health was improved, when compared with those reported before the surgery ($p \leq 0.05$). Joint work with Martin J. Moskovitz, Sherwood A. Baxt, and Robert E. Hausman.

Shidong Jiang

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

The Hyperpolarizabilities for One-Dimensional Infinite Single-Electron Periodic Systems

The analytical solutions for the general-four-wave-mixing hyperpolarizabilities on infinite chains under both Su-Shrieffer-Heeger and Takayama-Lin-Liu-Maki models of trans-polyacetylene are obtained through the scheme of dipole-dipole correlation. Analytical formats of DC Kerr effect, DC-induced second harmonic generation, optical Kerr effect and DC-electric-field-induced optical rectification are derived. By including or excluding the gradient terms in the calculations, comparisons show that the intraband contributions dominate the hyperpolarizabilities. The intraband transition leads to the break of the overall permutation symmetry in hyperpolarizabilities even for the low frequency and non-resonant region. Since the Kleinman symmetry is directly based on the overall permutation symmetry, our calculations provide a clear understanding about the break of the Kleinman symmetry that are widely observed in many experiments.

Ning Ju

Oklahoma State University, Department of Mathematics, 401 Mathematical Sciences, Stillwater, OK 74078

Vorticity Geometry and Regularity of 3D Incompressible Flow

New geometric constraints on vorticity are obtained which suppress possible development of finite-time singularity from the nonlinear vortex stretching mechanism. The continuity conditions of Constantin-Fefferman and Beirao da Veiga-Berselli for the direction of vorticity which yield the regularity of the solution to the 3D Navier-Stokes equations are further relaxed. The important role of the "critical plane" is recognized. This leads to two broad classes of geometric configurations which allow non-continuity of vorticity direction. This also leads to improvement of the results of Beirao da Veiga and Grujic and Ruzmaikina for 3D Navier-Stokes equations and a result of Constantin for 3D Euler equations.

Yeona Kang

SUNY at Stony Brook, Applied Mathematics and Statistics, Stony Brook, NY 11794

A Constrained Kinetic Approach to Fast Algorithms for the Protein Folding Dynamic

The Einstein Relation defines the ratio of diffusion constant to a field driven drift constant. A new Einstein Relation for charges constrained to move as an integral part of a shape changing protein structure was derived and applied to protein folding. The partitioning of free energy into structural and electrostatic (and/or any other easily mapped energy form) provides a constrained trajectory for environmentally induced protein shape change. Lagrange methods ensure that the simultaneous minimization of two or more energy forms is describable in terms of the free energies gradients or forces. Using an analysis based on gradients and measurable physical parameters such as: total free energy change, molecular mobility (and diffusion), and the steady-state key mechanisms of protein shape change were elucidated. Protein folding involves pivoting of large groups of atoms specific bonds. The functional atomic group of folding can be viewed as a wishbone comprised of a pivot bond and of two (or more) charged regions held distant from the pivot bond by inert lever arm regions. A computation using a Markov sequence of wishbone-based rotations was carried out. These considerations lead naturally to a description in which the cooperative motions of large numbers of protein atoms move simultaneously.

The goal of this work is to build upon known processes to provide and demonstrate a thermodynamic framework that simulates protein folding and elucidates the key mechanisms and structures related to protein folding. Furthermore another goal for the model is to wherever possible link input parameters to laboratory parameters using the mathematical description. In some cases this required the development of new relationships. For example, the well known Einstein relation links the diffusion constant to the drift mobility of a given species, thereby making it possible to link the potential energy gradient to the concentration gradient. Einstein relations are specifically developed for charged species constrained to move as part of a molecular chain that could itself store energy independently of the electric potential. Joint work with E. L. Jaen, J. H. Coleman, and C. M. Fortmann (SUNY at Stony Brook).

Said Kas-Danouche

Universidad de Oriente, Department of Mathematics, Cumana, Sucre 6101, Venezuela

A Mathematical Model for Core-Annular Flows with Surfactants and No Basic Flow

For oil companies, transport techniques are very important. One technique is transporting crude through water lubricated pipelinings. This is an example of core-annular flows. In our case, the annular fluid (water) lubricates the movement of the core fluid (crude). Instabilities occur at the interface of both fluids

giving rise to not wanted final products as, for example, emulsification. On the other hand, it is known that most of surfactants affect the interfacial dynamics since they reduce the interfacial tension and introduce the Marangoni force. We derive a mathematical model for core-annular flows without basic flow and with surfactants using asymptotic methods. We assume that the annulus is thin compared with the core radius. We obtain a coupled system of two nonlinear partial differential equations. One describes the evolution of the interface and the other the evolution of the concentration of surfactants at the interface. Joint work with Michael Siegel and Demetrios Papageorgiou (NJIT).

Aslan R. Kasimov

University of Illinois at Urbana-Champaign, Department of Theoretical and Applied Mechanics, 216 Talbot Lab, 104 S. Wright St., Urbana, IL 61801

Asymptotic Theory of Self-Sustained Detonations

A self-sustained detonation wave is a shock wave propagating in a reactive medium so that its motion is sustained only by the chemical energy released behind the shock. An important property that distinguishes self-sustained detonations from supported (also called overdriven) detonations is that in the former a sonic locus exists at the end of the reaction zone. We propose general sonic conditions for detonation waves with smoothly evolving reaction zone as compatibility conditions in the limiting forward characteristic surface, which defines a general sonic locus. Such a surface is an information boundary that isolates the lead shock from the influence of the downstream flow of burnt products. The sonic conditions are derived from the general system of reactive Euler equations. A numerical illustration of their properties in the simplest case of a pulsating one-dimensional detonation is given. The sonic conditions are essential ingredients of an asymptotic theory of a slowly evolving weakly curved detonation. We derive a reduced evolution equation for such detonations that relates the detonation-shock curvature, speed, and acceleration. Some properties of the evolution equation will be illustrated with an analysis of strong-blast initiation of a gaseous detonation.

Navodit Kaushik

University of Southern Mississippi, Department of Computer Science, Hattiesburg, MS 39406

Solving Second Order ODEs with the Local Discontinuous Galerkin Method: Effect of Stencil Selection on Accuracy

The Runge Kutta Discontinuous Galerkin (RKDG) method has been well established for compressible Euler equations by Cockburn, Shu and coworkers. Following the initial idea of Bassi and Rebay of writing the compressible Navier-Stokes equations as a system of first order equations, the Local Discontinuous Galerkin (LDG) method was also developed by Cockburn and Shu among others. Recent interests are in extending the Discontinuous Galerkin approach to elliptic equations and to the incompressible Navier Stokes equations.

We focus on the application of the LDG method to the following one-dimensional equation:

$$\frac{d^2 u}{dx^2} = f(x).$$

According to literature, one of the salient features of LDG method is that one can obtain high

orders of accuracy using a highly compact stencil. We show that the choice of stencil has a major effect on the accuracy of the method for the above model problem. In particular, it is shown that for a three point stencil, using Legendre polynomial bases, the resulting scheme is at best second order accurate, for the numerical fluxes considered in this work irrespective of the cell size and order of polynomials. The extension of the current work to two-dimensional cases is currently under study. Joint work with Beddhu Murali.

Adnan A. Khan

Rensselaer Polytechnic Institute, Mathematical Sciences, 8th Street, Troy, NY 12180

Turbulent Transport in the Presence of Periodic Fluctuations and Strong Mean Flow

The transport of passive scalars has been well studied using the advection-diffusion equation in the case of periodic fluctuations with a weak mean flow or a mean flow of equal strength through homogenization techniques. However, as the mean flow becomes stronger, homogenization seems to break down. We study the transport of passive scalars using Monte Carlo simulations for the stochastic differential equations describing the tracer trajectories, and study the transport properties as the mean flow is made stronger. We benchmark our numerical experiments by starting in a regime where homogenization works and agrees with our Monte Carlo results, and then we increase the mean flow strength and seek to describe how the breakdown in homogenization is manifested in the statistics of the tracer trajectories. Joint work with Peter R. Kramer (Rensselaer Polytechnic Institute).

Hafiz M. R. Khan

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Predictive Distributions for Responses from the Weibull Life Testing Model

The Weibull distribution has been widely used for modelling life data, and most studies on this distribution concentrate on inference about the parameters or on the reliability and hazard functions. This

presentation is concerned with predictive inference for future responses from a Weibull distribution in a Bayesian framework. It considers both the two-parameter and the three-parameter Weibull distributions. A numerical example is provided, and predictive bounds are determined for various values of the hyperparameters of the prior distribution.

Christopher E. Khedouri

Center for Drug Evaluation and Research (CDER), Food and Drug Administration (FDA), 9201 Corporate Blvd., Room S311, Rockville, MD 20850

Everyday Challenges of a Statistical Reviewer at the Food and Drug Administration (FDA)

This presentation provides a brief overview of the drug review process in the Center for Drug Evaluation and Research (CDER) at the Food and Drug Administration (FDA). The primary objective is to describe the everyday challenges that a statistical reviewer may face throughout the review process. Examples of actual statistical and/or design issues relevant to the Division of Anti-Infective Drug Products (DAIDP) at CDER will be discussed. (The views expressed are those of the author and not necessarily those of the FDA.)

Nickolas Kintos

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Modeling Actions of a Neuromodulator on a Rhythmic Neuronal Network

The stomatogastric ganglion (STG) of the crab, *Cancer borealis*, houses the rhythmic networks that control chewing (gastric mill) and filtering (pyloric) of food. These STG networks are modulated by several substances which are released by projection neurons. One such neuron is the modulatory commissural neuron 1 (MCN1), whose activity elicits a gastric mill rhythm. Previous modeling and experimental work studied the frequency regulation of the MCN1-elicited rhythm by the pyloric rhythm. Recent experimental work, however, showed that bath application of the neuropeptide, peppyrokinin elicits a gastric mill rhythm that is very similar to the MCN1-elicited rhythm. Using a 3D model of the MCN1-elicited rhythm, we reduce to 2D by exploiting the difference in time scales. Then, we investigate possible mechanisms by which the neuropeptide can elicit a gastric mill rhythm that mimics the MCN1-elicited rhythm. We also use a biophysically-detailed model to compare the MCN1-elicited and peptide-elicited rhythms.

Lou Kondic

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Thin Liquid Films with Contact Lines: Instabilities, Coalescence and Rupture

This contribution will concentrate on computational and experimental results involving dynamics of thin film flows on homogeneous and heterogeneous surfaces. In particular, I will present the dynamics of the fluid fronts, i.e., contact line. The presence of contact lines introduces microscales in a macroscale flow and therefore requires bridging the scales and careful modeling and numerical simulations. After presenting basic features of the flow, we will consider several flow configurations. One of these is an unstable configuration involving gravity driven flow on homogeneous and heterogeneous inclined solid surfaces, leading to pattern formation in the form of fingers and rivulets. In particular, the flow on heterogeneous surfaces is interesting since the effect of heterogeneity often competes with the basic instability mechanism, leading to an elaborate interplay of various sources of instability. The computational results are then related to the pattern formation process observed in the experiments performed at NJIT. Other topics of discussion include modeling of the problems involving topological changes, such as drop coalescence and formation of dry spots.

Gregory A. Kriegsmann

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Propagation in Periodic Dielectric Media

We have employed a homogenization procedure to describe the propagation of electromagnetic waves in a dielectric structure which is periodic in the X-Y plane and translationally invariant in the direction of propagation, Z. The fundamental cell is composed of an arbitrarily shaped pore filled with a dielectric and the host by another. The pore shape is allowed to depend upon Z.

Our analysis yields the structure of the electromagnetic fields at the micro level and gives an effective medium equation at the macro level. The latter contains a simple arithmetic average of the dielectric constants and a correction term which involves a line integral around the pore. The integrand of this integral depends upon the polarization of the wave and the solution to a canonical potential problem which has an equivalent variational formulation. We have used this to obtain a simple macroscopic description of the dielectric structure.

Stephen Kunec

Boston University, Center for Biodynamics and Department of Mathematics and Statistics, Boston, MA 02215

Encoding and Retrieval in the CA3 Region of the Hippocampus: A Model of Theta Phase Separation

Past research (Hasselmo et al. 2002) suggests that some fundamental tasks are better accomplished if memories are encoded and recovered during different parts of the theta cycle. A model of the CA3 subfield of the hippocampus is presented, using biophysical representations of the major cell types including pyramidal cells and two types of interneurons. Inputs to the network come from the septum and the entorhinal cortex (directly and via the dentate gyrus). A mechanism for parsing the theta rhythm into two epochs is proposed and simulated: in the first half, the strong, proximal input from the dentate to a subset of CA3 pyramidal cells and coincident, direct input from the entorhinal cortex to other pyramidal cells creates an environment for strengthening synapses between cells, thus encoding information. During the second half of theta, cueing signals from the entorhinal cortex, via the dentate, activate previously strengthened synapses, retrieving memories. Slow inhibitory neurons (O-LM cells) play a role in the disambiguation during retrieval. We compare and contrast our computational results with existing experimental data and other contemporary models. Joint work with M. Hasselmo and N. Kopell (Boston University).

Soumi Lahiri

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Linear and Log-Linear Models Based on Generalized Inverse Sampling Scheme

The linear and log-linear models based on generalized inverse sampling scheme, have a broad application to the research of biological and environmental sciences. Model parameters are estimated using maximum likelihood and modified minimum chi-square estimation methods. The model efficiency is verified based on Wald test. Some applications of these models will be demonstrated. Joint work with Sunil K. Dhar (NJIT).

Sanyogita Lakhera

University of Southern Mississippi, Department of Mathematics, 118 College Drive 5045, Hattiesburg, MS 39406-0001

Computing Numerical Derivatives Using Bernstein Functions

We examine the problem of estimating derivatives of grid functions based on non-compact, non-polynomial estimators built using stochastic interpolation and approximation methods, with particular attention to methods derived using Bernstein functions.

Bernstein functions arise naturally as an extension of the Bernstein polynomials in which the binomial probability density function (pdf) is replaced appropriately with a Gaussian pdf. These lead to row stochastic matrices which can be interpreted as convolution-deconvolution operators that can be used to construct Bernstein polynomial and Bernstein function interpolants.

These methods yield errors which are rapidly convergent (typically in the range of second to sixth order for smooth functions), however they appear to have advantages for non-smooth functions, providing robust interpolants even for very noisy data. Difficulties arise, though, in dealing with the boundaries of the domain as the errors typically decrease to first order near the boundary. Joint work with Joseph Kolibal (University of Southern Mississippi).

Joe Latulippe

Montana State University, Department of Mathematical Sciences, 2-214 Wilson, Bozeman, MT 59717-2400

Weakly Non-Autonomous Bursting Model for Visual Neurons

Certain visual system neurons exhibit On and Off responses when given a light stimulus. By drawing analogies to Hodgkin-Huxley type models, we systematically develop a phenomenological model which replicates On and Off behaviors. This model is also shown to (numerically) replicate the simultaneous stimuli case measured by Kuffler (1953). An examination of the fast-slow dynamics of the model is divided into two cases: constant and non-constant post-synaptic currents. In the constant stimuli case, singular perturbation methods are used to show the dependence of the cell's response for various combinations of stimulus intensity and duration. When the (slowly varying) synaptic current is non-constant the model is weakly non-autonomous. However, we show that by embedding the non-autonomous problem into a higher dimensional autonomous system, much of the analysis for the constant case extends. Preliminary results of this will be shown.

Eric Lauga

Harvard University, Division of Engineering and Applied Sciences, Cambridge, MA 02138

Bacteria Swimming in Circles

Near a solid boundary, *E. coli* does not swim in a straight line but performs a clockwise circular motion. We provide a hydrodynamic model for the motion of such bacteria near solid surfaces. We show that circular trajectories are natural consequences of force-free and torque-free swimming with account taken of hydrodynamic interactions with the boundary. Results of the model are compared to a new set of experimental data and reasonable agreement is obtained. Joint work with Willow DiLuzio, George M. Whitesides, and Howard A. Stone (Harvard University).

Kevin K. Lin

Courant Institute, New York University, 251 Mercer Street, New York, NY 10012

Entrainment and Chaos in the Pulse-Driven Hodgkin-Huxley Oscillator

The Hodgkin-Huxley system of differential equations was derived to model action potential generation in the squid giant axon, and has long served as a paradigm for quantitative models of neurons. I present numerical evidence that a surprisingly rich range of qualitatively distinct responses may be elicited from the Hodgkin-Huxley model by the addition of a periodic impulse train. These response types include

- * Stable entrainment to the input pulse train;
- * Transient chaos followed by entrainment;
- * Fully chaotic behavior characterized by a positive Lyapunov exponent, exponential decay of correlations, and the existence of a (unique) SRB measure.

These results are consistent with the predictions of Qiudong Wang and Lai-Sang Young, whose theory of kicked nonlinear oscillators motivated this work.

Xinfeng Liu

SUNY at Stony Brook, Applied Mathematics and Statistics, Stony Brook, NY 11794

Computational Algorithms for Dynamic Interface Tracking in Three Dimensions

A major challenge to this method is to handle changes in the interface topology. Two tracking methods, one named grid-free tracking and the other named grid-based tracking have been used to handle the three dimensional interface propagation and topological bifurcation. The former is a pure Lagrangian method in which the interface propagation and redistribution are independent of the underlying Eulerian grid. The detection and resolution of topological bifurcation is fully determined by the interface itself. This method is more accurate in the propagation of the interface position, but it is not robust in resolving the topological bifurcations. The latter is just the opposite. I developed locally grid based method, which uses Lagrangian propagation and redistribution, but applies Eulerian reconstruction for the bifurcation of Topology. This new method takes advantage of grid based method and grid free method, and reduces the use of Eulerian reconstruction to a minimum. It is particularly useful for the computation of an interface motion in which the interface forms sharp corners. It will also reduce the unphysical disappearance of the fragmented components of the material after bifurcation. The success of this improved algorithm is shown by simulations of turbulent mixing, such as the acceleration driven Rayleigh-Taylor instability, we found our simulations are now too fast, in disagreement with experiment, but when we include physical values of surface tension (we have implemented the revised calculation of curvature for a three dimensional surface into our code) and mass diffusion, we recover consistency with experimental values. Joint work with James Glimm and Xiaolin Li...

Xing Liu

University of Maryland, Department of Mathematics and Statistics, UMBC, 1000 Hilltop Circle, Baltimore, MD 21250

Optimization in 2D Gel Image Alignment

A new method for aligning families of two-dimensional polyacrylamide gel electrophoresis (2D-PAGE) images arising in proteomics studies is presented. Piecewise polynomial transformations via a multi-resolution approach is used to align the family of gels to an ideal gel. Both the ideal gel and the coefficients defining the transformations are obtained by solving a quadratic programming problem. Numerical results for a family of 123 gels are reported. Joint work with Florian Potra.

Dawn A. Lott

Delaware State University, Mathematics and Biotechnology, 1200 N. DuPont Hwy, Dover, DE 19901

The Configuration of the Aneurysm Neck and Proximal Dome Profoundly Affects Shear Stress and Flow Velocities within an Aneurysm and its Parent Vessel

Introduction: Flow characteristics of aneurysms depend upon the input flow and size of the parent vessel and aneurysm. Little is written describing the role of neck and dome configuration in establishing areas of shear stress as a potential for aneurysm growth. We developed a side-wall aneurysm model and compared different neck and dome configurations with regards to shear stress, velocity and pressure.

Methods: A finite-volume based package for modeling complex flow was utilized. A 5.3mm side-wall aneurysm with a 3.1mm neck was created and analyzed under pulsed-flow conditions. Shear stresses and velocities within the parent vessel, aneurysm, and interface were determined along several points of the cycle. Changes in neck and proximal dome geometry were made, and the analysis repeated. Comparisons in maximum velocities, pressures and shear stress were made.

Results: Changes in configuration of the neck, while maintaining its 3.1mm size altered the flow velocity, pressure and shear stress along the distal portion of the aneurysm, neck and parent vessel. Vortical flow patterns within the aneurysm dome changed, with flow velocities increasing at the interface between the aneurysm neck and parent vessel. The velocities at the distal neck (inflow zone) significantly increased when proximal dome configurations were modified as well.

Conclusion: The configuration of the neck and proximal dome, independent of the actual neck and aneurysm size, dramatically affects flow characteristics within the aneurysm and the aneurysm/parent vessel interface. Further benefits of the 2-D model as a possible predictor of aneurysm behavior will be explored. (Joint work with Hans Chaudhry and Michael Siegel (NJIT), and Charles J. Prestigiacomo (NJIT and UMDNJ).

Tianshi Lu

SUNY at Stony Brook, Department of Applied Math and Statistics, Stony Brook, NY 11794

Direct Numerical Simulation of Bubbly Flows and Application to the Mitigation of Cavitation Erosion

We have investigated the propagation of acoustic and shock waves in bubbly flows using direct numerical simulation. In this method, the liquid and the gas bubbles are represented by single-phase domains separated by free interfaces. FronTier, a front tracking hydro code was used for numerical simulations. It is capable of tracking simultaneously a large number of interfaces and resolving their topological changes in two- and three-dimensional spaces. We have compared results of our numerical simulations with theoretical predictions and experimental data on the propagation of linear sound waves and shock waves in bubbly fluids. The method has been applied to estimate the efficiency of gas bubble mitigation in reducing the cavitation erosion of the container of the Spallation Neutron Source liquid mercury target. Joint work with R. Samulyak (Brookhaven National Lab) and J. Glimm (SUNY at Stony Brook).

Jonathan H. C. Luke

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

An Effective Fluid Model for the Decay of Velocity Fluctuations in a Sedimenting Suspension

Effective fluid models for sedimentation have often neglected fluctuations in the particle density. For well-stirred suspensions, however, these fluctuations produce enormous velocity fluctuations in large containers. We use an effective fluid model to study the transition from the large fluctuations typical of well-stirred suspensions to the small-fluctuations seen in "steady" sedimentation. We present the time-scales of the fluctuation decay and an assessment of the short-time asymptotics of the system energy and velocity fluctuations.

Valeriy V. Lukyanov

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Analytical Modeling of Rayleigh-Bloch Surface Waves Along Metallic Rectangular Rods

We develop an analytical method to analyze and to study Rayleigh-Bloch surface waves propagating along a two-dimensional diffraction grating which again consists of a periodic array of rods with rectangular cross sections. The method is based on mode matching. By taking into account all propagating and only a finite number of evanescent modes in a specific portion of the waveguide, we show that the surface waves correspond to the zeros of the determinant of a Hermitian matrix. We demonstrate numerically that the method gives an accurate result if we take into account only several evanescent waves. Joint work with Gregory A. Kriegsmann (NJIT).

Marc Q. Ma

NJ Institute of Technology, Computer Science Department and Center for Applied Mathematics and Statistics, University Heights, Newark, NJ 07102

A Molecular Dynamics Study of the Effect of YC-1's Binding Modes on the Structure of Soluble Guanylyl Cyclase

Soluble guanylyl cyclase (sGC) is an enzyme that can be allosterically activated by synthetic compounds such as YC-1 and its derivatives. sGC catalyzes the cyclization of the substrate guanosine 5'-triphosphate (GTP) to guanosine 3',5'-cyclic monophosphate (cGMP). cGMP is a second messenger molecule that regulates lots of biological processes including vasodilation. The mechanisms regulating the catalytic activity of sGC remain unclear. Recently, however, a mutagenesis study has been reported, in which a variety of point mutations were made in wild-type (WT) sGC that affect the catalytic activity and YC-1 allosteric activation of sGC. These new data still do not immediately point to a unifying molecular mechanism to explain sGC's regulatory processes. We present preliminary results in using all-atom, classical molecular dynamics (MD) simulations to study the pre-chemistry conformational changes induced by YC-1 with the ultimate goal of establishing a valid detailed molecular model of allosteric regulation in sGC. The details of our simulation protocol and some novel discoveries will be presented. Joint work with Kentaro Sugino and Yu Wang (NJIT), and Annie V. Beuve (UMDNJ).

Marc Q. Ma

NJ Institute of Technology, Computer Science Department and Center of Applied Mathematics and Statistics, University Heights, Newark, NJ 07102

GO-LinkGenes, A Multiple Component Data Mining Framework for Knowledge Discovery in Microarrays

We present the design and evaluation of GO-LinkGenes, a data mining framework to facilitate exploration and knowledge discovery in DNA microarray experiments. GO-LinkGenes consists of two components, GO-Chart and LinkGenes. Each component can work independently of each other, or together in a unified fashion. GO-Chart is designed to visualize more conveniently the output from GOTM (GOTree Machine), while adjusting the p-values for multiple testing. GO-Chart allows users to analyze genes by Gene Ontology functional categories, occurrence, multiple parents-children relationships. LinkGenes is designed as a web-based tool for linking multiple genes to physical locations on human chromosomes and visualizing these genes. Feeding the output of genes identified using GO-Chart to LinkGenes, we are able to identify several human chromosomal regions that could be linked to either gene expression phenotypes or metabolic pathways. Genetic disease information associated with these genes is also obtainable through this tool. Joint work with Tongsheng Wang (NJIT and Public Health Research Institute, Newark, NJ 07103), Deepali Shah and Jason Wang (NJIT), Gerard Tromp (Wayne State University-School of Medicine), and Patricia Soteropoulos (Public Health Research Institute, Newark, NJ 07103).

Victor Matveev

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

A Bound-Calcium Mechanism of Synaptic Facilitation Revisited

Synaptic facilitation (SF) is a form of a transient increase in synaptic strength elicited with just one or several stimulation pulses, and decaying with time constants from tens to hundreds of milliseconds. At some synapses, SF may be caused by the increase in the activity-induced Ca^{2+} influx; however, in many other synaptic types SF is believed to result from the presynaptic accumulation of residual Ca^{2+} , under conditions of constant Ca^{2+} current from one pulse to the next. In the latter case, it is not known whether it is free or bound residual Ca^{2+} that underlies SF. Experimental work has demonstrated that an increase in intrasynaptic Ca^{2+} buffering capacity leads to a rapid reduction in both the baseline synaptic response and in the magnitude of SF; this is often viewed as a proof that SF is caused by the accumulation of Ca^{2+} in free form. In the past we have explored two variations of such free residual Ca^{2+} hypothesis of SF: the so-called two-site model, and the buffer saturation mechanism. However, here we show that a model including the contribution of bound Ca^{2+} to SF is also consistent with the observed effect of exogenous Ca^{2+} buffers on synaptic response, and thus represents a viable alternative to the two-site free residual Ca^{2+} model. In particular, we show that such hybrid free/bound Ca^{2+} model is not in contradiction with the Kamiya-Zucker protocol (1994, Nature 371:603), whereby the synaptic strength of the crayfish neuromuscular junction is seen to decrease within few milliseconds of a UV-flash photolysis liberating a Ca^{2+} buffering compound. While evidence indicates that buffer saturation may underlie SF at calbindin-positive central synapses, we conclude that SF at other synaptic types may well involve a slow Ca^{2+} unbinding step from a putative Ca^{2+} release sensor. Joint work with Richard Bertram (Florida State University) and Arthur Sherman (NIH).

Roberto Mauri

University of Pisa, Department of Chemical Engineering, Pisa, Italy

Mixing of Viscous Liquid Mixtures

We simulate the mixing process of a quiescent binary mixture that is instantaneously brought from the two- to the one-phase region of its phase diagram. Our theoretical approach follows the diffuse interface model, where convection and diffusion are coupled via a body force, expressing the tendency of the demixing system to minimize its free energy. In liquid systems, as this driving force induces a material flux which is much larger than that due to pure molecular diffusion, drops tend to coalesce and form larger domains which eventually must dissolve by diffusion. Therefore, in macroscopically quiescent mixtures, while convection speeds up phase separation, it effectively slows down mixing, which therefore is faster when the mixtures have large viscosities. In addition, the mixing rate is also influenced by the Margules parameter, which describes the relative weight of enthalpic versus entropic forces. In the late stage of the process, this influence can approximately be described assuming that mixing is purely diffusive and is therefore characterized by a self-similar solution of the governing equations, which leads to a $1/t$ power-law decay for the degree of mixing, i.e. the mean square value of the composition field. Joint work with A.G. Lamorgese (Cornell University).

Zoi-Heleni Michalopoulou

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Using Time-Frequency Analysis and the Bayesian Paradigm to Extract Modal Arrivals in Underwater Acoustics

Time-frequency analysis of signals that have propagated in an oceanic waveguide demonstrate modal dispersion in an informative manner. Identifying precise modal arrival times at specific frequencies can then be used for localization and geoacoustic inversion. Simple spectrograms, however, do not always allow accurate estimation of an arrival time - frequency pair for a given mode; especially in the presence of ambient noise, uncertainty characterizes such estimates. We here show how we can improve modal arrival estimates

using a Bayesian approach for time-frequency surface estimation as has been proposed for audio signals [Wolfe et al, Journal of the Royal Statistical Society, Series B, vol. 66, 2004].

Yuriy Mileyko

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Algorithms for Swept Surface Intersections

A new class of geometric objects called swept surfaces is studied and the problem of finding intersections of such objects is considered. Three algorithms for computing swept surface intersections are presented. The algorithms are based on ideas from existing intersection methods and utilize properties specific to swept surfaces to achieve the desired efficiency. Some of the employed properties of swept surfaces are also discussed.

Petronije Milojevic

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Fredholm Theory for Hammerstein Equations

Let X and Y be Banach spaces, $K : Y \rightarrow X$ be a linear map and $F : X \rightarrow Y$ be asymptotically positively homogeneous and odd, i.e., $F = F_1 + F_2$ with F_1 positively homogeneous and odd outside some ball and $\|F_2x\| \leq a \|x\| + b$ for large $\|x\|$ and some constants a and b . Assuming that $x + KF_1x = 0$ has only the trivial solution, we discuss various conditions on K and F that imply the solvability of the Hammerstein equation (*) $x + KF_x = f$ for each f in X (a generalized first Fredholm theorem). We require that $I + KF$ be a (pseudo) approximation-proper map. In particular, we consider the cases when K is compact, or monotone and possibly nonselfadjoint, while F is of ball contractive or monotone type. The number of solutions of (*) is discussed when $I + KF$ is approximation-proper. Under suitable conditions, we also prove a complete extension of the Fredholm alternative to these classes of maps. Applications to Hammerstein integral equations and to BV problems for differential equations are also given.

Victor A. Miroshnikov

College of Mount Saint Vincent, Department of Mathematics and Computer Science,
6301 Riverdale Avenue, Riverdale, NY 10471

The General Solution for the Two-Dimensional Poiseuille Flow

In two dimensions, general solutions of the unsteady Navier-Stokes equations are computed symbolically in the form of the Boussinesq-Rayleigh series and evaluated numerically. For finite Reynolds numbers, a nonlinear system of differential recurrent relations admits the following general solutions: the series solution for flows forced by the dynamic pressure and the series solution for freestreams. For generating functions, which are bounded together with their derivatives, the absolute convergence of the series solutions is shown by converting the differential recurrent relations into tensor recurrent relations and using the comparison and ratio tests. A triangular structure of three tensors of derivatives, which is employed in the tensor recurrent relations, is obtained by induction. It is shown that the general solutions away from boundaries are nonlinear superpositions of the Stokes flow, the Bernoulli flow, the Couette flow, and the Poiseuille flow. The general solution for the Poiseuille flow is specified by periodic generating functions, which model mixing of two-dimensional flows away from boundaries.

Milind Misra

NJ Institute of Technology, Department of Chemistry and Environmental Science, University Heights, Newark, NJ 07102

Fuzzy Relational Clustering of Molecular Conformations Using Novel Features Based on DNA Base-Pair Step Parameters

Six rigid-body parameters (Tilt, Roll, Twist, Shift, Slide, Rise) are commonly used to describe the relative orientation and positioning of any two base pairs in a nucleic acid structure. The present work generalizes the algorithms of the 3DNA software package (*Nucleic Acids Res.*, 31, 5108-21, 2001) to describe the relative orientation of any two planes in a molecule—for example, planes which contain important pharmacophore elements. Fuzzy relational clustering is used to classify molecular conformations using the six base-pair step parameters as features. This approach is applied to an analog of GBR 12909, a flexible inhibitor of the dopamine transporter potentially useful in the treatment of cocaine abuse. The results of this approach provide representative conformers to be used as templates for future 3D-QSAR (CoMFA) analysis. Joint work with Amit Banerjee, Deepa Pai, Rohan Woodley, Rajesh N. Davé, Liang-Yu Shih, Carol A. Venanzi (NJIT), and Xiang-Jun Lu, A. R. Srinivasan, and Wilma K. Olson (Rutgers University, Piscataway, NJ).

Robert M. Miura

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Modeling Nonlinear Waves of Spreading Depression

Slow chemical waves of spreading cortical depression (SD) have been observed during experiments in a variety of brain structures in different animals. Several mechanisms that are believed to be important in

modeling SD will be described, including ion diffusion, the spatial buffer mechanism, membrane ionic currents, osmotic effects, neurotransmitter substances, gap junctions, metabolic pumps, and synaptic connections. Ion diffusion and spatial buffering have been treated both theoretically and numerically, in simplified geometries, and several of the other mechanisms have been investigated numerically. In this talk, I will describe continuum models that consist of coupled nonlinear diffusion equations for the ion concentrations, and a discrete model that corresponds to treating the brain-cell microenvironment using a lattice Boltzmann method.

Brian E. Moore

McGill University, Mathematics and Statistics, 805 Sherbrooke Street W., Montreal, Quebec H3A 2K6 Canada

Modified Equations for Multi-Symplectic Integrators

A useful way to understand symplectic integration of Hamiltonian ODEs is through the system of equations, known as the modified equations, which are solved by the numerical solution. Here, the ideas of symplectic integration are extended to Hamiltonian PDEs, such that the symplectic structure in both space and time is exactly preserved. This paves the way for the development of a local modified equation analysis solely as a useful diagnostic tool for the study of these methods. In particular, the modified equations are used to derive modified conservation laws of energy and momentum that are preserved to higher order along the numerical solution. Results on the behavior of the modified equations are also demonstrated through an application to traveling wave solutions.

Richard O. Moore

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Thermally Induced Dynamics and Pattern Formation in Optical Parametric Oscillators

Optical parametric oscillators (OPOs) are an important source of laser-quality light in the far infrared. When operated at high average powers, absorption can lead to significant heating of the gain medium, changing the cavity properties and leading to thermal lensing and deformation of the transverse beam profile. To understand this process better, we consider the formation and evolution of patterns in reduced models of OPOs coupled to a diffusion equation for the temperature.

Pascal Moyal

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Stationarity of Some Queues with Impatient Customers

Consider a queueing system with one server and infinite capacity, where the customers enter following an ergodic sequence $(U_n, n \in \mathbb{N})$, requesting ergodic services $(\sigma_n, n \in \mathbb{N})$. The customers are furthermore impatient: they must meet their requirement before a fixed due-date, and leave the system if they did not reach the server before. Their *patiences* $(D_n, n \in \mathbb{N})$ are ergodic. We denote such a queue by $G/G/1+G$. Such a model has first been used to describe call centers with impatient customers, and applies furthermore in many time-sensitive computer networks or multimedia systems.

We investigate the stability of the queue: is 0 positive recurrent for the congestion process (*regenerativity*)? These questions have only been answered in the FIFO (*First in, First out*) context, where the arrival process is a renewal process, and the service durations as well as the patiences are i.i.d. ($G/G/1+GI$ system): a sufficient condition for the system to be regenerative is given by $P[\sigma < U] > 0$. But in a general $G/G/1+G$ system under any service discipline (as the EDF -*Earliest Deadline First-one*), the results are so far restricted, due to a higher stochastic instability. We prove the following more general statements:

1. any $G/G/1+GI$ system satisfying $P[D + \sigma < U] > 0$ is regenerative,
2. for any $G/G/1+G$ system, there exists a monodimensional description of the system $(Y_n, n \in \mathbb{N})$ tending to a stationary distribution Y as finite,
3. under further assumptions, Y is unique and the queue is regenerative.

Cyrill Muratov

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Signal Propagation and Failure in Discrete Autocrine Relays

This poster presentation is of my recent joint work with S. Shvartsman on modeling cell communication mechanisms in epithelial layers. The main mechanism at work here is ligand-induced ligand release coupled to the diffusion of ligands in the extracellular matrix. It is believed that under physiologically relevant range of parameters cell discreteness plays an important role in cell-to-cell communication. We formulated a fully discrete model of an autocrine relay, in which under physiologically reasonable assumptions it was possible to completely characterize propagation and failure. In particular, we obtained exact closed form discrete traveling wave solutions in this cell communication system.

Duane Nykamp

University of Minnesota, School of Mathematics, 206 Church Street SE, Minneapolis, MN 55455

Reconstructing Subpopulation Connectivity Within Neuronal Networks

To understand the function of neuronal networks within the brain, one would, at minimum, like to characterize the connectivity patterns that underlie these networks. Since one can simultaneously measure only a tiny fraction of neurons, the presence of vast numbers of unmeasured neurons can confound attempts to determine the subnetwork connecting the measured neurons.

A suitable mathematical framework can provide the structure to account for the presence of unmeasured neurons. Using a simple probabilistic model of a neuronal network, we demonstrate how to analyze connections among a subset of measured neurons embedded in the larger network. One obtains connectivity patterns in terms of certain neural subpopulations, which are groups of neurons with a similar response to a stimulus. This subpopulation connectivity can capture how network connections depend on the response properties of neurons.

Although the results are presented in terms of neuronal networks, the mathematical framework is applicable to other networks, such as gene regulatory networks.

Ozgur Ozen

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Experiments on Electrohydrodynamic Instability of Two-Layer Flow in a Square Channel

The stability of two-fluid flow in a channel is of importance in the design of microfluidic systems. Due to the low Reynolds numbers in micro-channels, it is relatively difficult to attain mixing in these geometries. Recent studies using miscible fluids have shown that applying electric fields to these systems drastically enhances mixing over a short distance and in short times. However, in a class of applications, the fluids in contact are not miscible, and the interfacial tension stabilizes the interface; an effect absent in the physics of miscible fluids.

We have performed experiments where two immiscible fluids of different electrical properties flowing in a square channel are subjected to an electrical field normal to the liquid-liquid interface. Our experiments show that electric fields indeed affect the stability of the interface between two immiscible fluids and careful application of electrohydrodynamic instability can be used to cause a controlled deflection of the interface.

The experiments are also compared qualitatively to the theoretical study of the problem where the linear stability analysis of the two-fluid flow subject to a normal electric field is carried out using the Chebyshev Spectral Tau method. Joint work with N. Aubry, D. T. Papageorgiou, and P. Petropoulos (NJIT).

Demetrios T. Papageorgiou

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Nonlinear Dynamics of a Leaky Dielectric Fluid Sheet Under Horizontal Electric Fields

In related work we have shown that a horizontal electric field applied to a fluid sheet comprised of a leaky dielectric can become unstable. This instability is not present for perfect dielectrics and arises due to the build-up of charge at the interface and the ensuing modification of the tangential stress balance there. In the linear regime we identify a canonical asymptotic limit that yields a rational long wave nonlinear model. This limit occurs as the electric field parameter, E_b say, tends to zero and a band of unstable waves is seen with a cut-off wavenumber $E_b^{1/2}$ and growth rate E_b . This leads to nonlinear long waves of size $E_b^{1/2}$ which is a measurable physical parameter (in general long wave theories assume the existence of an arbitrary long wavelength). A nonlinear system of evolution equations is derived asymptotically and it is found that the electric field in the surrounding medium enters in a local manner due to the smallness of E_b . The equations are studied for nonlinear traveling waves and the initial value problem is addressed to study possible rupture of the sheet. Joint work with Ozgur Ozen and Peter G. Petropoulos (NJIT).

John A. Pelesko

University of Delaware, Department of Mathematical Sciences, Ewing Hall Room 406, Newark, DE 19716

The MEC Lab, Hands-On Mathematics at the University of Delaware

Just about everyone expects to find mathematics being used in the laboratory, but few of us expect to find mathematicians in the laboratory. Well, at the University of Delaware in the Department of Mathematical Sciences, finding mathematicians in the laboratory is quickly becoming a commonplace event. This was made possible with the establishment of the MEC Lab in the fall of 2002. This experimental laboratory, housed in the Department of Mathematical Sciences, takes its name from the words Modeling, Experiment, and Computation. These three words capture the hands-on philosophy of the lab; in the MEC Lab, math is meant to be experienced from every possible direction. This means carrying out real-world, often-dirty, hands-on experiments, constructing mathematical models, and analyzing them with the aid of

computers. In this poster, we showcase the many ways the MEC Lab has been incorporated into teaching and research in the Department of Mathematical Sciences at the University of Delaware. Joint work with Louis Rossi.

Mark Pernarowski

Montana State University, Mathematical Sciences, Bozeman, MT 59717

Return Map Characterizations for a Model of Bursting with Two Slow Variables

Many neurons and endocrine cells exhibit periodic bursting oscillations in their transmembrane electrical potential. The fast subsystems of the corresponding models exhibit bistability between stable equilibria and periodic orbits. Slow variables evolve in a manner which causes the solutions to switch between pseudo-stationary and oscillatory states resulting in a characteristic bursting pattern.

Most recent models involve two slow variables which tends to complicate their analyses. Using singular perturbation techniques we show that bursting solutions of such models correspond to fixed points of a one dimensional map constructed from the fast and slow subsystems. We further show that for some parameter values, bistability between bursting solutions and stable equilibria is possible. Joint work with Roger Griffiths (Mercyhurst College).

Peter G. Petropoulos

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Chemical Oscillations & Waves in the Undergraduate Applied Mathematics Laboratory: The Belousov-Zhabotinsky (BZ) Reaction

By considering the modeling of the BZ reaction in a closed reactor (glass beaker) with continuous stirring (spatially homogeneous reaction modeled by a 3x3 system of nonlinear ordinary differential equations) the students make the connection between topics from nonlinear dynamics theory (stability of steady-states and birth/death of limit cycles through super/sub-critical bifurcations) and a real-world problem where chemical oscillations arise. Then, the students perform the actual experiment where potentiometric methods are employed to measure the two most important chemical concentrations as the reaction proceeds to equilibrium. Finally, the students identify the theoretical attributes of the BZ mathematical model in the measured time traces by constructing the phase plane from the measurements.

By considering the modeling of the BZ reaction in Petri dishes (spatially inhomogeneous reaction) the students study how systems of parabolic reaction-diffusion partial differential equations arise and how traveling waves are then possible in such systems. By performing the actual experiment and capturing sequences of images the students experimentally measure the speed of propagation of reaction fronts and compare to the theoretically obtained results for the speed of traveling waves.

A web diary of our progress can be found at <http://web.njit.edu/~peterp/Capstone.html>.

Valentin Polishchuk

SUNY at Stony Brook, Applied Mathematics and Statistics, Stony Brook, NY 11794-3600

Touring Convex Polytopes – A Conic Programming Solution

We study the problem of finding a shortest tour visiting a given sequence of convex bodies in \mathbb{R}^d . To our knowledge, this is the first attempt to attack the problem in its full generality: we investigate high-dimensional case ($d \geq 2$); we consider convex bodies bounded by (hyper)planes and/or (hyper)spheres; we allow a different cost of travel through each of the bodies; we do not restrict the start and the goal positions of the path to be single points, etc. Formulating the problem as a second order cone program (SOCP) makes it possible to incorporate distance constraints, which cannot be handled by a purely geometric algorithm.

Next, we focus on the planar case ($d = 2$), with the convex bodies being just straight line segments – adjacent edges in a triangulation of a polygonal domain. In this setting we introduce a set of separation constraints on the path and observe that the new constraints can naturally be handled by the conic program. We also consider the case in which each face of the triangulation is assigned a weight, and the length of the path is measured according to the weighted region metric.

We implemented the SOCP program in MATLAB and obtained its solution with the SeDuMi package. We ran computational experiments, which suggest that the proposed solution is practical. Finally, we present NP-hardness results, showing that the assumptions we make in the statement of our problems are crucial for the problems to be tractable. Joint work with Joseph S. B. Mitchell (Stony Brook University).

Keywords: Computational Geometry, Weighted Region Metric, Conic Programming, Algorithms Complexity.

Joshua L. Proctor

University of Washington, Department of Applied Mathematics, Box 352420, Seattle, WA 98195

Theory of Q-Switching in Actively Mode-Locked Lasers

An analytic theory is proposed which characterizes Q-switching in an active mode-locked cavity as the nonlinear interaction of two unstable modes: one symmetric, another anti-symmetric. The phase difference between these modes generates a nonlinear beating interaction which gives rise to quasi-periodic behavior in the laser cavity. This quasi-periodic behavior is responsible for the Q-switching phenomena and is controlled by the interaction and overlap between neighboring pulses. Using a linear stability analysis, a

simple qualitative description of the Q-switching phenomena is given which is verified with numerical simulations of the governing active mode-locked equations. This model characterizes the Q-switching as a function of the physical parameters of the laser cavity and elucidates the mechanisms for controlling its behavior.

Christopher Raymond

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Effect of Reversible Chemistry on Immunocolloid Labeling

We investigate the effect of reversible chemistry on the time course of a surface-volume biochemical reaction. We show how to modify asymptotic techniques developed for the irreversible case to produce approximate analytical solutions for arbitrarily strong reversibility and validate these approximate solutions by comparison to full numerical solutions.

Gregory T. Reeves

Princeton University, Chemical Engineering Department, Princeton, NJ

Computational Analysis of EGFR Inhibition by Argos

Argos, a secreted inhibitor of Drosophila epidermal growth factor receptor, and the only known secreted inhibitor of receptor tyrosine kinases, acts by sequestering the EGFR ligand Spitz. We use computational modeling to show that the biochemically-derived mechanism of Argos action is consistent with genetic data on EGFR/Spitz/Argos interactions in vivo. We find that efficient Spitz sequestration by Argos is key for explaining the existing data and for providing a robust feedback loop that modulates the Spitz gradient in patterning of the embryonic ventral ectoderm. Our analysis of the EGFR/Spitz/Argos module shows that Argos need not be long-ranged in order to account for genetic data in the ventral ectoderm. In our models, Argos action over both long and short length scales can effectively limit the range of secreted Spitz. Thus, the spatial range of Argos does not have to be tightly regulated or may depend on the developmental context. In addition, we verify the experimentally observed robustness of the wild type patterning of the ventral ectoderm and show that negative feedback control through Argos serves to impart this robustness.

Joint work with Rachel Kalifa and Stanislav Y. Shvartsman (Princeton University) and Daryl E. Klein and Mark A. Lemmon (University of Pennsylvania School of Medicine).

Max Roman

NJ Institute of Technology, Mechanical Engineering Department, University Heights, Newark, NJ 07102

Modeling, Design, and Fabrication of Pulsed Micro-Jet Actuators

The forced vibration of a thin flexible plate or membrane in a sealed cavity with a small opening can cause fluid to be pumped into and out-of the cavity. If the frequency and amplitude of vibration are large enough, a streaming of vortex rings occurs near the orifice. Moving under their own self-induced momentum, downstream of the opening these vortex pairs ultimately break up to form a fully developed jet. This phenomenon has been shown to be effective for a multitude of applications, including mixing, cooling of electronic components, micro-propulsion and flow control, such as the suppression of vortex shedding. Our own preliminary studies measuring the heat dissipation of dense circuitry by direct impingement with a pulsed jet have shown great promise.

Microfabrication (MEMS) offers a platform to build miniaturized inexpensive, reliable, light-weight, and low power actuators and sensors. Such small actuators can have a very unique function in microfluidics, where they can serve as micromixers, pumps, and non-invasive cell manipulators. This work is dedicated to the analysis, design, and fabrication of electrostatically actuated pulsed microjet actuators. This work is unique in that theoretical modeling, computer simulation (CFD-Computational Fluid Dynamics) and experiments are conducted in unison. A low dimensional theoretical model takes into account the coupling between the electrostatic actuation, the solid deformation of the membrane, and the squeeze flow in the cavity. Computational fluid dynamics allows us to better understand the formation of the vortex rings both inside and outside the cavity and how the jet develops. Experiments, aided by the use of PIV (Particle Image Velocimetry) validate both the model and CFD results. It is our goal that by understanding fundamentally the coupled physics of the actuator, the performance, in terms of exiting volume flux and dynamic response, can be precisely controlled and, therefore, applied to many diverse applications. Joint work with Nadine Aubry (NJIT).

Satrajit Roychoudhury

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Family of Probability Generating Functions Induced by Shock Model

The question investigated here can be simply posed as follows: Under what conditions is the function defined by:

$$\int_0^{\infty} p^z Q(dp),$$

a *probability generating function* (p.g.f) of an non-negative integer valued random variable N . The answer is clearly affirmative if the support of the mixing distribution Q is no larger than $(0; 1]$. This paper explores necessary conditions for which the function is a p.g.f. The motivation of this problem comes from some nonparametric aging properties of reliability methods. Joint work with M.C. Bhattacharjee (NJIT).

Keywords: Probability generating function, completely monotone functions, Essay Marshall and Proschan (EMP) shock model.

Roman Samulyak

Brookhaven National Laboratory, Computational Science Center, Upton, NY 11973

MHD of Multiphase Flows at Low Magnetic Reynolds Numbers

We have developed mathematical models, numerical algorithms, and computational software for the study of magnetohydrodynamics (MHD) at low magnetic Reynolds numbers of multiphase flows with phase transitions and free surfaces. The numerical models are based on the front tracking method for interfaces, a liquid-vapor phase transition and ablation models for phase boundaries, and high performance solvers for coupled hyperbolic - elliptic systems in geometrically complex and moving domains. Numerical simulations of hydro- and MHD processes in liquid mercury targets for future accelerators, and processes associated with the refueling of tokamaks through the injection of frozen deuterium pellets will be discussed.

Tobias Schaefer

CUNY, Mathematics Department, 2800 Victory Boulevard, Staten Island, NY 10314

Impact of Microstructures on Macroscopic Observables in Nonlinear Systems

We present three approaches to understand the relation of microscopic and macroscopic scales in nonlinear systems. Each approach is discussed with an example that is of its own interest:

- (a) Method of multiple scales: Nonlinear Schroedinger vs. Shortpulse equation
- (b) Coarse-graining randomness in nonlinear systems: Random susceptibility and random dispersion
- (c) Chorin's Optimal Prediction: Averaging of Hamiltonian Systems

Eric Shea-Brown

Courant Institute, New York University, 251 Mercer St., New York, NY 10012

How Architecture Restricts Spiking Patterns in Networks of Phase Oscillators

We study networks of coupled phase oscillators and show how their architecture shapes their dynamics, forcing subsets to have the same oscillation numbers and interleaving spiking times. Our analysis follows the theory of coupled systems of ODEs developed by Stewart, Golubitsky, Pivato, and Torok, developing surprising new consequences for phase systems.

We say that two oscillators coevolve if the space on which their phases are equal is dynamically invariant and prove the results about oscillation numbers and spike times for these pairs. We then introduce a partition of oscillators in a network into 'collections' with closely related dynamics. We illustrate all of our results using the well-known 'theta neuron' model; implications for integrate and fire neurons are also discussed. Joint work with Martin Golubitsky and Kresimir Josic (University of Houston).

Asya Shpiro

New York University, Center for Neural Science, 4 Washington Place, New York, NY 10003

Dynamics of Neuronal Competition Models for Binocular Rivalry

We consider several reduced population firing rate models to describe oscillation dynamics during neuronal competition, in particular, during binocular rivalry - an alternation of percepts when different steady images are presented to the two eyes. These models, by Laing and Chow (2002) and Wilson (2003), involve two neuronal populations, that correspond to the neural representations of competing percepts. They include recurrent excitation, cross-inhibition, and a slow negative feedback process. We focus on examining the effect of the input strength (e.g., contrast) to the two populations on the rate (and existence) of oscillations. In all models considered, there is a range of parameters when five distinct regimes of behavior are observed: as

the stimulus strength, common to both populations, decreases, the system's dynamics changes from both populations being highly active (fusion regime) to oscillations with the period increasing with decreasing input, followed by a winner-take-all regime, then by the oscillatory regime again, but with the period of oscillations decreasing as input decreases, and, finally, by the fusion at low activity level.

This result contradicts the classical view, based on many psychophysics experiments, in which the oscillation's period increases with decreasing input (this constitutes Levelt's Proposition I, 1968). We use phase plane methods to analyze the system's dynamics in a particular case of a Laing-Chow-like model with spike frequency adaptation, and show that different period vs. stimulus strength dependencies correspond to different mechanisms of establishing the alternating behavior: "release" and "escape" (Wang and Rinzal, 1992). We calculate the values of the input where the transitions between regimes occur, as functions of the model's parameters. Finally, we use phase plane arguments to analyze Levelt's Proposition II, which states

that if the stimulus strength to one population is changed, it is the dominance duration of the other population will be affected. We show that satisfaction of Levelt's Proposition II depends on whether the Levelt's Proposition I is satisfied or not. Joint work with Rodica Curtu (Transylvania University of Brasov, Romania), Nava Rubin (New York University), and John Rinzel (Courant Institute of Mathematical Sciences and Center for Neural Science).

Michael Siegel

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Steady Deformation and Tip-Streaming of a Slender Bubble with Surfactant in an Extensional Flow

We consider steady-state deformation and time-dependent evolution of an inviscid, axisymmetric bubble in zero-Reynolds-number extensional flow when an insoluble and immobile surfactant is present on the bubble surface. Asymptotic solutions of slender-body theory show (i) steady ellipsoidal bubbles covered with surfactant, and (ii), at sufficiently large deformation, bubbles with a cylindrical surfactant-free central part and ellipsoidal surfactant-covered end caps. These bubble shapes are rounded at their end-points, in contrast to the non-rounded shapes found for entirely surfactant-free bubbles.

Simple expressions are found that relate the capillary number to the bubble slenderness ratio, and these show a critical capillary number above which steady solutions cease to exist. Beyond the critical capillary number, slender-body model equations show time-dependent spindle shapes at the bubble end-points with tip-streaming filaments. Joint work with M.R. Booty (NJIT).

Linda Smolka

Bucknell University, Department of Mathematics, Lewisburg, PA 17837

Getting Your Hands Wet in Mathematics

In recent years there has been increasing interest and support for physical labs in mathematics departments. The newest of these labs is being built at Bucknell University. Bucknell is a liberal arts college of 3,350 undergraduate and 150 graduate students. Bucknell is distinct among liberal arts colleges in that it has several professional programs including an engineering school.

I'll explain the role and goals of the lab at Bucknell within the context of the mathematics department, the university, and my own research program. I'll also discuss the process of designing a lab; including issues of funding, renovating, and selecting/purchasing equipment for the lab.

Seongho Song

University of Connecticut, Department of Statistics, 215 Glenbrook Rd. U-4120, Storrs, CT 06269-4120

Genetic Diversity of Microsatellite Loci in Hierarchically Structured Populations

Patterns of genetic variation within and among populations are determined by population sizes, migration rates, and mutation rates. We provide exact expressions for the first two moments of a stochastic model with hierarchically structured migration, mutation, and drift. Specifically, we consider the symmetric stepwise mutation model for microsatellite evolution under the assumption that the number of repeats is finite in Wright's finite island model. Further, we suggest two measures, $\theta^{(b)}$ and $\theta^{(m)}$, as an analogue of Wright's F -Statistics and compared them to R_{ST} as discussed in Slatkin(1995). As a result, θ 's and R_{ST} have the same properties under some conditions. Through numerical study we observe that R_{ST} measure is quite robust for the stepwise mutation model and the two-phase mutation model (Di Rienzo et al, 1994).

Keywords: F -statistics, Finite Island Model, Genetic Drift, Hierarchical Population Structure, Microsatellite Loci, Migration, Mutation, Singular Value Decomposition, Stepwise Mutation Model. Joint work with Dipak K. Dey and Kent E. Holsinger (University of Connecticut).

David Stickler

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Layered Media

Ray theoretic methods are very useful in determining high frequency solutions to the scalar Helmholtz wave equation. This poster concerns the solution of the eiconal equation for an almost stratified medium. For a stratified medium the index of refraction depends only on Z , i.e., $n = n(Z)$. For an almost stratified medium $n = n(Z - E h(xy))$ where E is a small parameter and $h(xy)$ are given. With minor assumptions on $n(Z)$ and $h(xy)$, a solution to the eiconal equation to order E^2 is obtained which is uniform in E , i.e., it contains no secular terms.

Louis Tao

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Visual Cortical Orientation Selectivity by Fluctuation-Controlled Criticality

We examine how synaptic fluctuations modify the effects of strong recurrent network amplification to produce orientation selectivity in a large-scale neuronal network model of the macaque primary visual cortex. Typically, strong cortical amplification leads to network instabilities and unrealistically high firing rates even in the presence of strong cortical inhibition. In this poster, we show that large fluctuations in the cortico-cortical conductances can stabilize the network, allowing strong cortical gain and the emergence of orientation selective neurons. By increasing the strength of synaptic fluctuations, say, through sparsifying the network connectivity, we identify a transition between two types of dynamics, mean- and fluctuation-driven. In a network with strong recurrent excitation, this fluctuation-controlled transition is signified by a near hysteretic behavior and a rapid rise of network firing rates as the synaptic drive or stimulus input is increased. Finally, we demonstrate that network sparsity leads naturally to a recently observed invariance of orientation selectivity across the cortical network, even in the presence of orientation hypercolumns.

Burt S. Tilley

Franklin W. Olin College of Engineering, Department of Mathematics, 1000 Olin Way, Needham, MA 02492-1200

Electrokinetic Instabilities in Thin Microchannels

An important class of electrokinetic, microfluidic devices aims to pump and control electrolyte working liquids that have spatial gradients in conductivity. These high-gradient flows can become unstable under the application of a sufficiently strong electric field. In many of these designs, flow channels are thin in the direction orthogonal to the main flow and the conductivity gradient. Viscous stresses due to the presence of these walls introduce a stabilizing force that plays a major role in determining the overall instability. A thin channel model for fluid flow is developed and shown to provide good agreement with a complete three-dimensional model for channel aspect ratio less than approximately one-tenth. Joint work with Brian D. Storey (Franklin W. Olin College of Engineering) and Hao Lin and Juan G. Santiago (Stanford University).

Natalia Toporikova

Florida State University, Department of Biological Science and Mathematics and Institute of Molecular Biophysics, Tallahassee, FL 32306

Mathematical Model of Neuronal Network Regulating Prolactin Release in Rats

The mating stimulus of the rat uterine cervix (CS) induces two daily surges of the anterior pituitary hormone prolactin (PRL), which lasts for 12 days. PRL secretion occurs in response to relief from hypothalamic dopaminergic (DA) inhibition and stimulation by releasing oxytocin (OT). We have proposed a mathematical model to explain the origin of rhythmic PRL secretion. The model consists of a system of delay differential equations (DDE) and includes variables for the activity level of OT, PRL and DA cell populations. The oscillatory mechanism in the model is the mutual interaction between DA neurons and lactotrophs. Prior to CS, the system is in a stable state with a low basal level of PRL and a high level of DA. Following CS or injection of OT, PRL and the DA levels start to oscillate. The rhythm termination factors (PL or some other uterus factors) are modeled as an increasing function of time with a positive connection to the DA neurons. After 12 days this factor suppresses the oscillation by stimulation DA neurons and the system returns to the steady state. Joint work with M. Egli (Florida State University and Space Biology Group, ETH Zurich, Switzerland), and R. Bertram and M. E. Freeman (Florida State University).

A. David Trubatch

United States Military Academy, Department of Mathematical Sciences, West Point, NY 10996

Soliton Dynamics in the Integrable Discrete Vector Nonlinear Schrödinger Equation

The fundamental solitons of the integrable discrete vector nonlinear Schrödinger equation (IDVNLS) retain many of the interesting properties of their continuous counterpart (i.e., the solitons of Vector NLS, also known as the Manakov equation). In particular, the collision-induced phase shifts of the vector solitons can be described by fractional linear transformations and constitute a solution of the set-theoretical Yang-Baxter equation. Moreover, as in the continuous case, logical operations can be encoded in the vector-soliton interaction such that it is possible to carry out general computations with the discrete vector solitons. On the other hand, the discrete system contains composite soliton states that have no counterpart in the Manakov system. These traveling breathers are related to a class of traveling breathers of the scalar integrable discrete NLS (also known as the Ablowitz-Ladik equation). However, unlike their scalar, discrete counterparts the composite solitons have a minimal nonlinear spectrum as solutions of the vector system.

Dmitri Tseluiko

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Nonlinear Stability of the Solutions of Modified Kuramoto-Sivashinsky Equations

We study equations that arise in the modelling of the wave motion in a perfectly conducting viscous thin film falling down an inclined plane under gravity in the presence of an electric field which is uniform in its undisturbed state, and normal to the plate at infinity. These are modified Kuramoto-Sivashinsky equations with an additional non-local term due to the Maxwell stresses exerted at the interface by the electric field. The numerical results show that the solutions of these equations are nonlinearly stable

and exhibit a complicated behavior including chaotic oscillations as in the case of the usual Kuramoto-Sivashinsky equation; the basins of attraction of chaotic dynamics is significantly affected by the presence of the electric field, however, and this is seen as significantly larger windows in phase space where the global attracting solutions exhibit complicated dynamics. The nonlinear stability of the solutions is also proved analytically. The proof also leads to an upper bound estimate of the 2-norm of the solution in terms of the length of the system and the electric field intensity parameter. Joint work with Demetrios T. Papageorgiou (NJIT).

A. Kerem Uguz

University of Florida, Chemical Engineering Department, Gainesville, FL 32611

An Experimental Study on the Stability of Elliptical Liquid Bridges

The break-up point of an elliptical liquid bridge was investigated in a Plateau chamber using two density-matched liquids. The objective was to show that an elliptical liquid bridge is more stable than a companion circular bridge where the elliptical end plates are slight deviations from the circular end plates. The semi-major axis of the ellipse was roughly 20% larger than the radius of the circle yet maintaining the areas the same for both geometries. The volume in the elliptical liquid bridge was fixed to be the same as the volume of the companion cylindrical bridge at its critical point and the end plates were parallel and oriented so that they were not twisted with respect to one another. In this study, two different sizes of circular end plates were used to confirm our results. The distortion amount, which was roughly 20%, was kept constant for both cases and elliptical liquid bridge experiments were performed. It was found that the elliptical liquid bridge break-up point is about 3% longer than its corresponding cylindrical liquid bridge demonstrating that an elliptical liquid bridge is more stable than its companion circular liquid bridge. A theory showing that the elliptical end plates do not break the classical bifurcation is presented. This supports the view that the elliptical liquid bridge ought to be more stable than its circular counterpart. Joint work with N. J. Alvarez and R. Narayanan (University of Florida).

A. Kerem Uguz

University of Florida, Chemical Engineering Department, Gainesville, FL 32611

Rayleigh-Taylor Instability with Shear-Induced Flow

It is known in the Rayleigh Taylor problem that there is a decrease in stability when the liquid is sheared with a constant stress. This decrease in the stability limit has been explained with the symmetry breaking effect of the shear. In this study, we prove that the fluid mechanics of the light fluid (vapor or liquid) is important and it changes the characteristics of the problem. Shear induced Rayleigh-Taylor instability in an open-channel flow and in a closed container is studied in this paper. For both cases, at the base state, we satisfied the conditions for a flat interface between the two liquids and studied the stability of this base state to small disturbances via linear stability analysis. In the open channel flow, the critical point remained unchanged compared to the classical Rayleigh-Taylor instability, but the critical point exhibits oscillations and the frequency of the oscillations depends on the wall speed. On the other hand, in a closed geometry, moving the wall stabilizes the classical Rayleigh-Taylor instability and it does not show any oscillation at the onset of break-up. We also applied weakly nonlinear analysis to study the nature of the bifurcation via a dominant balance method and concluded that the problem shows a backward pitchfork bifurcation, as does the classical Rayleigh-Taylor instability problem. Joint work with Ranga Narayanan (University of Florida).

Suleyman Ulusoy

Georgia Institute of Technology, School of Mathematics, Atlanta, GA 30332-0160

An Entropy Dissipation-Entropy Estimate for a Thin Film Type of Equation

We prove a lower bound on the rate of relaxation to equilibrium in the H^1 norm for a thin film equation. We find a two stage relaxation, with power law decay in an initial interval, followed by exponential decay, at an essentially optimal rate, for large times. The waiting time until the exponential decay sets in is explicitly estimated. Joint work with E.A. Carlen. Keywords: thin film, Lyapunov functional, entropy dissipation.

Vianey Villamizar

Brigham Young University, Mathematics Department, Provo, UT 84602

Grid Generation with Curve Control and Its Application in Acoustic Scattering

A new algorithm to automatically generate two-dimensional boundary conforming coordinates, with grid curve control, on multiply connected regions, is devised. The technique is based on the numerical solution of the widely used Poisson grid generation equations. The physical domain D is transformed into a topologically equivalent connected rectangular domain D' by defining a branch cut inside D . Grid curve control over the multiply connected regions with one or more interior holes is established by appropriately distributing grid points with the desired spacing on the branch cut, and on other boundary curves. From this initial distribution of points, control functions present in the Poisson system are defined, creating a natural link between clustering properties and the control functions. An iterative smoothing process relocating the branch cut is described. As a result, a smooth grid retaining the spacing of the non-smooth initial grid is obtained.

Tested hole shapes include a rose, an epicycloid, an astroid, and a pacman. These grids are used to numerically solve acoustic scattering problems. Approximations of the pressure field and the scattered cross section for arbitrary shape obstacles are obtained. Joint work with Joseph Mabey (Brigham Young University).

X. Sheldon Wang

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

From Immersed Boundary Method to Immersed Continuum Method

In this talk, a new modeling method, the immersed continuum method (ICM), is proposed for the coupling of different continua (fluid-fluid, fluid-solid, or solid-solid). This new method retains the same strategies employed in the immersed boundary (IB) method [1] and the recent extensions, namely, the extended immersed boundary method (EIBM) [2] and the immersed finite element method (IFEM) [3]. The immersed continuum method deals with submerged continua (compressible or incompressible) which occupy finite volumes within the surrounding fluid (or solid) medium. In the context of fluid-solid interactions [4] [5], like EIBM and IFEM, in ICM, an independent solid mesh moves on top of a fixed or prescribed arbitrary Lagrangian-Eulerian (ALE) background fluid mesh. The procedure of constructing an entire fluid domain over both fluid and solid domains is similar to that of the fictitious domain method [6]. However, unlike the fictitious domain method which is designed for immersed rigid bodies; ICM handles arbitrary immersed deformable solids/structures. Furthermore, in order to handle submerged compressible continua and to circumvent the stringent requirement for the time step size as demanded in the explicit time marching schemes in the IB method, EIBM, and IFEM, we introduce in ICM an implicit formulation along with the combination of the Newton-Raphson and GMRES iterations. The new attributes in ICM enable a wide variety of applications in aerospace, mechanical, civil, and in particular bio-engineering fields. Several numerical examples in micro- and capillary vessel hemodynamics are also presented for illustrative purposes.

Reference:

- [1] C.S. Peskin, The immersed boundary method, *Acta Numerica*, 11: 479–517, 2002.
- [2] X. Wang and W.K. Liu, Extended immersed boundary method using FEM and RKPM, *Computer Methods in Applied Mechanics and Engineering*, 193:1305–1321, 2004.
- [3] L. Zhang, A. Gerstenberger, X. Wang, and W.K. Liu, Immersed finite element method, *Computer Methods in Applied Mechanics and Engineering*, 193:2051–2067, 2004.
- [4] T.E. Tezduyar, Finite element methods for flow problems with moving boundaries and interfaces, *Archives of Computational Methods in Engineering*, 8(2):83–130, 2001.
- [5] K.J. Bathe and H. Zhang, "Finite element developments for general fluid flows with structural interactions", *International Journal for Numerical Methods in Engineering*, 60: 213–232, 2004.
- [6] R. Glowinski, T.W. Pan, T.I. Hesla, D.D. Joseph, and J. Periaux, A fictitious domain approach to the direct numerical simulation of incompressible viscous flow past moving rigid bodies: Application to particulate flow, *Journal of Computational Physics*, 169:363–426, 2001.

Xinli Wang

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Influence of Surfactant on Air Entrainment at a Contact Line

We present a model for the rolling motion of a viscous fluid onto a rigid substrate, in the case when insoluble surfactant is present at the fluid interface. The model is used to examine the influence of surfactant on air entrainment at the line where the fluid contacts the wall. In the absence of surfactant, an analytical solution has been obtained by Benney and Timson (*Stud. Appl. Math* 1980) for the local flow field near the point of steady attachment. They find that a steady local solution exists for all capillary numbers. In the presence of surfactant, we find that there is a critical capillary above which steady shapes no longer exist. This suggests that the presence of surfactant can lead to air entrainment for large enough capillary number. Joint work with Michael Siegel (NJIT).

Wonsuk Yoo

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Model Selection Method using Reversible Jump MCMC Algorithm for Longitudinal Biomarkers

Markov chain Monte Carlo (MCMC) methods provide a very effective sampling base by generating from extremely complex distributions but they can be used only within fixed dimensions. We consider a model within a fully Bayesian framework which postulates a mixture for longitudinal data, for which one component contains a change point and the other does not. While MCMC methods cannot be used in this trans-dimensional situation which needs to move between different states or dimensions, the reversible jump Markov chain Monte Carlo (Green 1995) allows a Markov transition between models with parameter spaces of different dimensions, but holds the aperiodicity, irreducibility, and detailed balance conditions necessary for MCMC convergence. It is particularly useful in the model selection procedure within the Bayesian context. This paper focuses developing the algorithms of the reversible jump MCMC, and evaluating them on the basis of the early detection. This paper can be valuable in Bayesian model determination using reversible jump method since the algorithms are proposed based on usage of various auxiliary variables for dimension-matching. The results from these proposed algorithms can be more reasonable because they allow various

and realistic situation on the relationship between parameters of both models. We provide following steps: birth, death, update and move steps on each iteration. We apply this reversible jump algorithm to Nutritional prevention of Cancer Trial (Clark 1996).

Yun Yoo

Drexel University, Department of Mathematics, Korman Center 252, 3141 Chestnut Street, Philadelphia, PA 19104

Periodic Orbits Near a Saddle-Focus in Systems with Strong Contraction

We study a one-parameter family of three-dimensional flows near an Andronov-Hopf bifurcation (AHB). We specify conditions on the global vector field that guarantee the existence of a rich family of multimodal periodic orbits passing close to a saddle-focus. We identify two bifurcation scenarios for periodic orbits depending on the character of the AHB and analyze the properties of the periodic orbits using asymptotic techniques for ordinary differential equations. This work is motivated by the numerical results for a finite-dimensional approximation of a free boundary problem modeling solid combustion.

Yuan-Nan Young

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Effect of Capillary Network Hemodynamics on Tumor Growth

A two-dimensional model of tumor necrosis, neo-vascularization and tissue invasion was recently developed by Zheng, Wise and Cristini. This model describes the angiogenesis and vascular nutrient delivery that occur during tumor growth; however, blood flow and other physiological details of transport in the capillary network are not included. To improve this model, a two-phase (red cells and plasma) continuum model of blood flow is used to estimate the local capacity for nutrient delivery within the network. This model includes empirical relations describing known nonlinear effects (Fahraeus, Fahraeus-Lindqvist, and plasma skimming) and allows calculation of the steady-state blood flow distribution within the network. Using calculated capillary network blood flows to modulate nutrient delivery, it is shown how tumor growth and steady-state patterns differ from those obtained assuming constant nutrient delivery throughout the network. Currently, blood flows are calculated only at selected times during tumor growth, however in the future the blood flow pattern will be updated simultaneously with tumor evolution. This is a joint collaboration with Daniel Goldman (NJIT), and V. Cristini and X. Zhang (UC Irvine).

Yuan-Nan Young

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Effect of Surfactant on Bubble Deformation

The effect of surfactant on bubble deformation is investigated using direct numerical simulations. In particular we focus on the formation of long, thin thread due to the surfactant effect, and investigate the possibility of local similarity solution that leads to pinch-off of the thin thread in the presence of insoluble surfactant. In the second part of this investigation we include soluble surfactant and report how the formation of the thin thread can be altered. This is a joint work with Jie Li (Cambridge University), and Michael Siegel and Demetrios T. Papageorgiou (NJIT).

Hui Zhang

SUNY at Stony Brook, Department of Mechanical Engineering, Stony Brook, NY 11794-2300

Adaptive Level Set Method for Droplet Dynamics and Solidification in Sprayed Coating Applications

Sprayed coatings are formed by injecting a powder feedstock through a thermal plasma or combustion flame, where the powder is melted. The resulting molten droplets are accelerated to high velocity and impact on a substrate. The droplets spread into disk-like pancake shapes called splats and rapidly solidified. Agglomerating splats form a thick film coating with a distinctive lamellar microstructure, characterized by the splat interfaces or interphases. In this presentation, current status of thermal spray research will be reviewed. An advanced numerical model based on the adaptive level set method is developed that is capable of solving the deformation of the free surface, fluid instability, and evolution of solidification interface simultaneously. In the new model conservation of mass is guaranteed and the solution domain can be adaptively deformed depending on the solution process. Theoretical and numerical analyses are performed to predict the splat-flattening ratio and related the splat geometry with the droplet Reynolds, Weber, Jakob numbers and substrate conditions. The numerical predictions are in good agreement with the experimental data. Through studies, good understanding has been achieved on the single splat formation and the roles of kinetic, dissipation and surface energies on the droplet spreading and solidification process. Numerical and experimental studies are also performed to study the splashing phenomena. Different splashing mechanisms are proposed and confirmed with the experimental data. The fragmentation degree is related to surface wetting, rapid solidification, substrate roughness and surface chemistry. The ability to control and manipulate the splat formation will significantly advance the spray coating technology.

Kai Zhang

TIMDA: A Toolkit for Integrated Genotyping Microarray Data Analysis

Background: TIMDA (Toolkit for Integrated Genotyping Microarray Data Analysis) is designed for integrated data analysis for spotted single nucleotide polymorphism (SNP) genotyping microarrays.

Results: We have implemented TIMDA using Matlab, a powerful and popular engineering computing programming language which integrates seamlessly the mathematical computation, analysis, visualization and algorithm development. The framework of TIMDA consists of several function modules for microarray image processing which implements precise gridding and robust segmentation techniques, data preprocessing, genotype-calling which implements novel machine learning-based algorithms and Loss-of-Heterozygosity which is used to study the genetic basis of certain diseases, Graphical User Interface and text and graphical outputs. Each individual module works seamlessly with each other. Some modules, such as the image-processing module, can also be singled out to work on their own. A user can also bypass some modules by supplying with information generated from other software, which makes TIMDA a flexible framework.

Conclusions: We have done extensive testing of TIMDA, which shows that TIMDA is a powerful framework for performing robust data analysis in genotyping microarrays.

Joint work with Li Jia, Marc Ma, Frank Shih, Yu Wang, and I-Jen Yeh (NJIT), Honghua Li and Hui-Yun Wang (UMDNJ, Robert Wood Johnson Medical School), Tongsheng Wang (NJIT and Public Health Research Institute), and Patricia Soteropoulous (Public Health Research Institute).

Yili Zhang

Rutgers University, Biological Sciences Department, Newark, NJ 07102

Modeling Recovery of Rhythmic Activity: Hypothesis for the Role of a Calcium Pump

The rhythmic activity produced by the pyloric network of crustaceans depends on the release of neuromodulatory substances by axon terminals from adjacent ganglia. After these terminals are destroyed or action potential transmission along these axons is inhibited (decentralization), the rhythmic pyloric activity recovers spontaneously, but the process of activity recovery follows a very complex temporal dynamics that involves the alternating turning on and off of the pyloric rhythm (that we term 'bouts'). This bouting period lasts several hours after which a stable pyloric rhythm emerges.

We have built a model of the activity of a pyloric neuron to study the recovery of rhythmic activity after decentralization. Our model is based on a dynamic oscillation of intracellular Ca^{2+} due to the interaction of Ca^{2+} influx, Ca^{2+} diffusion, Ca^{2+} pump activity, and IP3 receptor activity at the endoplasmic reticulum (ER). The model assumes that activity is monitored by sensors of intracellular Ca^{2+} concentration changes. The three Ca^{2+} sensors represent biochemical pathways sensitive tuned to different frequencies of Ca^{2+} change. They act as feedback regulators of Ca^{2+} and K^{+} conductances, and of Ca^{2+} pump activity. Our model reproduces qualitatively accurately the dynamics of recovery after decentralization, particularly the transition to a stable rhythm after a period of bouting. We observe that the regulation of the activity of the ER Ca^{2+} pump is key in the generation of this rich dynamics. Supported by NIH grant MH-64711 (J.G.). Joint work with Jorge Golowasch (NJIT and Rutgers University).

Yongmin Zhang

SUNY at Stony Brook, Applied Mathematics and Statistics, Stony Brook, NY 11794

Modeling and Simulation of Fluid Mixing for Laser Experiments and Supernova

Recently, laboratory astrophysics has been playing an important role in the study of astrophysical systems, especially in the case of supernova explosions through the creation of scaled reproductions of astrophysical systems in the laboratory. In collaboration with a team centered at U. Michigan and LLNL, we have conducted front tracking simulations for axisymmetrically perturbed spherical explosions relevant to supernovae as performed on NOVA laser experiments, with excellent agreement with experiments [1]. We have extended the algorithm and its physical basis for preshock interface evolution due to radiation preheat [2]. The preheat simulations motivate direct experimental measurements of preheat as part of any complete study of shock-driven instabilities by such experimental methods.

Our second focus is to study turbulent combustion in a type Ia supernova (SN Ia) which is driven by Rayleigh-Taylor mixing. We have extended our front tracking to allow modeling of a reactive front in SN Ia. Our 2d axisymmetric simulations show a successful level of burning [3]. Our front model contains no adjustable parameters so that variations of the explosion outcome can be linked directly to changes in the initial conditions.

References:

- [1] R. P. Drake, H. F. Robey, O. A. Hurricane, Y. Zhang, B.A. Remington, J. Knauer, J. Glimm, D. Arnett, J. O. Kane, K. S. Budil, and J. W. Grove. Experiments to produce a hydrodynamically unstable spherically diverging system of relevance to instabilities in supernovae. *Astrophysical Journal*, 564:896-906, 2002.
- [2] Y. Zhang, P. Drake, J. Glimm, J. Grove, and D. H. Sharp. Radiation coupled front tracking simulations for laser driven shock experiments. *J Nonlinear Analysis*, 2005. In press. LANL report No. LA-UR-04-2381.
- [3] Y. Zhang, J. Glimm, and S. Dutta. Tracked flame simulation for Type Ia supernova. In *Proceeding of Third MIT Conference*. Elsevier, 2005. accepted.

Joint work with James Glimm, Paul Drake, Srabasti Dutta, John W. Grove, and David H. Sharp (State University of New York, Stony Brook).

Lin Zhou

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Complete Transmission Through a Periodically Perforated Solid Slab

The propagation of a normal incident plane acoustic wave through a 3-dimensional rigid solid slab with periodically placed holes is modeled and analyzed. The period of the structure S , the wavelength and the thickness of the slab L are order one parameters compared to the characteristic size of the holes R , which is a small quantity. Scattering matrix techniques are used to derive expressions for the transmission and reflection coefficients of the lowest mode. These expressions depend on only the transmission coefficient of an infinitely long slab with the same configuration. An infinite system of algebraic equations of the transmission coefficients is derived and solved approximately by exploiting the small parameter R/S . It turns out that the structure is transparent at certain frequencies which could prove useful in narrow band filters and resonators. Joint work with Gregory A. Kriegsmann (NJIT).

Ivan Zorych

NJ Institute of Technology, Mathematical Sciences Department, University Heights, Newark, NJ 07102

Bayesian Models for Location Estimation in Wireless Networks

The Bayesian modeling approach is used to study location problems in wireless networks using Markov Chain Monte Carlo methods in DAGs (directed acyclic graphs). Nonhierarchical and hierarchical Bayesian models are investigated, as well as penalized Bayesian splines and bivariate splines in combination with a particle approach. Two different data sets are used to illustrate proposed models. (Joint work with D. Madigan, Rutgers University.)