

# DETOXIFICATION OF CHEMICAL WARFARE AGENTS BY THE PLANT CHOLINERGIC SYSTEM

Suman S. Thakur, Gregory E. Garcia, Haim N. Leader, Deborah Moorad-Doctor, Rajendra Gupta\*, Richard K. Gordon, and Bhupendra P. Doctor

Walter Reed Army Institute of Research, Division of Biochemistry,  
Department of Biochemical Pharmacology Silver Spring, MD 20910-7500, USA  
and \*University of Delhi, Department of Botany, Delhi, India

## ABSTRACT

Plants have cholinesterases (pChEs), anti-ChEs, and activators of ChEs. We have isolated pChE from mung bean sprout. We investigated 300 plants and found that 75% of them contained anti-ChE that inhibited ChEs. Thirty-five percent contained activators of ChEs. An activator of ChE from wheat leaf, “Tritiacche-T123”, activates fetal bovine serum AChE, equine BChE, and pChE. This non-oxime natural plant product may offer a new approach to the reactivation of OP-inhibited ChEs. pChE showed substrate differences from human acetylcholinesterase (AChE) or butyrylcholinesterase, hydrolyzing propionylthiocholine > acetylthiocholine > butyrylthiocholine. pChEs might also be useful as bioscavengers of OP nerve agents.

## INTRODUCTION

Organophosphate chemical warfare agents (CWA) and pesticides are extremely poisonous compounds. Nerve agents irreversibly inhibit acetylcholinesterase (AChE) activity at synapses, stop neuronal signal transmission, and destroy the cholinergic system machinery and can ultimately cause death. Reactivation of inhibited AChE by traditional oxime reactivators (for example 2-PAM, LÜH6, TMB4, and HI-6,) is an efficient way to attenuate toxicity and thus plays a key role in the treatment of nerve agent poisoning. However, their efficacies can be limited due to either poor or non-reactivation of the CWA-inhibited cholinesterases (ChEs), most notably in the case where ChEs have been exposed to soman (GD) because it “ages” rapidly. New activator/reactivators<sup>1</sup> could be useful to overcome these difficulties.

Plants also contain anti-ChEs, for example physostigmine and huperzine A<sup>1,2</sup>. Pyridostigmine is a synthetic analogue of physostigmine<sup>3</sup>, a natural product from the plant *Physostigma venenosum*, and is used therapeutically for a variety of diseases including myasthenia gravis.

Among the enzymes being considered and evaluated as bioscavengers of organophosphate (OP) nerve agents are the cholinesterases (ChEs), specifically human serum butyrylcholinesterase<sup>4</sup>, the OP hydrolases (bacterial sources), and carboxylesterase. An effort has been made to find new ChE candidates as bioscavenger. Interestingly, plants have been found to have cholinesterase(s)<sup>5</sup>. Therefore we have isolated and partially purified a plant ChE from mung bean sprouts. These biological compounds from plants may be useful as bioscavengers and protectants of OPs, and activators of OP-poisoned enzymes.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>01 OCT 2005</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Detoxification Of Chemical Warfare Agents By The Plant Cholinergic System</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Walter Reed Army Institute of Research, Division of Biochemistry, Department of Biochemical Pharmacology Silver Spring, MD 20910-7500, USA</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM001851, Proceedings of the 2003 Joint Service Scientific Conference on Chemical &amp; Biological Defense Research, 17-20 November 2003. , The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## METHODS

Isolation and Characterization of plant activator(s): Wheat (*Triticum aestivum*) leaves were grown in laboratory. The methanol extract of wheat leaves was analyzed for its various chemical constituents by thin layer chromatography (TLC), normal phase column chromatography (silica), and HPLC. TLC plates (20 x 20 cm), coated with silica gel G (0.25 mm thickness), were pre-dried and then loaded with the methanolic solution. The plates were developed in a variety of solvent systems. Fractions were analyzed for their ability to increase (activate) AChE activity in the Ellman assay<sup>6</sup>. The band that activated AChE the most had an R<sub>f</sub> of 0.53 in chloroform: methanol (1:4) system. This material exhibited bright blue fluorescence under UV illumination. The activity was concentrated under vacuum, and then re-chromatographed on a silica gel column in water: methanol (1:3). The active fraction (R<sub>f</sub> 0.94) was once again re-re chromatographed on the silica column in water: methanol (1:3). These fractions are being evaluated for purity and structural elucidation by HPLC, NMR, and MS.

Purification of plant cholinesterase (pChE): pChE was extracted from the roots of mung bean sprouts (*Phaseolus aureus*). The roots were crushed and homogenized in 0.1 M sodium phosphate buffer pH 8; 2:1, (v/w), and extracted by stirring for 1 h in 1% Triton X and 1 M NaCl. The remaining insoluble material was removed by filtration through a nylon mesh. The filtrate was centrifuged at 15,000 x g for 30 min at 4 C. The resulting supernatant was dialyzed against 5 mM sodium phosphate buffer pH 8. The protein fraction was concentrated by using polyethylene glycol. Next, the enzyme was applied to a procainamide affinity gel column for purification. Active fractions (1 ml each) were collected, and pooled prior to use.

## RESULTS AND DISCUSSION

ChEs inhibited by certain OPs can become refractory to spontaneous or oxime induced reactivation because the enzyme-OP has undergone dealkylation or “aging”. Alternatively, some enzyme: OP respond very slowly to oxime reactivation. For these reasons, there has been a renewed interest in novel oxime or non-oxime reactivators for nerve agent poisoning.

Anti-cholinesterase compounds (anti-ChEs) are widespread in plants as shown in Table 1. On the other hand, plants were also found to be a rich source of compounds that can act as ChE activators as shown in Table 2. Purified plant derived activators may be of value as a treatment against organophosphate poisoned ChEs.

Plant Activators of Cholinesterase: We found that an extract from wheat leaves, such as “Tritiacche-T123” and other compounds from plants activated fetal bovine serum AChE up to three fold (Fig.1), and also activated equine BChE and pChE cholinesterases. Two novel partially purified compounds that activate eel AChE were isolated from wheat leaves, Triticheac<sup>1</sup> and Whecheac<sup>1</sup>. A third activator of ChE from wheat leaf, “Tritiacche-T123” is three-fold more potent than Triticheac. These non-oxime natural plant products may offer a new approach to the reactivation of OP-inhibited ChEs. The mechanism by which Tritiacche-T123 activated purified fetal bovine serum AChE three-fold is currently being investigated.

<b>Table 1. Survey of plants for anti-ChE Activity</b>				
<b>Division</b>	<b>Families</b>	<b>Families tested positive</b>	<b>Species</b>	<b>Species tested positive</b>
<b>Algae</b>	<b>8</b>	<b>6</b>	<b>19</b>	<b>10</b>
<b>Fungi</b>	<b>4</b>	<b>4</b>	<b>13</b>	<b>12</b>
<b>Lichens</b>	<b>4</b>	<b>4</b>	<b>6</b>	<b>6</b>
<b>Bryophytes</b>	<b>11</b>	<b>3</b>	<b>18</b>	<b>4</b>
<b>Pteridophytes</b>	<b>12</b>	<b>12</b>	<b>38</b>	<b>35</b>
<b>Gymnosperms</b>	<b>10</b>	<b>10</b>	<b>18</b>	<b>16</b>
<b>Angiosperms</b>	<b>112</b>	<b>86</b>	<b>191</b>	<b>145</b>
<b>Total</b>	<b>161</b>	<b>125</b>	<b>303</b>	<b>228</b>

<b>Table 2. Survey of plants for activator(s) of AChE [Acti-ChE]</b>				
<b>Division</b>	<b>Families</b>	<b>Families tested positive</b>	<b>Species</b>	<b>Species tested positive</b>
<b>Algae</b>	<b>8</b>	<b>7</b>	<b>19</b>	<b>9</b>
<b>Fungi</b>	<b>4</b>	<b>1</b>	<b>13</b>	<b>3</b>
<b>Lichens</b>	<b>4</b>	<b>0</b>	<b>6</b>	<b>0</b>
<b>Bryophytes</b>	<b>11</b>	<b>10</b>	<b>18</b>	<b>14</b>
<b>Pteridophytes</b>	<b>12</b>	<b>6</b>	<b>38</b>	<b>9</b>
<b>Gymnosperms</b>	<b>10</b>	<b>3</b>	<b>18</b>	<b>2</b>
<b>Angiosperms</b>	<b>112</b>	<b>38</b>	<b>191</b>	<b>69</b>
<b>Total</b>	<b>161</b>	<b>65</b>	<b>303</b>	<b>106</b>

Plant cholinesterase: Interestingly, plants have been found to contain ChEs, and these enzymes have been proposed as a bioscavenger for OPs<sup>4</sup>. We have examined ChE activity derived from mung bean roots. This partially purified pChE showed substrate specificity different than mammalian acetylcholinesterase, hydrolyzing thiocholine substrates in the following order: propionylthiocholine > acetylthiocholine > butyrylthiocholine (Fig. 2). Based on acryl amide gel electrophoresis analysis, the pChE appears to be a dimer (~120 kD).

These natural pChEs might be useful for immobilizing on polyurethane sponge used for decontamination of skin and wounds and as a pretreatment for OP poisoning, if it is compatible with human system.





Fig. 1. Effect of Tritiacche-T123 compound on FBS AChE

**Fig. 2. Plant ChE (pChE, from mung bean)**  
**Hydrolysis of ATC, BTC, & PTC and protein concentration after**  
**procainamide column affinity purification**



## CONCLUSION

- Plants have cholinesterase and anti-cholinesterase and novel activator(s) of ChEs.
- A partially purified compound from wheat, “Tritiacche-T123”, activated fetal bovine serum AChE up to three-fold.
- These non-oxime natural plant AChE activating products may offer a new approach to the activation/reactivation of OP-inhibited ChEs.
- Based on acryl amide gel electrophoresis analysis, the pChE appears to be a dimer (~120 kD).
- This partially purified pChE showed substrate specificity different than mammalian acetylcholinesterase, hydrolyzing thiocholine substrates in the following order: propionylthiocholine > acetylthiocholine > butyrylthiocholine.
- pChE may be useful for the decontamination of OPs and as a bioscavengers for OPs, thereby preventing or decreasing OP poisoning.

## REFERENCES

1. **Thakur, S.S.** 2001. Studies on Cholinesterases and Anticholinesterases in Plants, and the First Isolation, Purification and Characterization of Naturally Occurring Activators of Acetylcholinesterase: Triticheac and Whecheac. Ph.D Thesis, University of Delhi, Delhi, India.
2. **Taylor, P.** 2001. Anticholinesterase Agents. In: Goodman and Gilman's, *The Pharmacological Basis of Therapeutics*, J.G. Hardman and L.E. Limbird, eds., Tenth Edition, pp. 175-192.
3. FDA News, PO3-08 February 5, 2003 <http://www.fda.gov/bbs/topics/NEWS/2003/NEW00870.html>
4. **Doctor, B. P.** 2003. Butyrylcholinesterase: its use for prophylaxis of organophosphate exposure. In: Giacobini, E (ed.). Butyrylcholinesterase: its function and inhibitors. Pp. 163-177. Martin Dunitz, UK.
5. **Miura, G.A.**, Broomfield, C.A., Lawson, M.A. and Worthley, E.G. 1982. Widespread occurrence of cholinesterase activity in plant leaves. *Physiol. Plant.* **56**: 28-32.
6. **Ellman, G.L.**, Courtney, K.D., Andres, V., Jr. and Featherstone, R.M. 1961. A new rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **7**: 88-95.