

Cold Injury of Amputated Digits

Uros Ahcan¹, Bostjan Luzar², Fajko Bajrovic² & Igor Mekjavic³

¹Department of Plastic Surgery and Burns, Clinical Centre Ljubljana, Zaloska 7, SI-1000 Ljubljana;

²Institute of Pathophysiology, Faculty of Medicine, University of Ljubljana, SI-1000 Ljubljana;

³Department of Automation, Biocybernetics and Robotics, Jozef Stefan Institute, Jamova 39, SI-1000, Ljubljana, Slovenia.

Contact person: uros.ahcan@guest.arnes.si

ABSTRACT

Cooling of amputated parts during transportation delays the onset of ischaemic tissue damage and contributes to successful replantation. The most common error in preservation of amputated parts is exposing them to temperatures, which may cause cold injury, and thus render them unusable. Two case reports will be presented, which illustrate how freezing cold injury (FCI) of amputated digits may decrease the viability of such digits following replantation, and discuss the potential benefit of hyperbaric oxygen therapy (HBOT) in treating FCI of amputated digits following replantation. During the period 1998 to 2002, 124 injured individuals with severed digits were admitted to the Department of Plastic Surgery and Burns at the Ljubljana Clinical Centre. During this period, the number of attempted replantations increased from 29% in 1998 to 48% in 2002. The success rate of these replantations was 81%. Cold injury to the amputated parts was a significant factor affecting the replantation success rate in six patients. In all these cases, all fingers with the exception of the thumb were severed, and placed on ice for preservation during the transportation to the hospital. In all cases, the amputated digits suffered FCI, which was noted after successful replantation. In one patient we assessed the benefit of administering HBOT to treat FCI of the replanted digits. Signs of FCI appeared three days after replantation. One patient received HBOT 7 days after replantation, whereas the others did not. Due to oedema and progressive necrosis observed in the replanted fingers of these patients, reamputation was necessary in most cases. The detrimental consequences of inappropriate preservation of the severed fingers at subzero temperatures were: non-freezing tissue damage, arterial and venous thrombosis during the microsurgical procedure, and longer operations. Post-operatively, improper preservation during transport caused complications associated with freezing- and non-freezing cold injury, despite successful reinstatement of perfusion to the replanted digits. Proper preservation of amputated body parts is essential for successful replantation. Hypothermic preservation is appropriate, if conducted properly. Since digits contain no muscle tissue, irreversible damage appears after significantly longer periods of ischaemia than in other types of tissue. Although we did not observe any benefit of HBOT in one patient, this is most likely due to the delay in initiating the treatment. HBOT should be administered immediately post-surgery, both to minimise reperfusion injury and freezing/non-freezing cold injury.

1.0 INTRODUCTION

Cooling of amputated parts during transportation delays the onset of ischaemic tissue damage and contributes to successful replantation. Current guidelines for a doctor, who refers a patient to a hospital for replantation surgery, are strictly defined. The amputated part should be wrapped in sterile moist (using Ringer lactate) gauze, then placed in a clean plastic bag and floated in a mixture of water and ice. The temperature of the mixture must be between 4 and 6 degrees Celsius. The stump should be protected with moist gauze and bandaged. The limb should be immobilized and elevated (Janezic and Arnez, 1998). Optimal conditions for hypothermic preservation of amputated digits are hard to attain in the field even

Ahcan, U.; Luzar, B.; Bajrovic, F.; Mekjavic, I. (2005) Cold Injury of Amputated Digits. In *Prevention of Cold Injuries* (pp. 10-1 – 10-4). Meeting Proceedings RTO-MP-HFM-126, Paper 10. Neuilly-sur-Seine, France: RTO. Available from: <http://www.rto.nato.int/abstracts.asp>.

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 01 MAY 2005	2. REPORT TYPE N/A	3. DATES COVERED -			
4. TITLE AND SUBTITLE Cold Injury of Amputated Digits		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Plastic Surgery and Burns, Clinical Centre Ljubljana, Zaloska 7, SI-1000 Ljubljana		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001854, Prevention of Cold Injuries (Prevention des accidents dus au froid) ., The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	UU	4	

when all the necessary equipment is available and the above guidelines are followed (Sapega et al., 1988; VanGiesen et al., 1983). The knowledge of the professional and lay public is insufficient. The most common error in preservation of amputated parts is exposing them to temperatures, which may cause cold injury, and thus render them questionable for replantation. The extent of tissue damage due to freezing is difficult to evaluate clinically. In the literature, hypothermic tissue injury is not listed as a contraindication to replantation surgery (Soucacos, 2001; Pederson, 2001; Gold and Lee, 1981). The surgeon is faced with a dilemma whether to replant or not. Most surgeons will attempt to replant the digit despite clinically obvious freezing injuries. A higher incidence of intra- and postoperative complications is noted in these cases. The replanted digit may be lost due to arterial and venous thrombosis in the immediate postoperative period. In this article, six cases will be presented, which illustrate how freezing cold injury (FCI) of amputated digits may decrease the viability of such digits following replantation, and discuss the potential benefit of hyperbaric oxygen therapy (HBOT) in treating FCI of amputated digits following replantation.

2.0 METHODS

During the period 1998 to 2002, in 6 of 124 admitted injured individuals the digits were brought frozen to the Department of Plastic Surgery and Burns at the Ljubljana Clinical Centre. Upon admission, the type of injury was assessed, and 18 of the 22 severed digits were replanted.

Due to the reported decrease in reperfusion injury and edema, and enhancement of rate of healing of problem wounds with hyperbaric oxygen (HBO) therapy, it was considered appropriate to include this therapy as an adjunct treatment, while one patient was hospitalised post-replantation. This patient received five daily 90 minute HBO treatments at 2.5 ATA in a multiplace hyperbaric chamber. The treatment was initiated 7 days after replantation.

3.0 RESULTS

In all cases, the amputated digits suffered FCI, which was noted after successful replantation. Signs of FCI appeared three days after replantation. Five of the 18 replanted digits remained viable, 13 had to be reamputated later.

In one patient we assessed the benefit of administering HBOT to treat FCI of the replanted digits. This patient received HBOT 7 days after replantation, whereas the others did not. Due to oedema and progressive necrosis observed in the replanted fingers of these patients, reamputation was necessary in most cases.

The detrimental consequences of inappropriate preservation of the severed fingers at subzero temperatures were: non-freezing tissue damage, arterial and venous thrombosis during the microsurgical procedure, and longer operations. Post-operatively, improper preservation during transport caused complications associated with freezing- and non-freezing cold injury, despite successful reinstatement of perfusion to the replanted digits.

4.0 DISCUSSION

We have found that success rate of replantation of amputated digits is significantly reduced by incorrect cooling of amputated digits during preoperative preservation. Because amputated digits are not warmed by inflow of arterial blood, they become frozen more quickly than perfused digits and are more susceptible to frostbite. Other factors that promote heat loss are contact with moisture or bare metal, high wind velocities, immobility and the small size of the amputated digit (Molnar et al., 1973; House and Fidler,

1993). A frozen digit is pale, hard and cool. More subtle hypothermic damage to the small blood vessels cannot be determined in the preoperative period. Additional tissue damage occurs during reperfusion (reperfusion injury). Vascular endothelium, bone and skin are most vulnerable. Vascular changes lead to thrombotic occlusion of arteries and veins, causing vascular spasms and subsequent diminished blood supply to the replanted digit (Mowlavi et al., 2003; Nakazato et al., 1986). In children, bone damage becomes evident 1-2 years after exposure to severe cold as stunted growth and destruction of epiphyseal growth cartilage, leading to flexion deformities of the fingers, radial deviation and anomalous fingernail growth (Nakazato and Ogin, 1986; Wenzel et al., 1967). Signs of oedema and epidermolysis are seen on the skin and subcutaneous tissue. Progressive necrosis (demarcation) affects the skin, blood vessels, soft tissues and bones (VanGiesen et al., 1983; House and Fidler, 1993).

Therefore, proper preservation of amputated body parts is essential for successful replantation. Contrary to our previous experience with treating freezing cold injury with HBO, we did not observe any benefit of HBOT in the patient with replanted cold-injured digits. Most likely, this was due to the delay in initiating the treatment. Also, HBO therapy should be perhaps applied more aggressively (2 treatments daily) in the early phase of the healing process, and should include a much greater number of treatments. The protocol used for treating acute traumatic peripheral ischaemias with HBO therapy should be evaluated in future studies. HBOT should be considered as an adjunct therapy and administered immediately post-surgery, both to minimise reperfusion injury and freezing/non-freezing cold injury.

5.0 ACKNOWLEDGEMENTS

This work was supported, in part, by a Science for Security and Peace grant (project no. M2-0018) from the Ministry of Defence and Ministry of Education, Science and Sport of the Republic of Slovenia.

6.0 REFERENCES

- [1] Janezic T, Arnez ZM (1998). Indikacije za replantacijo. *Zdrav. Vestn.* 67: 241 – 244.
- [2] Soucacos PN (2001). Indications and selection for digital amputation and replantation. *J. Hand Surg.*, 26B: 572 – 581.
- [3] Pederson WC. Replantation. *Plastic and Reconstr. Surg.*, March 2001: 823 – 841.
- [3] Gold AH, Lee GW (1981). Upper extremity replantation: Current concepts and patient selection. *J. Trauma* 21: 551 – 557.
- [4] Sapega AA et al. (1988). The bioenergetics of preservation of limbs before replantation. *J. Bone Joint Surg.* 70-A: 1500 – 1513.
- [5] VanGiesen PJ, Seaber AV, Urbaniak JR. (1983). Storage of amputated parts prior to replantation— an experimental study with rabbit ears. *J. Hand Surg.* 8: 60 – 65.
- [6] Molnar GW, Hughes AL, Wilson O, Goldman RF (1973). Effect of skin wetting on finger cooling and freezing. *J. Appl. Physiol.* 35: 205 – 207.
- [7] House JH, Fidler MO (1993). Frostbite. In: Green DP (ed.): *Operative Hand Surgery*, 3rd ed., Vol. 2; Churchill Livingstone: New York; pp. 2033 – 2041.
- [8] Mowlavi A, Neumeister MW, Wilhelmi BJ, Song YH, Suchy H, Russell RC (2003). Local hypothermia during early reperfusion protects skeletal muscle from ischaemia – reperfusion injury. *Plast. Reconstr. Surg.* 111; 242 – 250.

Cold Injury of Amputated Digits

- [9] Nakazato T, Ogino T (1986). Epiphyseal destruction of children's hands after frostbite: A report of two cases. *J. Hand Surg.*, 11A: 289 – 292.
- [10] Wenzel JE, Burke EC, Bianco AJ (1967). Epiphyseal destruction from frostbite of the hands. *Am J Dis. Child* 114: 668 – 70.