

AD_____

AWARD NUMBER: W81XWH-04-1-0050

TITLE: Investigating the Role of Hepatocyte Nuclear Factor-3 (HNF-3) Alpha and Beta in Prostate Cancer and Cellular Differentiation

PRINCIPAL INVESTIGATOR: Janni Mirosevich, Ph.D.

CONTRACTING ORGANIZATION: Beckman Research Institute of the City of Hope
Duarte, California 91010

REPORT DATE: January 2006

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) January 2006		2. REPORT TYPE Annual Summary		3. DATES COVERED (From - To) 1 Jan 04 – 31 Dec 05	
4. TITLE AND SUBTITLE Investigating the Role of Hepatocyte Nuclear Factor-3 (HNF-3) Alpha and Beta in Prostate Cancer and Cellular Differentiation				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-04-1-0050	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Janni Mirosevich, Ph.D. E-Mail: jmirosevich@coh.org				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Beckman Research Institute of the City of Hope Duarte, California 91010				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT No abstract provided.					
15. SUBJECT TERMS No subject terms provided.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

Introduction.....	1
Task 1 - Research Performed2
Key Findings	3
Task 2 - Research Performed	4
Key Findings	4
Task 3 - Research Performed	7
Conclusion	8
Training9
Reportable Outcomes.....	9
References	10

Introduction:

The Hepatocyte Nuclear Factor-3 (HNF-3) family of *forkhead* transcription factors (HNF-3 α , β and γ) are essential for endoderm development (Kaestner et al., 1994). HNF-3 proteins, along with other cell specific transcription factors, have been shown to activate the transcription of numerous genes important for hepatocytic and respiratory epithelial cell function (Kaestner et al., 2000). We have recently demonstrated that HNF-3 α is a transcriptional potentiator for androgen-regulated prostatic gene expression in epithelial cells and is also necessary for normal murine prostate development (Gao et al., 2003; Mirosevich et al., 2005; Gao et al., 2005). Further, I have obtained significant correlative evidence for the role of HNF-3 α and β proteins in oncogenic involvement in tumor initiation and/or progression, in addition to prostate epithelial differentiation (Mirosevich et al., In Press). HNF-3 α and β protein expression is altered in both the LADY transgenic mouse prostate cancer model, in human prostate adenocarcinomas and is differentially expressed in a variety of well established human prostate cancer cell lines (Mirosevich et al., In Press). The precise role that HNF-3 proteins play in normal prostate epithelial cell function, prostate cancer development and progression is unknown.

The signal transducer and activator of transcription-3 (Stat3) protein belongs to the Stat family of transcription factors and regulates numerous genes involved in normal cellular processes including cell proliferation, differentiation and apoptosis (Yu & Jove, 2004). In normal cells Stat3 phosphorylation is tightly regulated, however during cellular transformation Stat3 expression becomes altered and is frequently observed to be up-regulated and constitutively activated, i.e. phosphorylated, in many tumors, including prostate carcinomas (Mora et al., 2001). HNF-3 α and Stat3 have previously been shown to interact directly and bind to the core enhancer region of the hepatitis B virus to regulate viral gene transcription (Waris & Siddiqui, 2002). I have recently identified and confirmed that Stat3 transcription factor binding sites are located adjacent to HNF-3 sites in the promoters of several prostate specific genes including the Prostate Specific Antigen (PSA) gene. Currently, the roles that HNF-3 α and Stat3 protein interactions play in regulating normal prostate specific gene transcription, epithelial cell function and tumor initiation/progression are unknown.

The overall objective of this proposal is to elucidate the roles that HNF-3 and Stat3 proteins play in normal and cancerous prostate epithelial cells by employing several molecular biological techniques; *in vitro* and *in vivo* models. The first goal is to establish the functional significance of HNF-3 α and β proteins, in normal and cancerous prostatic epithelial cells. The central hypothesis of this project is that HNF-3 α and β are involved in prostatic epithelial differentiation and that altered expression of these genes is important in development and/or progression of prostate adenocarcinoma. The second goal is to establish the functional significance of HNF-3 α and Stat3 interactions in normal and cancerous prostatic epithelial cell gene regulation. The hypothesis is that HNF-3 α and Stat3 interact to regulate genes involved in normal prostatic epithelial differentiation and that the expression and interaction of these transcription factors is important in the development and/or progression of prostate adenocarcinoma. In summary, the proposed research will define the importance of HNF-3 *forkhead* proteins and Stat3 proteins in prostate epithelial cells and the development/progression of prostate tumors.



Task 1.

Research Performed

- While transient transfection experiments, using a tetracycline inducible system and tetracycline regulated HNF-3 α siRNAs, produced adequate protein knockdown, similar experiments performed to test siRNAs towards HNF-3 β failed to produce a decrease in protein levels. As such, experiments using Dharmacon RNAis directed towards HNF-3 were performed. Unfortunately, these siRNAs also failed to adequately reduce HNF-3 β protein levels (results not shown).
- Due to the problems associated with the *in vitro* RNAi knockdown experiments, Task 1.5: Vary the ratios of HNF-3 α and β in epithelial cells for characterization, was not pursued further.
- DU-145 and BPH-1 prostate cancer cells lines were retrovirally infected with pLZRS-CMV-HNF-3 α or HNF-3 β expression vectors to produce HNF-3 overexpressing prostate cancer cells (Figure 1). LNCaPs could not be infected with the HNF-3 β expression construct. As such, a lentiviral expression construct is being created to infect these cells.
- The proliferation rates of control and infected DU-145 and BPH-1 cell lines were measured using the cell titer 96 Aqueous One solution cell proliferation assay. The apoptosis assay has yet to be performed.
- Western blot analysis was performed on the cell lines to examine *in vitro* androgen receptor, PSA and synaptophysin expression.

Key Research Findings

- Overexpression of HNF-3 α and β (Figure1), had no effect on the *in vitro* proliferative index of DU-145 cells, but did increase proliferation of BPH-1/HNF-3 infected cells (Figure 2). BPH-1 cells overexpressing HNF-3 proteins were chosen for subsequent tissue recombinant experiments.
- No differences in androgen receptor, PSA and synaptophysin expression were observed between experimental and control DU-145 or BPH-1 cell lines (results not shown).



Task 2.

Research Performed

- HNF-3 α or HNF-3 β overexpressing BPH-1 cell lines were recombined with rat embryo 18 day urogenital sinus mesenchymal cells and placed under the kidney capsule of athymic nude mice.
- Eight mice were used for 16 experimental grafts, containing cells over expressing HNF-3 α or HNF-3 β , in the left kidney. Twelve BPH-1 tissue recombination controls were grafted into the right kidney.
- Mice were sacrificed at six weeks and three months after implantation, the grafts harvested and analyzed for histology, HNF-3 expression, proliferation and expression of various differentiation markers, i.e. androgen receptor, PSA and synaptophysin. Immunohistochemical staining for apoptosis has yet to be performed.
- The proposed tumorigenicity studies have not been performed due to my recent move from the Moffitt Cancer Center in Tampa, Florida to the Beckman Research Institute of the City of Hope in Duarte, California. (see Additional Comments, page 10)

Key Research Findings

- Histologically, the HNF-3 α and HNF-3 β infected BPH-1 tissue recombinant grafts appeared similar to control BPH-1 cells. All tissue recombinants appeared to form canalized glandular epithelial structures with defined eosinophilic basement membrane surrounded by stroma. (Figure 3).
- As expected, BPH-1 infected cells expressed their respective HNF-3 proteins (Figure 4A&B). No differences were observed in the number of proliferating epithelial cells between experimental and control groups (results not shown). Interestingly, the control non-HNF-3 infected BPH-1 tissue recombination grafts also expressed HNF-3 α (Figure 4C), but not HNF-3 β (results not shown).
- When BPH-1 cells were grafted under the renal capsules without mesenchymal cells, the SV-40 transformed BPH-1 cells expressed T-antigen, however, HNF-3 α expression was not detected by immunohistochemistry (Figure 5B).




Figure 3. Histology of BPH-1 and BPH-1 HNF-3 tissue recombinants. The epithelial glandular structures which formed in the (B) BPH-1/HNF-3 β and (C) BPH-1/HNF-3a tissue recombinants were similar to the glandular structures which formed in the (A) BPH-1 control tissue recombinants. x10




Figure 4. Immunohistochemical (IHC) analysis for HNF-3 expression in BPH-1 tissue recombinants following 6-weeks of sub-renal grafting. (B) Strong HNF-3 β expression was detected in BPH-1 epithelial cells infected with HNF-3a, while (C) strong HNF-3a expression was detected in BPH-1/ HNF-3a infected cells. (A) Interestingly, HNF-3a expression was also detected in epithelial cells of non-infected control BPH-1 tissue recombinants. x20 All cells expressed T-antigen while BPH-1/ HNF-3a and BPH-1/ HNF-3 β also expressed GFP (results not shown).






Figure 5. Immunohistochemical (IHC) analysis for T-antigen and HNF-3 expression in BPH-1 cells following 4-weeks of sub-renal kidney capsule grafting. Immunohistochemical staining of serial sections derived from kidney capsule grafts of BPH-1 cells without urogenital sinus mesenchyme tissue recombination showed; (A) Strong T-antigen expression was observed in BPH-1 epithelial cells, however, HNF-3a expression was not detected in these grafts (B). x5


- We have previously demonstrated that HNF-3 β is expressed in synaptophysin expressing neuroendocrine cells of normal mouse (Figure 6A) and human prostates; neuroendocrine prostate tumors from the highly metastatic androgen independent 12T-10 LADY prostate cancer model (Figure 6B) and small cell carcinomas from human prostate biopsy samples (Mirosevich et al., In Press).
- Synaptophysin expression was examined in BPH-1/HNF-3 β tissue recombination experiments by immunohistochemistry, however, no synaptophysin expression was detected in the tissue recombinant samples (Figure 7).



Task 3.

Research Performed

- I have recently shown, by electromobility shift assay (EMSA), that Stat3 binding sites exist adjacent to HNF-3 α on prostate specific promoters. As such, I plan to perform experiments to determine how these two proteins interact to regulate prostate gene transcription.
- I have acquired both full-length and various truncated HNF-3 α -V5 tagged expression constructs that have previously been used to determine HNF-3 and androgen receptor interactions (Figure 8A; Gao et al., 2003).
- DU-145 prostate cancer cell lines, which constitutively expresses activated Stat3, were transfected with these HNF-3 α -V5 tagged expression constructs. A preliminary co-immunoprecipitation experiment showed that Stat3 interacts with the *forkhead* DNA binding domain of HNF-3 α .
- Recently, full length and truncated Stat3-GST tagged constructs have been kindly donated by Dr. Xinmin Cao (Figure 8B; Zhang et al., 2002). These constructs will be transcribed and translated *in vitro* and used in GST-pull down experiments. Protein interactions will be determined by western blot analysis using the V5 antibody (Invitrogen, Carlsbad, USA).



Additional Comments

- In September of this year I moved with Dr. Jove from the Moffitt Cancer Center in Tampa, Florida to the Beckman Research Institute of the City of Hope in Duarte, California. Subsequently, I received a promotion from a post-doctoral fellow to research assistant scientist.

Conclusions

The findings from these studies have provided interesting insights into the biological and molecular roles of HNF-3 and Stat3 proteins in prostate cancer cells. Results from the last six months have demonstrated several key findings. Firstly, *in vitro* cell culture conditions are quite different to the *in vivo* conditions that can be achieved by techniques such as tissue recombination. These differences were highlighted when the BPH-1 cell *in vitro* proliferation results were compared with the *in vivo* tissue recombination experiments; in addition to the re-expression of HNF-3 α following tissue recombination and sub-renal grafting. The re-expression of HNF-3 α following tissue recombination with urogenital sinus mesenchymal cells also confirms that epithelial-stromal interactions play an important and necessary role in prostate organogenesis (Marker et al., 2003). Classical tissue recombination experiments by Cunha and colleagues have established that epithelial and stromal cells produce reciprocal paracrine signaling molecules to induce normal prostate development (Cunha et al., 1987). The findings from my work show, for the first time, that mesenchymal/stromal cells produce signaling molecules that stimulate expression of HNF-3 α , an important transcription factor for both normal prostate development and cancer. The signaling molecules, which are involved in stimulating the *in vivo* HNF-3 α epithelial cell expression, are currently unknown and are actively being investigated.

In my initial research proposal, I expected that the expression of HNF-3 β may cause luminal epithelial cells to differentiate into a neuroendocrine phenotype. However, the tissue recombination results in this study demonstrated that HNF-3 β alone is insufficient to promote a prostatic neuroendocrine epithelial cell phenotype, suggesting that other factors are required. Since, BPH-1 cells do not express the androgen receptor (Wang et al., 2001), all the BPH-1 cell sub-lines produced thus far have been infected with androgen receptor expressing constructs. These cells will be used in subsequent tissue recombination experiments to see the effects of HNF-3 and androgen receptors on prostate epithelial cell biology/phenotype.

Finally, the preliminary immunoprecipitation experiments performed show that the *forkhead* DNA binding domain directly interacts with Stat3. This interaction is similar to our previously published work demonstrating that the androgen receptor also interacts with the DNA binding domain of HNF-3 α (Gao et al., 2003), and suggests that the androgen receptor/HNF-3 α /Stat3 proteins associate within close proximity to form a prostate tissue specific transcriptional unit.

The initiation and progression of prostate cancer is considered a multi-step process involving, primarily, dysregulated gene expression. Currently, very little is known about the genes which are necessary for this process. Elucidating the role of HNF-3 and Stat3 protein interactions in both prostate gene regulation and tumor initiation and/or progression will define the importance of HNF-3/Stat3 transcription factors in prostate biology, and will verify that these protein complexes are potential therapeutic targets for the treatment of advanced prostate cancer.

Training

So far this year I have attended two national meetings.

- American Association for Cancer Research 96th Annual Meeting, Anaheim, CA, April 2005.
- Society for Basic Urologic Research Spring 2005 Meeting, San Antonio, TX, May 2005.

I will also attend and participate in a third national meeting in December.

- Society for Basic Urologic Research Fall 2005 Meeting, Miami, FL, December 2005.

In addition, I gave an oral presentation to the Department of Pathology at Royal Perth Hospital in Perth, Western Australia, during a visit to Australia in February.

Lunchtime and evening seminars and mini symposiums also contributed to an overall training strategy in place within the Moffitt Cancer Center and the Department of Molecular Oncology. Finally, laboratory meeting discussions and journal club presentations are integral to the running of the department. I attended and presented within these forums.

Reportable outcomes

Publications

Mirosevich J, Gao N, Gupta A, Shappell SB, Jove R & Matusik RJ. 2005 The expression and role of Foxa proteins in prostate cancer. *The Prostate*, In Press.

Nam S, Kim D, Cheng JQ, Zhang S, Lee J, Buettner R, **Mirosevich J**, Lee FY & Jove R. 2005 Action of the Src family kinase inhibitor, Dasatinib (BMS-354825), on human prostate cancer cells. *Cancer Research*, **65** 9185-9189.

Gao N, Ishii K, **Mirosevich J**, Kuwajima S, Oppenheimer SR, Roberts RL, Jiang M, Shappell SB, Caprioli RM, Stoffel M, Hayward SW & Matusik RJ. 2005 Forkhead box A1 regulates prostate ductal morphogenesis and promotes epithelial cell maturation. *Development*, **132** 3431-3443.

Yi Y, **Mirosevich J**, Shyr Y, Matusik R & George AL Jr. 2005 Coupled analysis of gene expression and chromosomal location. *Genomics*, **85** 401-412.

Klezovitch O, Chevillet J, **Mirosevich J**, Roberts RL, Matusik RJ & Vasioukhin V. 2004 A causal role for hepsin in prostate cancer metastasis. *Cancer Cell*, **6** 185-195.

Poster Abstracts

Mirosevich J, Shappell SB, Jove R & Matusik RJ. Prostatic neuroendocrine cells express variable levels of androgen receptor in normal and tumorigenic cell populations. *Society for Basic Urologic Research Fall 2005 Meeting*, Miami, FL, December 2005.

References

- Cunha GR, Donjacour AA, Cooke PS, Mee S, Bigsby RM, Higgins SJ, Sugimura Y. The endocrinology and developmental biology of the prostate. *Endocr Rev* 1987;8:338-362.
- Gao N, Ishii K, Mirosevich J, Kuwajima S, Oppenheimer SR, Roberts RL, Jiang M, Yu X, Shappell SB, Caprioli RM, Stoffel M, Hayward SW, Matusik RJ. Forkhead box A1 regulates prostate ductal morphogenesis and promotes epithelial cell maturation. *Development* 2005;132(15):3431-3443.
- Gao N, Zhang J, Rao MA, Case TC, Mirosevich J, Wang Y, Jin R, Gupta A, Rennie PS, Matusik RJ. The role of hepatocyte nuclear factor-3 alpha (Forkhead Box A1) and androgen receptor in transcriptional regulation of prostatic genes. *MolEndocrinol* 2003;17(8):1484-1507.
- Kaestner KH, Hiemisch H, Luckow B, Schutz G. The HNF-3 gene family of transcription factors in mice: gene structure, cDNA sequence, and mRNA distribution. *Genomics* 1994;20(3):377-385.
- Kaestner KH, Knochel W, Martinez DE. Unified nomenclature for the winged helix/forkhead transcription factors. *Genes Dev* 2000;14(2):142-146.
- Marker PC, Donjacour AA, Dahiya R, Cunha GR. Hormonal, cellular, and molecular control of prostatic development. *Dev Biol* 2003;253:165-174.
- Mirosevich J, Gao N, Gupta A, Shappell SB, Jove R, Matusik RJ. Expression and role of Foxa proteins in prostate cancer. *The Prostate; In Press*.
- Mirosevich J, Gao N, Matusik RJ. Expression of foxa transcription factors in the developing and adult murine prostate. *The Prostate* 2005;62:339-352.
- Mora LB, Buettner R, Ahmad N, Bassel Y, Jove R, Seigne JD. Prostate adenocarcinoma: cellular and molecular abnormalities. *Cancer Control* 2001;8(6):551-561.
- Wang Y, Sudilovsky D, Zhang B, Haughney PC, Rosen MA, Wu DS, Cunha TJ, Dahiya R, Cunha GR, Hayward SW. A human prostatic epithelial model of hormonal carcinogenesis. *Cancer Res*. 2001;61:6064-6072.
- Waris G, Siddiqui A. Interaction between STAT-3 and HNF-3 leads to the activation of liver-specific hepatitis B virus enhancer 1 function. *J Virol* 2002;76(6):2721-2729.
- Yu H, Jove R. The STATs of cancer--new molecular targets come of age. *Nat Rev Cancer* 2004;4(2):97-105.
- Zhang T, Seow KT, Ong CT, Cao X. Interdomain interaction of Stat3 regulates its Src homology 2 domain-mediated receptor binding activity. *J Biol Chem* 2002;277(20):17556-17563.