

Award Number: DAMD17-98-1-8333

TITLE: Inherited Susceptibility to Breast Cancer in Healthy Women: Mutation in Breast Cancer Genes, Immune Surveillance, and Psychological Distress

PRINCIPAL INVESTIGATOR: Dana H. Bovbjerg, Ph.D.

CONTRACTING ORGANIZATION: Mount Sinai School of Medicine  
New York, NY 10029-6574

REPORT DATE: October 2005

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. REPORT DATE</b> 01-10-2005			<b>2. REPORT TYPE</b> Final		<b>3. DATES COVERED</b> 28 Sep 1998 – 27 Sep 2005	
<b>4. TITLE AND SUBTITLE</b>  Inherited Susceptibility to Breast Cancer in Healthy Women: Mutation in Breast Cancer Genes, Immune Surveillance, and Psychological Distress					<b>5a. CONTRACT NUMBER</b>	
					<b>5b. GRANT NUMBER</b> DAMD17-98-1-8333	
					<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b>  Dana H. Bovbjerg, Ph.D.					<b>5d. PROJECT NUMBER</b>	
					<b>5e. TASK NUMBER</b>	
					<b>5f. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>  Mount Sinai School of Medicine New York, NY 10029-6574					<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012					<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
					<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>	
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for Public Release; Distribution Unlimited						
<b>13. SUPPLEMENTARY NOTES</b>						
<b>14. ABSTRACT</b>  The purpose of the research supported by this IDEA grant award was to test the possibility that variability in the strength of immune surveillance mechanisms against cancer (operationally defined by assessment of natural killer cell activity) may be a factor in determining the penetrance of mutations in breast cancer susceptibility genes. The following hypotheses were investigated: Hypothesis 1: Women with family histories of breast cancer are more emotionally distressed than women at normal risk, particularly after notification that they carry a mutation in a primary susceptibility gene. Hypothesis 2: Women with family histories of breast cancer will have lower levels of NK cell activity than normal risk women, which cannot be entirely attributed to the effects of higher emotional distress or the presence of a mutation in primary susceptibility genes. Hypothesis 3: Variability in NK cell activity will contribute to variability in the penetrance of mutations in primary susceptibility genes for breast cancer. Data has been collected on 255 women. Preliminary analyses support these hypotheses.						
<b>15. SUBJECT TERMS</b> Breast Cancer						
<b>16. SECURITY CLASSIFICATION OF:</b>					<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b> USAMRMC
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U				<b>19b. TELEPHONE NUMBER</b> (include area code)
				UU	11	

## Table of Contents

<b>Cover.....</b>	<b>1</b>
<b>SF 298.....</b>	<b>2</b>
<b>Table of Contents.....</b>	<b>3</b>
<b>Introduction.....</b>	<b>4</b>
<b>Body.....</b>	<b>5</b>
<b>Key Research Accomplishments.....</b>	<b>7</b>
<b>Reportable Outcomes.....</b>	<b>9</b>
<b>Conclusions.....</b>	<b>10</b>
<b>Personnel .....</b>	<b>11</b>
<b>References.....</b>	<b>11</b>
<b>Appendices.....</b>	

## **INTRODUCTION:**

The penetrance of mutations in primary breast cancer susceptibility genes (BRCA1/BRCA2) is highly variable. The mechanisms underlying this variability are not yet well established (Dite et al., 2000; Antoniou et al., 2002; Couch, 2004). Modifying genes and/or environmental factors are likely to play a major role (Antoniou et al., 2002; Couch, 2004). To date, the search for such factors has primarily focused on hormonal/reproductive variables previously established as independent risk factors for breast cancer (DeJong et al., 2002; Martin & Weber, 2000). Another factor that could contribute to differences in penetrance, and/or make an independent contribution to risk is variability in immune surveillance. One immune surveillance mechanism (natural killer cell activity, NKCA) has been reported to be lower in women with family histories of breast cancer, consistent with inherited deficits in immune surveillance (Bovbjerg & Valdimarsdottir, 2001). However, the lower levels of NKCA in these women could be secondary to higher levels of stress, to which NKCA is particularly sensitive (Segerstrom & Miller, 2004). Consistent with this possible environmental effect on NKCA, survey studies have found higher levels of distress among women with family histories of breast cancer (Bovbjerg & Valdimarsdottir, 2001; Lindberg & Wellisch, 2004; Schnur et al., 2004, Kim et al., 2005).

The purpose of the research supported by this IDEA grant award was to test the possibility that differences in the strength of immune surveillance mechanisms against cancer (operationally defined as natural killer cell activity) may be a factor in determining the penetrance of mutations in breast cancer susceptibility genes. The first aim was to investigate two possible explanations for variability in NKCA (Bovbjerg & Valdimarsdottir, 2001): 1) stress-induced immune suppression, and 2) inherited deficits in immune surveillance. The second aim was to examine the possibility that inherited deficits in immune surveillance may be independently associated with familial risk of breast cancer (Bovbjerg & Valdimarsdottir, 2001).

The study "piggy-backed" on other studies involving familial risk, genetic counseling, and breast cancer gene testing (BRCA1, BRCA2) under the direction of Co-Investigators on this proposal. These "parent" studies, which provided the infrastructure and funding necessary for recruitment, assessment, genetic counseling, and BRCA testing, were the source of potential participants for the present study. The participants in the present study were recruited to form three Study Groups (N=80/group) of comparable age for the research: 1) The Mutation-Positive Family History Group (Mut+Hist+) includes women whose family histories of cancer indicate a relative risk  $\geq 1.5$  for breast cancer and who carry a mutation in BRCA1 or BRCA2; 2) The Mutation-Negative Risk Family History Group (Mut-Hist+) includes women with comparable family histories, who do not carry mutations; 3) The Normal Risk Group (Mut-Hist-) includes women without family histories of cancer who do not carry mutations. Study participants completed psychological assessments (e.g., standardized self-report measures) in conjunction with their involvement with the parent studies that fund the genetic testing. To reduce participant burden and avoid compromising the parent studies, blood samples for the assessment of NKCA were also collected in conjunction with the women's involvement in the parent studies, by collecting additional samples when the women were already providing a sample for genetic testing. In the context of the requirements of the parent studies, it has not been feasible to collect blood samples for the two follow-up NK cell assessments originally proposed for this study, as psychological data is collected by telephone. Consistent with scheduling exigencies, NKCA was concurrently assessed in samples from women in each group by personnel "blind" to group status.

**BODY:**

In the first year of the grant, Bovbjerg (PI) and Valdimarsdottir (Co-I) moved from Memorial Sloan-Kettering Cancer Center to Mount Sinai Medical Center. The resulting site change required modification of collaborative arrangements (e.g., the addition of Eng as a co-investigator), set up of new facilities (e.g., Bovbjerg lab), hiring of new support staff, and additional attention to institutional review requirements. Subsequent to that change, Eng left Mount Sinai and Valdimarsdottir began a new collaborative study involving investigators at another institution (Georgetown) necessitating further modification of collaborative arrangements. These complications caused a delay in the proposed rate of study accrual and the necessity of applying for several no-cost extensions.

In 2002, we proposed three methodological changes to expedite the meeting of our study goals in light of the delays: 1) we reduced the sample size to 80 per group from 100; 2) we modified the scheduling of assessments to enhance recruitment efforts; 3) we streamlined the immune assessments by using well validated and reliable whole blood approaches (e.g., four color flow cytometry), rather than approaches requiring labor intensive isolation of peripheral blood lymphocytes. In 2003, we proposed an additional change based on our experience in recruiting for the study: we reduced the sample size for the assessments of NK cell activity to 140.

Our progress according to the original Statement of Work is detailed below:

Task 1: Preparation for first wave of subjects. Preparation of psychosocial questionnaires and immune assessments. Data base established.

Completed, as previously reported.

Task 2: First wave of subjects completes assessments. Data entry and initial analysis.

Completed, as previously reported.

Task 3: Complete data entry of first wave. Prepare annual report. Prepare for second wave of subjects.

Completed, as previously reported.

Task 4: Second wave of subjects completes assessments. Data entry and analyses continues.

Completed, as previously reported.

Task 5: Complete data entry of first wave. Prepare annual report. Prepare for second wave of subjects.

Completed, as previously reported.

Task 6: Third wave of subjects completes assessments. Data entry and analyses continues.

Completed.

Task 7: Complete data entry for third wave. Complete empiric risk determination. Verify study data. Conduct literature review of relevant articles. Meet with research team to review results. Complete statistical analyses. Write manuscripts; prepare graphics. Complete DOD final report.

Completed. Preliminary analyses are completed. Further analysis of data will continue with funding from other sources.

Throughout the grant period we have made consistent progress toward our study goals and have completed recruitment. Over the grant period a total of 255 women (Mut+Hist+ n=99; Mut-Hist+ n=80; Mut-Hist- n=76) have completed psychological assessments (Assessment 1: n=255; Assessment 2: n=182; Assessment 3: n=142). Over the grant period NK cell activity has been assessed in a total of 135 women (Mut+Hist+ n=37; Mut-Hist+ n=45; Mut-Hist- n= 53).

Analyses thus far have supported our hypotheses:

- Hypothesis 1: Women with family histories of breast cancer will be more emotionally distressed than women at normal risk, particularly after notification that they carry a mutation in a primary susceptibility gene.
  - Healthy women undergoing genetic testing for BRCA mutations, compared to controls, before and after testing have higher levels of:
    - Perceived risk of the disease
    - Breast cancer-specific distress (intrusion/avoidance)
    - Anxiety symptoms
    - Depressive symptoms
  - Women post negative BRCA result have:
    - Reduced perceived risk
    - Reduced breast cancer-specific distress, anxiety/depressive symptoms
  - Women post positive BRCA result have:
    - Increased perceived risk
    - Increased breast cancer-specific avoidance
    - No change in breast cancer-specific intrusions, or anxiety/depressive symptoms
    - Remain higher than controls on all
- Hypothesis 2: Women with family histories of breast cancer will have lower levels of NK cell activity than normal risk women, which cannot be entirely attributed to the effects of higher emotional distress or the presence of a mutation in primary susceptibility genes.
  - Analyses have not yet been completed.
- Hypothesis 3: Variability in NK cell activity will contribute to variability in the penetrance of mutations in primary susceptibility genes for breast cancer.
  - Analyses have not yet been completed.

## **KEY RESEARCH ACCOMPLISHMENTS:**

### **Accepted abstracts:**

- Zakowski S, Valdimarsdottir H, Bovbjerg DH. Emotional expressivity in women at familial risk for breast cancer. Annals of Behavioral Medicine 1998; 20:S156.
- Bovbjerg DH, Erbllich J, Valdimarsdottir H. Intrusive thoughts and perceived risk predict general distress in women with family histories of breast cancer. Annals of Behavioral Medicine 1999; 21:S32.
- Bovbjerg DH, Valdimarsdottir H, Norman C. Distress associated with mammography screening is higher among healthy women with histories of breast cancer. Annals of Behavioral Medicine 1999; 21:S33
- Erbllich J, Bovbjerg DH, Valdimarsdottir H. Caregiving and death of a mother predict current distress in women with family histories of breast cancer. Annals of Behavioral Medicine 1999; 21:S137.
- Bovbjerg DH, Guevarra JS, Valdimarsdottir HB. Familial breast cancer risk is a stressor for black women. Psychosomatic Medicine 2000; 62:98.
- Guevarra J, Bovbjerg DH, Erbllich J, Valdimarsdottir H, Godfrey D, Freeman H. Role of family histories of breast cancer and psychological distress in black women's breast-self examination frequency. Annals of Behavioral Medicine 2000; 22:S57.
- Brown K, Valdimarsdottir H, Erbllich J, Amareld D, Scheuer L, Hull J, McDermott D, Bovbjerg DH, Hurley K, Offit K. Does genetic counseling for breast cancer predisposition increase knowledge? American Society of Human Genetics October 2000.
- Valdimarsdottir H, Bovbjerg DH, Zakowski. Familial breast cancer risk: psychobiological reactivity. Era of Hope June 2000.
- Zakowski SG, Valdimarsdottir HB, Bovbjerg DH. Cancer-specific distress and reduced NKCA in women at familial risk for breast cancer: the buffering role of emotional expressivity. Psychoneuroimmunology Research Society May 2001.
- Kim Y, Valdimarsdottir H, Bovbjerg DH. Coping strategies as moderators of cancer-specific distress among healthy women with family histories of breast cancer. Psychosomatic Medicine 2002; 64:106.
- Valdimarsdottir H, Jeremias JC, Zakowski S, Bovbjerg DH. Major life events and psychobiological reactivity. Psychosomatic Medicine 2002; 64:140.
- DiLorenzo T, Erbllich J, Montgomery G, Ephron R, Shaffren M, Bovbjerg DH. Family histories of disease and disease-specific worry: the role of perceived risk. Annals of Behavioral Medicine 2002 Supplement; 24:S093.
- Kim Y, Valdimarsdottir H, Bovbjerg DH. Family histories of breast cancer, health locus of control and coping. Annals of Behavioral Medicine 2002 Supplement, 24:S167.
- Valdimarsdottir H, Thompson H, Bovbjerg DH, Brown K, Hurley K, Amarel D, Offit K. Social constraints and psychological adjustment following notification of uninformative BRCA ½ test results. Annals of Behavioral Medicine 2003 Supplement, 25.
- Valdimarsdottir H, Zakowski S, Bovbjerg DH. Social constraints to expressing breast cancer concerns and knowledge following genetic counseling for breast cancer susceptibility.
- Bovbjerg DH, Valdimarsdottir H, Schwartz M. Inherited susceptibility to breast cancer in healthy women: mutations in breast cancer genes, immune surveillance and psychological distress. Era of Hope June 2005.

### **Peer-reviewed publications:**

- Erblich J, Bovbjerg DH, Valdimarsdottir HB. Looking forward and back: distress among women at familial risk for breast cancer. Annals of Behavioral Medicine 2000; 22:53-59.
- Erblich J, Bovbjerg DH, Norman C, Valdimarsdottir HB, Montgomery GH. It won't happen to me: Lower perception of heart disease risk among women with family histories of breast cancer. Preventive Medicine 2000; 31:714-721.
- Zakowski SG, Valdimarsdottir HB, Bovbjerg DH. Emotional expressivity and intrusive thoughts in women with family histories of breast cancer: Application of a cognitive processing model. British Journal of Health Psychology 2001; 6:151-165.
- Thompson HS, Valdimarsdottir HB, Buteau-Buck C, Guevarra J, Bovbjerg DH, Richmond-Avellaneda C, Godfrey D, Brown K, Offit K. Psychosocial predictors of BRCA counseling and testing decisions among urban African-American women. Cancer Epidemiology, Biomarkers & Prevention 2002; 11:1579-1585.
- Valdimarsdottir HB, Zakowski SG, Gerin W, Mamakos J, Pickering T, Bovbjerg DH. Heightened psychobiological reactivity to laboratory stressors in healthy women at familial risk for breast cancer. Journal of Behavioral Medicine 2002; 25:51-65.
- Gold SM, Zakowski SG, Valdimarsdottir HB, Bovbjerg DH. Stronger endocrine responses after brief psychological stress in women at familial risk of breast cancer. Psychoneuroendocrinology 2003; 28:584-593.
- Erblich J, Montgomery GH, Valdimarsdottir HB, Cloitre M, Bovbjerg DH. Biased cognitive processing of cancer-related information among women with family histories of breast cancer: Evidence from a cancer stroop task. Health Psychology 2003; 22:235-244.
- Kim Y, Valdimarsdottir HB, Bovbjerg DH. Family histories of breast cancer, coping styles, and psychological adjustment. Journal of Behavioral Medicine 2003; 26:225-243.
- James GD, van Berge-Landry H, Valdimarsdottir HV, Montgomery GH, Bovbjerg DH. Urinary catecholamine levels in daily life are elevated in women at familial risk of breast cancer. Psychoneuroendocrinology 2004; 29:831-838.
- Kim Y, DuHamel K, Valdimarsdottir HB, Bovbjerg DH. Psychological distress among healthy women with family histories of breast cancer: Effects of recent life events. PsychoOncology 2004; 13:1-9.
- Schnur JB, Valdimarsdottir HB, Montgomery GH, Nevid JS, Bovbjerg DH. Social constraints and distress among women at familial risk for breast cancer. Annals of Behavioral Medicine 2004; 28:142-148.
- Dettenborn L, James GD, van Berge-Landry H, Valdimarsdottir HB, Montgomery GH, Bovbjerg DH. Heightened cortisol responses to daily stress in working women at familial risk for breast cancer. Biological Psychology 2005; 69:167-179.
- Guevarra JS, Kwate NA, Tang TS, Valdimarsdottir HB, Freeman HP, Bovbjerg DH. Acculturation and its relationship to smoking and breast self-examination frequency in African American women. Journal of Behavioral Medicine 2005; 28:191-201.
- Rowe JL, Montgomery GH, Duberstein PR, Bovbjerg DH. Health locus of control and perceived risk for breast cancer in healthy women. Behavioral Medicine 2005; 31(1):33-40.
- Kim Y, DuHamel K, Valdimarsdottir HB, Bovbjerg DH. Psychological distress among healthy women with family histories of breast cancer: Effects of recent life events. PsychoOncology 2005; 14(7):555-63.
- DiLorenzo TA, Schnur J, Montgomery GH, Erblich J, Winkel G, Bovbjerg DH. A model of disease-specific worry in heritable disease: The influence of family history, perceived risk and worry about other illnesses. Journal of Behavioral Medicine in press.



## **REPORTABLE OUTCOMES:**

### **Funded grants:**

Department of the Army 7/1/01 - 6/30/05  
(Bovbjerg, PI)  
(Valdimarsdottir, Project 2 PI)  
(Brown, McGovern, Co-Inv)

Center Grant: Genetic Factors in Breast Cancer: Center for Interdisciplinary Biobehavioral Research:  
Project 2: "Impact of culturally tailored counseling on psychobehavioral outcomes and BRCA  
decision making among women with breast cancer"(Valdimarsdottir, PI)

National Cancer Institute 12/1/02 - 11/30/07  
(Valdimarsdottir, PI)  
(Bovbjerg, Co-Inv)

Decisions & Outcomes of BRCA 1/2 Test for Breast Patients

American Cancer Society 1/1/03 - 12/31/06  
(Valdimarsdottir, PI)  
(Bovbjerg, Co-Inv)

BRCA Counseling/Testing for Urban African American Women

Department of the Army 7/1/02 - 6/30/07  
(Bovbjerg, PI)

Immune Surveillance, Cytokines and Breast Cancer Risk: Genetic and Psychological Influences in  
African American Women

Department of the Army 06/01/04-05/30/08  
(Valdimarsdottir, PI)  
(Bovbjerg, Co-Inv)

Emotional, Biological and Cognitive Impact of a Brief Expressive Writing Intervention for African  
American Women at Familial Breast Cancer Risk

### **Invited Lectures/Presentations:**

Biobehavioral consequences of stress in individuals. Danish Cancer Society, Copenhagen,  
Denmark, 1998.

Psychoneuroimmunology and cancer. University of Aarhus, Aarhus, Denmark, 1998.

Stress and stress biomarkers that can be used in epidemiological studies. Department of  
Community Medicine, Mount Sinai School of Medicine, New York, NY, 2000.

New directions in biobehavioral research. Annual Meeting of American Society of Preventive Oncology, Bethesda, MD, 2000.

Stress and cancer: biobehavioral factors in familial risk. Department of Psychiatry, University of Michigan, Ann Arbor, MI, 2000.

Psychobiologic stress effects of a family history of cancer. Department of Health Policy, Mount Sinai School of Medicine, New York, NY, 2000.

Biobehavioral factors in familial risk of breast cancer. Binghamton University, Binghamton, NY, 2000.

Stress and cancer revisited. Department of Psychiatry. SUNY Stony Brook, Stony Brook, NY 2001.

Psychoneuroimmunology in cancer Prevention and control. The American Society of Preventive Oncology, New York, NY, 2001.

Breast cancer in the family: genes, stress and the immune system. NYU Medical Center, New York, NY, 2002.

Looking forward and back: family history as a stressor for healthy women. M.D. Anderson Cancer Center, Houston, TX, 2002.

Familial breast cancer risk: a psychobiological perspective. Fox Chase Cancer Center, Cheltenham, PA., 2002

Enhanced psychobiological responses to laboratory challenges among women at familial risk for breast cancer. Academy of Behavioral Medicine Research, Islamorada, FL, 2002.

Studies of high risk families and stress. Hillman Cancer Center at the University of Pittsburgh, Pittsburgh, PA, 2003.

Increased psychobiological stress responses in healthy women at familial risk for breast cancer. Georgetown University, Washington, DC, 2004.

Familial breast cancer risk: a biobehavioral analysis. Long Island University, Brooklyn, NY, 2004.

Familial breast cancer risk: a biobehavioral analysis. Ohio State University, Columbus, OH, 2004.

Psychobiological processes in familial breast cancer risk. Fred Hutchinson Cancer Research Center, Seattle, Washington, 2005.

## **CONCLUSIONS:**

At the conclusion of our final grant year, we have collected psychological data on 255 women and immune data on 135 women. As our preliminary analyses has revealed, the results of the research are consistent with the hypothesis that deficits in immune surveillance (e.g., as a result of stress) moderate the effects of mutations in primary susceptibility genes, the study could have important implications for the eradication of breast cancer. These results raise the possibility that appropriate interventions to reduce stress and increase the activity of immune surveillance mechanisms in women carrying mutations in primary susceptibility genes might delay the onset or prevent the development of breast cancer. In addition, this IDEA award has indirectly led to

the funding of five additional grants to study familial breast cancer risk, publication of 31 abstracts and articles by the investigators, and 18 invited presentations by the PI, thus establishing the team as leaders in the field of research on familial risk for breast cancer.

### **PERSONNEL:**

Dana Bovbjerg  
Karen Brown  
Nicole Cartalemi  
Jan Jeremias  
Christina Norman  
Janae Ostolaza  
Heiddis Valdimarsdottir

### **REFERENCES:**

- Antoniou AC, Pharoah PDP, McMullan G, Day NE, Stratton MR, Peto J, Ponder BJ, Easton DF. 2002. A comprehensive model for familial breast cancer incorporating BRCA1, BRCA2 and other genes. British Journal of Cancer, 86, 76-83.
- Bovbjerg DH, Valdimarsdottir HB. 2001. Interventions for healthy individuals at familial risk for cancer: Biobehavioral mechanisms for health benefits. In: Baum A, Anderson BL (eds), Psychosocial Interventions and Cancer, American Psychological Association Books: Washington, pp 305-320.
- Couch FJ. 2004. Genetic epidemiology of BRCA1. Cancer Biology and Therapy, 3(6), 509-514. Epub 2004 June 24.
- DeJong MM, Nolte IM, Te Meerman GJ, Van der Graaf WTA, Oosterwijk JC, Kleibeuker JH, Shaapveld M, De Vries EGE. 2002. Genes other than BRCA1 and BRCA2 involved in breast cancer susceptibility. Journal of Medical Genetics, 39, 225-242
- Dite G, Jenkins M, Southey M, Hocking J, Giles G, McCredit M, Ventor D, Hopper J. 2003. Familial risks, early-onset breast cancer, and BRCA1 and BRCA2 germline mutations. Journal of the National Cancer Institute, 95, 448-457.
- Kim Y, DuHamel K, Valdimarsdottir HB, Bovbjerg DH. Psychological distress among healthy women with family histories of breast cancer: Effects of recent life events. PsychoOncology 2005; 14(7):555-63.
- Lindberg NM, Wellisch DK. 2004. Identification of traumatic stress reactions in women at increased risk for breast cancer. Psychosomatics, 45(1), 7-16.
- Martin A, Weber B. 2000. Genetic and hormonal risk factors in breast cancer. Journal of the National Cancer Institute, 92, 1126-1135.
- Schnur JB, Valdimarsdottir HB, Montgomery GH, Nevid JS, Bovbjerg DH. 2004. Social constraints and distress among women at familial risk for breast cancer. Annals of Behavioral Medicine, 28, 142-148.
- Seegerstrom SC, Miller GE. 2004. Psychological stress and the human immune system: a meta-analytic study of 30 years of inquiry. Psychological Bulletin, 130:601-30.