

AD\_\_\_\_\_

AWARD NUMBER: DAMD17-02-1-0239

TITLE: Naked DNA Immunization for Prevention of Prostate Cancer in a Dunning Rat Prostate Tumor Model

PRINCIPAL INVESTIGATOR: Milcho Mincheff, M.D., Ph.D.

CONTRACTING ORGANIZATION: The George Washington University  
Washington, DC 20052

REPORT DATE: June 2006

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. REPORT DATE (DD-MM-YYYY)</b> 01-06-2006		<b>2. REPORT TYPE</b> Final		<b>3. DATES COVERED (From - To)</b> 1 Jun 2002 – 31 May 2006	
<b>4. TITLE AND SUBTITLE</b>  Naked DNA Immunization for Prevention of Prostate Cancer in a Dunning Rat Prostate Tumor Model				<b>5a. CONTRACT NUMBER</b>	
				<b>5b. GRANT NUMBER</b> DAMD17-02-1-0239	
				<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b>  Milcho Mincheff, M.D., Ph.D  E-Mail: <a href="mailto:mcamsm@gwumc.edu">mcamsm@gwumc.edu</a>				<b>5d. PROJECT NUMBER</b>	
				<b>5e. TASK NUMBER</b>	
				<b>5f. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>  The George Washington University Washington, DC 20052				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
				<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>	
<b>12. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for Public Release; Distribution Unlimited					
<b>13. SUPPLEMENTARY NOTES</b>					
<b>14. ABSTRACT</b>  After cloning a "truncated" (no peptide leader sequence) versions of the human prostate acid phosphatase (H-PAP-T), the human prostate-specific antigen (H-PSA-T) and the rat analogue of the human PSMA (R-"PSMA"-S), all plasmids were produced under GLP conditions. Their safety was tested in rats and the efficacy to stimulate T cells and to prevent development of transplantable tumors was shown in a rat model. Best protection was obtained following immunization with a cocktail containing all plasmids and a plasmid encoding rat GM-CSF. None of the animals immunized with the truncated plasmids developed antibodies against the native antigen while immunization with a plasmid encoding for an antigen that was secreted did. The antibodies were of mixed (Th1 and Th2) type (IgG1 and IgG2a). When priming was performed with the "truncated" version of the vaccines (HPSMA-T or HPSA-T), however and boosting with the "secreted" ones, the antibodies were mainly of the Th1 (complement-binding) type (IgG2a and IgG2b). The best protection was achieved when priming was performed with a plasmid encoding a xenogeneic protein and boosting with a plasmid encoding a syngeneic one. Genetic modification of rat tumor cells that led to expression of human (xeno) antigens, made them immunogenic, identifying other possible practical applications for vaccine design.					
<b>15. SUBJECT TERMS</b> plasmid DNA vaccine. PSMA, PAP, PSA, AT3B-1, GM-CSF, tumor protection, rat model					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b>
<b>a. REPORT</b> U	<b>b. ABSTRACT</b> U	<b>c. THIS PAGE</b> U			USAMRMC
			UU	93	<b>19b. TELEPHONE NUMBER (include area code)</b>

## TABLE OF CONTENTS

<b>Cover</b> .....	<b>1</b>
<b>SF 298</b> .....	<b>2</b>
<b>Introduction</b> .....	<b>4</b>
<b>Body</b> .....	<b>4</b>
<b>Key Research Accomplishments</b> .....	<b>16</b>
<b>Reportable Outcomes</b> .....	<b>17</b>
<b>Conclusions</b> .....	<b>18</b>
<b>References</b> .....	<b>20</b>
<b>Appendices</b> .....	<b>21</b>

## **INTRODUCTION:**

Recent research indicated that loss of expression of certain antigens was a common phenomenon during cancer progression and, therefore, targeting a single, defined tumor antigen would be disadvantageous to inducing an immune response against a broad spectrum of potential targets. Furthermore, the generation of various peptides when the immunization was performed against different tissue-specific antigens would eliminate possible differences in the affinity for peptide binding of different MHC alleles thus increasing the probability of creating a universal tissue-specific vaccine for the highly polymorphic human population. For this reason, we speculated that a cancer vaccine targeting a multitude of tissue-specific antigens was more likely to raise an effective immune response. We enlarged the target panel to include the prostate-specific membrane antigen (PSMA), the prostate acidic phosphatase (PAP) and the prostate-specific antigen (PSA). We planned to test the efficacy of immunization in an in vitro human model and the safety and efficacy for tumor protection in a Copenhagen rat prostate tumor model.

## **BODY OF THE REPORT**

### **1. Cloning of H-PAP, H-PSA and R-PSMA.**

The human prostate specific antigen and the human prostate acidic phosphatase were cloned by RT-PCR of total RNA from LNCaP using specific primers.

The rat equivalent of the extracellular domain of the human prostate specific membrane antigen (R-PSMA) was cloned by RT-PCR of total RNA isolated from rat brain tissue. It has 85.66% homology with the human PSMA at the nucleotide level.

For the SH-PSMA and SR-PSMA plasmids, the extracellular portion of the human PSMA or its rat equivalent respectively, were cloned into pSecTag2 vectors (Invitrogen), which provided the murine Ig k-chain leader sequence. The inserts from the obtained clones were then sub-cloned into pVAX-1 expression vectors (Invitrogen).

A stable transfectant of the AT3B1 cell line, co-expressing the truncated H-PSMA and the secreted H-PSA, was obtained by standard technique with the commercially available reagent Fugene. The level of expression was tested and confirmed by Western blot for PSMA and PSA and by ELISA for the PSA.

### **2. Transfection of AT3B-1 cells**

The rat AT3B-1 cell line expresses PAP. We have not been able to confirm expression at the protein level for PSMA and PSA in rat AT3-B1 cells. Since future immunotherapy experiments would use PSMA and PSA as targets for tumor development prevention, tumor cells needed to be stably transfected with both PSMA and PSA. A stable transfectant of the AT3B1 cell line, co-expressing the truncated H-PSMA and the secreted H-PSA, was obtained by standard technique with the commercially available reagent Fugene. Cells were initially transfected with PSA and selected with the antibiotic G418 (neomycin). The level of expression was tested and confirmed by Western blot and ELISA. The cells were then transfected for HS-PSMA and selected with the antibiotics zeocin and neomycin.

### **3. GLP-level production and purification of the plasmid vectors**

In order to proceed with the animal experiments as detailed in the Statement of Work of the original proposal, the plasmid vectors needed to be produced and purified under good laboratory practice

(GLP) conditions. This was achieved using standard Qiagen technology. The following plasmids were produced and purified: H-PSMA-T, R-“PSMA”-T, H-PSA, H-PSA-T, H-PAP-T and R”PSMA”-S.

We performed the following DNA Quality Control:

- 1 Endotoxin: Limulus amoebocyte lysate test: <100 E.U./mg DNA
- 2 DNA homogeneity: >90% on agarose gel assay.
- 3 RNA – no contamination on Agarose Gel and HPLC assays.
- 4 E.Coli Genomic DNA contamination: <50 ug/mg DNA by Southern blot.
- 5 Protein: Spectrophotometric scans between A220 and A320. Quantification is performed by use of the BCA protein test.
- 6 Bioburden assay (for DNA of a final concentration not more than 1 mg/ml); 1 colonies following United States Pharmacopeia (USP 23) or Ph. Eur. Suppl 3,1998.

The structure of the DNA was monitored during production by restriction endonuclease analysis and following production by direct sequencing. It was not monitored prior to application.

### **Characteristic Specifications Assay**

- 1 Appearance: Clear, colorless solution on visual inspection
- 2 Homogeneity of plasmid DNA >90% co-valently closed supercoiled circle using agarose gel
- 3 Sterility Sterile Assay according to USP 23
- 4 Presence of host DNA <5% (<0.05 mg/mg plasmid) by blot hybridization
- 5 Presence of Proteins <1%(<0.01 mg/mg plasmid) by BCA colorimetric assay
- 6 Presence of RNA and ssDNA: Not detectable on agarose gel and HPLC
- 7 Presence of endotoxins <0.1 EU/μg plasmid using LAL test according to USP 23
- 8 Plasmid identity: exclusive presence of target plasmid by restriction digest, sequencing

## **4. Development of the Copenhagen rat prostate tumor model**

Preliminary studies using the Copenhagen rat tumor prostate model showed uniform tumor development in rats that were injected subcutaneously with 100 000 AT3B-1<sup>PSMA,PSA</sup> cells. Healthy, Copenhagen 2331 male retired rat breeders were injected sub-cutaneously in the right flank with 1x10<sup>5</sup> tumor cells from the AT3B-1<sup>PSMA,PSA</sup> rat prostate carcinoma cell line. Injections were performed using 25-ga needles. Tumor size was evaluated every other day by measuring two perpendicular diameters by a caliper. The animals were observed twice daily for morbidity and mortality, and once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the whole duration of the study. Two weeks after tumor inoculation all rats were sacrificed, tumor were excised and measured. Necropsy was performed, tumor and sera was collected and liver and lungs were inspected macroscopically for metastases.

Good size (~2 cm<sup>3</sup>) subcutaneous tumors were established two weeks following injection. Tumors did not invade muscle and there were no skin ulcerations.

## **5. In vitro transfection of human dendritic cells with PSMA**

We next needed to test the efficacy of transfection of human dendritic cells with PSA and PAP and the ability of such transfected Using a commercially available transfection device from Amaxa, we could transfect differentiated dendritic cells with 20-40% efficiency. Such transfected dendritic cells stimulated in vitro autologous T cells to PSMA. T cells cytotoxicity was then tested against tumor cells or peptide-

pulsed T2 target cells. Both H-PSMA-T DCs and S-PSMA DCs generated antigen-specific cytotoxic T cell responses. The immune response was restricted towards one of four PSMA derived epitopes when priming and boosting was performed with S-PSMA. In contrast, T-PSMA transfected DCs primed T cells towards several PSMA derived epitopes. Subsequent repeated boosting with transfected DCs restricted the immune response to a single immunodominant epitope. The results have been incorporated in a manuscript which has been submitted for publication in Cancer Gene Therapy [1].

## **6. In vitro transfection of human dendritic cells with PAP and PSA**

In the last year's annual report we presented evidence that human dendritic cells, genetically engineered to express human prostate-specific membrane antigen, stimulate cytotoxic T cell response in autologous T cells [1]. We next needed to test the efficacy of transfection of human dendritic cells with PSA and PAP and the ability of such transfected cells to stimulate autologous T cells to lyse the target LNCaP cell line. Using a commercially available transfection device from Amaxa, we could transfect differentiated dendritic cells with 20-40% efficiency. Such transfected dendritic cells stimulated in vitro autologous T cells to PAP or PSA. T cells cytotoxicity was then tested against tumor cells or peptide-pulsed T2 target cells. Both T-PSMA H (tPSA)-DCs and S-PSA H (sPSA)-DCs generated antigen-specific cytotoxic T cell responses. The immune response was restricted towards one of four PSA derived epitopes when priming and boosting was performed with S-PSMA. In contrast, T-PSMA or T-PAP transfected DCs primed T cells towards several PSMA derived epitopes. Subsequent repeated boosting with transfected DCs restricted the immune response to a single immunodominant epitope. The immunodominance could be alleviated by depletion of CD25+ T cells prior to, or by GITR-L co-expression during, priming. The results have been incorporated in a manuscript which has been submitted for publication in Cancer Gene Therapy [2].

## **7. Titration of AT3B-1 and AT3B-1PSA for tumor cell development**

Preliminary studies using the Copenhagen rat tumor prostate model showed uniform tumor development in rats that were injected subcutaneously with 100 000 AT3B-1 cells. We needed to define the tumorigenicity of the PSA-transfected cell line and compare it to the parental cell line. To do that, healthy, Copenhagen 2331 male retired rat breeders were injected sub-cutaneously in the right flank with different numbers of tumor cells from the AT3B-1 or the AT3B-1<sup>PSA</sup> rat prostate carcinoma cell line (see below). For the AT3B-1 cell line, cells (passage 85, delivered 9/03) were purchased from the ATCC. Three rats per group were injected. Injections were performed using 25-ga needles. The animals were observed twice daily for morbidity and mortality, and once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the whole duration of the study. Tumor size was evaluated every other day by measuring two perpendicular diameters by a caliper. Three days after tumor formation or five weeks after tumor inoculation, all rats were sacrificed, tumor when present were excised and measured. Necropsy was performed, tumor and sera was collected and liver and lungs were inspected macroscopically for metastases.

1. Titration of AT3B-1<sup>PSA</sup> for tumor induction
  - $5 \times 10^6$  – tumor development in 10-12 days
  - $1 \times 10^6$  – tumor development in 15-19 days
  - $0.500 \times 10^6$  – tumor development 27-29 days
  - $0.250 \times 10^6$  – no tumors
  - $0.1 \times 10^6$  – no tumors

2. Titration of AT3B-1 cells for tumor induction
  - $1 \times 10^6$  – tumor development day 10-12
  - $0.5 \times 10^6$  – tumor development day 10-12
  - $0.25 \times 10^6$  – tumor development day 19-25
  - $0.1 \times 10^6$  – tumor development day 23-25

**Conclusion:**

1. Injection of the parent cell line (AT-3B1) leads to uniform development of tumors and there is a relationship between number of cell injected and the time of tumor appearance.
2. Injection of  $0.25$  and  $0.1 \times 10^6$  of PSA-transfected AT3B-1 cells does not result in tumor development.
3. Rats that have received  $0.25$  and  $0.1 \times 10^6$  of PSA-transfected AT3B-1 cells become immunized to human PSA and reject the transfected AT3B-1 cells
4. Rats, rejecting the AT3B-1<sup>PSA</sup> cells become immunized to other tumor associated antigens due to “antigen epitope spread”.
- 5.
6. Injection of a larger dose of the transfected cells leads to tumors, but their development is delayed.

For future experiments all immunized rats will be challenged with  $1 \times 10^6$  AT3B-1<sup>PSA</sup> cells.

**8. In vivo testing in a rat model the safety and efficacy of a combination of gene-based vaccines encoding human PSA, PAP and PSMA (months 10-30)**

The usual dose for DNA immunization of small rodents is 100 ug of plasmid DNA. Safety studies in mice have shown no toxicity in a dose range between 1ug and 100 ug plasmid DNA. After adjusting the dose per body weight, we assumed that injecting of 150 ug plasmid DNA per rat per immunization will have no toxic effect on the animal.

1. **Immunization cocktail A:** 50 ug H PSMA-T plasmid + 50 ug H PSA-T plasmid + 50 ug H PAP-T plasmid
2. **Immunization cocktail G:** 150 ug empty plasmid backbone
3. **Immunization cocktail H:** saline

Healthy, Copenhagen 2331 male retired rat breeders were used. All rats were immunized three times at 10-day intervals. The immunizations were intradermal and were performed by a standard intradermal injection technique with a small gauge (25-27 gauge) needle and an intradermal bevel. The needle was advanced into the intradermal region and the material was slowly injected while observing the formation of a “bleb” indicative of a proper injection. Volumes of up to 0.1 ml per site were injected.

A total of 104 animals were injected with the immunization cocktails (35 rats with cocktail A, 39 with cocktail G and 30 with cocktail H).

The animals were observed once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the study. When re-immunizations were performed, mice were re-immunized at weekly intervals. Two weeks after the last immunization, all mice either received a tumor inoculation or were sacrificed, blood and sera were collected and spleens harvested.

All immunizations were tolerated very well and there were no signs of toxicity after immunization. There were no differences in the overall condition of the animals between those who received the gene-based vaccines, the empty plasmid backbone or the saline injections.

Two weeks after the last immunization, all rats were injected sub-cutaneously in the right flank with  $1 \times 10^6$  tumor cells from the AT3B-1<sup>PSA</sup> rat prostate carcinoma cell line. Injections were performed using 25-ga needles. The AT3B-1<sup>PSA</sup> prostate cancer cell line is a non-metastatic cell line. Tumor size was evaluated every other day by measuring two perpendicular diameters by a caliper. The animals were observed twice daily for morbidity and mortality, and once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the whole duration of the study. Two weeks after tumor inoculation all rats were sacrificed, tumor were excised and measured. Necropsy was performed and kidney, especially proximal tubule, intestines, lungs, liver, prostate and brain were examined by histology. Spleen cells and sera were collected and tested for cytotoxicity against AT3B-1<sup>PSA</sup> cells or for antibody against the target antigen.

None of the 35 animals immunized with the immunization cocktail A (a mixture of HPSMA-T; HPAP-T and HPSA-T plasmids) developed tumors. In contrast, 37 of the 39 control rats immunized with the empty backbone and all 30 animals immunized with saline developed tumors 14-20 days after AT3B-1<sup>PSA</sup> cell inoculation. None of the tumor bearing rats showed distant metastases.

There were no macroscopical or histological changes in the proximal kidney tubule, intestines, lungs, liver, prostate or brain tissues of the immunized rats.

Spleen cells from the immunized animals were cytotoxic against the parent (AT3B-1) and the transfected cell line. The spleens were processed individually. A single cell suspension was obtained cutting and filtering the organ through a sterile cell strainer (70  $\mu\text{m}$ ; Becton Dickinson, Franklin Lakes, NJ). Thirty million splenocytes were resuspended in 10 ml RPMI 1640 containing 10% heat inactivated FCS, 50  $\mu\text{M}$  2-ME, 2 mM L-glutamine, 10 mM HEPES, 1 mM sodium pyruvate, 100 U/ml penicillin, and 100  $\mu\text{g}/\text{ml}$  streptomycin (culture medium) and cultured in a T25 flask with  $3 \times 10^6$  of mitomycin-c-treated of AT3B-1PSA cells in 10 ml culture medium. After 4 days, blasts were isolated on a density gradient, cultured for an additional day in medium supplemented with 20 IU/ml human rIL-2, and tested for cytolytic activity in a 4-hour 3H thymidine labeled DNA fragmentation assay [3] using the respective targets.

Interestingly, spleen cells from 18 of the 39 animals who were immunized with the empty backbone, and from 19 of the 30 animals who received saline, were cytotoxic against the transfected but not against the parent cell line. We interpret this as evidence that animals become immunized against PSMA after inoculation with the AT3B-1<sup>PSA</sup> cells. Immunization against PSA could also explain the reduced tumorigenicity of the PSA-transfected cell line (see 2. of the present report) but additional experiments are necessary to clarify this hypothesis.

The presence of anti-PSMA or anti-PSA antibodies in serum pre- and post-vaccination was analyzed by Western blot and ELISA. Anti-human PSA monoclonal antibodies sc-7316 and sc-7638 were obtained from Santa Cruz Biotechnologies, Inc. (Santa Cruz, CA, USA). Purified PSA protein was obtained from International Immuno-Diagnostics, Foster City, CA. Anti-human PSMA monoclonal antibodies Y-PSMA1 and Y-PSMA2 were obtained from Yes Biotech Laboratories Ltd (Mississauga, Ontario, Canada). Immunoblotting and electrophoretic assays of cell lysates and culture supernatants were performed using NuPage BioTris electrophoretic system (Invitrogen). Protein samples in loading buffer were heated at 70°C for 10 min and loaded on 10% Bis-Tris gels. After electrophoresis and electro-transfer, the nitrocellulose membrane was blocked with 1% casein in TBS/T for 40 min. The membranes were probed with poly- or monoclonal anti-PSMA Abs (see below) for 1 h at RT. Detection was



performed using goat anti-rabbit or anti-mouse IgG-HRP conjugates (Sigma) and visualized with WestPico Super Signal Chemo luminescent Substrate (Pierce) in accordance with the manufacturer's recommendations. Polyvinyl chloride 96-well microtiter plates (Dynatech Laboratories, Chantilly, VA) were incubated overnight at 4°C with a purified preparation of PSA (Vitro Diagnostics, Littleton, CO), as well as BSA or human serum albumin at 100 ng/well in 50 µl of PBS (pH 7.2). The wells were blocked for 1 h with PBS containing 1% BSA (assay buffer). Tested serum and control pooled rat serum were diluted in assay buffer and added to wells in triplicate in a volume of 50 µl/well. Purified human antimurine PSA-specific IgG antibody (Fitzgerald Industries, Concord, MA) was used as a positive control for PSA binding. Purified murine IgG was used as a negative control. After incubation overnight at room temperature, the wells were washed four times with assay buffer, and 50 µl of a 1:4000 dilution of peroxidase-conjugated goat antihuman IgG (Kirkegaard & Perry Laboratories, Gaithersburg, MD) were added to each well. A 1:2000 dilution of peroxidase-conjugated goat antimurine IgG (Kirkegaard & Perry) was used for the PSA antibody control. After incubation at 37°C for 1 h, wells were washed four times with assay buffer, and 100 µl each of the chromogen O-phenylenediamine dihydrochloride (Sigma, St. Louis, MO) and hydrogen peroxide was added to each well. After a 10-min incubation in the dark, the reaction was stopped with 25 µl of 4N NH<sub>2</sub>SO<sub>4</sub>. The absorbance of each well was measured at 490 nm using an ELISA microplate autoreader (Bio-Tek Instruments, Winooski, VT).

Sera from 17 of the 39 rats animals immunized with cocktail A (HPSMA-T; HPAP-T; HPSA-T) contained very low titer of antibodies against the target antigens that could be detected on Western against a lysate of COS-1 cells transfected with the respected target antigen. No antibodies against recombinant PSMA or PAP could be detected on ELISA. We interpret this as evidence that no antibodies against the native molecule of the target antigen develop when immunization is performed with the “truncated” (HPSMA-T and HPAP-T) vectors whose products are not secreted but proteasomally degraded. In contrast, however, all rats developed antibodies against the native PSA (detectable both on Western blots and ELISA). A possible explanation for this is that the rats become immunized and develop antibodies against the native PSA only after they are inoculated with the transfected tumor cells that secrete human PSA (AT3B-1<sup>PSA</sup>). Additional support for this hypothesis came from the fact that 22 of the 39 animals immunized with the empty backbone and 14 of the 30 who received saline, but were all inoculated with AT3B-1<sup>PSA</sup>, had antibodies against the native PSA. To confirm this observation, the following experiment was performed. Five rats received 3 immunizations with HPSA-T and were then sacrificed and their sera tested against PSA by Western blot and ELISA. None of the rats developed antibodies against native PSA detected on ELISA. Two of them had low titer of antibodies when tested against a lysate of COS-1 cells on Western blot.

### **Conclusion:**

1. Immunization with the mixture of the three plasmids (cocktail A) is safe and well tolerated.
2. Three immunizations with this cocktail provide excellent protection against tumor development induced by inoculation of  $1 \times 10^6$  AT3B-1<sup>PSA</sup> cells.
3. Spleen cells from the immunized rats are cytotoxic to both the parental and transfected cell lines.
4. Spleen cells from some control rats that have been inoculated with AT3B-1<sup>PSA</sup> cells are cytotoxic to the transfected but no to the parental cell line.
5. Inoculation of rats with AT3B-1<sup>PSA</sup> cells immunizes them against human PSA. The immune response could be detected as cytotoxicity against the transfected cell line or as antibodies against the native PSA detected by ELISA. This could explain the reduced tumorigenicity of the transfected cell line.
6. Some of the animals immunized with the “truncated” vectors whose products are retained in the cytosol and degraded in the proteasome, contain antibodies to the non-glycosylated forms of the antigen (protein core), but not against the natively folded glycoprotein.

Currently, results from the above experiments are being processed and a manuscript is in preparation.

## **9. In vivo testing the safety and efficacy of naked DNA immunization when single tissue-specific antigens (PSA, PSMA or PAP) are targeted for immunotherapy (months 18-36)**

Immunization cocktails:

- 1 Immunization cocktail C: 50 ug H PSA-T plasmid + 100 ug empty plasmid backbone
- 2 Immunization cocktail D: 50 ug H PAP-T plasmid + 100 ug empty plasmid backbone
- 3 Immunization cocktail F: 50 ug R PSMA-T plasmid + 100 ug empty plasmid backbone

The usual dose for DNA immunization of small rodents is 100 µg of plasmid DNA. Safety studies in mice have shown no toxicity in a dose range between 1ug and 100 µg plasmid DNA. After adjusting the dose per body weight, we assumed that injecting of 150 µg plasmid DNA per rat per immunization will have no toxic effect on the animal.

Healthy, Copenhagen 2331 male retired rat breeders were used. All rats were immunized three times at 10-day intervals. The immunizations were intradermal and were performed by a standard intradermal injection technique with a small gauge (25-27 gauge) needle and an intradermal bevel. The needle was advanced into the intradermal region and the material was slowly injected while observing the formation of a “bleb” indicative of a proper injection. Volumes of up to 0.1 ml per site were injected.

A total of 54 animals were injected with the immunization cocktails (18 rats per each cocktail).

The animals were observed once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the study.

Two weeks after the last immunization, all rats were injected sub-cutaneously in the right flank with  $1 \times 10^6$  tumor cells from the AT3B-1<sup>PSA</sup> rat prostate carcinoma cell line. Injections were performed using 25-ga needles. The AT3B-1<sup>PSA</sup> prostate cancer cell line is a non-metastatic cell line. Tumor size was evaluated every other day by measuring two perpendicular diameters by a caliper. The animals were observed twice daily for morbidity and mortality, and once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the whole duration of the study. Two weeks after tumor inoculation all rats were sacrificed, tumor were excised and measured. Necropsy was performed and kidney, especially proximal tubule, intestines, lungs, liver, prostate and brain were examined by histology. Spleen cells and sera were collected and tested for cytotoxicity against AT3B-1<sup>PSA</sup> cells or for antibody against the target antigen.

All immunizations were tolerated very well and there were no signs of toxicity after immunization. There were no differences in the overall condition of the animals between those who received the gene-based vaccines, the empty plasmid backbone or the saline injections.

Five control animals that were immunized with an empty plasmid + GM-CSF developed tumors 16 days following tumor inoculation. Immunization with any of the vectors that encoded a target antigen led to protection of animals against development of tumors. None of the 18 animals immunized with the

immunization cocktail C (H PSA-T plasmid) developed tumors. In contrast, 4 of the 18 rats immunized with cocktail D and 2 of the 18 rats immunized with cocktail F developed tumor. The difference between group C and D is statistically significant ( $\chi^2 = 4.5$ ;  $p < 0.05$ ) while between C and F is insignificant ( $\chi^2 = 2.11$ ;  $p < 0.2$ ). None of the tumor bearing rats showed distant metastases. All animals developed antibodies against PSA that were detected by ELISA, probably as a result of the inoculation of the PSA-expressing cell line. From experimentation performed last year (year 2 of the award) we found that AT3B-1<sup>PSA</sup> cells were immunogenic and injection of low numbers (less than  $0.5 \times 10^6$ ) of cells per rat did not lead to tumor formation, but to immunity against PSA which spread to other tumor antigens so that animals became protected against the non-transfected parental cell line. A no-cost extension of the award to study this phenomenon was requested and granted.

None of the animals immunized with the H PAP-T and H PSMA-T constructs developed antibodies against the target antigen (testing by Western blot and ELISA) – see # 4.

### **Conclusions:**

1. Immunization against the tissue-specific antigens PSMA, PSA and PAP is safe and well tolerated.
2. All 5 control animals immunized with the empty plasmid backbone-rat GM-CSF cocktail develop tumors 16 days after AT3B-1<sup>PSA</sup> cell inoculation.
3. None of the 18 animals immunized with the immunization cocktail C (H PSA-T plasmid) developed tumors.
4. The difference between the rate of tumor development in rats immunized against PSA, PSMA or PAP was statistically insignificant at the current number of rats tested.
5. All animals developed anti-PSA antibodies as a result of the inoculation of the AT3B-1<sup>PSA</sup> cells.
6. None of the animals immunized with H PSMA-T or H PAP-T developed antibodies against the target antigen.

### **10. Effectiveness of immunotherapy after immunization with plasmids encoding either syngeneic (R-“PSMA”-T) or xenogeneic (H-PSMA-T) proteins (months 24-36)**

Ten healthy, Copenhagen 2331 male retired rat breeders were immunized three times at 10-day intervals with a plasmid encoding the rat equivalent of the extracellular portion of the human PSMA (R PSMA-T plasmid) and rat recombinant GM-CSF. All immunizations were tolerated very well and there were no signs of toxicity after immunization. There were no differences in the overall condition of the animals between those who received the gene-based vaccines, the empty plasmid backbone or the saline injections.

Two weeks after the last immunization, all rats were injected sub-cutaneously in the right flank with  $1 \times 10^6$  tumor cells from the AT3B-1<sup>PSA</sup> rat prostate carcinoma cell line. Injections were performed using 25-ga needles. Tumor size was evaluated every other day by measuring two perpendicular diameters by a caliper. The animals were observed twice daily for morbidity and mortality, and once daily for clinical signs of toxicity (cage side). A more thorough examination was performed at weekly intervals. Animals were monitored for weight on weekly intervals throughout the whole duration of the study. Two weeks after tumor inoculation all rats were sacrificed, tumor were excised and measured. Necropsy was performed and kidney, especially proximal tubule, intestines, lungs, liver, prostate and brain were examined by histology. Spleen cells and sera were collected and tested for cytotoxicity against AT3B-1<sup>PSA</sup> cells or for antibody against the target antigen. The results from this study were compared with the results in which the animals were immunized prior to tumor cell inoculation with the H PSMA-T plasmid (see #2).

None of the animals developed tumors until day 16 after tumor inoculation and were sacrificed. In contrast, 2 of the 18 rats, immunized with H PSMA-T developed tumors, but the difference between the two groups is not statistically significant. Spleen cells from animals from both groups were cytotoxic against the AT3B-1<sup>PSA</sup> rat prostate carcinoma cell line. No antibodies against PSMA were detected by ELISA or Western blot.

### **Conclusions:**

1. Immunization with the R PSMA-T is safe and well tolerated.
2. Animals, immunized with the R PSMA-T plasmid are protected against development of AT3B-1<sup>PSA</sup>-induced tumors.
3. No statistically significant difference in tumor protection is observed when animals immunized with H PSMA-T are compared to animals immunized with the R PSMA-T.
4. Rats immunized with the R PSMA-T do not develop antibodies against the target antigen.

## **11. To compare different priming and boosting strategies for tumor prevention when immunizing against PSMA**

Since the AT3B-1 cells that were transfected with human the PSA elicited a tumor response, initially we decided to work for these experiments with a non-transfected parental cell line. We found, however, that both rat prostate tissue and AT3B-1 cell lysates expressed low level of PSMA when tested by PCR (data not shown). Since PSMA expression is up-regulated in human prostate cancer cells, we decided to transfect the AT3B-1 cells with the HPSMA-S, a vector that contains the murine Ig k-chain leader sequence so that the extracellular portion of the human PSMA is glycosylated and secreted<sup>1</sup>. We did not transfect the cells with the full length PSMA since our goal was to study T cell-mediated and not antibody-mediated tumor protection and we felt that if PSMA was expressed as a membrane protein by the transfected AT3B-1 cells, some of the protection might have been antibody mediated. This would have been true especially when immunization was performed with DNA constructs that led to anti-PSMA antibody development (see below).

### **3.1 AT3B-1 transfection with HPSMA-S.**

Monolayers of AT3B-1 cells (CRL-2375; ATCC, Manassas, VA) were transfected with the HPSMA-S using the FuGENE6 transfection reagent (Roche Diagnostics Corporation, Indianapolis, IN) and assayed for PSMA production by Western blot. The cells were seeded in 6-well tissue culture plates (Nunc, Denmark) at  $1.5 \times 10^5$  cells per well and grown to 50-70% confluence in DMEM supplemented with 25 mM HEPES (pH 7.5), 1 mM sodium pyruvate, 3.7 g/L sodium bicarbonate, 100 µg/ml penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin B and 10% (v/v) of fetal bovine serum. COS-1 or AT3B-1 cells were transfected with 1.5 µg of plasmid DNA pre-condensed with 4.5 µl of FuGENE 6 reagent in serum-free DMEM for 30 min at room temperature. Cells were then grown for 72 h in complete DMEM and then harvested.

The transfected cells with stable expression of HPSMA-S were selected and cloned after growth in zeocin-containing medium (0.1 mg/ml). We obtained two clones of stably transfected AT3B-1 cells – YM1 and YM2, both of which are high expressors and PSMA can be detected in the culture medium. A significant portion of the expressed product is degraded in the proteasome and could be detected in cell

lysates only in the presence of proteasomal inhibitor lactacystin. All further experimentation was performed with the YM2 clone.

All non-immunized rats developed tumors 12 days after inoculation of  $1 \times 10^5$  YM2 cells. We found that the protection of the immunized animals from development of tumor, however, depended on the amount of YM2 cells inoculated during tumor challenge. Partial tumor protection could be seen when  $2 \times 10^5$  or  $3 \times 10^5$  tumor cells were injected. All immunized animals developed tumors when the tumor dose per animal was higher than  $4 \times 10^5$  cells. In an attempt to define the best strategy of vaccination, we explored different priming and boosting strategies. A total of 40 rats were included in these experiments.

All immunizations were performed with plasmid DNA cocktails that contained rat GM-CSF. All animals were challenged with  $2 \times 10^5$  cells. Rats were sacrificed on the day tumor was detected. All tumor-free animals were sacrificed at day 16 following tumor cell inoculation. Ten rats were included per group. The best protection following immunization involved priming with H PSMA-T and boosting with R PSMA-T and all the animals were tumor free 16 days after tumor inoculation (figure 4; circles). In contrast, when priming was performed with HPSMA-S and boosting with R PSMA-T, three rats developed tumors on day 12, while the rest of the rats (66%) remained tumor free. The difference between these two groups was not statistically significant ( $\chi^2 = 3.52$  at  $p < 0.1$ ). Spleen cells from the HPSMA-S-immunized animals, however, gave the strongest proliferative response after stimulation with recombinant PSMA (data not shown). Priming and boosting with R PSMA-T resulted in only partial protection – three rats developed tumor on day 10, and 3 rats on day 11. Four of those rats (34%) remained tumor free until day 16 (figure 4, diamonds). The difference between this group and the control group, though, was statistically significant ( $\chi^2 = 5$  at  $p < 0.05$ ) since all control rats developed tumors by day 12 after tumor inoculation.

## Conclusions:

1. Priming with a plasmid encoding a xenogeneic construct and boosting with a plasmid encoding a syngeneic protein is safe and well tolerated.
2. Priming with a plasmid encoding a xenogeneic protein followed by re-immunization with a plasmid encoding a syngeneic leads to the best protection against AT3B-1<sup>PSMA</sup>-induced tumors.
3. Spleen cells from animals immunized with the HPSMA-S plasmid give the strongest proliferative response to PSMA.

## 12. To compare the development of anti-PSMA antibodies in a Dunning rat prostate cancer model between groups of animals that are immunized with a truncated R"PSMA"-T plasmid and R-"PSMA"-S plasmid

Healthy, Copenhagen 2331 male retired rat breeders were used. All rats were immunized three times at 10-day intervals. The immunizations were intradermal and were performed by a standard intradermal injection technique with a small gauge (25-27 gauge) needle and an intradermal bevel. The needle was advanced into the intradermal region and the material was slowly injected while observing the formation of a "bleb" indicative of a proper injection. Volumes of up to 0.1 ml per site were injected.

A total of 20 animals were injected with an immunization cocktail that contained plasmid DNA and  $9 \mu\text{g}/\text{m}^2$  recombinant rat GM-CSF. Ten rats were immunized with R PSMA-S and ten rats with R PSMA-T. Two weeks following the last immunization, the rats were inoculated with the with  $1 \times 10^5$  YM2 cells. None of the immunized animals developed tumors.

As observed in earlier studies, rats immunized with “truncated” constructs (H PSMA-T or R PSMA-T) did not develop antibodies to PSMA as detected by ELISA or Western blot, no matter whether GM-CSF was present in the immunization cocktail or not. No antibodies against PSMA (detection by Western blot or ELISA) developed following immunization with plasmid, encoding the secreted PSMA (R PSMA-S), or with a recombinant protein, when GM-CSF was absent from the immunization cocktail. Immunization with R PSMA-S in the presence of GM-CSF led to formation of antibodies against the native protein (detected by ELISA). Antibodies, developing following immunization with R PSMA-S or GCPII, were of mixed Th2 and Th1 type, since both IgG1 and IgG2a subtypes were detected. On the contrary, priming with H PSMA-T or R PSMA-T and boosting with R PSMA-S led to formation of Th1, cytotoxic antibodies (IgG2a and IgG2b) with relatively high titer.

**Conclusions:**

1. Immunization with R PSMA-T does not lead to development of antibodies against the target antigen.
2. Immunization with R PSMA-S results in the development of antibodies to the target antigen when GM-CSF is used as an adjuvant. The antibodies are of mixed Th2 and Th1 type.
3. Priming with H PSMA-T or R PSMA-T and boosting with R PSMA-S led to formation of Th1, cytotoxic antibodies (IgG2a and IgG2b) with relatively high titer.

**13. To compare the development of anti-PSA antibodies in a Dunning rat prostate cancer model with already established tumors after immunization with full length or human truncated PSA constructs (months 24-36)**

In experiments performed in year two of the award, we found that AT3B-1<sup>PSA</sup> cells were immunogenic and that their inoculation per se led to development of anti-PSA antibodies. As a result of this, the experiments listed in Task 6 were modified and no tumor cell inoculation prior to immunization was performed. Healthy, Copenhagen 2331 male retired rat breeders were used. All rats were immunized three times at 10-day intervals. The immunizations were intradermal and were performed by a standard intradermal injection technique with a small gauge (25-27 gauge) needle and an intradermal bevel. The needle was advanced into the intradermal region and the material was slowly injected while observing the formation of a “bleb” indicative of a proper injection. Volumes of up to 0.1 ml per site were injected.

A total of 57 animals (19 rats per group) were immunized 3 times at 10-day intervals with either H PSA-T, HPSA-S or empty plasmid cocktails. All immunizations were tolerated very well and there were no signs of toxicity after immunization. There were no differences in the overall condition of the animals between those who received the gene-based vaccines or the empty plasmid backbone. Two weeks after the last immunization, all animals were sacrificed, spleen cells and sera were collected and frozen until further use. The sera were tested for anti-PSA antibodies by Western blot and ELISA. None of the rats immunized with H PSA-T or the empty plasmid developed antibodies against PSA. In contrast, all animals immunized with HPSA-S developed antibodies against the target antigen that were of mixed Th1-Th2 type (IgG1 and IgG2a).

**Conclusion:**

1. No anti-PSA antibodies are detected in the sera of animals immunized with the H PSA-T plasmid.
2. Immunization with HPSA-S plasmid in the presence of rat GM-CSF leads to development of antibodies against PSA that are of mixed Th1-Th2 type.

**14. To study antigenic spread during inoculation of rats with “xenogenized” tumor cell lines**

In experiments performed in year two of the award, we found that that injection into Copenhagen rats of low numbers of AT3B-1<sup>PSA</sup> cells (AT3B-1 cells transfected with the human prostate-specific antigen, also called “xenogenized” tumor cells) led to development of anti-PSA antibodies. Such rats reject the xenogenized cells and did not form tumors. During rejection however, this immunity spread to other tumor-associated antigens present in the parental (non-transfected) cell line. As a consequence rats, that had once rejected the xenogenized tumor cells, rejected the parental cells also. This discovery opened new venues for designing tumor vaccines.

## KEY RESEARCH ACCOMPLISHMENTS:

During the period covered by the award we have:

- 1 Cloned of H-PAP, H-PSA and R-PSMA.
- 2 Designed and production of constructs for H-PAP and H-PSA that have no leader sequence and whose expression products are cytosolically retained and degraded in the proteasome (H-PSA-T and H-PAP-T)
- 3 Designed and production of constructs that lead to secreted H-PSMA (SHPSMA) and secreted R-PSMA (SR-PSMA).
- 4 Produced and purified GLP-grade gene-based vaccines for use in the Copenhagen rat prostate tumor protection model (H-PSMA-T, R-“PSMA”-T, H-PSA, H-PSA-T, H-PAP-T and R”PSMA”-S)
- 5 Obtained a stable transfectant of the AT3B-1 cell line co-expressing the human PSA and the human PSMA (AT3B-1<sup>PSA,PSMA</sup>)
- 6 Developed a tumor development model in a Copenhagen rat with the stably-transfected tumor cells AT3B-1<sup>PSA,PSMA</sup>
- 7 Studied the efficacy of human dendritic cells transfected with the H-PSMA plasmid to stimulate autologous T cells in vitro; tested their ability to lyse target tumor cells or HLA-identical cells that have been pulsed with PSMA plasmids
- 8 Proven in an animal model the safety of naked DNA immunization with a mixture of three “truncated” plasmid vectors that encode for the human PSMA, PSA and PAP that are not glycosylated but are degraded in the proteasome of the cells that express them.
- 9 Proven in rats the efficacy of immunization with the mixture of the gene-based vaccines to prevent AT3B-1<sup>PSA</sup> induced tumor development.
- 10 Shown that spleen cells derived from rats previously immunized with the mixture of the three “truncated” plasmid vectors are cytotoxic to both the parental (AT3B-1) and the transfected (AT3B-1<sup>PSA</sup>) cell lines.
- 11 Shown that AT3B-1 cells stably transfected with the human PSA (AT3B-1<sup>PSA</sup>) have decreased tumorigenicity in rats when compared to the parental cell line.
- 12 Shown that sub-cutaneous inoculation of control (non-immunized) rats with the AT3B-1<sup>PSA</sup> immunizes them against human PSA. Immune responses can be detected by cytotoxicity against the AT3B-1<sup>PSA</sup> cells or by ELISA with recombinant PSA.
- 13 Shown that immunization with the “truncated” vectors whose products after expression are degraded in the proteasome lead to development of low titer antibodies against the protein core.
- 14 Shown that injection of small numbers of AT3B-1<sup>PSA</sup> cells leads to anti-PSA immunity and rejection of the AT3B-1<sup>PSA</sup> cells
- 15 Shown that immunity to human PSA spreads to other tumor-associated antigens during rejection and such animals become immune against the parental, non-transfected cell line.



## REPORTABLE OUTCOMES:

1. Gene-based vaccines for immunotherapy of cancer:
  - H-PSA – a plasmid vector that contains an expression cassette for the human PSA
  - H-PSA-T – a plasmid vector that contains an expression cassette for the human PSA with no signal peptide
  - H-PAP – a plasmid vector that contains an expression cassette for the human PAP
  - H-PAP-T – a plasmid vector that contains an expression cassette for the human PAP with no signal peptide
  - R-PSMA – a plasmid vector that contains an expression cassette for the mouse analogue of the extracellular domain of the human PSMA
  - SR-PSMA – a plasmid vector that contains an expression cassette for the mouse analogue of the extracellular domain of the human PSMA with a signal peptide
  - SH-PSMA – a plasmid vector that contains an expression cassette for the extracellular domain of the human PSMA with a signal peptide
2. AT3B-1<sup>PSMA,PSA</sup> – a rat AT3B-1 cell line that secretes the extracellular domain of the human PSMA and the human PSA.
3. AT3B-1<sup>PSMA</sup> (YM2) cell line which is the AT3B-1 cell line transfected to secrete the extracellular portion of the human prostate specific membrane antigen
4. Mincheff M, Zoubak S, Altankova I, Tchakarov S, Makogonenko Y, Botev C, Ignatova I, Dimitrov R, Madarzhieva K, Hammett M, Pomakov Y, Meryman H, Lissitchkov T. Human dendritic cells genetically engineered to express cytosolically retained fragment of prostate-specific membrane antigen prime cytotoxic T-cell responses to multiple epitopes. *Cancer Gene Ther.* 10:907;2003
5. Mincheff M, Zoubak S, Altankova I, Tchakarov S, Pogribnyy P, Makogonenko Y, Botev C, Meryman, HT. Depletion of CD25+ cells from human T-cell enriched fraction eliminates immunodominance during priming and boosting with genetically modified dendritic cells. *Cancer Gene Therapy* 12:185;2005
6. Mincheff M, Makogonenko Y, Zoubak S. Immune responses against PSMA after gene-based vaccination for immunotherapy: A. Results from immunizations in animals. *Cancer Gene Therapy* 2005 (in press)

## CONCLUSIONS:

1. The research has progressed according to the approved Statement of Work. There have been no problems so far connected with the experimental design except for the fact that PSMA expression in the AT3B-1 cell line has been detected only at the mRNA but not protein level. All tumor protection studies will be performed with the AT3B-1<sup>PSMA</sup> cell line.
2. Dendritic cells transfected with a construct whose product is retained in the cytosol and degraded in the proteasome, prime to both dominant and subdominant epitopes.
3. Early CD25+ cell depletion during priming in vitro enhances priming to sub-dominant epitopes.
4. Co-expression of GITR-L during priming may alleviate immunodominance.
5. Injection of the parent cell line (AT-3B1) leads to uniform development of tumors and there is a relationship between number of cell injected and the time of tumor appearance.
6. Injection of 0.25 and 0.1 x10<sup>6</sup> of PSA-transfected AT3B-1 cells does not result in tumor development.
7. Injection of a larger dose of the transfected cells leads to tumors, but their development is delayed.
8. Immunization with the mixture of the three plasmids (cocktail A) is safe and well tolerated.
9. Three immunizations with this cocktail provide excellent protection against tumor development induced by inoculation of 1x10<sup>6</sup> AT3B-1<sup>PSA</sup> cells.
10. Spleen cells from the immunized rats are cytotoxic to both the parental and transfected cell lines.
11. Spleen cells from some control rats that have been inoculated with AT3B-1<sup>PSA</sup> cells are cytotoxic to the transfected but no to the parental cell line.
12. Inoculation of rats with AT3B-1<sup>PSA</sup> cells immunizes them against human PSA. The immune response could be detected as cytotoxicity against the transfected cell line or as antibodies against the native PSA detected by ELISA. This could explain the reduced tumorigenicity of the transfected cell line.
13. None of the animals immunized with H PSMA-T or H PAP-T developed antibodies against the target antigen. Immunization with the R PSMA-T is safe and well tolerated.
14. Animals, immunized with the R-PSMA plasmid are protected against development of T3B-1<sup>PSA</sup>-inuced tumors.
15. No statistically significant difference in tumor protection is observed when animals immunized with H PSMA-T are compared to animals immunized with the R PSMA-T.
16. Rats immunized with the R PSMA-T do not develop antibodies against the target antigen.
17. Priming with a plasmid encoding a xenogeneic construct and boosting with a plasmid encoding a syngeneic protein is safe and well tolerated.
18. Priming with a plasmid encoding a xenogeneic protein followed by re-immunization with a plasmid encoding a syngeneic leads to the best protection against AT3B-1<sup>PSMA</sup>-induced tumors.
19. Spleen cells from animals immunized with the HPSMA-S plasmid give the strongest proliferative response to PSMA.
20. Immunization with R PSMA-T does not lead to development of antibodies against the target antigen.
21. Immunization with R PSMA-S results in the development of antibodies to the target antigen when GM-CSF is used as an adjuvant. The antibodies are of mixed Th2 and Th1 type.
22. Priming with H PSMA-T or R PSMA-T and boosting with R PSMA-S led to formation of Th1, cytotoxic antibodies (IgG2a and IgG2b) with relatively high titer.
23. No anti-PSA antibodies are detected in the sera of animals immunized with the H PSA-T plasmid. A "truncated" version of gene-based vaccines, therefore, can be safely used when targeting tumor markers that are used for monitoring disease such as PSA and PAP.
24. Immunization with HPSA-S plasmid in the presence of rat GM-CSF leads to development of antibodies against PSA that are of mixed Th1-Th2 type.

**The “so what” section:** Loss of expression of certain antigens is a common phenomenon during cancer progression and, therefore, targeting a single, defined tumor antigen will be disadvantageous to inducing an immune response against a broad spectrum of potential targets. Our results showed that a combination of vaccines that target three different auto-antigens was safe to administer and very effective in prevention of tumor development. Since immunization with “truncated” DNA sequence did not lead to formation of antibodies against the native antigens, DNA vaccines encoding truncated PSA could be used for immunotherapy since no antibodies that would interfere with PSA testing would be developed. Genetic modification of rat tumor cells that led to expression of human (xeno) antigens, made them immunogenic. Due to antigen epitope spread during elimination of these cells, the inoculated animals became immunized to antigens present in the parental cell line, thus identifying possible practical applications for vaccine design.

## REFERENCES:

1. Mincheff M *et al.* Human dendritic cells genetically engineered to express cytosolically retained fragment of prostate-specific membrane antigen prime cytotoxic T-cell responses to multiple epitopes. *Cancer Gene Ther* 2003; **10**: 907-917.
2. Mincheff M *et al.* Depletion of CD25+ cells from human T-cell enriched fraction eliminates immunodominance during priming with dendritic cells genetically modified to express a secreted protein. *Cancer Gene Ther* 2005; **12**: 185-197.
3. Mincheff M *et al.* Immune responses against PSMA after gene-based vaccination for immunotherapy-A: results from immunizations in animals. *Cancer Gene Ther* 2006;13:436-44

## **APPENDICES:**

### **A. Manuscript published in *Cancer Gene Therapy*, 10:907; 2004**

Human Dendritic Cells Genetically Engineered to Express Cytosolically-Retained Fragment of Prostate-Specific Membrane Antigen Prime Cytotoxic T Cell Responses to Multiple Epitopes

Milcho Mincheff<sup>1,4</sup>, M.D., Ph.D. Serguei Zoubak<sup>1</sup>, Ph.D., Iskra Altankova<sup>2</sup>, M.D., D.Sci., Stoyan Tchakarov<sup>3</sup>, M.D., Ph.D., Yevgen Makogonenko<sup>1</sup>, D.Sci., Chavdar Botev<sup>4</sup>, M.D., Irena Ignatova<sup>4</sup>, Rosen Dimitrov<sup>4</sup>, M.D., Kalina Madarzhieva<sup>4</sup>, M.D., Michael Hammett<sup>1</sup>, Yavor Pomakov<sup>2</sup>, M.D., Harold T. Meryman<sup>1</sup>, M.D., T. Lissitchkov<sup>4</sup>, M.D.

The George Washington University Medical Center, Washington DC  
Central Laboratory for Immunology, St. Ivan Rilski Hospital, Sofia, Bulgaria  
Department of Urology, St. Ann University Hospital, Sofia, Bulgaria  
National Institute of Hematology and Transfusion Medicine, Sofia, Bulgaria

Corresponding author:

Milcho Mincheff

Head, Tumor Immunology Laboratory

Department of Medicine

The George Washington Medical Center

2300 Eye Street, N.W., Ross Hall 705

Washington, DC 20037

Telephone 202 994 7765; fax 202 994 0465

E-mail: [mcamsm@gwumc.edu](mailto:mcamsm@gwumc.edu)

Running title: Human Dendritic Cells Genetically Engineered

Keywords: PSMA. Gene-based vaccine, immunodominance, CTLA-4

Supported in part by grant N00014-00-1-0787 from the Office of Naval Research.

Supported in part by award No DAMD17-02-1-0239. The U.S. Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick, MD 21702-5014 is the awarding and administering acquisition office.

The content of the information does not necessarily reflect the position or the policy of the Government, and no official endorsement should be inferred. For purpose of this article, information includes news releases, articles, manuscripts, brochures, advertisements, still and motion pictures, speeches, trade association proceedings etc.

## **ABSTRACT**

The ability of two plasmid DNA vaccines to stimulate lymphocytes from normal human donors and to generate antigen-specific responses is demonstrated. The first vaccine (truncated; tPSMA) encodes for only the extracellular domain of PSMA. The product, expressed following transfection with this vector, is retained in the cytosol and degraded by the proteasomes. For the “secreted” (sPSMA) vaccine, a signal peptide sequence is added to the expression cassette and the expressed protein is glycosylated and directed to the secretory pathway. Monocyte-derived dendritic cells (DCs) are transiently transfected with either sPSMA or tPSMA plasmids. The DCs are then used to activate autologous lymphocytes in an in vitro model of DNA vaccination. Lymphocytes are boosted following priming with transfected DCs or with peptide-pulsed monocytes. Their reactivity is tested against tumor cells or peptide-pulsed T2 target cells. Both tPSMA DCs and sPSMA DCs cells generate antigen-specific cytotoxic T cell responses. The immune response is restricted towards one of the four PSMA derived epitopes when priming and boosting is performed with sPSMA. In contrast, tPSMA transfected DCs prime T cells towards several PSMA derived epitopes. Subsequent repeated boosting with transfected DCs, however, restricts the immune response to a single epitope due to immunodominance.

## INTRODUCTION

Immunotherapy of prostate cancer could be a safe, non-invasive, relatively inexpensive procedure that can avoid bowel and bladder injury and impotence often resulting from surgical, cryosurgical or radiation therapy. Several groups have recently reported on the safety of DNA vaccines for immunizations against tumor antigens. {Mincheff, 2000 #698;Mincheff, 2001 #697;Weber, 1998 #1237} A possible target for prostate cancer immunotherapy is the prostate specific membrane antigen (PSMA), also known as glutamate carboxypeptidase II (GCPII). {Israeli, 1993 #189;Wright, 1996 #238;Chang, 2000 #220;Chang, 1999 #219;Heston, 1996 #234} PSMA expression is normally restricted to the prostate gland, brain tissue, jejunum and proximal kidney tubules. {Silver, 1997 #208;Shneider, 1997 #1249;Halsted, 2000 #1248} Its expression is increased nearly 10-fold in prostate cancer cells and is also found in tumor but not normal neovasculature. {Chang, 1999 #219;Chang, 1999 #213}

The main effectors in anti-tumor immunity are CD8<sup>+</sup> cytotoxic T cells that recognize tumor-associated antigen-derived peptides in association with major histocompatibility complex (MHC) class I molecules. {Sprent, 1989 #1245;Bennink, 1993 #1246;Restifo, 1993 #1247} Gene-based vaccination in its current mode of application is effective in breaking tolerance to a self-antigen, but the response is narrow and is restricted to few of the potential epitopes. This presents a problem in vaccinology since loss of an MHC haplotype that participates in the conformation of the T cell antigen, or point mutation in the recognized sequence would result in ineffective immune surveillance. {Dunn, 2002 #1276;Schreiber, 2002 #1085;Ikeda, 2002 #1277} New vaccines and/or new methods of immunizations need to be developed for those instances. These hopefully will raise responses to subdominant determinants so that the selection of tumor escape variants that fail to express immunodominant epitopes will be prevented. {Dunn, 2002 #1276}

Numerous factors combine to establish an immunodominance hierarchy, {Yewdell, 1999 #1096} among them the ineffective generation and transport of sub-dominant epitopes by antigen-presenting cells (APCs). Since proteasomal degradation is the main source of antigenic fragments destined for MHC presentation, {Bennink, 1993 #1246} we speculated that purposeful cytosolic retention of newly synthesized tumor-associated antigens in genetically manipulated antigen presenting cells may increase both the quantity and the diversity of such fragments. PSMA is a type II transmembrane glycoprotein which is comprised of 750 amino acids. {Pinto, 1996 #216;Rawlings, 1997 #686} It lacks a signal peptide sequence and we speculated that elimination of sequences for its transmembrane region might impede the translocation of the encoded product to the endoplasmic reticulum. Such product should not be N-glycosylated, should be retained in the cytosol and rapidly degraded in the proteasome. In theory, DC transfected with such “truncated” sequences, may have an advantage of presenting “subdominant” antigenic determinants that otherwise may not be generated at sufficient density to prime antigen-specific cytotoxic T cell responses. The following experiments were designed to test this hypothesis in an in vitro immunization system with human cells.

## MATERIALS AND METHODS

All human cellular material used in these experiments was obtained following informed consent through protocols approved by the local Committee for Bioethics (Bulgaria) or the Investigational Review Board (IRB) at George Washington University Medical Center in Washington, DC.

### *tPSMA and sPSMA plasmids construction*

The cDNA encoding the extracellular portion (AA 44-750) of the human PSMA (XC-PSMA) was cloned into the pCR2.1 vector (Invitrogen, Carlsbad, CA) after RT-PCR of total mRNA from the human prostate cancer cell line LNCaP (CRL 1740, ATCC). The forward primer introduced a NotI cloning site and a Kozak sequence with a Met codon (GCCACCATG) into the 5'-end of XC-PSMA. The tPSMA plasmid for the transfection experiments was obtained by NotI-XhoI sub-cloning of XC-PSMA into a pVAX1 mammalian expression vector (Invitrogen, Carlsbad, CA). A secretable variant of the tPSMA plasmid – the sPSMA plasmid - was obtained by fusion of the XC-PSMA with a murine Ig k-chain leader sequence. The insert from tPSMA plasmid was sub-cloned by BamHI-XhoI into the mammalian expression vector pSecTag2A (Invitrogen, Carlsbad, CA) providing the murine Igk-chain leader sequence (clone 96). The single SfiI cloning site of the vector was used to fuse the XC-PSMA in-frame with the leader. The 5'-portion of the XC-PSMA between start Met and a single HpaI site was re-amplified in order to introduce SfiI site (Met codon was not included). The SfiI-HpaI fragment of clone 96 was replaced with the PCR product pre-digested with the same restriction endonucleases and the NruI-XhoI fragment from obtained construct was moved to the pVAX1 vector. The inserts in both constructs are under the regulation of a human cytomegalovirus (CMV) immediate-early promoter/enhancer and a bovine growth hormone polyadenylation signal. The plasmid DNA specifications include endotoxin content below 0.1 EU per microgram of DNA; lack of detectable amounts of bacterial RNA, genomic DNA or ssDNA as determined by agarose-gel electrophoresis; less than 10 microgram of protein per 1 mg of plasmid DNA as determined by colorimetric assay (Bio-Rad, Hercules, CA).

### COS-1 transfection

Expression of PSMA constructs was performed in Cos-1 cells (ATCC). Monolayers were transfected with FuGENE 6 transfection reagent (Roche) and assayed for PSMA production by Western blot. Cos-1 cells were seeded in 6-well tissue culture plates (Nunc, Denmark) at  $1.5 \times 10^5$  cells per well and grown to 50-70% confluence in DMEM supplemented with 25 mM HEPES (pH 7.5), 1 mM sodium pyruvate, 3.7 g/L sodium bicarbonate, 100 µg/ml penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin B and 10% (v/v) of heat inactivated fetal bovine serum. Cos-1 cells were transfected with 1.5 µg of plasmid DNA pre-condensed with 4.5 µl of FuGENE 6 reagent in serum-free DMEM for 30 min at room temperature. Cells were then grown for 72 h in supplemented DMEM and then harvested.

In the proteasome inhibition studies, lactacystin (Sigma) was added to the culture media (final concentration 10 µM) 24 h before harvesting.

Cells were harvested by gentle scraping after double wash with 2 ml of cold PBS and adding to each well of 0.25 ml of cold lysis buffer (0.5 M NaCl, 1% triton X-100, 0.2% Tween 20, 50 mM HEPES, pH 7.0). Lysates were transferred to Eppendorf tubes and homogenized by repeated pipetting on ice.

For detection of secreted PSMA, the serum containing DMEM was removed 48 h after transfection, the cells were washed twice with 2 ml of PBS, serum free DMEM (2 ml per well) was added and cells were incubated for additional 24 h in the 6-well plates. After collection of the medium, the cell debris was removed by centrifugation (35,000 g, 20 min) and supernatants were concentrated with Centricon-50 centrifuge filtering device (Millipore) and stored at -30°C until further use.

### Deglycosylation of PSMA



The deglycosylation of PSMA proteins was carried out by treatment of 25 mg protein samples with 5 units of glycopeptidase F from *Chryseobacterium Meningosepticum* (Sigma) in 50 mM phosphate buffer (pH 7.5) containing 0.1% (w/v) SDS, 50 mM  $\beta$ -mercaptoethanol, 0.75% (v/v) Triton X-100 for 4 h at 37°C. {Holmes, 1996 #298}

#### Electrophoresis and immunoblotting

Immunoblotting and electrophoretic assays of cell lysates and culture supernatants were performed using NuPage BioTris electrophoretic system (Invitrogen). Protein samples in loading buffer were heated at 70°C for 10 min and loaded on 10% Bis-Tris gels. After electrophoresis and electro-transfer, the nitrocellulose membrane was blocked with 1% casein in TBS/T for 40 min. The membranes were probed with poly- or monoclonal anti-PSMA Abs (see below) for 1 h at RT. Detection was performed using goat anti-rabbit or anti-mouse IgG-HRP conjugates (Sigma) and visualized with WestPico Super Signal Chemo luminescent Substrate (Pierce) in accordance with the manufacturer's recommendations.

Anti-human PSMA monoclonal antibodies Y-PSMA1 and Y-PSMA2 were obtained from Yes Biotech Laboratories Ltd (Mississauga, Ontario, Canada).

#### Cell culture and generation of monocytes-derived dendritic cells

Cell cultures from peripheral blood mononuclear cells were maintained in research grade serum-free AIM-V medium (Invitrogen, Carlsbad, CA). The human prostate cancer cell line LNCaP was purchased from ATCC and was maintained in RPMI supplemented with 10% FCS (Life Technologies Inc., Rockville, MD), 2 mM L-glutamine, 50 units/ml penicillin and 50 mg/ml streptomycin. The human T2 cell line is transporter associated with antigen processing (TAP) deficient, resulting in inefficient loading of leukocyte antigen class I molecules with endogenous peptides. {DeMars, 1984 #1278} As a consequence, the HLA-A0201 molecules of T2 cells can be efficiently loaded with exogenous peptides. The T2 cell line was purchased from ATCC and maintained in IMDM supplemented with 20% FBS (Life Technologies Inc., Rockville, MD).

For T cell stimulation, the leukocyte fraction was collected by leukapheresis and mononuclear cells were separated on a density gradient. Cells were resuspended in serum-free AIM-V medium at  $2 \times 10^7$ /ml and incubated for 2 hrs in a humidified incubator at 37°C. The non-adherent T-cell enriched fraction and part of the adherent cells were harvested and frozen for future use.

The rest of the adherent cells were differentiated into DC by culture in serum-free AIM-V medium supplemented with IL-4 (PeproTech, Rocky Hill, NJ) and GM-CSF (Oncology Supplies Inc. (Dothan, AL) for 6 days. The non-adherent cells were harvested and used for transfection. Transfection was performed using the Nucleofector<sup>TM</sup> device and corresponding transfection kit (Amaxa GmbH, Cologne, Germany). The efficiency of Nucleofector<sup>TM</sup> transfection of human DCs was  $32 \pm 8.8\%$  (n=5) as estimated by control transfection with green fluorescence protein (GFP). Following transfection, dendritic cells were resuspended in serum-free AIM-V medium and matured with TNF- $\alpha$  (Becton Dickinson Inc., Bedford, MA) at 37°C for 24 hours. Following maturation, the DCs were resuspended in AIM-V medium at  $1 \times 10^5$  cells/ml.

#### In Vitro Generation of CTL Responses

For T cell stimulation, the T-cell enriched fraction was thawed, washed, resuspended in AIM-V medium and added to the DC suspension so that the final concentration of the T-cell enriched fraction was  $1 \times 10^6$  cells/ml. The cell suspension was then distributed into 24-wells plates (1ml/well) and cultured at 37°C. Three days later, the medium was replaced with serum free AIV-M medium supplemented with human IL-2 (20 U/ml) and human IL-7 (10 U/ml) (PeproTech, Rocky Hill, NJ). Cells were additionally stimulated with autologous PSMA-transfected DCs (stimulator:effector ratio of 1:10) or with peptide-pulsed autologous monocytes (stimulator:effector ratio of 1:1) twice, 8 days apart.

For pulsing with peptides, monocytes were resuspended at  $10^6$ /ml in serum-free RPMI-1640 medium supplemented with L-glutamine and penicillin/streptomycin. Peptide was added to a final concentration of 0.05 mg/ml and the cells were incubated for 4 hours at 37°C in a controlled CO<sub>2</sub> humidified incubator. The cells were then washed twice with serum-free medium, irradiated (1500 rads) and resuspended in IL-2 and IL-7 containing medium.

For cytotoxicity assays, target T2 cells were pulsed for 6 hours with 10 mg/ml peptide and 1mCi/well <sup>3</sup>H-thymidine (ICN Biomedical Inc., Irvine, CA).

After 20 days of culture, effector T cells were harvested without further separation for microcytotoxicity assays. The cells were analyzed by flow cytometry and 83±10% of them were CD3<sup>+</sup> and ~45% or ~60% of them were CD3<sup>+</sup>CD8<sup>+</sup> when primed with sPSMA DCs or tPSMA DCs respectively (data not shown).

For CTLA-4 inhibition experiments, Fab-fragments, prepared (see below) from anti-CD152 (HB-12319, ATCC) monoclonal antibody, were added (0.02 mg/ml final concentration) to the T cells at the initiation of culture or 6, 10 or 18 hours after that. Cultures were incubated for 3 additional days and the cells were washed and re-suspended in IL-2 and IL-7 containing AIV-M medium.

#### *Fab-fragment preparation of anti-CD152 monoclonal antibodies*

Purified IgG (2 mg/ml in PBS) was cleaved with papain (Sigma) at a ratio 1:25 (w/w) of papain to IgG in the presence of 0.01 mM L-cystein at 37°C for 5.5 h. The reaction was stopped by the addition of N-ethylmaleimide to a final concentration of 30 mM. The Fab fragment was isolated by gel filtration with a Superdex 75 column (Pharmacia) followed by ion exchange column (ResourceQ, Pharmacia) chromatography. The buffer for the eluted protein was changed to 10 mM Tris-HCl (pH 7.4) by gel filtration and the pooled protein was concentrated to 15 mg/ml with centricon-10 (Millipore). The protein was tested for purity at each step by SDS-PAGE under reduced and non-reduced conditions.

#### Cytotoxicity testing

Cytotoxicity against LNCaP cells or against peptide pulsed T2 cells (both HLA-A2-positive) was tested after 20 days of culture and compared to a control cell line that did not express PSMA. Cytotoxicity was tested using the JAM test. {Matzinger, 1991 #1269} Briefly, target or control cells were grown overnight with <sup>3</sup>H-thymidine, then washed, resuspended in complete RPMI-1640 medium and used in 4-hour cytotoxicity test. The killing was detected as a fall in counts per minute in cell samples undergoing apoptosis due to DNA fragmentation. All of the E:T ratios were tested in triplicate. Spontaneous cytotoxicity was determined in medium alone without effector cells.

Unlabelled K562 cells (no MHC expression, sensitive to natural killer cell-mediated lysis) were included at 50 x the target cell number to inhibit nonspecific lysis. Control experiments involved the Malme M3 melanoma cell line, which is also HLA-A2 positive.

#### Cell Lines

The human LNCaP (CRL-1740), T2 (CRL-1992) and Malme 3M (HTB-64) cell lines were purchased from ATCC and were maintained according to ATCC instructions.

#### Flow cytometry analysis

Antibodies used to phenotype the cells were anti-CD1a, anti-HLA-DR-PE, anti-CD80-PE, anti-CD86-PE, anti-CD83-FITC, anti-CD54-PE, anti-HLA-ABC-FITC, anti-CD14-FITC, anti-CD8-PE, anti-CD4-PE, anti-CD69-FITC (PharMingen, San Diego, CA), and anti-CD3-PerCP (Becton Dickinson, San Jose, CA). For HLA-A2 Typing, aliquots of peripheral blood mononuclear cells (PBMC) of healthy donors were tested with the FITC-labeled anti-HLA-A2 antibody BB7.2 (Becton Dickinson). For staining,  $10^5$  cells were suspended in 100 µl of PBS and were incubated with 10 µl of the antibodies for 20 min on ice. Flow cytometry analysis was performed on a FACS-Calibur (Becton Dickinson). GM-CSF was

purchased from Oncology Supplies Inc. (Dothan, AL), IL-4 and IL-7 from PeproTech Inc. (Rocky Hill, NJ), IL-2 and TNF- $\alpha$  from Becton Dickinson Inc. (Bedford, MA).  $^3\text{H}$ -thymidine was purchased from ICN Biomedical Inc (Irvine, CA).

### Statistics and Epitope Binding Predictions

Analysis of cytotoxicity data were performed using two-tailed Student's t tests assuming equal variance. We used the predictive algorithm from the Bioinformatics and Molecular Analysis Section of the NIH ("BIMAS") that was developed by Parker et al{Parker, 1992 #1270}, ranking potential MHC binders according to the predictive one-half-time disassociation of peptide/MHC complexes.

### Peptide synthesis and purification

Peptides were custom synthesized and purified by Sigma Genosys (The Woodlands, Texas).

## RESULTS

### Selection of HLA-A2-binding PSMA peptides

The aminoacid sequence of the extracellular domain of PSMA was analyzed for the existence of 9-aminoacid peptides predicted to bind to HLA-A0201, the most common human MHC class I allele. Using the computer-based algorithm (<http://bimas.cit.nih.gov/>) four 9-mer peptides that contains binding motifs for the HLA-A0201 class I molecule were identified (table 1).

Table 1. **HLA-A0201-restricted PSMA-derived peptides**

Rank	Position at PSMA molecule*	Peptide Sequence	Score**
1	663	MMNDQLMFL	1360
2	711	ALFDIESKV	1055
3	668	LMFLERAFI	261
4	707	GIYDALFDI	251


\* positions determined using the numbering from Israeli et al{Israeli, 1993 #189}.

\*\* score corresponds to the estimated half-time of dissociation of complexes containing the peptide at 37°C at pH

6.5


### COS-1 cells transfected with tPSMA or sPSMA plasmids express the encoded product

Following transfection with sPSMA plasmid, the encoded product is N-glycosylated (fig.1) and could be detected intra- and extracellularly – i.e. is secreted (data not shown). The product expressed following transfection with the tPSMA plasmid is not glycosylated, but is retained in the cytosol and rapidly degraded in the proteasome. It could be detected following proteasomal inhibition with lactacystin (fig.2).



**Fig.1.** COS-1 cells were transfected with empty (pVAX), tPSMA or sPSMA plasmids, grown for 72 hours and the cell lysates were analyzed on Western blot with (+G) or without (-G) prior treatment with glycopeptidase F. Unlike the

product obtained following transfection with tPSMA, the product obtained following transfection with sPSMA is N-glycosylated. Bands from recombinant PSMA or from cell lysates from LNCaP cells are shown for comparison. Legend – pVAX, tPSMA and sPSMA – lysates from COS-1 cells transfected with the respected plasmids; RP – recombinant PSMA expressed in insect cells, LNCaP – cell lysate form LNCaP.




**Fig 2.** COS-1 cells were transfected with the tPSMA plasmids and the cells were grown for 72 hours, the last 24 hours in the presence of lactacystin (10  $\mu$ M). Unlike the product expressed following transfection with sPSMA (left panel), most of the product expressed following transfection with tPSMA (right panel) is degraded in the proteasome and could be detected following proteasomal inhibition. Legend –LNCaP- LNCaP cell lysate; +/- L – in the presence (+) or absence (-) of the proteasomal inhibitor lactacystin.

### Generation of mature DCs from peripheral blood monocytes


Immature DC were generated from PBMC after 6 days of culture in GM-CSF- and IL-4- conditioned medium. DC appeared as non-adherent cells with the typical DC morphology and high expression of CD1a (80-90%). Treatment of tPSMA- and sPSMA-transfected DCs with TNF- $\alpha$  triggers a coordinate series of phenotypic changes, resulting in an up-regulation of co-stimulatory molecules (CD80, CD86, CD40) and HLA class II antigens (fig.3).

**Fig.3.** Change in cell surface antigen expression in PSMA-transfected DCs following maturation with TNF- $\alpha$ . Since surface antigens shows a heterogeneous baseline expression, the up-regulation of those molecules is presented as a ratio of mean fluorescent intensity (MFI activated/MFI resting). Data from 5 experiments with different donors are shown.




Both tPSMA DCs and sPSMA DCs prime and support development of T cells that are cytotoxic against LNCaP cells

The T cell enriched fraction from each leukapheresis was primed and then boosted twice, at 8-day intervals with autologous PSMA-transfected DCs. Their cytotoxicity was then tested against LNCaP cells or control Malme M3 melanoma cells. Both tPSMA DCs and sPSMA DCs primed and supported development of T cells that are cytotoxic against LNCaP cells (fig.4).



**Fig. 4.** Generation of CTL by tPSMA DCs (squares) or sPSMA DCs (triangles). Non-adherent (T cell-enriched) HLA A2 (+) peripheral blood mononuclear cells were primed and then boosted twice at 8-day intervals with autologous monocytes-derived dendritic cells that were transiently transfected with either tPSMA plasmid or sPSMA plasmid. The medium was changed, initially 72 hours following priming, and then during boosting. The cells were grown in IL-2 and IL-7 medium for 20 days and the specific cytotoxicity tested against LNCaP. Each point represents the mean and SD of triplicate experiments. Significant differences ( $p < 0.01$ ) in specific cytotoxicity at the 30:1 E/T ratio are indicated (xx). Control experiments involved priming and boosting of the T-cell enriched fraction with DCs transfected with empty plasmid (diamonds) or testing of cytotoxicity against the Malme-3M melanoma cell line (circles).



**Fig.5.** Induction of CTL responses with autologous dendritic cells that have been transfected with the extracellular domain of PSMA. Peripheral blood mononuclear cells that had been depleted of monocytes were primed with autologous DCs transfected with either sPSMA (triangles), tPSMA (squares) or control (mock, empty) plasmids (vertical bar). Responding cultures were then boosted with monocytes pulsed with one of four HLA-A2 restricted PSMA-derived peptides (table 1) and their cytotoxicity tested against T2 cells pulsed with the same peptide that was used during boosting, or with a control (influenza derived) peptide. Data points for the control peptide are not shown but are identical to those obtained with empty plasmid-transfected DCs. Each point represents the mean and SD of triplicate experiments. Significant differences ( $p < 0.01$ ) in cytotoxicity against specific and control targets at the 30:1 E/T ratio are indicated (**xx**).

*tPSMA DCs, but not sPSMA DCs prime T cells that are reactive to sub-dominant PSMA epitopes*


Since proteasomal degradation is the main source of antigenic fragments destined for MHC class I presentation, {Bennink, 1993 #1246} and since purposeful cytosolic retention of non-glycosylated and misfolded newly synthesized tumor-associated antigens enhances such degradation (fig.2), we reasoned that tPSMA transfected DCs will have a greater potential to

prime T cells specific for sub-dominant PSMA epitopes. To check this hypothesis, we primed the T cell enriched fraction from each leukapheresis with mature tPSMA DCs or sPSMA DCs and boosted them twice, at 8-day intervals, with autologous monocytes pulsed with one of several PSMA derived peptides (table 1). Twenty days after priming, cytotoxicity was tested against T2 cells pulsed with the same peptide used for boosting. Unlike the dendritic cells pulsed with the sPSMA plasmid, the tPSMA DCs prime T cells to all four PSMA-derived peptides in four of the five donors (fig.5).

*CTLA-4 inhibition during sPSMA DC-priming leads to stimulation of T cells reactive to sub-dominant epitopes*

Lack of responses to the subdominant PSMA epitopes when DCs are transfected with sPSMA may result from complete lack of such epitopes on the membrane of the APCs. Alternatively, the epitopes may be generated but factors other than T-cell receptor (TCR) signaling such as the CTLA-4/B7 pathway, could be contributing to the ineffective proliferation of T cell to subdominant epitopes.


Currently, there is a general consensus that effective activation of naive T cells requires two signals: one dependent on the engagement of the TCR by peptide-MHC complexes, and the second co-stimulatory signal that is provided by interactions between cell surface molecules on the T cell and the APC. {Kaye, 1995 #1354} Numerous studies have indicated that the CD28 molecule, expressed on T cells, provides a potent co-stimulatory signal following engagement with its ligands, B7-1 and B7-2. {Lenschow, 1996 #1353} The co-stimulatory function of CD28 is counterbalanced by the existence of a second higher-affinity receptor for B7, termed CTLA-4. {Walunas, 1994 #1352} The latter inhibits T cell activation at instances of weak T cell receptor engagement {Manzotti, 2002 #1330} and we speculated that we could enhance T cell stimulation to existing subdominant epitopes if we inhibited CTLA-4 signaling with anti-CTLA-4 Fab fragments. Indeed, CTLA-4 inhibition during sPSMA DC-priming led to stimulation of T cells reactive to sub-dominant epitopes. We interpret this as evidence that sub-dominant epitopes are generated in DC transfected with the sPSMA plasmid but the response can be detected only after CTLA-4 inhibition (fig.6).



**Fig. 6.** HLA A2 (+) peripheral blood mononuclear cells that had been depleted of monocytes were primed with autologous dendritic cells transfected with either sPSMA plasmid (triangles) or control (mock, empty) plasmid (diamonds) in the presence (open symbols) or absence (closed symbols) of blocking anti-CD152 Fab fragments. The Fab fragments (0.02 mg/ml) were added at the initiation of the culture. Three days later, and then 2 more times at weekly intervals, the responding cultures were boosted with monocytes pulsed with one of two HLA-A2 restricted, PSMA-derived peptides (PSMA<sub>707</sub> or PSMA<sub>668</sub>). Cytotoxicity

was tested against T2 cells pulsed with the same peptide that was used during boosting, or with a control (influenza) peptide. Data points for the control peptide are not shown but are identical to those obtained with empty plasmid-transfected DCs. Results from 3 independent experiments are shown. Significant differences ( $p < 0.01$ ) in cytotoxicity against specific and control targets at the 30:1 E/T ratio are indicated (xx).

Since CTLA-4 was not expressed in resting T cells, we determined the time points during the 72-hour priming phase that the inhibitory control by CTLA-4 became apparent. The anti-CD152 antibody Fab fragments (20  $\mu\text{g/ml}$ ) were added either at the beginning, or 6, 10 or 18 hours after the start of the culture. CTLA-4 inhibition was effective when performed early (0 or 6 hours) after initiation of culture (fig.7).



**Fig. 7.** HLA A2 (+) peripheral blood mononuclear cells depleted of monocytes were primed in the presence of blocking anti-CD152 Fab fragments with autologous dendritic cells that had been transfected with sPSMA plasmid. The Fab fragments (0.02 mg/ml) were added either at the beginning (diamonds), or 6 (squares), 10 (triangles) or 18 (circles) hours after initiation of culture. Lymphocytes were then cultured for 3 days, then washed, resuspended in IL-2 and IL-7 AIM-V medium and boosted twice at 1-week interval with autologous PSMA<sub>707</sub>-pulsed monocytes. Cytotoxicity was tested against T2 cells pulsed with the same peptide. Results from 3 independent experiments are shown. Significant differences ( $p < 0.01$ ) in cytotoxicity against specific and control targets at the 30:1 E/T ratio are indicated (xx).



*Repeated boosting with transfected DCs restricts the response towards one immunodominant epitope*

Maturation of potent CTL effectors requires repetitive boosting with target antigen. To determine the effect of a prime/boost vaccination strategy on the clonality of the T cell response, tPSMA DCs-primed cultures, known to contain CTLs to subdominant epitopes (fig.5), were boosted with transfected or peptide pulsed dendritic cells or monocytes, and their cytotoxicity was tested against PSMA-peptide pulsed T2 target cells. Boosting with either antigen presenting cells that express multiple PSMA-derived epitopes (transfected DCs, or DCs or monocytes pulsed with multiple peptides) restricts the immune response towards one immunodominant epitope (table 2). A subdominant T cell response could only be preserved if boosting is performed with an APC pulsed with the particular subdominant epitope.


**Table 2.** Boosting with dendritic cells that express multiple antigenic epitopes restricts the response to an immunodominant epitope

Prime/Boost Strategy*	Number of Patients Developing CTL Activity against PSMA Peptide-Pulsed T2 cells			
	PSMA <sub>663</sub>	PSMA <sub>711</sub>	PSMA <sub>668</sub>	PSMA <sub>707</sub>
Prime: tPSMA DCs. 2 Boosts tPSMA DCs	5 of 5	1 of 5	0 of 5	0 of 5
Prime: tPSMA DCs. 2 Boosts sPSMA DCs	5 of 5	1 of 5	0 of 5	0 of 5
Prime: tPSMA DCs. 2 Boosts 1P DCs	5 of 5	4 of 5	5 of 5	4 of 5
Prime: tPSMA DCs. 2 Boosts 4P Mos	5 of 5	0 of 5	0 of 5	0 of 5
Prime: tPSMA DCs. 2 Boosts 1P Mos	5 of 5	4 of 5	5 of 5	4 of 5

\* 1P – pulsed with single peptide, 4P – pulsed with all four peptides, Mos- monocytes

Anti-tumor reactivity of PSMA-specific CTLs

The ability of the PSMA-reactive T cell clones to recognize tumor cells that express PSMA was tested in an in vitro cytotoxicity assay. In all 5 donors, only two of the peptide-specific CTL clones (PSMA<sub>663</sub> and PSMA<sub>711</sub>) were cytotoxic against non-modified LNCaP cells (fig.8).



**Fig. 8.** Anti-tumor reactivity of PSMA peptide-specific CTLs. Peptide-reactive CTL lines shown in fig. 5 were tested for their ability to recognize and kill prostate tumor cells. LNCaP cells (filled symbols) or

Malme-3M melanoma cells were used as targets. Each point represents the mean and SD of triplicate experiments. Significant differences ( $p < 0.01$ ) in cytotoxicity against specific and control targets at the 30:1 E/T ratio are indicated (xx). A representative result from five tested donors is shown.

## **DISCUSSION**

Despite being effective in breaking tolerance to a self-antigen or tumor-associated antigen, anti-tumor vaccination in its current mode of application leads to a narrow, directed to few of the potential epitopes response. Immunodominance is a natural control mechanism that ensures the tight specificity of the immune reaction and prevents untoward autoimmunity. {Yewdell, 1999 #1096} Immunodominance, however, presents a problem in vaccinology since loss of an MHC haplotype that participates in the conformation of the T cell antigen, or point mutation in the recognized sequence, would result in ineffective immune surveillance. {Dunn, 2002 #1276; Schreiber, 2002 #1085} We have explored possibilities to extend the target epitope specificity of the immune response and the following study was designed to test the ability of dendritic cells transfected with different plasmid constructs to prime and boost cytotoxic T cell responses.

A critical requirement for T cell activation is the engagement of TCRs with major histocompatibility complex molecules presenting antigenic peptides on the surface of an antigen-presenting cell. {Davis, 1989 #1303} Efficient priming of multi-specific CTL responses is limited by the poor immunogenicity of subdominant MHC class I-binding epitopes. {Yewdell, 1999 #1096} T cells binding with high affinity to the antigen compete with low affinity T cells and inhibit their interaction with the same antigen-presenting cell by down-modulation of peptide-MHC complexes on the APC. {Kedl, 2002 #1317} Inhibition of responses to sub-dominant epitopes, therefore, could be partially overcome by simply increasing the amount of antigen present on the APCs. {Mitchison, 1992 #1311; Lightstone, 1992 #1312; Clark, 1981 #1313} We reasoned that, since proteasomal degradation is the main source of antigenic fragments destined for MHC class I presentation, {Bennink, 1993 #1246} purposeful cytosolic retention of newly synthesized tumor-associated antigens in genetically manipulated antigen presenting cells may enhance their proteasomal degradation and thus increase the quantity of such fragments. The transport from the cytosol into the endoplasmic reticulum (ER) is an important step in the biogenesis of many proteins, including secretory proteins and proteins of the plasma membrane. It is triggered by a signal sequence, which is normally located at the amino terminus of those polypeptides. For many membrane proteins, the signal sequence is identical with the first membrane anchor (signal-anchor sequence). {Rapoport, 1996 #1301} Since PSMA is a type II, bitopic integral membrane protein, we speculated that elimination of the sequence that encodes for its transmembrane domain will lead to expression of a product that will not be translocated to the ER, but will be degraded by the proteasomes, presumably to peptides that will access the class I presentation pathway. Indeed, the product expressed following transfection of COS-1 cells with such "truncated" construct is not glycosylated, but is retained in the cytosol and rapidly degraded by the proteasomes. This leads to generation of PSMA peptides that are expressed via the class I MHC presentation pathway at density sufficient to prime CTLs to all four PSMA epitopes tested.

Contrary to that, dendritic cells transfected with DNA encoding for a product, which is translocated to the ER, then glycosylated and secreted, do not prime against subdominant epitopes (fig.5). Although all four PSMA epitopes are generated, T cells

reactive to the subdominant ones could only be primed if CTLA-4 signaling is inhibited (fig.6). Since high-affinity receptor T cells down-modulate peptide-MHC complexes from the APC membrane, {Kedl, 2000 #1314} increased production of peptides after tPSMA transfection favors sub-dominant CTL priming. Lower concentration of MHC-peptide complexes on the APC membrane following sPSMA transfection, however, requires CTLA-4 inhibition. A homologue of CD28, CTLA-4 also binds to the B-7 family members {Greene, 1996 #1318; van der Merwe, 1997 #1319} but inhibits T cell activation. {Thompson, 1997 #1130} Mice lacking CTLA-4 reveal a striking phenotype of polyclonal T cell activation and tissue infiltration which results in death by 3-4 weeks of age, indicating a powerful regulatory role for CTLA-4. {Waterhouse, 1995 #1320; Tivol, 1995 #1321}

Our finding that priming to subdominant responses is enhanced by CTLA-4 inhibition may seem contradictory to the model proposed by others. {Egen, 2002 #1128} In that model CTLA-4 signaling and not inhibition allows for a greater diversity in the T cell response. Our results, however, support the model originally proposed by Manzotti et al. {Manzotti, 2002 #1330} The following operational mechanisms are, therefore, possible, but their clarification requires additional experimentation:

First, the intensity of TCR stimulation seems to be important with weak signals being prompt to inhibition. {Manzotti, 2002 #1330} In other words, weak signals being overwhelmed by inhibition through CTLA-4 will have no chance to prime CTL responses.

Second, CTLA-4 may act as a non-signaling "decoy" receptor reducing the available ligand for CD28 costimulation. {Masteller, 2000 #1323; Doyle, 2001 #1324}

There is also a third possibility {Read, 2000 #1328; Takahashi, 2000 #1327} that a small number of CTLA-4-expressing and -stimulated T cells exerts a suppressive or regulatory effect on other T cells. These cells appear to be similar if not identical to T regulatory cells. {Shevach, 2000 #1331; Groux, 1999 #1332; Sakaguchi, 2003 #1333}

Experiments are currently in progress to validate any of these hypotheses.

Priming with tPSMA DCs leads to polyclonal CTL stimulation, but boosting with APCs that express both dominant and sub-dominant epitopes narrows the immune response to the dominant ones. Research from other groups has gained similar results. {Smith, 2001 #1174; Mateo, 1999 #1334; Firat, 1999 #1335; Loirat, 2000 #1336} In all these instances, boosting with polyepitope constructs has resulted in failure to expand polyepitope CTLs. A likely explanation is that competition between T cells for antigen on individual APC leads to obscuring of responses to sub-dominant epitopes when both the dominant and subdominant epitopes are present on the same APC. {Palmowski, 2002 #1102; Kedl, 2003 #1337} New vaccines or new strategies employing modulation of co-stimulatory networks during boosting may need to be developed if the polyepitope response against a target is to be maintained.

Finally, not all PSMA-peptide specific CTL clones exhibited tumor cell killing (fig.8). A likely explanation for the lack of cytotoxicity is the downregulation of MHC - unlike other tumor cells, LNCaP cells express low levels of MHC class I molecules. {Lu, 2002 #1355} However, neither the level of class I MHC expression, nor the percentage of killing, was influenced by pre-treatment of the target cells with  $\gamma$ -IFN (data not shown). Alternatively, these subdominant epitopes may not be generated at sufficient quantity by LNCaP cells and further experimentation is necessary to test this possibility.

In conclusion, we have shown that:

1. Dendritic cells transfected with plasmid DNA can successfully prime CTL responses in vitro.
2. Dendritic cells transfected with a construct whose product is retained in the cytosol and degraded in the proteasome (tPSMA), prime to both dominant and subdominant epitopes. In contrast, sPSMA DCs prime to dominant epitopes only.
3. CTLA-4 inhibition during priming in vitro enhances priming to subdominant epitopes generated following transfection of DCs with sPSMA.
4. In vitro boosting with APCs that express both dominant and subdominant epitopes narrows the immune response to the dominant ones.

If confirmed in animal studies, these results will pose important questions on the design of vaccines and methods for re-immunization if a polyepitope response is to be maintained.

## REFERENCES

1. Mincheff M, Tchakarov S, Zoubak S, et al. Naked DNA and adenoviral immunizations for immunotherapy of prostate cancer: a phase I/II clinical trial. *Eur Urol.* 2000;38(2):208-217.
2. Mincheff M, Altankova I, Zoubak S, et al. In vivo transfection and/or cross-priming of dendritic cells following DNA and adenoviral immunizations for immunotherapy of cancer--changes in peripheral mononuclear subsets and intracellular IL-4 and IFN-gamma lymphokine profile. *Crit Rev Oncol Hematol.* 2001;39(1-2):125-132.
3. Weber LW, Bowne WB, Wolchok JD, et al. Tumor immunity and autoimmunity induced by immunization with homologous DNA. *J Clin Invest.* 1998;102(6):1258-1264.
4. Israeli RS, Powell CT, Fair WR, et al. Molecular cloning of a complementary DNA encoding a prostate-specific membrane antigen. *Cancer Res.* 1993;53(2):227-230.
5. Wright GL, Jr., Grob BM, Haley C, et al. Upregulation of prostate-specific membrane antigen after androgen-deprivation therapy. *Urology.* 1996;48(2):326-334.
6. Chang SS, Gaudin PB, Reuter VE, et al. Prostate-specific membrane antigen: present and future applications. *Urology.* 2000;55(5):622-629.
7. Chang SS, Gaudin PB, Reuter VE, et al. Prostate-Specific Membrane Antigen: Much More Than a Prostate Cancer Marker. *Mol Urol.* 1999;3(3):313-320.
8. Heston WD. [Significance of prostate-specific membrane antigen (PSMA). A neurocarboxypeptidase and membrane folate hydrolase]. *Urologe A.* 1996;35(5):400-407.
9. Silver DA, Pellicer I, Fair WR, et al. Prostate-specific membrane antigen expression in normal and malignant human tissues. *Clin Cancer Res.* 1997;3(1):81-85.
10. Shneider BL, Thevananther S, Moyer MS, et al. Cloning and characterization of a novel peptidase from rat and human ileum. *J Biol Chem.* 1997;272(49):31006-31015.
11. Halsted CH, Ling E, Luthi-Carter R, et al. Folylpoly-gamma -glutamate carboxypeptidase from pig jejunum. Molecular characterization and relation to glutamate carboxypeptidase II. *J Biol Chem.* 2000;275(39):30746.
12. Chang SS, O'Keefe DS, Bacich DJ, et al. Prostate-specific membrane antigen is produced in tumor-associated neovasculature. *Clin Cancer Res.* 1999;5(10):2674-2681.
13. Sprent J, Schaefer M. Antigen-presenting cells for unprimed T cells. *Immunol Today.* 1989;10(1):17-23.
14. Bennink JR, Anderson R, Bacik I, et al. Antigen processing: where tumor-specific T-cell responses begin. *J Immunother.* 1993;14(3):202-208.
15. Restifo NP, Kawakami Y, Marincola F, et al. Molecular mechanisms used by tumors to escape immune recognition: immunogenethrapy and the cell biology of major histocompatibility complex class I. *J Immunother.* 1993;14(3):182-190.
16. Dunn GP, Bruce AT, Ikeda H, et al. Cancer immunoediting: from immunosurveillance to tumor escape. *Nat Immunol.* 2002;3(11):991-998.
17. Schreiber H, Wu TH, Nachman J, et al. Immunodominance and tumor escape. *Semin Cancer Biol.* 2002;12(1):25-31.

18. Ikeda H, Old LJ, Schreiber RD. The roles of IFN gamma in protection against tumor development and cancer immunoediting. *Cytokine Growth Factor Rev.* 2002;13(2):95-109.
19. Yewdell JW, Bennink JR. Immunodominance in major histocompatibility complex class I-restricted T lymphocyte responses. *Annu Rev Immunol.* 1999;17:51-88.
20. Pinto JT, Suffoletto BP, Berzin TM, et al. Prostate-specific membrane antigen: a novel folate hydrolase in human prostatic carcinoma cells. *Clin Cancer Res.* 1996;2(9):1445-1451.
21. Rawlings ND, Barrett AJ. Structure of membrane glutamate carboxypeptidase. *Biochim Biophys Acta.* 1997;1339(2):247-252.
22. Holmes EH, Greene TG, Tino WT, et al. Analysis of glycosylation of prostate-specific membrane antigen derived from LNCaP cells, prostatic carcinoma tumors, and serum from prostate cancer patients. *Prostate Suppl.* 1996;7:25-29.
23. DeMars R, Chang CC, Shaw S, et al. Homozygous deletions that simultaneously eliminate expressions of class I and class II antigens of EBV-transformed B-lymphoblastoid cells. I. Reduced proliferative responses of autologous and allogeneic T cells to mutant cells that have decreased expression of class II antigens. *Hum Immunol.* 1984;11(2):77-97.
24. Matzinger P. The JAM test. A simple assay for DNA fragmentation and cell death. *J Immunol Methods.* 1991;145(1-2):185-192.
25. Parker KC, Bednarek MA, Hull LK, et al. Sequence motifs important for peptide binding to the human MHC class I molecule, HLA-A2. *J Immunol.* 1992;149(11):3580-3587.
26. Kaye PM. Costimulation and the regulation of antimicrobial immunity. *Immunol Today.* 1995;16(9):423-427.
27. Lenschow DJ, Walunas TL, Bluestone JA. CD28/B7 system of T cell costimulation. *Annu Rev Immunol.* 1996;14:233-258.
28. Walunas TL, Lenschow DJ, Bakker CY, et al. CTLA-4 can function as a negative regulator of T cell activation. *Immunity.* 1994;1(5):405-413.
29. Manzotti CN, Tipping H, Perry LC, et al. Inhibition of human T cell proliferation by CTLA-4 utilizes CD80 and requires CD25+ regulatory T cells. *Eur J Immunol.* 2002;32(10):2888-2896.
30. Davis MM, Chien YH, Bjorkman PJ, et al. A possible basis for major histocompatibility complex-restricted T-cell recognition. *Philos Trans R Soc Lond B Biol Sci.* 1989;323(1217):521-524.
31. Kedl RM, Schaefer BC, Kappler JW, et al. T cells down-modulate peptide-MHC complexes on APCs in vivo. *Nat Immunol.* 2002;3(1):27-32.
32. Mitchison NA. Specialization, tolerance, memory, competition, latency, and strife among T cells. *Annu Rev Immunol.* 1992;10:1-12.
33. Lightstone E, Marvel J, Mitchison A. Memory in helper T cells of minor histocompatibility antigens, revealed in vivo by alloimmunizations in combination with Thy-1 antigen. *Eur J Immunol.* 1992;22(1):115-122.
34. Clark EA, Lake P, Favila-Castillo L. Modulation of Thy-1 alloantibody responses: donor cell-associated H-2 inhibition and augmentation without recipient Ir gene control. *J Immunol.* 1981;127(5):2135-2140.
35. Rapoport TA, Rolls MM, Jungnickel B. Approaching the mechanism of protein transport across the ER membrane. *Curr Opin Cell Biol.* 1996;8(4):499-504.

36. Kedl RM, Rees WA, Hildeman DA, et al. T cells compete for access to antigen-bearing antigen-presenting cells. *J Exp Med.* 2000;192(8):1105-1113.
37. Greene JL, Leytze GM, Emswiler J, et al. Covalent dimerization of CD28/CTLA-4 and oligomerization of CD80/CD86 regulate T cell costimulatory interactions. *J Biol Chem.* 1996;271(43):26762-26771.
38. van der Merwe PA, Bodian DL, Daenke S, et al. CD80 (B7-1) binds both CD28 and CTLA-4 with a low affinity and very fast kinetics. *J Exp Med.* 1997;185(3):393-403.
39. Thompson CB, Allison JP. The emerging role of CTLA-4 as an immune attenuator. *Immunity.* 1997;7(4):445-450.
40. Waterhouse P, Penninger JM, Timms E, et al. Lymphoproliferative disorders with early lethality in mice deficient in CtlA-4. *Science.* 1995;270(5238):985-988.
41. Tivol EA, Borriello F, Schweitzer AN, et al. Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity.* 1995;3(5):541-547.
42. Egen JG, Allison JP. Cytotoxic T lymphocyte antigen-4 accumulation in the immunological synapse is regulated by TCR signal strength. *Immunity.* 2002;16(1):23-35.
43. Masteller EL, Chuang E, Mullen AC, et al. Structural analysis of CTLA-4 function in vivo. *J Immunol.* 2000;164(10):5319-5327.
44. Doyle AM, Mullen AC, Villarino AV, et al. Induction of cytotoxic T lymphocyte antigen 4 (CTLA-4) restricts clonal expansion of helper T cells. *J Exp Med.* 2001;194(7):893-902.
45. Read S, Malmstrom V, Powrie F. Cytotoxic T lymphocyte-associated antigen 4 plays an essential role in the function of CD25(+)CD4(+) regulatory cells that control intestinal inflammation. *J Exp Med.* 2000;192(2):295-302.
46. Takahashi T, Tagami T, Yamazaki S, et al. Immunologic self-tolerance maintained by CD25(+)CD4(+) regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4. *J Exp Med.* 2000;192(2):303-310.
47. Shevach EM. Regulatory T cells in autoimmunity\*. *Annu Rev Immunol.* 2000;18:423-449.
48. Groux H, Powrie F. Regulatory T cells and inflammatory bowel disease. *Immunol Today.* 1999;20(10):442-445.
49. Sakaguchi S. Regulatory T cells: mediating compromises between host and parasite. *Nat Immunol.* 2003;4(1):10-11.
50. Smith SG, Patel PM, Porte J, et al. Human dendritic cells genetically engineered to express a melanoma polyepitope DNA vaccine induce multiple cytotoxic T-cell responses. *Clin Cancer Res.* 2001;7(12):4253-4261.
51. Mateo L, Gardner J, Chen Q, et al. An HLA-A2 polyepitope vaccine for melanoma immunotherapy. *J Immunol.* 1999;163(7):4058-4063.
52. Firat H, Garcia-Pons F, Tourdot S, et al. H-2 class I knockout, HLA-A2.1-transgenic mice: a versatile animal model for preclinical evaluation of antitumor immunotherapeutic strategies. *Eur J Immunol.* 1999;29(10):3112-3121.
53. Loirat D, Lemonnier FA, Michel ML. Multiepitopic HLA-A\*0201-restricted immune response against hepatitis B surface antigen after DNA-based immunization. *J Immunol.* 2000;165(8):4748-4755.
54. Palmowski MJ, Choi EM, Hermans IF, et al. Competition between CTL narrows the immune response induced by prime-boost vaccination protocols. *J Immunol.* 2002;168(9):4391-4398.

55. Kedl RM, Kappler JW, Marrack P. Epitope dominance, competition and T cell affinity maturation. *Curr Opin Immunol.* 2003;15(1):120-127.
56. Lu J, Celis E. Recognition of prostate tumor cells by cytotoxic T lymphocytes specific for prostate-specific membrane antigen. *Cancer Res.* 2002;62(20):5807-5812.



## **A. Manuscript published in Cancer Gene Therapy**

### **Human Dendritic Cells Genetically Engineered to Express Cytosolically-Retained Fragment of Prostate-Specific Membrane Antigen Prime Cytotoxic T Cell Responses to Multiple Epitopes**

Milcho Mincheff<sup>1,4</sup>, M.D., Ph.D. Serguei Zoubak<sup>1</sup>, Ph.D., Iskra Altankova<sup>2</sup>, M.D., D.Sci., Stoyan Tchakarov<sup>3</sup>, M.D., Ph.D., Yevgen Makogonenko<sup>1</sup>, D.Sci., Chavdar Botev<sup>4</sup>, M.D., Irena Ignatova<sup>4</sup>, Rosen Dimitrov<sup>4</sup>, M.D., Kalina Madarzhieva<sup>4</sup>, M.D., Michael Hammett<sup>1</sup>, Yavor Pomakov<sup>2</sup>, M.D., Harold T. Meryman<sup>1</sup>, M.D., T. Lissitchkov<sup>4</sup>, M.D.

- 1- The George Washington University Medical Center, Washington DC
- 2- Central Laboratory for Immunology, St. Ivan Rilski Hospital, Sofia, Bulgaria
- 3- Department of Urology, St. Ann University Hospital, Sofia, Bulgaria
- 4- National Institute of Hematology and Transfusion Medicine, Sofia, Bulgaria

Corresponding author:

Milcho Mincheff

Head, Tumor Immunology Laboratory

Department of Medicine

The George Washington Medical Center

2300 Eye Street, N.W., Ross Hall 705

Washington, DC 20037

Telephone 202 994 7765; fax 202 994 0465

E-mail: [mcamsm@gwumc.edu](mailto:mcamsm@gwumc.edu)

Running title: Human Dendritic Cells Genetically Engineered

Keywords: PSMA. Gene-based vaccine, immunodominance, CTLA-4

Supported in part by grant N00014-00-1-0787 from the Office of Naval Research.

Supported in part by award No DAMD17-02-1-0239. The U.S. Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick, MD 21702-5014 is the awarding and administering acquisition office.

The content of the information does not necessarily reflect the position or the policy of the Government, and no official endorsement should be inferred. For purpose of this article, information includes news releases, articles, manuscripts, brochures, advertisements, still and motion pictures, speeches, trade association proceedings etc.

## **ABSTRACT**

The ability of two plasmid DNA vaccines to stimulate lymphocytes from normal human donors and to generate antigen-specific responses is demonstrated. The first vaccine (truncated; tPSMA) encodes for only the extracellular domain of PSMA. The product, expressed following transfection with this vector, is retained in the cytosol and degraded by the proteasomes. For the “secreted” (sPSMA) vaccine, a signal peptide sequence is added to the expression cassette and the expressed protein is glycosylated and directed to the secretory pathway. Monocyte-derived dendritic cells (DCs) are transiently transfected with either sPSMA or tPSMA plasmids. The DCs are then used to activate autologous lymphocytes in an in vitro model of DNA vaccination. Lymphocytes are boosted following priming with transfected DCs or with peptide-pulsed monocytes. Their reactivity is tested against tumor cells or peptide-pulsed T2 target cells. Both tPSMA DCs and sPSMA DCs cells generate antigen-specific cytotoxic T cell responses. The immune response is restricted towards one of the four PSMA derived epitopes when priming and boosting is performed with sPSMA. In contrast, tPSMA transfected DCs prime T cells towards several PSMA derived epitopes. Subsequent repeated boosting with transfected DCs, however, restricts the immune response to a single epitope due to immunodominance.

## INTRODUCTION

Immunotherapy of prostate cancer could be a safe, non-invasive, relatively inexpensive procedure that can avoid bowel and bladder injury and impotence often resulting from surgical, cryosurgical or radiation therapy. Several groups have recently reported on the safety of DNA vaccines for immunizations against tumor antigens.<sup>1-3</sup> A possible target for prostate cancer immunotherapy is the prostate specific membrane antigen (PSMA), also known as glutamate carboxypeptidase II (GCPII).<sup>4-8</sup> PSMA expression is normally restricted to the prostate gland, brain tissue, jejunum and proximal kidney tubules.<sup>9-11</sup> Its expression is increased nearly 10-fold in prostate cancer cells and is also found in tumor but not normal neovasculature.<sup>7,12</sup>

The main effectors in anti-tumor immunity are CD8<sup>+</sup> cytotoxic T cells that recognize tumor-associated antigen-derived peptides in association with major histocompatibility complex (MHC) class I molecules.<sup>13-15</sup> Gene-based vaccination in its current mode of application is effective in breaking tolerance to a self-antigen, but the response is narrow and is restricted to few of the potential epitopes. This presents a problem in vaccinology since loss of an MHC haplotype that participates in the conformation of the T cell antigen, or point mutation in the recognized sequence would result in ineffective immune surveillance.<sup>16-18</sup> New vaccines and/or new methods of immunizations need to be developed for those instances. These hopefully will raise responses to subdominant determinants so that the selection of tumor escape variants that fail to express immunodominant epitopes will be prevented.<sup>16</sup>

Numerous factors combine to establish an immunodominance hierarchy,<sup>19</sup> among them the ineffective generation and transport of sub-dominant epitopes by antigen-presenting cells (APCs). Since proteasomal degradation is the main source of antigenic fragments destined for MHC presentation,<sup>14</sup> we speculated that purposeful cytosolic retention of newly synthesized tumor-associated antigens in genetically manipulated antigen presenting cells may increase both the quantity and the diversity of such fragments. PSMA is a type II transmembrane glycoprotein which is comprised of 750 amino acids.<sup>20,21</sup> It lacks a signal peptide sequence and we speculated that elimination of sequences for its transmembrane region might impede the translocation of the encoded product to the endoplasmic reticulum. Such product should not be N-glycosylated, should be retained in the cytosol and rapidly degraded in the proteasome. In theory, DC transfected with such "truncated" sequences, may have an advantage of presenting "subdominant" antigenic determinants that otherwise may not be generated at sufficient density to prime antigen-specific cytotoxic T cell responses. The following experiments were designed to test this hypothesis in an in vitro immunization system with human cells.

## MATERIALS AND METHODS

All human cellular material used in these experiments was obtained following informed consent through protocols approved by the local Committee for Bioethics (Bulgaria) or the Investigational Review Board (IRB) at George Washington University Medical Center in Washington, DC.

### *tPSMA and sPSMA plasmids construction*

The cDNA encoding the extracellular portion (AA 44-750) of the human PSMA (XC-PSMA) was cloned into the pCR2.1 vector (Invitrogen, Carlsbad, CA) after RT-PCR of total mRNA from the human prostate cancer cell line LNCaP (CRL 1740, ATCC). The forward primer introduced a NotI cloning site and a Kozak sequence with a Met codon (GCCACCATG) into the 5'-end of XC-PSMA. The tPSMA plasmid for the transfection experiments was obtained by NotI-XhoI sub-cloning of XC-PSMA into a pVAX1 mammalian expression vector (Invitrogen, Carlsbad, CA). A secretable variant of the tPSMA plasmid – the sPSMA plasmid - was obtained by fusion of the XC-PSMA with a murine Ig k-chain leader sequence. The insert from tPSMA plasmid was sub-cloned by BamHI-XhoI into the mammalian expression vector pSecTag2A (Invitrogen, Carlsbad, CA) providing the murine Igk-chain leader sequence (clone 96). The single SfiI cloning site of the vector was used to fuse the XC-PSMA in-frame with the leader. The 5'-portion of the XC-PSMA between start Met and a single HpaI site was re-amplified in order to introduce SfiI site (Met codon was not included). The SfiI-HpaI fragment of clone 96 was replaced with the PCR product pre-digested with the same restriction endonucleases and the NruI-XhoI fragment from obtained construct was moved to the pVAX1 vector. The inserts in both constructs are under the regulation of a human cytomegalovirus (CMV) immediate-early promoter/enhancer and a bovine growth hormone polyadenylation signal. The plasmid DNA specifications include endotoxin content below 0.1 EU per microgram of DNA; lack of detectable amounts of bacterial RNA, genomic DNA or ssDNA as determined by agarose-gel electrophoresis; less than 10 microgram of protein per 1 mg of plasmid DNA as determined by colorimetric assay (Bio-Rad, Hercules, CA).

### **COS-1 transfection**

Expression of PSMA constructs was performed in Cos-1 cells (ATCC). Monolayers were transfected with FuGENE 6 transfection reagent (Roche) and assayed for PSMA production by Western blot. Cos-1 cells were seeded in 6-well tissue culture plates (Nunc, Denmark) at  $1.5 \times 10^5$  cells per well and grown to 50-70% confluence in DMEM supplemented with 25 mM HEPES (pH 7.5), 1 mM sodium pyruvate, 3.7 g/L sodium bicarbonate, 100 µg/ml penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin B and 10% (v/v) of heat inactivated fetal bovine serum. Cos-1 cells were transfected with 1.5 µg of plasmid DNA pre-condensed with 4.5 µl of FuGENE 6 reagent in serum-free DMEM for 30 min at room temperature. Cells were then grown for 72 h in supplemented DMEM and then harvested.

In the proteasome inhibition studies, lactacystin (Sigma) was added to the culture media (final concentration 10  $\mu$ M) 24 h before harvesting.

Cells were harvested by gentle scraping after double wash with 2 ml of cold PBS and adding to each well of 0.25 ml of cold lysis buffer (0.5 M NaCl, 1% triton X-100, 0.2% Tween 20, 50 mM HEPES, pH 7.0). Lysates were transferred to Eppendorf tubes and homogenized by repeated pipetting on ice.

For detection of secreted PSMA, the serum containing DMEM was removed 48 h after transfection, the cells were washed twice with 2 ml of PBS, serum free DMEM (2 ml per well) was added and cells were incubated for additional 24 h in the 6-well plates. After collection of the medium, the cell debris was removed by centrifugation (35,000 g, 20 min) and supernatants were concentrated with Centricon-50 centrifuge filtering device (Millipore) and stored at -30°C until further use.

### **Deglycosylation of PSMA**

The deglycosylation of PSMA proteins was carried out by treatment of 25 mg protein samples with 5 units of glycopeptidase F from *Chryseobacterium Meningosepticum* (Sigma) in 50 mM phosphate buffer (pH 7.5) containing 0.1% (w/v) SDS, 50 mM  $\beta$ -mercaptoethanol, 0.75% (v/v) Triton X-100 for 4 h at 37°C.<sup>22</sup>

### **Electrophoresis and immunoblotting**

Immunoblotting and electrophoretic assays of cell lysates and culture supernatants were performed using NuPage BioTris electrophoretic system (Invitrogen). Protein samples in loading buffer were heated at 70°C for 10 min and loaded on 10% Bis-Tris gels. After electrophoresis and electro-transfer, the nitrocellulose membrane was blocked with 1% casein in TBS/T for 40 min. The membranes were probed with poly- or monoclonal anti-PSMA Abs (see below) for 1 h at RT. Detection was performed using goat anti-rabbit or anti-mouse IgG-HRP conjugates (Sigma) and visualized with WestPico Super Signal Chemo luminescent Substrate (Pierce) in accordance with the manufacturer's recommendations.

Anti-human PSMA monoclonal antibodies Y-PSMA1 and Y-PSMA2 were obtained from Yes Biotech Laboratories Ltd (Mississauga, Ontario, Canada).

### **Cell culture and generation of monocytes-derived dendritic cells**

Cell cultures from peripheral blood mononuclear cells were maintained in research grade serum-free AIM-V medium (Invitrogen, Carlsbad, CA). The human prostate cancer cell line LNCaP was purchased from ATCC and was maintained in RPMI supplemented with 10% FCS (Life Technologies Inc., Rockville, MD), 2 mM L-glutamine, 50 units/ml penicillin and 50 mg/ml streptomycin. The human T2 cell line is transporter associated with antigen processing (TAP) deficient, resulting in inefficient loading of leukocyte antigen class I molecules with endogenous peptides.<sup>23</sup> As a consequence, the HLA-A0201 molecules of T2 cells can be efficiently loaded with exogenous peptides. The T2 cell line was purchased from ATCC and maintained in IMDM supplemented with 20% FBS (Life Technologies Inc., Rockville, MD).

For T cell stimulation, the leukocyte fraction was collected by leukapheresis and mononuclear cells were separated on a density gradient. Cells were resuspended in serum-free AIM-V medium at  $2 \times 10^7$ /ml and incubated for 2 hrs in a humidified incubator at 37°C. The non-adherent T-cell enriched fraction and part of the adherent cells were harvested and frozen for future use.

The rest of the adherent cells were differentiated into DC by culture in serum-free AIM-V medium supplemented with IL-4 (PeproTech, Rocky Hill, NJ) and GM-CSF (Oncology Supplies Inc. (Dothan, AL) for 6 days. The non-adherent cells were harvested and used for transfection. Transfection was performed using the Nucleofector<sup>TM</sup> device and corresponding transfection kit (Amaxa GmbH, Cologne, Germany). The efficiency of

Nucleofector™ transfection of human DCs was  $32 \pm 8.8\%$  (n=5) as estimated by control transfection with green fluorescence protein

**B. Manuscript published in Cancer Gene Therapy, 12, 185, 2005**

Depletion of CD25<sup>+</sup> Cells from Human T-Cell Enriched Fraction Eliminates Immunodominance during Priming with Dendritic Cells Genetically Modified to Express a Secreted Protein

Milcho Mincheff<sup>1,4</sup>, M.D., Ph.D. Serguei Zoubak<sup>1</sup>, Ph.D., Iskra Altankova<sup>2</sup>, M.D., D.Sci., Stoyan Tchakarov<sup>3</sup>, M.D., Ph.D., Petro Pogribnyy<sup>1</sup>, D.Sci., Yevgen Makogonenko<sup>1</sup>, D.Sci., Chavdar Botev<sup>4</sup>, M.D., Harold T. Meryman, M.D.<sup>1</sup>

- 1 The George Washington University Medical Center, Washington DC
- 2 Central Laboratory for Immunology, St. Ivan Rilski Hospital, Sofia, Bulgaria
- 3 Department of Urology, St. Ann University Hospital, Sofia, Bulgaria
- 4 National Institute of Hematology and Transfusion Medicine, Sofia, Bulgaria

Corresponding author:

Milcho Mincheff

Head, Tumor Immunology Laboratory

Department of Medicine

The George Washington Medical Center

2300 Eye Street, N.W., Ross Hall 705

Washington, DC 20037

Telephone 202 994 7765; fax 202 994 0465

E-mail: [mcamsm@gwumc.edu](mailto:mcamsm@gwumc.edu)

Running title: Depletion of CD25<sup>+</sup> cells, GITR-L co-transfection and immunodominance

Keywords: PSA, PAP, Gene-based vaccine, immunodominance, CD4+CD25+ cells, T regulatory cells, GITR-L

Supported in part by grant N00014-00-1-0787 from the Office of Naval Research.

Supported in part by award No DAMD17-02-1-0239. The U.S. Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick, MD 21702-5014 is the awarding and administering acquisition office.

The content of the information does not necessarily reflect the position or the policy of the Government, and no official endorsement should be inferred. For purpose of this article, information includes news releases, articles, manuscripts, brochures, advertisements, still and motion pictures, speeches, trade association proceedings etc.



## ABSTRACT

The ability of dendritic cells (DCs), genetically modified with one of two types of plasmid DNA vaccines to stimulate lymphocytes from normal human donors and to generate antigen-specific responses, is compared. The first type, also called “secreted” vaccine (sVac), encodes for the full length of the human prostate-specific antigen (PSA) with a signal peptide sequence so that the expressed product is glycosylated and directed to the secretory pathway. The second type, truncated vaccines (tVacs), encodes for either hPSA or human prostate acidic phosphatase (hPAP), both of which lack signal peptide sequences and are retained in the cytosol and degraded by the proteasomes following expression. Monocyte-derived dendritic cells are transiently transfected with either sVac or one of two tVacs. The DCs are then used to activate CD25<sup>+</sup>-depleted or non-depleted autologous lymphocytes in an *in vitro* model of DNA vaccination. Lymphocytes are boosted following priming with transfected DCs, peptide pulsed DCs or monocytes. Their reactivity is tested against tumor cells or peptide-pulsed T2 target cells. Both tVacDCs and sVacDCs generate antigen-specific cytotoxic T cell responses. The immune response is restricted towards one of three antigen-derived epitopes when priming and boosting is performed with sVacDCs. In contrast, tVac transfected DCs prime T cells towards all antigen-derived epitopes. Subsequent repeated boosting with transfected DCs, however, restricts the immune response to a single epitope due to immunodominance. While CD25<sup>+</sup> cell depletion prior to priming with sVacDCs alleviates immunodominance, co-transfection of dendritic cells with GITR-L does so in some but not all cases.

## INTRODUCTION

In a previous study,<sup>1</sup> we found that dendritic cells transfected with products that were cytosolically retained and degraded in the proteasomes, primed autologous T cells to multiple epitopes. Priming with sVacDCs, however, restricted the immune response to one of the epitopes due to immunodominance and the latter was alleviated if anti-CTLA-4 antibodies were present. We wanted to extend these observations to other tumor-associated antigens such as prostate-specific antigen (PSA) and prostate acid phosphatase (PAP), as well as to study the effect that the removal of CD4<sup>+</sup>CD25<sup>+</sup> cells prior to priming may have on immunodominance development. Both PSA and PAP are currently used as targets for immunotherapy of cancer.<sup>2-4</sup> On the other hand, gene-based vaccination in its current mode of application is effective in breaking tolerance to a self-antigen, but the response appears to be narrow and restricted to few of the potential epitopes. For example, the post-vaccination T cell response of some of the HLA A2 patients from the clinical trial performed by us<sup>5</sup> was directed against only two of the potential 4 PSMA peptide motifs that had high affinity for binding [M. Mincheff, unpublished]. Immunodominance is a natural mechanism for control that ensures the tight specificity of the immune reaction and prevents untoward autoimmunity, but it also carries the risk of inefficient immune surveillance in cases such as cancer where mutations of the epitope or downregulation of MHC alleles occur.<sup>6-8</sup> Malignant transformation and tumor progression are frequently associated with loss of HLA class I antigens. For example, a recent review of the literature<sup>9</sup> has reported that ~ 15% and 55% of surgically removed primary and metastatic melanoma lesions, respectively, are not stained in immunohistochemical reactions by monoclonal antibodies to monomorphic determinants of HLA class I antigens. Loss or reduced HLA class I antigen expression enables tumor cells to evade the host's immune response<sup>6,7,10-12</sup> and downregulation of HLA class I antigens in metastases from patients with malignant melanoma has been associated with poorer prognosis.<sup>13</sup> Immunodominance, therefore, presents a problem in vaccinology.<sup>8,14,15</sup> New vaccines and/or new methods of immunizations need to be developed for those instances. These hopefully will raise responses to subdominant determinants so that the selection of tumor escape variants that fail to express immunodominant epitopes will be prevented.<sup>8</sup>

Numerous factors combine to establish an immunodominance hierarchy.<sup>16</sup> Preliminary results from our laboratory suggest that the enhanced priming to subdominant epitopes by CTLA-4 inhibition is at least partially mediated through the inhibition of CD4<sup>+</sup>CD25<sup>+</sup> T cell function. These CD4<sup>+</sup> T cells are a minor subpopulation (10%) that co-expresses the IL-2 receptor  $\alpha$ -chain (CD25)<sup>17</sup> and they can prevent both the induction and effector function of autoreactive T cells.<sup>18-20</sup> Additionally, they suppress polyclonal T cell activation *in vitro* by inhibiting IL-2 production.<sup>21</sup> Very little is known of the physiologic regulation of CD4<sup>+</sup>CD25<sup>+</sup> T cells *in vivo*.<sup>22</sup> Recent reports suggest that glucocorticoid-induced tumor necrosis factor receptor (GITR), also known as TNFRSF18 – a member of the TNF-nerve growth factor receptor gene superfamily – is predominantly expressed on CD4<sup>+</sup>CD25<sup>+</sup> T cells<sup>22,23</sup> and stimulation of GITR abrogates CD4<sup>+</sup>CD25<sup>+</sup> T cell-mediated suppression.<sup>23</sup> The gene encoding the natural ligand of human GITR has been cloned and characterized. It is called GITR-L, a human activation-inducible TNF receptor (AITR) ligand, or TL6. Expression of the GITR-L is detected in immature and mature splenic dendritic cells. GITR-L binds GITR expressed on HEK 293 cells and triggers NF-kappa B activation. Functional studies

reveal that soluble CD8-GITR-L prevents CD4+CD25+ regulatory T-cell-mediated suppressive activities<sup>24</sup>. Would CD25+ T cell depletion prior to priming alleviate immunodominance? If CD25+ cells suppress priming of sub-dominant T cell clone, could immunodominance be restricted by GITR signaling? Could this be achieved by enhanced GITR-L co-expression during re-immunization? The following experiments were designed to test these hypotheses in an *in vitro* immunization system with human cells.

## MATERIALS AND METHODS

All human cellular material used in these experiments was obtained following informed consent through protocols approved by the local Committee for Bioethics (Bulgaria) or the Investigational Review Board (IRB) at George Washington University Medical Center in Washington, DC. Use of recombinant DNA was approved by the local IRAC committees.

### htPAP-, hGITR-L-, hsPSA- and htPSA plasmids construction

*hPAP* was obtained by RT-PCR of total RNA from LNCaP cells using TriPure RNA/DNA isolation reagent and Titan RT-PCR kit (Roche). The first cDNA strand was synthesized using hPAP-specific reverse primer (5'-GAGATCTCTGTGCACACTAATCTGTA-3'). Amplification was performed using the direct primer 5'-TCCTAACTCCTGCCAGAAACAGCTCT-3' and the same reverse primer. After initial denaturation, step (2 min, 94°C) and the first 10 cycles (30 sec at 94°C, 25 sec at 60°C, 45 sec at 72°C) the extension time was progressively incremented by 15 sec after each of the additional 20 cycles. The gel purified PCR product was cloned into a pCR2.1 vector (Invitrogen) and several clones were sequenced. All of them contained silent or non-silent substitutions. Two clones were selected for further work: 150.16 (no substitution in the 5'-region) and 150.20 (no substitution in the 3'-region).

*Truncated hPAP (H PAP-T)* was obtained by PCR of the clone 150.16 containing no substitution in the region between the end of signal peptide (position 97 on cds) and a single SfuI site (position 311 on cds). The direct primer (CGGCGGGGTACCATGGAGTTGAAGTTTGTGACTTTGGTG) introduces a KpnI site and a Kozak sequence (underlined). The first amino acid next to the signal peptide (Lys) was replaced by Met (bold italic). The reverse primer (5'-GGCTGCCAGAGTAGGATAGGATTC-3') anneals to the region downstream of a single SfuI site. The PCR product was digested with KpnI and SfuI endonucleases, gel purified, and sub-cloned by KpnI-SfuI sites into the clone 150.20 replacing its 5'-region with substitutions and signal peptide sequence. The correctness of the resulting clone was checked by sequencing.

Finally, the *H PAP-T* sequence was transferred by BamHI-XbaI sites from the pCR2.1 vector to the mammalian expression vector p147 (see below).

Human GITR-L was obtained by RT-PCR of total RNA from HuVec cells using specific primers (direct: 5'-GGTACCATGTGTTTGAGCCACTTGGAAAATATGCC-3'; reverse: 5'-CTAGGAGATGAATTGGGGATTTGC-3'). The PCR product (544 bp long) was cloned into a pCR2.1 vector and several clones were sequenced. All of them contained mutations. hGITR-L insert from one of the clones was transferred by HindIII-XhoI sites into the vector p147 and the mutations were corrected using QuickChange Site

Directed Mutagenesis kit (Stratagen). The sequence of the final clone is identical to hGTR-L genebank entry AF125303.

The *hPSA* containing plasmid was kindly provided by Jan Geliebter, New York Medical College, Department of Microbiology and Immunology, Valhalla, NY, USA. The clone represents the complete *hPSA* cDNA sequence included in a pCDNA3 (Invitrogen) vector. The clone contains two non-silent nucleotide substitutions: C to A at cds position 289, and G to A at cds position 406 leading to substitutions Pro97 by Thr and Val136 by Met respectively. These mutations do not affect the generation of PSA-derived 9-mer epitopes with high affinity for binding to HLA A201 as predicted by the computer-based algorithm available at <http://bimas.cit.nih.gov/>.

To obtain a 5'-truncated form of *hPSA* (*H PSA-T*) that lacks a signal peptide, a PCR of the *hPSA* region from the end of signal peptide (cds position 52) to the end of cds was performed. A direct primer (GCGGCCGCCGCCACCAATGGCACCCCTCATCCTGTCTCGG) introduced a NotI recognition site and a Kozak sequence (underlined) as well as the start methionine (bold italic). A reverse primer (5'-GTTTAAACTCAGGGGTTGGCCACGATGGTGTC-3') introduced the PmeI site (underlined) just next to stop codon. The PCR product was cloned into a pCR2.1 vector and sequenced. The *H PSA-T* was then excised from the pCR2.1 vector by cutting with NotI+PmeI, gel purified and ligated with pVAX-1 mammalian expression vector (Invitrogen) by NotI and XhoI sites. The last one was blunted with Klenow enzyme in the presence of excess of dNTP. Consequently, *H PSA-T* was transferred by BamHI-XbaI sites to the mammalian expression vector p147.

The mammalian expression vector p147 represents a modified pCDNA3 vector (Invitrogen) in which the ampicillin resistance gene was replaced with kanamycin resistance gene. The replacement was performed by ligation of BclI-BspHI fragment from pVAX-1 plasmid with BglII-BspHI fragment of plasmid pCDNA3. The former fragment contains kanamycin resistance gene, the last one includes all elements of pCDNA3 vector except the ampicillin resistance gene.

The plasmid-DNA product specifications include endotoxin content below 0.1 EU per microgram of DNA; >90% of covalently closed circle DNA, lack of detectable amounts of bacterial RNA, genomic DNA or ssDNA as determined by agarose-gel electrophoresis; less than 10 microgram of protein per 1 mg of plasmid DNA as determined by colorimetric assay (Bio-Rad, Hercules, CA).

### **COS-1 transfection**

Expression of PSA and PAP constructs (sVacs or tVacs) was performed in COS-1 cells (ATCC). A stably transfected AT3B-1 (ATCC) cell line which secretes PSA was obtained. Monolayers were transfected with FuGENE 6 transfection reagent (Roche) and assayed for PSA or PAP production by Western blot. COS-1 or AT3B-1 cells were seeded in 6-well tissue culture plates (Nunc, Denmark) at  $1.5 \times 10^5$  cells per well and grown to 50-70% confluence in DMEM supplemented with 25 mM HEPES (pH 7.5), 1 mM sodium pyruvate, 3.7 g/L sodium bicarbonate, 100 µg/ml penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin B and 10% (v/v) of heat inactivated fetal bovine serum. COS-1 or AT3B-1 cells were transfected with 1.5 µg of plasmid DNA pre-condensed with 4.5 µl of FuGENE 6 reagent in serum-free DMEM for 30 min at room temperature. Cells were then grown for 72 h in complete DMEM and then harvested.

In the proteasome inhibition studies, lactacystin (Sigma) was added to the culture media (final concentration 10  $\mu$ M) 24 h before harvesting.

Cells were harvested by gentle scraping, washed twice with 2 ml of cold PBS and 0.25 ml cold lysing buffer (0.5 M NaCl, 1% triton X-100, 0.2% Tween 20, 50 mM HEPES, pH 7.0) was added to each well. Lysates were transferred to Eppendorf tubes and homogenized by repeated pipetting on ice.

For detection of secreted PSA, the serum containing DMEM was removed 48 h after transfection, the cells were washed twice with 2 ml of PBS, serum free DMEM (2 ml per well) was added and cells were incubated for additional 24 h in the 6-well plates. After collection of the medium, the cell debris was removed by centrifugation (35,000 g, 20 min) and supernatants were concentrated with Centricon centrifuge filtering device (Millipore) and then stored at -30°C until further use.

### **Electrophoresis and immunoblotting**

Immunoblotting and electrophoretic assays of cell lysates and culture supernatants were performed using NuPage BioTris electrophoretic system (Invitrogen). Protein samples in loading buffer were heated at 70°C for 10 min and loaded on 10% Bis-Tris gels. After electrophoresis, the proteins were electro-transferred onto nitrocellulose membrane and blocked with 1% casein in TBS/T for 40 min. The membranes were probed with poly- or monoclonal anti-PSA or PAP Abs (see below) for 1 h at RT. Detection was performed using goat anti-rabbit or anti-mouse IgG conjugated with HRP (Sigma) and visualized with WestPico Super Signal Chemo luminescent Substrate (Pierce) in accordance with the manufacturer's recommendations.

Anti-human PSA monoclonal antibodies sc-7316 and sc-7638 were obtained from Santa Cruz Biotechnologies, Inc. (Santa Cruz, CA, USA). Purified PSA protein was obtained from International Immuno-Diagnostics, Foster City, CA. Anti-human PAP monoclonal antibody (mouse, IgG1), clone Pase/4LJ, was obtained from Acris Antibodies GmbH (Germany).

### **Flow cytometry analysis**

Antibodies used to phenotype the cells were anti-CD1a, anti-HLA-DR-PE, anti-CD80-PE, anti-CD86-PE, anti-CD83-FITC, anti-CD54-PE, anti-HLA-ABC-FITC, anti-CD14-FITC, anti-CD8-PE, anti-CD4-PE, anti-CD69-FITC (PharMingen, San Diego, CA), and anti-CD3-PerCP (Becton Dickinson, San Jose, CA). Anti-GITR-L antibodies (clones 109114 and 109117) and anti-GITR (clone 110416) were supplied by R&D Systems Inc.; Minneapolis, MN 55413, USA. For HLA-A2 Typing, aliquots of PBMC from buffy coats of healthy donors were tested with the FITC-labeled anti-HLA A2 antibody BB7.2 (Becton Dickinson). For staining,  $10^5$  cells were suspended in 100  $\mu$ l of PBS and were incubated with 10  $\mu$ l of the antibodies for 20 min on ice. Flow cytometric analysis was performed on a FACS Calibur (Becton Dickinson).

### **Phenotyping of CD25+ regulatory T cells (TR)**

Briefly, fresh or frozen PBMC were washed once in phosphate-buffered saline (PBS) containing 1% bovine serum albumin (BSA) (Sigma-Aldrich, St. Louis, Mo.) and stained with antibodies for CD3 (peridinin chlorophyll protein; PerCP), CD4 (fluorescein isothiocyanate; FITC), CD8 (allophycocyanin; APC), CD25-FITC or -phycoerythrin (PE), CD38-APC or -PE, and/or HLA-DR—APC (BD BioSciences, San Jose, Calif., and BD BioSciences PharMingen, San Diego, Calif.) for 20 min at 4°C. The cells were then

washed twice with PBS containing 1% BSA, fixed in 1% paraformaldehyde, acquired on a flow cytometer (FACSCalibur; BD BioSciences), and analyzed using FlowJo software (Tree Star, San Carlos, Calif.).

### **Cell cultures**

Cell cultures from peripheral blood mononuclear cells were maintained in research grade serum-free AIM-V medium (Invitrogen, Carlsbad, CA). The human prostate cancer cell line LNCaP was purchased from ATCC and was maintained in RPMI supplemented with 10% FCS (Life Technologies Inc., Rockville, MD), 2 mM L-glutamine, 50 units/ml penicillin and 50 mg/ml streptomycin (complete medium; CM). The human T2 cell line is transporter associated with antigen processing (TAP) deficient, resulting in inefficient loading of human leukocyte antigen class I molecules with endogenous peptides<sup>25</sup>. As a consequence, the HLA-A0201 molecules of T2 cells can be efficiently loaded with exogenous peptides. The T2 cell line was purchased from ATCC and maintained in IMDM supplemented with 20% FBS (Life Technologies Inc., Rockville, MD).

For T cell stimulation, the leukocyte fraction was collected by leukapheresis and mononuclear cells were separated on a density gradient. Cells were resuspended in serum-free AIM-V medium at  $2 \times 10^7$ /ml in culture flasks for 2 hrs in a humidified incubator at 37°C. The non-adherent T-cell enriched fraction (later in the text referred to as “T cells”) and part of the adherent cells were harvested and frozen for future use.

### **Depletion of CD25<sup>+</sup> cells**

CD25<sup>+</sup> cells were purified with MACS CD25 MicroBeads (Miltenyi Biotec, Auburn, Calif.). Briefly, the non-adherent T-cell enriched fraction (T cells) was washed twice in PBS containing 0.5% BSA and 2 mM EDTA, resuspended in 80  $\mu$ l of PBS containing 0.5% BSA-2 mM EDTA and 20  $\mu$ l of MACS CD25 MicroBeads per  $10^7$  total PBMC, and incubated for 15 min at 6 to 12°C. The cells were washed twice in PBS containing 0.5% BSA and 2 mM EDTA and applied to a magnetic column on a MidiMACS separation unit (Miltenyi Biotec). CD25<sup>+</sup> and CD25<sup>-</sup> T-cell fractions were collected. The CD25<sup>+</sup> cell fraction contained >90% CD4<sup>+</sup> T cells. In some experiments the CD25<sup>+</sup> cell fraction was purified to >99% CD4<sup>+</sup> T cells by cell sorting after staining with monoclonal antibodies to CD3 and CD4 (FACSVantage; BD Biosciences) and frozen for add-back experiments. In other experiments, CD4<sup>+</sup>CD25<sup>+</sup> T cells were stimulated for 48 hours with plate-bound anti-CD3 monoclonal antibody and IL-2 and used for staining with anti-GITR antibody.

### **Generation of monocytes-derived dendritic cells**

The rest of the adherent cells were differentiated into DC by culture in serum-free AIM-V medium with IL-4 (PeproTech, Rocky Hill, NJ) and GM-CSF (Oncology Supplies Inc. (Dothan, AL) for 6 days. The non-adherent cells were harvested at that time and used for transfection. Transfection was performed using the Nucleofector<sup>TM</sup> device and transfection kit (Amaxa GmbH, Cologne, Germany). The green fluorescence protein (GFP) transfection efficiency of human DCs after Nucleofector<sup>TM</sup> transfection was 32 $\pm$ 8.8 % (n=5). Following transfection, dendritic cells were resuspended in serum-free AIM-V medium and matured with TNF- $\alpha$  (Becton Dickinson Inc., Bedford, MA) at 37°C for 24 hours. Following maturation, the DCs were resuspended in AIM-V medium at  $1 \times 10^5$  cells/ml. Treatment of tPSMA- and sPSMA-transfected DCs with TNF alpha

triggers a coordinate series of phenotypic changes, resulting in an up-regulation of co-stimulatory molecules (CD80, CD86, CD40) and HLA class II antigens<sup>1</sup>.

### **In Vitro Generation of CTL Responses**

For T cell stimulation, the T cells with or without prior CD25<sup>+</sup>-depletion, were thawed, washed, resuspended in AIM-V medium and added to the DC suspension so that the final concentration of the T cells was 1x10<sup>6</sup> cells/ml (T cell/DC ratio = 10:1). The cell suspension was then distributed into 24-wells plates (1ml/well; Costar plates) and cultured at 37°C.

In some experiments CD4<sup>+</sup>CD25<sup>+</sup> T cells were added back to the stimulated T cells. In those, 0.25x10<sup>6</sup> CD4<sup>+</sup>CD25<sup>+</sup> T cells were added to each well (responder T cells/CD4<sup>+</sup>CD25<sup>+</sup> T cell ratio = 4:1) at different time points following priming.

Three days later, the medium was removed and the T cells were cultured from that moment in serum free AIV-M medium supplemented with human IL-2 (20 U/ml) and human IL-7 (10 U/ml) (PeproTech, Rocky Hill, NJ). Cells were additionally stimulated with autologous PSMA-transfected DCs (stimulator:effector ratio of 1:10) or with peptide-pulsed autologous monocytes (stimulator:effector ratio of 1:1) twice, 8 days apart.

For pulsing with peptides, monocytes were resuspended at 10<sup>6</sup>/ml in serum-free RPMI-1640 with L-glutamine and penicillin/streptomycin. Peptide was added to a final concentration of 0.05 mg/ml and the cells were incubated for 4 hours at 37°C in a controlled CO<sub>2</sub> humidified incubator. The cells were then washed twice with serum-free medium, irradiated (1500 rads) and used for boosting in IL-2 and IL-7 containing medium.

For peptide pulsing of T2 cells, we initially performed peptide-binding assays. For that purpose, T2 cells were incubated (10<sup>5</sup> cells/well) overnight in 96-well plates with serial dilutions of peptides in RPMI 1640/10% boiled fetal calf serum (to prevent protease activity), and then analyzed by FACS analysis for surface expression of HLA-A0201. Mean fluorescence intensities at varying concentrations of peptide were compared (data not shown). In subsequent experiments, T2 cells were pulsed for 6 hours with 0.01mg/ml peptide and 1mCi/well <sup>3</sup>H thymidine (ICN Biomedical Inc., Irvine, CA).

After 20 days of culture, effector cells were harvested without further separation for micro-cytotoxicity assays. The cells were analyzed by flow cytometry and 83% +/- 10% of them were CD3<sup>+</sup> and ~ 45% of them (when primed with the sPSA DCs) and ~60% of them (when primed with the tPSA DCs) were CD3<sup>+</sup>CD8<sup>+</sup> (data not shown).

### **Selection of HLA-A2-binding PSA- and PAP- peptides**

For the present study, the aminoacid sequence of the truncated (no-leader sequence) PSA (AA 25-161) and PAP (AA 33-386) were analyzed for the existence of 9-aminoacid peptides predicted to bind to HLA-A201, the most common human MHC class I allele, using the computer-based algorithm (<http://bimas.cit.nih.gov/>) (table1). Three 9-mer peptides from each sequence that contained peptide-binding motifs for the HLA A201 class I molecule were identified:

### **Cytotoxicity testing**

Cytotoxicity against LNCaP cells or against peptide pulsed T2 cells (both HLA A2-positive) was tested after 20 days of culture and compared to a control cell line that did not express PSA or PAP. Cytotoxicity was tested using the JAM test<sup>26</sup>. Briefly, target or control cells were grown overnight with <sup>3</sup>H-thymidine, then washed, resuspended in

CM and used in 4-hour cytotoxicity test. The killing was detected as a fall in counts per minute due to DNA fragmentation in cell samples undergoing apoptosis. All of the E:T ratios were tested in triplicate. Spontaneous cytotoxicity was determined in medium alone without effector cells.

Unlabelled K562 cells (no MHC expression and sensitive to natural killer cell-mediated lysis) were included at 50 x the target cell number to inhibit nonspecific lysis. Control experiments involved the Malme M3 melanoma cell line, which is also HLA A2 positive.

### **Cell Lines**

The human LNCaP (CRL-1740), T2 (CRL-1992), Malme 3M (HTB-64), COS-1 (CRL-1650) and AT3B-1 (CRL-2375) cell lines were purchased from ATCC and were maintained according to ATCC instructions.

### **Cytokines and 3H-thymidine**

GM-CSF was purchased from Oncology Supplies Inc. (Dothan, AL), IL-4 and IL-7 from PeproTech Inc. (Rocky Hill, NJ), IL-2 and TNF- $\alpha$  from Becton Dickinson Inc. (Bedford, MA). 3H-thymidine was purchased from ICN Biomedical Inc (Irvine, CA).

### **Statistics and Epitope Binding Predictions**

Analysis of cytotoxicity data were performed using two-tailed Student's t tests assuming equal variance. We used the predictive algorithm from the Bioinformatics and Molecular Analysis Section of the NIH ("BIMAS") that was developed by Parker et al <sup>27</sup>, ranking potential MHC binders according to the predictive one-half-time disassociation of peptide/MHC complexes for epitope binding prediction.

### **Peptide synthesis and purification**

Peptides were custom synthesized and purified by Sigma Genosys (The Woodlands, Texas).

## **RESULTS**

### **COS-1 cells transfected with tPSA, sPSA or GITR-L plasmids express the encoded product**

Following transfection with sPSA plasmid, the encoded product is N-glycosylated and could be detected intra- and extracellularly – i.e. is secreted (fig.1 A and B). The product expressed following transfection with the tPSA plasmid is not glycosylated, but is retained in the cytosol and rapidly degraded in the proteasome. Similarly to transfection with another “truncated” gene-based vaccine <sup>1</sup>, the product could be detected following proteasomal inhibition with lactacystin. COS-1 cells transfected with GITR-L stain positively with antibodies specific for the GITR-L (fig.2).

### **tVacDCs ( tPSA DCs, tPAPDCs) and sVac DCs (sPSA DCs) prime and support development of T cells that are cytotoxic against LNCaP cells**

The T cell enriched fraction from each leukapheresis was primed and then boosted twice, at 8-day intervals with autologous PSA-transfected DCs. Their cytotoxicity was then tested against LNCaP cells or control Malme M3 melanoma cells.



Both tVacDCs and sVacDCs primed and supported development of T cells that are cytotoxic against LNCaP cells (fig. 3).

### **tVacDCs but not sVacDCs prime T cells that are reactive to sub-dominant PSA or PAP epitopes**

Recently, we found that dendritic cells transfected with a fragment of the human prostate-specific membrane antigen that was cytosolically retained and degraded in the proteasome (tPSMA DCs), primed autologous T cells to multiple epitopes<sup>1</sup>. To extend these observations to other prostate antigens such as PSA and PAP, we primed the T cell enriched fraction from leukapheresis with either tPSA DCs, sPSA DCs or tPAP DCs and boosted them twice, at 8-day intervals, with autologous monocytes pulsed with one of several PSA or PAP derived peptides (table 1). Twenty days after priming, cytotoxicity was tested against T2 cells pulsed with the same peptide used for boosting. Unlike dendritic cells pulsed with the sPSA plasmid, tVac DCs prime T cells to all PSA- or PAP-derived peptides in all five donors tested (fig. 4). DCs or monocytes, loaded with PSA peptides, support development of T cell effectors with similar efficacy (fig. 4).

### **Repeated boosting with transfected DCs restricts the response towards one immunodominant epitope**

Previously, we found that boosting with polyepitope expressing DCs restricts the immune response towards a single immunodominant epitope. To determine the effect of a prime/boost vaccination strategy on the clonality of the T cell response, tVac DCs-primed cultures, known to contain CTLs to sub-dominant epitopes (fig. 4), were boosted with transfected or peptide pulsed dendritic cells or monocytes, and their cytotoxicity was tested against PSA- or PAP-peptide pulsed T2 targets. Boosting with antigen presenting cells that express multiple PSA- or PAP-derived epitopes (transfected DCs, or DCs or monocytes pulsed with multiple peptides) restricts the immune response towards one immunodominant epitope (table 2), a finding that has already been observed by others.<sup>28</sup> A subdominant T cell response could only be preserved if boosting is performed with an APC (DC or monocyte) pulsed with the particular sub-dominant epitope (table 2; fig.4).

### **Depletion of CD25+ cells prior to initial exposure to antigen leads to generation of T cells reactive to both dominant and sub-dominant epitopes**

Subdominant epitopes are generated when dendritic cells are transfected with sVacDCS, but factors other than TCR signaling such as the CTLA-4/B7 pathway, are contributing to the ineffective proliferation of T cell to subdominant epitopes<sup>1</sup>. Since non-activating anti-CTLA4 antibodies block the suppressor activity of regulatory cells in vitro<sup>29</sup>, we decided to explore the effect of CD25+ cell depletion prior to priming of peripheral blood T cells with genetically modified autologous dendritic cells. Similarly to CTLA-4 inhibition<sup>1</sup>, CD25+ cell depletion prior to priming with sPSA DCs led to stimulation of T cells reactive to sub-dominant PSA-derived epitopes. We interpret this as evidence that T cell responses to sub-dominant epitopes are generated following priming with sPSA DCs but are inhibited by CD25+ cells present in the T cell-enriched fraction (fig.4). Both peptide pulsed DCs and peptide-pulsed MCs support development of T cells effectors in cultures depleted of CD25+ cells prior to priming with sPSA DCs (fig.5).

### **Addition of CD25+ cells back to T cells within the first hour of priming with sPSA DCs reverses immunodominance.**

Suppression of T cells reactive to sub-dominant epitopes by CD25+ cells occurs early during T cell priming. Addition of CD25+ T cell to T cells that are primed by sPSA DCs is suppressive only if CD4+CD25+ T cells are added within the first hour after initiation of culture (figs.6 and 7). No suppression is seen if CD4+CD25+ T cells are added 8 hours after initiation of priming.

### **sPSA DCs co-transfected with the human GITR-L may support priming and development of T cells reactive to sub-dominant epitopes**

Murine CD25+ T regulatory cells expressed high levels of GITR. We find that human CD4+CD25+ cells also express high levels of GITR in an activation-dependent manner (fig.8), similarly to human cytotoxic T cells isolated from tumor lesions<sup>30</sup>. Since signaling through GITR has been found to downregulate the function of T regulatory cells and enhance the development of autoimmunity,<sup>22,23,31,32</sup> we decided to explore the effect of co-transfection of DCs with sPSA and GITR-Ligand. In two out of five separate experiments (donor B and C), such DCs primed T cells to the sub-dominant epitopes (fig.9).

## **DISCUSSION**

T cells that are specific for PSA- or PAP-derived peptides exist in the adult male since both sVac DCs and tVacDCs prime T cells that are cytotoxic to LNCaP cells in vitro (fig.3). Gene-based vaccination in its current mode of application is effective in breaking tolerance to a self-antigen, but the response is narrow and is restricted to few of the potential epitopes. This presents a problem in vaccinology since loss of an MHC haplotype that participates in the conformation of the T cell antigen, or point mutation in the recognized sequence would result in ineffective immune surveillance.<sup>8,14,15,33</sup>

Unlike sVacDCs, tVacDCs prime cytotoxic T cells that are specific for both dominant and sub-dominant epitopes (fig.4). Numerous factors combine to establish an immunodominance hierarchy,<sup>16</sup> among them the ineffective generation and transport of sub-dominant epitopes by antigen-presenting cells (APCs). Since proteasomal degradation is the main source of antigenic fragments destined for MHC presentation,<sup>34</sup> purposeful cytosolic retention of newly synthesized tumor-associated antigens in genetically manipulated antigen presenting cells increases both the quantity and the diversity of such fragments. Dendritic cells, transfected to synthesize such products, clearly have the advantage to prime to both dominant and sub-dominant epitopes (fig.4).<sup>1</sup>

Similarly to boosting with tPSMA transfected DCs,<sup>1</sup> boosting with polyepitope expressing DCs or monocytes, restricts the immune response to the dominant epitope (table 2; fig.4). New vaccines and/or new methods of immunizations need to be developed for those instances. These hopefully will preserve responses to subdominant determinants during re-immunization so that the selection of tumor escape variants that fail to express immunodominant epitopes will be prevented.<sup>8</sup>

In a previous study we found that, under conditions that favor priming of T cells to dominant epitopes, CTLA-4 inhibition alleviates immunodominance. Possible operational mechanisms involved CTLA-4 acting as a non-signaling "decoy" receptor reducing the available ligand for CD28 costimulation<sup>35,36</sup> or creating opportunities for weak signals coming from sub-dominant epitopes otherwise prompt to inhibition by CTLA-4.<sup>37</sup>

A third possibility also existed that needed to be explored.<sup>1</sup> Since non-activating anti-CTLA4 antibodies have been found to block the suppressor activity of regulatory cells *in vitro*<sup>29</sup>, a possibility existed<sup>38,39</sup> that a small number of CTLA-4-expressing and -stimulated T cells exerted a suppressive or regulatory effect on other T cells. These cells appeared to be similar if not identical to T regulatory cells.<sup>40-42</sup> In murine models, suppression of auto-reactive T cells has been attributed to a population of spontaneously occurring CD4+CD25+ T cells.<sup>43</sup> Cells with similar phenotype and function have been found in healthy humans.<sup>44-47</sup> Of interest for cancer immunotherapy is the fact that depleting these cells results in the induction of anti-tumor immune responses, particularly after tumor specific vaccination.<sup>48,49</sup> One hypothesis is that depleting these CD4+CD25+ regulatory T cells in humans enhances a polyclonal T cell response.<sup>21,50</sup>

Removal of CD25+ T cells from the T cell reactive pool prior to priming does result in elimination of immunodominance so that T cells are primed by sVac DCs to both the dominant and the sub-dominant epitopes (fig.5). Similarly to CTLA-4 inhibition,<sup>1</sup> CD25+ cell depletion is effective if performed early during priming – actually within the first hour after initiation of culture (fig.6). Whether alleviation of immunodominance through CTLA-4 inhibition acts through T regulatory cell suppression needs additional experimentation. One way would be to look whether CD25+ cell removal and CTLA-4 inhibition have a synergistic effect. A synergism of cytotoxic T lymphocyte-associated antigen 4 blockade and depletion of CD25(+) regulatory T cells in antitumor therapy has already been described<sup>51</sup> and this suggests that these interventions may act through different pathways. Unfortunately, our current experimentation system does not permit quantitative analysis of responses to sub-dominant epitopes under different conditions.

Finally, the glucocorticoid-induced TNFR (GITR) is expressed at high levels on resting CD4 +CD25+ T regulatory cells and regulates their suppressive phenotype.<sup>32</sup> Antibodies to GITR abrogate suppression, demonstrating a functional role for this receptor in regulating the CD4+CD25+ T cell subset.<sup>22</sup> In our hands, co-transfection of DCs with the natural ligand for GITR leads to T cell priming to sub-dominant epitopes in 2 out of 5 experiments (fig.7). This could be the result of T regulatory cell suppression by GITR signaling.<sup>22,23,31,32</sup> On the other hand, GITR-GITR-L interaction could also provide a costimulatory signal for the antigen-driven proliferation of naive T cells.<sup>52,53</sup> No matter what the mechanism is, GITR-L co-expression during gene-based vaccination may lead to enhancement of the immune response and alleviation of immunodominance. All of these results, however, have been obtained in an *in vitro* experimentation system. Additional *in vivo* experimentation is necessary to validate their significance in immunocompetent hosts.

In conclusion, we have shown that:

1. Dendritic cells transfected with a construct whose product is retained in the cytosol and degraded in the proteasome, prime to both dominant and subdominant epitopes.
2. Early CD25+ cell depletion during priming *in vitro* enhances priming to sub-dominant epitopes.
3. Co-expression of GITR-L during priming may alleviate immunodominance.

## References

1. Mincheff M, Zoubak S, Altankova I, et al. Human dendritic cells genetically engineered to express cytosolically retained fragment of prostate-specific membrane antigen prime cytotoxic T-cell responses to multiple epitopes. *Cancer Gene Ther.* 2003;10(12):907-917.
2. Fong L, Brockstedt D, Benike C, et al. Dendritic cell-based xenoantigen vaccination for prostate cancer immunotherapy. *J Immunol.* 2001;167(12):7150-7156.
3. Eder JP, Kantoff PW, Roper K, et al. A phase I trial of a recombinant vaccinia virus expressing prostate-specific antigen in advanced prostate cancer. *Clin Cancer Res.* 2000;6(5):1632-1638.
4. Wolchok JD, Gregor PD, Nordquist LT, et al. DNA vaccines: an active immunization strategy for prostate cancer. *Semin Oncol.* 2003;30(5):659-666.
5. Mincheff M, Tchakarov S, Zoubak S, et al. *Naked DNA immunizations for immunotherapy of prostate cancer.* in *Annual Meeting of the American Urological Association.* 1999. Dallas TX.
6. Hicklin DJ, Wang Z, Arienti F, et al. beta2-Microglobulin mutations, HLA class I antigen loss, and tumor progression in melanoma. *J Clin Invest.* 1998;101(12):2720-2729.
7. Hiraki A, Kaneshige T, Kiura K, et al. Loss of HLA haplotype in lung cancer cell lines: implications for immunosurveillance of altered HLA class I/II phenotypes in lung cancer. *Clin Cancer Res.* 1999;5(4):933-936.
8. Dunn GP, Bruce AT, Ikeda H, et al. Cancer immunoediting: from immunosurveillance to tumor escape. *Nat Immunol.* 2002;3(11):991-998.
9. Ferrone S, Marincola FM. Loss of HLA class I antigens by melanoma cells: molecular mechanisms, functional significance and clinical relevance. *Immunol Today.* 1995;16(10):487-494.
10. Cordon-Cardo C, Fuks Z, Drobnjak M, et al. Expression of HLA-A,B,C antigens on primary and metastatic tumor cell populations of human carcinomas. *Cancer Res.* 1991;51(23 Pt 1):6372-6380.
11. Rivoltini L, Barracchini KC, Viggiano V, et al. Quantitative correlation between HLA class I allele expression and recognition of melanoma cells by antigen-specific cytotoxic T lymphocytes. *Cancer Res.* 1995;55(14):3149-3157.
12. de la Salle H, Zimmer J, Fricker D, et al. HLA class I deficiencies due to mutations in subunit 1 of the peptide transporter TAP1. *J Clin Invest.* 1999;103(5):R9-R13.
13. van Duinen SG, Ruiter DJ, Broecker EB, et al. Level of HLA antigens in locoregional metastases and clinical course of the disease in patients with melanoma. *Cancer Res.* 1988;48(4):1019-1025.
14. Schreiber H, Wu TH, Nachman J, et al. Immunodominance and tumor escape. *Semin Cancer Biol.* 2002;12(1):25-31.
15. Ikeda H, Old LJ, Schreiber RD. The roles of IFN gamma in protection against tumor development and cancer immunoediting. *Cytokine Growth Factor Rev.* 2002;13(2):95-109.
16. Yewdell JW, Bennink JR. Immunodominance in major histocompatibility complex class I-restricted T lymphocyte responses. *Annu Rev Immunol.* 1999;17:51-88.
17. Sakaguchi S, Sakaguchi N, Asano M, et al. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol.* 1995;155(3):1151-1164.

18. Suri-Payer E, Amar AZ, Thornton AM, et al. CD4+CD25+ T cells inhibit both the induction and effector function of autoreactive T cells and represent a unique lineage of immunoregulatory cells. *J Immunol.* 1998;160(3):1212-1218.
19. Shevach EM. Certified professionals: CD4(+)CD25(+) suppressor T cells. *J Exp Med.* 2001;193(11):F41-46.
20. Levings MK, Sangregorio R, Roncarolo MG. Human cd25(+)cd4(+) t regulatory cells suppress naive and memory T cell proliferation and can be expanded in vitro without loss of function. *J Exp Med.* 2001;193(11):1295-1302.
21. Thornton AM, Shevach EM. CD4+CD25+ immunoregulatory T cells suppress polyclonal T cell activation in vitro by inhibiting interleukin 2 production. *J Exp Med.* 1998;188(2):287-296.
22. McHugh RS, Whitters MJ, Piccirillo CA, et al. CD4(+)CD25(+) immunoregulatory T cells: gene expression analysis reveals a functional role for the glucocorticoid-induced TNF receptor. *Immunity.* 2002;16(2):311-323.
23. Shimizu J, Yamazaki S, Takahashi T, et al. Stimulation of CD25(+)CD4(+) regulatory T cells through GITR breaks immunological self-tolerance. *Nat Immunol.* 2002;3(2):135-142.
24. Kim JD, Choi BK, Bae JS, et al. Cloning and characterization of GITR ligand. *Genes Immun.* 2003;4(8):564-569.
25. DeMars R, Chang CC, Shaw S, et al. Homozygous deletions that simultaneously eliminate expressions of class I and class II antigens of EBV-transformed B-lymphoblastoid cells. I. Reduced proliferative responses of autologous and allogeneic T cells to mutant cells that have decreased expression of class II antigens. *Hum Immunol.* 1984;11(2):77-97.
26. Matzinger P. The JAM test. A simple assay for DNA fragmentation and cell death. *J Immunol Methods.* 1991;145(1-2):185-192.
27. Parker KC, Bednarek MA, Hull LK, et al. Sequence motifs important for peptide binding to the human MHC class I molecule, HLA-A2. *J Immunol.* 1992;149(11):3580-3587.
28. Palmowski MJ, Choi EM, Hermans IF, et al. Competition between CTL narrows the immune response induced by prime-boost vaccination protocols. *J Immunol.* 2002;168(9):4391-4398.
29. Boden E, Tang Q, Bour-Jordan H, et al. The role of CD28 and CTLA4 in the function and homeostasis of CD4+CD25+ regulatory T cells. *Novartis Found Symp.* 2003;252:55-63; discussion 63-56, 106-114.
30. Wang HY, Lee DA, Peng G, et al. Tumor-specific human CD4+ regulatory T cells and their ligands: implications for immunotherapy. *Immunity.* 2004;20(1):107-118.
31. Kwon B, Kim BS, Cho HR, et al. Involvement of tumor necrosis factor receptor superfamily(TNFRSF) members in the pathogenesis of inflammatory diseases. *Exp Mol Med.* 2003;35(1):8-16.
32. Kohm AP, Williams JS, Miller SD. Cutting Edge: Ligation of the Glucocorticoid-Induced TNF Receptor Enhances Autoreactive CD4(+) T Cell Activation and Experimental Autoimmune Encephalomyelitis. *J Immunol.* 2004;172(8):4686-4690.
33. Draenert R, Le Gall S, Pfafferott KJ, et al. Immune Selection for Altered Antigen Processing Leads to Cytotoxic T Lymphocyte Escape in Chronic HIV-1 Infection. *J Exp Med.* 2004;199(7):905-915.

34. Bennink JR, Anderson R, Bacik I, et al. Antigen processing: where tumor-specific T-cell responses begin. *J Immunother.* 1993;14(3):202-208.
35. Masteller EL, Chuang E, Mullen AC, et al. Structural analysis of CTLA-4 function in vivo. *J Immunol.* 2000;164(10):5319-5327.
36. Doyle AM, Mullen AC, Villarino AV, et al. Induction of cytotoxic T lymphocyte antigen 4 (CTLA-4) restricts clonal expansion of helper T cells. *J Exp Med.* 2001;194(7):893-902.
37. Manzotti CN, Tipping H, Perry LC, et al. Inhibition of human T cell proliferation by CTLA-4 utilizes CD80 and requires CD25+ regulatory T cells. *Eur J Immunol.* 2002;32(10):2888-2896.
38. Read S, Malmstrom V, Powrie F. Cytotoxic T lymphocyte-associated antigen 4 plays an essential role in the function of CD25(+)CD4(+) regulatory cells that control intestinal inflammation. *J Exp Med.* 2000;192(2):295-302.
39. Takahashi T, Tagami T, Yamazaki S, et al. Immunologic self-tolerance maintained by CD25(+)CD4(+) regulatory T cells constitutively expressing cytotoxic T lymphocyte-associated antigen 4. *J Exp Med.* 2000;192(2):303-310.
40. Shevach EM. Regulatory T cells in autoimmunity\*. *Annu Rev Immunol.* 2000;18:423-449.
41. Groux H, Powrie F. Regulatory T cells and inflammatory bowel disease. *Immunol Today.* 1999;20(10):442-445.
42. Sakaguchi S. Regulatory T cells: mediating compromises between host and parasite. *Nat Immunol.* 2003;4(1):10-11.
43. Morse MA, Clay TM, Mosca P, et al. Immunoregulatory T cells in cancer immunotherapy. *Expert Opin Biol Ther.* 2002;2(8):827-834.
44. Sakaguchi S. Naturally Arising CD4+ Regulatory T Cells for Immunologic Self-Tolerance and Negative Control of Immune Responses. *Annu Rev Immunol.* 2004;22:531-562.
45. Jonuleit H, Schmitt E, Stassen M, et al. Identification and functional characterization of human CD4(+)CD25(+) T cells with regulatory properties isolated from peripheral blood. *J Exp Med.* 2001;193(11):1285-1294.
46. Misra N, Bayry J, Lacroix-Desmazes S, et al. Cutting Edge: Human CD4(+)CD25(+) T Cells Restrain the Maturation and Antigen-Presenting Function of Dendritic Cells. *J Immunol.* 2004;172(8):4676-4680.
47. Baecher-Allan C, Viglietta V, Hafler DA. Human CD4+CD25+ regulatory T cells. *Semin Immunol.* 2004;16(2):89-98.
48. Casares N, Arribillaga L, Sarobe P, et al. CD4+/CD25+ regulatory cells inhibit activation of tumor-primed CD4+ T cells with IFN-gamma-dependent antiangiogenic activity, as well as long-lasting tumor immunity elicited by peptide vaccination. *J Immunol.* 2003;171(11):5931-5939.
49. Manjili MH, Wang XY, Chen X, et al. HSP110-HER2/neu chaperone complex vaccine induces protective immunity against spontaneous mammary tumors in HER-2/neu transgenic mice. *J Immunol.* 2003;171(8):4054-4061.
50. Piccirillo CA, Shevach EM. Cutting edge: control of CD8+ T cell activation by CD4+CD25+ immunoregulatory cells. *J Immunol.* 2001;167(3):1137-1140.
51. Suttmuller RP, van Duivenvoorde LM, van Elsas A, et al. Synergism of cytotoxic T lymphocyte-associated antigen 4 blockade and depletion of CD25(+) regulatory T cells in antitumor therapy reveals alternative pathways for suppression of autoreactive cytotoxic T lymphocyte responses. *J Exp Med.* 2001;194(6):823-832.

52. Tone M, Tone Y, Adams E, et al. Mouse glucocorticoid-induced tumor necrosis factor receptor ligand is costimulatory for T cells. *Proc Natl Acad Sci U S A*. 2003;100(25):15059-15064.
53. Ronchetti S, Zollo O, Bruscoli S, et al. GITR, a member of the TNF receptor superfamily, is costimulatory to mouse T lymphocyte subpopulations. *Eur J Immunol*. 2004;34(3):613-622.


**Table 1. HLA A201-restricted PSA- and PAP-derived peptides**

Rank	Start position	Subsequence residue listing	BIMAS Score (estimate of half-time of disassociation of a molecule containing this sequence)
1 PSA	170	KLQCVDLHV	243
2 PSA	52	GVLVHPQWV	124
3 PSA	53	VLVHPQWVL	123
1 PAP	135	ILLWQPIPV	437
2 PAP	112	TLMSAMTNL	182
3 PAP	33	KELKFVTLV	153

**Table 2.** Priming with tVac DCs stimulates T cells to all 3 PSA- or PAP-derived epitopes. Boosting with dendritic cells and monocytes that express multiple antigenic epitopes restricts the response to an immunodominant epitope

	Number of Patients Developing CTL Activity against PSA or PAP Peptide-Pulsed T2 cells					
	PSA <sub>170</sub>	PSA <sub>52</sub>	PSA <sub>53</sub>	PAP <sub>135</sub>	PAP <sub>112</sub>	PAP <sub>33</sub>
Prime: tVac (PSA or PAP) DCs 2 Boosts tVac(PSA or PAP) DCs	5 of 5	1 of 5	0 of 5	5 of 5	0 of 5	1 of 5
Prime: tVac (PSA or PAP) DCs 2 Boosts 1P (PSA- or PAP-derived) MCs	5 of 5	4 of 5	4 of 5	5 of 5	3 of 5	5 of 5
Prime: tVac (PSA or PAP) DCs 2 Boosts 3P (PSA- or PAP-derived) MCs	5 of 5	0 of 5	0 of 5	5 of 5	0 of 5	0 of 5
Prime: sVac (PSA) DCs 2 Boosts sVac (PSA) DCs	5 of 5	0 of 5	0 of 5	5 of 5	0 of 5	0 of 5
Prime: s Vac (PSA) DCs. 2 Boosts 1P (PSA-derived) MCs	5 of 5	0 of 5	0 of 5	5 of 5	0 of 5	1 of 5

\* *tVac* – “truncated” (no signal sequence) plasmid DNA vaccine; *sVac* – plasmid vaccine that encodes for the mature form of human PSA with a signal sequence; *1P* – pulsed with single either PSA-, or PAP-derived peptide; *3P* – pulsed with all three, either PSA- or PAP-derived, peptides;; *MCs* – monocytes



**FIGURES AND LEGENDS TO FIGURES**




Fig. 1. A. Detection of human PSA in cell lysates. Line 1: The parent AT3B-1; Line 2: AT3B-1 transfected with hPSA plasmid, mass culture; Line M: molecular weight marker; Line 3: D12/1, a PSA producing AT3B-1 clone (reduced conditions); Line 4: The D12/1 clone (non-reduced conditions); Lines 5 and 6: Recombinant PSA protein, reduced and non-reduced conditions respectively, loading 200 ng per line.

Fig. 1. B. Detection of secreted human PSA in the culture medium. M molecular weight marker; Line 1: culture medium from the clone D12/1; Line 2: culture medium from the parent AT3B-1 cell line; Line 3: culture medium from a sham transfected AT3B-1 cell line. Cells were grown during 60h in serum free medium, then media were collected, concentrated by filtration and normalized by protein mass.






Fig.2. Staining COS-1 cells with FITC-labeled monoclonal antibody (clone 109114) against the GITR-L. A – prior to transfection; B – following transfection with plasmid encoding for the human GITR-L.




Fig.3. Generation of CTLs by:

- A. PSA DCs (tPSA – squares; sPSA - triangles)
- B. tPAP DCs (squares).

Non-adherent (T cell-enriched) HLA A2(+) peripheral blood mononuclear cells were primed and then boosted twice at 8-day intervals with autologous monocytes-derived dendritic cells that were transiently transfected with either sPSA, tPSA or tPAP plasmid. The medium was changed, initially 72 hours following priming, and then during boosting. The cells were grown in IL-2 and IL-7 medium for 20 days and the specific cytotoxicity was tested against LNCaP cells. Each point represents the mean and SD of

triplicate experiments. Control experiments involved priming and boosting of the T-cell enriched fraction with DCs transfected with empty plasmid (diamonds) or testing of cytotoxicity against the Malme-3M melanoma cell line (circles).




Fig. 4. tVacDCs but not sVacDCs prime T cells that are reactive to sub-dominant PSA or PAP epitopes. Induction of CTL responses with autologous dendritic cells that have been transfected with either the secreted or truncated version of the human PSA. Peripheral blood mononuclear cells that had been depleted of monocytes were primed with autologous DCs transfected with either sPSA (first column) or tPSA (second and third column). Responding cultures were then boosted with monocytes pulsed with one of three HLA-A2 restricted PSA-derived peptides (table 1) and their cytotoxicity was tested against T2 cells pulsed with the same peptide that was used during boosting (PSA<sub>170</sub> – squares; PSA<sub>51</sub> – triangles and PSA<sub>53</sub> – circles). Data points for the control (influenza) peptide are not shown but are identical to those obtained with empty plasmid-transfected DCs (donor A) in all donors tested. Each point represents the mean and SD from three different experiments. Both tPSA DCs- and sPSA DCs-primed T cells are cytotoxic against T2 cells pulsed with PSA<sub>170</sub> peptide. Values for cytotoxicity of either tPSA DCs- or sPSA DCs-primed T cells against T2 targets pulsed with PSA<sub>170</sub> and PSA<sub>51</sub> or PSA<sub>53</sub> peptides were compared. Significant differences at the 30:1 E/T ratio are indicated with xx (p<0.01). Level of cytotoxicity for sPSA DCs primed T cells against T2 cells pulsed with either PSA<sub>51</sub> or PSMA<sub>53</sub> for all 5 donors are identical to controls (T cells primed with empty plasmid transfected DCs).




Fig. 5. Depletion of CD25<sup>+</sup> cells prior to initial exposure to antigen leads to generation of T cells reactive to both dominant and sub-dominant epitopes. HLA A2(+) peripheral blood mononuclear cells that had been depleted of monocytes, with or without additional removal of CD25<sup>+</sup> cells, were primed with autologous dendritic cells transfected with the sPSA plasmid. Responding cultures were then boosted twice with monocytes pulsed with one of three HLA-A2 restricted PSA-derived peptides (PSA<sub>170</sub> – squares; PSA<sub>51</sub> – triangles or PSA<sub>53</sub> – diamonds) (table 1) and their cytotoxicity was tested against T2 cells pulsed with the same peptide that was used during boosting. Each point represents the mean and SD from three different experiments. Values for cytotoxicity of sPSA DCs-

primed T cells against T2 targets pulsed with PSA<sub>170</sub>, PSA<sub>51</sub> or PSA<sub>53</sub> peptides were compared. Significant differences at the 30:1 E/T ratio are indicated with xx (p<0.01).




Fig.6. Addition of CD4<sup>+</sup>CD25<sup>+</sup> cells back to T cells within the first hour of priming with sPSA DCs reverses immunodominance. HLA A2(+) peripheral blood mononuclear cells that had been depleted of monocytes and CD25<sup>+</sup> cells were primed with autologous dendritic cells transfected with the sPSA plasmid. Purified CD4<sup>+</sup>CD25<sup>+</sup> T cells were added back at different time points following the initiation of the cultures (responder T cells/CD4<sup>+</sup>CD25<sup>+</sup> T cell ratio = 4:1). Responding cultures were then boosted twice with

monocytes pulsed with one of three HLA-A2 restricted PSA-derived peptides (PSA<sub>170</sub> – squares; PSA<sub>51</sub> – triangles or PSA<sub>53</sub> – diamonds) and their cytotoxicity was tested against T2 cells pulsed with the same peptide that was used during boosting. Each point represents the mean and SD from three different experiments. Values for cytotoxicity of sPSA DCs-primed T cells against T2 targets pulsed with PSA<sub>170</sub>, PSA<sub>51</sub> or PSA<sub>53</sub> peptides were compared. Significant differences at the 30:1 E/T ratio are indicated with xx (p<0.01).




Fig.7. CD4<sup>+</sup>CD25<sup>+</sup> T cells act early during priming to establish immunodominance to PSA<sub>52</sub>. HLA A2(+) peripheral blood mononuclear cells that had been depleted of monocytes and CD25<sup>+</sup> cells were primed with autologous dendritic cells transfected with the sPSA plasmid. Purified CD4<sup>+</sup>CD25<sup>+</sup> T cells were added back at different time points (triangles – before; squares – 1 hour after; diamonds – 8 hours after) onset of the (responder T cells/CD4<sup>+</sup>CD25<sup>+</sup> T cell ratio = 4:1). Responding cultures were then boosted twice with PSA52-pulsed monocytes and their cytotoxicity was tested against T2 cells pulsed with the same peptide.




Fig. 8. Activation of human CD4<sup>+</sup>CD25<sup>+</sup> T cells increases the membrane GITR expression. Freshly isolated (1) or stimulated human CD4<sup>+</sup>CD25<sup>+</sup> T cells stained with anti-GITR antibody. FITC conjugated mouse IgG1 was used as a control.






Fig.9. sPSA DCs co-transfected with the human GITR-L may support priming and development of T cells reactive to sub-dominant epitopes. HLA A2(+) peripheral blood mononuclear cells that had been depleted of monocytes were primed with sPSA plasmid-transfected autologous dendritic cells that had (filled in symbols) or had not been (open

symbols) transfected with the human GITR-L. Responding cultures were then boosted twice with monocytes pulsed with one of three HLA-A2 restricted PSA-derived peptides (PSA<sub>170</sub> – squares; PSA<sub>51</sub> – triangles or PSA<sub>53</sub> – diamonds) and their cytotoxicity was tested against T2 cells pulsed with the same peptide that was used during boosting. Each point represents the mean and SD from three different experiments. Values for cytotoxicity of sPSA DCs-primed T cells against T2 targets pulsed with PSA<sub>51</sub> or PSA<sub>53</sub> peptides were compared (third column). Significant differences at the 30:1 E/T ratio are indicated with xx (p<0.01; reactivity against PSA<sub>53</sub> for donor B and PSA<sub>51</sub> or PSA<sub>53</sub> for donor C).

**C. Manuscript published in *Cancer Gene Therapy* 2006 Apr;13(4):436-44.**

Immune responses against PSMA after gene-based vaccination for immunotherapy: A. Results from immunizations in animals

Milcho Mincheff, Serguei Zoubak, Yevgen Makogonenko

The George Washington University Medical Center, Washington, DC

Corresponding author:

Milcho Mincheff

Head, Tumor Immunology Laboratory

Department of Medicine

The George Washington Medical Center

2300 Eye Street, N.W., Ross Hall 705

Washington, DC 20037

Telephone 202 994 7765; fax 202 994 0465

E-mail: [mcamsm@gwumc.edu](mailto:mcamsm@gwumc.edu)

Running title: Anti-PSMA gene-based vaccination in animals

Keywords: PSMA, Gene-based vaccine, prostate cancer, cytotoxicity, antibody production

Supported in part by grant N00014-00-1-0787 from the Office of Naval Research.

Supported in part by award No DAMD17-02-1-0239. The U.S. Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick, MD 21702-5014 is the awarding and administering acquisition office.

The content of the information does not necessarily reflect the position or the policy of the Government, and no official endorsement should be inferred. For purpose of this article, information includes news releases, articles, manuscripts, brochures, advertisements, still and motion pictures, speeches, trade association proceedings etc.

## **ABSTRACT**

Two plasmid DNA vaccines, encoding either products that are retained in the cytosol and degraded in the proteasome (tVacs; hPSMA<sub>t</sub>), or secreted proteins (sVacs; hPSMA<sub>s</sub>) were evaluated for stimulation of cytotoxic cell or antibody responses. Immunization with both vectors led to generation of cell cytotoxicity providing GM-CSF was administered with the vaccine. Spleen cells from animals immunized with hPSMA<sub>t</sub> demonstrated stronger cytotoxicity to the target cells. Priming with a vector that encoded a xenogeneic protein (hPSMA<sub>t</sub>; “xenogeneic” construct) and boosting with a vector that encoded an autologous protein (rPSMA<sub>t</sub>; “autologous” construct) gave the best protection against tumor challenge. Immunization with tVacs did not lead to formation of antibodies to the target protein as detected by Western blot or ELISA, while immunization with sVacs or with the protein did. Antibodies were of mixed Th1-Th2 isotype. Priming with tVacs and boosting with protein also resulted in antibody formation, but in this case the antibodies were from the cytotoxic, Th1 isotype. The best strategy to obtain a strong cellular cytotoxic response, therefore, seems to be gene-based vaccinations with tVacs, priming with the “xenogeneic” and boosting with the “autologous” constructs. When cytotoxic antibody production is the goal, priming should be performed with the tVacs while boosting with the protein.

## INTRODUCTION

Several groups have recently reported on the safety and biologic effectiveness of DNA vaccines for immunization against pathogen or tumor antigens.<sup>1-4</sup> The use of PSA-recombinant vaccinia virus has also been found to be safe and to lead to an immune response.<sup>5,6</sup> The main effectors in anti-tumor immunity after DNA immunization are CD8+ cytotoxic T cells that recognize tumor or tumor-associated antigen-derived peptides expressed in association with MHC class I molecules.<sup>7</sup> A rate limiting step in loading the latter with peptides seem to be the antigen supply.<sup>8</sup> We have found that newly synthesized proteins that are purposefully retained in the cytosol are targeted for proteasomal degradation.<sup>9</sup> Transfection with truncated DNA constructs (tVacs) that lack sequences encoding the leader peptide of a secreted protein, or the membrane domain of type II protein, lead to expression of such products. In our immunotherapeutic studies, we have targeted the prostate-specific membrane antigen (PSMA; gamma carboxyl peptidase II, GCPII). PSMA expression is normally restricted to the prostate gland, brain tissue, jejunum and proximal kidney tubules.<sup>10-12</sup> Its expression is increased nearly 10-fold in prostate cancer cells and is also found in tumor but not normal neovasculature.<sup>13,14</sup> Elevated PSMA serum levels have been observed in healthy males and females, as well as in patients with benign prostate hypertrophy, prostate and breast cancer.<sup>15-17</sup> It is a type II integral membrane protein<sup>18,19</sup> whose main portion is extracellular. We have cloned the cDNA encoding the extracellular portion (AA 44-750) of the human PSMA (XC-PSMA) and included it in a mammalian expression vector (hPSMA<sub>t</sub>). Dendritic cells, transfected with hPSMA<sub>t</sub>, stimulate in vitro autologous T cell proliferation and cytotoxicity against the target antigen.<sup>9</sup> We speculate that, similarly to results from in vitro experiments, in vivo immunization with this vector will result in enhanced cytotoxicity and protection from tumor development.<sup>9,20</sup>

Additionally, gene-based vaccines encoding for a xenogeneic rather than autologous protein have been shown to more likely break existing immunological tolerance.<sup>3,21-24</sup> We tested this approach and compared immunizations in rats with tVacs encoding the human PSMA or its rat equivalent. Additionally, since transfection with hPSMA<sub>t</sub> does not result in the expression of a full length glycosylated and folded protein,<sup>9</sup> it is of academic interest whether immunization with hPSMA<sub>t</sub>s will lead to formation of antibody against the native antigen, against carbohydrate residues or against linear epitopes. To study this, we have developed a transplantable tumor rat model in which animals are immunized against PSMA with different DNA vectors with or without recombinant rat GM-CSF, and then inoculated with syngeneic tumor cells (AT3B-1 cells) transfected to express the target protein.

## MATERIALS AND METHODS

### *Rats and Immunizations and Tumor Cell Inoculation*

All animal experiments were approved by the George Washington University Medical Center (GWUMC) Institutional Animal Care and Use Committee. Male retired breeders (Copenhagen rats) were purchased from Harlan Sprague Dawley Laboratories (Indianapolis, IN), maintained at the GWUMC Animal Research Facility (ARF), and fed a standard diet. The ARF has maintained continuous accreditation by the Association for Assessment and Accreditation of Laboratory Animal Care since 1974, and has an

assurance statement on file with the Office Laboratory Animal Welfare, thus meeting all Public Health Service guidelines for animal use and care.

For immunizations, 100 µg plasmid DNA with or without rat recombinant GM-CSF (R&D Systems, Minneapolis, MN), were brought to 100 µl with sterile saline for injection. The immunizations were performed at 10 day-intervals at the right rear flank using insulin syringes. All injections were intradermal, taking caution that a blister was formed and the infusate was retained at the epidermal-dermal junction.

For tumor cell inoculation,  $10^5$  of AT3B-1<sup>PSMA</sup> cells (the YM2 clone, see “AT3B-1 transfection with hPSMAs”) were resuspended in 100 µl sterile saline and injected subcutaneously in the opposite rear flank.

### *Plasmid DNA Vaccines*

The H PSMA-K plasmid (hPSMA<sub>H</sub>) was constructed on the basis of pVAX1 vector (Invitrogen, Carlsbad, CA) and contained a pUC derived pMB1 origin of replication, a kanamycin resistance gene and the mammalian transcriptional unit with DNA encoding for the extracellular portion of the human PSMA under the regulation of the immediate-early CMV promoter/enhancer and a bovine growth hormone polyadenylation signal. The plasmid size is 5.1 kb.

For the R (rat) “PSMA”-K plasmid (rPSMA<sub>R</sub>), cDNA encoding for a rat analogue of the human extracellular PSMA was cloned from rat brain tissue and included in a pVAX1 backbone. The cDNA had 85.66% homology with the human PSMA at the nucleotide level. The plasmid size was 5.1 kb. Products expressed from hPSMA<sub>H</sub> or rPSMA<sub>R</sub> plasmids were retained in the cytosol and degraded in the proteasome<sup>9</sup>.

For the S-PSMA plasmid (hPSMA<sub>S</sub>), the extracellular portion of the human PSMA was cloned into a pSecTag2 vector (Invitrogen), which provided the murine Ig k-chain leader sequence. The insert from the obtained clone, which represents human XC-PSMA fused with leader sequence, was then sub-cloned into a pVAX1 expression vector (Invitrogen, Carlsbad, CA). The plasmid size was 5.1 kb. Following transfection with this vector, the expressed product was translocated to the endoplasmic reticulum (ER), glycosylated and secreted<sup>9</sup>. Cells, transfected with hPSMA<sub>H</sub> or hPSMA<sub>S</sub> synthesize proteins that have identical aminoacid sequences. Due to post-translational modification in the ER, however, the secreted protein is N-glycosylated and secreted, while the other is not transported to the ER and not glycosylated, but remains in the cytosol and is degraded in the proteasome.<sup>9</sup>

For transfection of the AT3B-1 cells with the human PSMA, the extracellular portion of the human PSMA was cloned into a pSecTag2 vector containing zeocin resistance gene.

### *AT3B-1 transfection with hPSMAs*

Monolayers of AT3B-1 cells (CRL-2375; ATCC, Manassas, VA) were transfected with the hPSMA<sub>S</sub> using the FuGENE6 transfection reagent (Roche Diagnostics Corporation, Indianapolis, IN) and assayed for PSMA production by Western blot. The cells were seeded in 6-well tissue culture plates (Nunc, Denmark) at  $1.5 \times 10^5$  cells per well and grown to 50-70% confluence in DMEM supplemented with 25 mM HEPES (pH 7.5), 1 mM sodium pyruvate, 3.7 g/L sodium bicarbonate, 100 µg/ml

penicillin, 100 µg/ml streptomycin, 0.25 µg/ml amphotericin B and 10% (v/v) of fetal bovine serum. COS-1 or AT3B-1 cells were transfected with 1.5 µg of plasmid DNA pre-condensed with 4.5 µl of FuGENE 6 reagent in serum-free DMEM for 30 min at room temperature. Cells were then grown for 72 h in complete DMEM and then harvested.

The transfected cells with stable expression of hPSMAs were selected and cloned after growth in zeocin-containing medium (0.1 mg/ml).

### *Electrophoresis and immunoblotting*

For electrophoresis and Western blotting, cells were harvested by gentle scraping, washed twice with 2 ml of cold PBS and lysed with 0.25 ml cold lysing buffer (0.5 M NaCl, 1% triton X-100, 0.2% Tween 20, 50 mM HEPES, pH 7.0). Lysates were transferred to Eppendorf tubes and homogenized by repeated pipetting on ice. Immunoblotting and electrophoretic assays of cell lysates and culture supernatants were performed using NuPage BisTris electrophoretic system (Invitrogen, Carlsbad, CA). Protein samples in loading buffer were heated at 70°C for 10 min and loaded on 10% BisTris gels. After electrophoresis and electro-transfer, the nitrocellulose membrane was blocked with 1% casein in TBS/T for 40 min. The following primary antibodies were used: Y-PSMA1 and Y-PSMA-2 (Yes Biotech Laboratories Limited, Mississauga, Ontario, Canada), rabbit polyclonal anti-GCPII antibodies, anti-GCPII monoclonal antibodies A5, E1, E7, E12 and H2 (all developed in the Institute of Organic Chemistry and Biochemistry, Prague, the Czech Republic). The membranes were probed with poly- or monoclonal anti-PSMA Abs for 1 h at RT. Detection was performed using goat anti-rabbit or anti-mouse IgG-HRP conjugates (Sigma, St. Louis, MO) and visualized with WestPico Super Signal Chemo luminescent Substrate (Pierce, Rockford, IL) in accordance with the manufacturer's recommendations.

For detection of secreted PSMA, the serum containing DMEM was removed 48 h after transfection, the cells were washed twice with 2 ml of PBS, serum free DMEM (2 ml per well) was added and the cells were incubated for additional 24 h in the 6-well plates. After collection of the medium, the cell debris was removed by centrifugation (35,000 g, 20 min) and supernatants were concentrated with Centricom centrifuge filtering device (Millipore, Bedford, MA) and then stored at -30°C until further use.

For a control, an extracellular portion of the human glutamate carboxypeptidase II (amino acids 44-750) expressed in *Drosophila Schneider's* cells, purified to homogeneity (GCP II) was used.<sup>25</sup> This material, together with the rabbit anti-PSMA serum and some of the monoclonal antibodies were provided to us by Dr. Jan Konvalinka, Institute of Organic Chemistry and Biochemistry, Czech Academy of Science, Prague, The Czech Republic.

### *Cell Proliferation*

Standard [<sup>3</sup>H]thymidine incorporation assays were performed to assess spleen cell proliferative responses. Splenocytes from immunized rats were resuspended at 5 x 10<sup>6</sup> cells/ml in RPMI 1640 containing 10% FBS. One hundred microliters of cell suspension were added to each well in 96-well plates with 20, 10, 5, or 0 µg/ml of recombinant PSMA. After 4 days of culture, 1 µCi [<sup>3</sup>H]thymidine (Amersham Biosciences, Piscataway, NJ) was added to each well. Following a 16-h incubation, cells were harvested on glass filter paper and radioactivity was measured in a Beckman LS 2501

liquid scintillation counter. The stimulation index (SI) was calculated as: counts per minute (cpm) with Ag stimulation/background cpm without Ag.

### *Cell Cytotoxicity*

Cell Cytotoxicity was determined as previously described.<sup>26</sup> Briefly, target cells (AT3B-1 or YM2 cells) were grown overnight in complete medium supplemented with 1  $\mu$ Ci [<sup>3</sup>H]thymidine (Amersham Biosciences, Piscataway, NJ) then washed and resuspended in complete medium. Cells in different target to effector cell ratio were plated in 96 round bottom well plates for 6 hours at 37°C in a humidified incubator. Cells were then harvested on glass filter paper, and radioactivity was measured in a liquid scintillation counter. The killing was detected as a fall in counts per minute due to DNA fragmentation in cell samples undergoing apoptosis. All of the E:T ratios were tested in triplicate. Cell cytotoxicity was normalized to control samples that contained target cells only and expressed as specific lysis. Spontaneous cytotoxicity was determined in medium alone without effector cells.

### *Tumor Development*

Animal weight and tumor development were monitored daily. Tumor size was measured after the animals were euthanized and the tumors were excised.

### *Antibody determination*

For Western blot, SDS gel electrophoresis of cell lysates, cell supernatants or GCP II were performed as described above. After electrotransfer and blocking, the nitrocellulose membranes were stained with sera from immunized rats or anti-PSMA antibodies for an hour. Secondary staining with antibodies conjugated to horseradish peroxidase (Sigma, St. Louis, MO; dilution 1:10,000; for one hour) was performed after three washes with TBS, 0.05% Tween20. Finally, the blots were developed using SuperSignal Chemiluminescent Substrate (Pierce, Rockford, IL) and exposed to BioMax MR film (Kodak, New Haven, CT).

In some cases, serum anti-PSMA antibody titers from immunized rats were measured by a direct ELISA. Ninety-six-well plates coated overnight with 100  $\mu$ l/well of 10  $\mu$ g/ml recombinant GCPII in PBS were blocked for 2 hours with PBS containing 2% BSA and 0.05% Tween20. Sera were then added in serial dilutions and incubated for 1 hour. The plates were washed three times with PBS containing 0.05% Tween20 and incubated for 1 hour with 1/5000 dilution of a peroxidase conjugated anti-rat kappa-lambda chain secondary antibody (Jackson Laboratories, Bar Harbor, ME). The plates were then washed three times, developed with tetramethylbenzidine (Kirkegaard & Perry, Gaithersburg, MD), stopped with 1% HCl, and analyzed at 450



nm with a Dynatech MR5000 ELISA plate reader (Dynatech Laboratories, Alexandria, VA).

### *Statistical Analyses*

Significant differences in cytotoxicity of spleen cells or antibody titers were assessed by Student's *t* test. The difference between groups was considered statistically significant when the *P* was lower than 0.05. A non-parametric test of statistical significance for bivariate tabular analysis (chi-square) was performed for data accumulated when tumor development was assayed.

## **RESULTS**

### *Transfection of AT3B-1 cells with hPSMAs*

We find that both rat prostate tissue and AT3B-1 cell lysates express low level of PSMA when tested by PCR (data not shown). Since PSMA expression is up-regulated in human prostate cancer cells, we decided to transfect the AT3-B1 cells with the hPSMAs, a vector that contains the murine Ig k-chain leader sequence so that the extracellular portion of the human PSMA is glycosylated and secreted<sup>9</sup>. We did not transfect the cells with the full length PSMA since our goal was to study T cell-mediated and not antibody-mediated tumor protection and we felt that if PSMA was expressed as a membrane protein by the transfected AT3B-1 cells, some of the protection might have been antibody mediated. This would have been true especially when immunization was performed with DNA constructs that led to anti-PSMA antibody development (see below). We obtained two clones of stably transfected AT3B-1 cells – YM1 and YM2, both of which are high expressors and PSMA can be detected in the culture medium (figure 1). A significant portion of the expressed product is degraded in the proteasome and could be detected in cell lysates only in the presence of proteasomal inhibitor lactacystin (figure 1). All further experimentation was performed with the YM2 clone.

### *Addition of rat GM-CSF to the immunization cocktail increases the efficacy of immunization*

Male Copenhagen rats were immunized with the plasmid vaccines alone (figure 2A), or with a cocktail that, in addition to the plasmid vaccine, contained 9  $\mu\text{g}/\text{m}^2$  recombinant rat GM-CSF (R&D, Minneapolis, MN) (figure 2B and C). In separate experiments, rats were primed with the hPSMA<sub>1</sub> GM-CSF, but boosted with the rPSMA<sub>1</sub> GM-CSF cocktail (figure 2D). Addition of GM-CSF to the immunization cocktail increased the efficacy of the immunization (figure 2B and C). Rats immunized with hPSMA<sub>1</sub> showed stronger spleen cell cytotoxicity to YM2 compared to those immunized with rPSMA<sub>1</sub>, but the best results were obtained when priming was performed with hPSMA<sub>1</sub>, while boosting with rPSMA<sub>1</sub> (figure 2D).

Slightly stronger cytotoxicity to YM2 cells was observed in spleen cells from animals immunized with hPSMA<sub>t</sub> when compared to spleen cells from animals immunized with hPSMA<sub>s</sub> (figure 3).

*In the absence of GM-CSF, immunization with a construct encoding either human or rat PSMA partially protects the rats from developing tumors*

Copenhagen rats, inoculated sub-cutaneously with  $10^5$  syngeneic AT3B-1<sup>PSMA</sup> YM2 tumor cells, develop detectable tumors by day 10-12 following injection. To explore the role of pre-immunization on tumor development, rats were immunized three times at 10-day intervals with either saline, empty plasmid vector, hPSMA<sub>t</sub> or rPSMA<sub>t</sub> (table 1). No GM-CSF was included in the immunization cocktail. Fourteen days after the last immunization, all animals received  $10^5$  of YM2 tumor cells in the right rear flank. Tumors, when developing, were palpable at day 12 following tumor cell inoculations. All animals were sacrificed 15 days following tumor inoculations and tumors, when present, excised and measured. Only two of the ten animals immunized against human PSMA and five of the 10 animals immunized against the rat analogue developed tumors. In contrast, 17 of the 20 animals in the control groups developed tumors. The tumors, developing in the group, immunized with the human or the rat construct, were smaller than the ones in the control group (table 1).

*Addition of rat GM-CSF to the immunization cocktail enhances gene-based vaccination-induced tumor protection*

From an earlier clinical trial,<sup>4</sup> we found that the magnitude of the response could be increased if the plasmid vaccine is co-injected with GM-CSF, a cytokine that has been known for its adjuvant properties.<sup>27</sup> The above experiment was repeated, but this time recombinant rat GM-CSF ( $9 \mu\text{g}/\text{m}^2$ ) was added to the immunization cocktail. In this study, full protection was observed when recombinant rat GM-CSF was used as an ingredient of the immunization cocktail (table 2), no matter whether animals were vaccinated with the vaccine encoding the human or the rat protein.

*In the presence of GM-CSF, priming with a construct encoding a xenogeneic protein and boosting with a construct encoding an autologous protein lead to the best protection*

The protection of the immunized animals from development of tumor depends on the amount of cells inoculated during tumor challenge. Partial tumor protection could be seen when  $2 \times 10^5$  or  $3 \times 10^5$  tumor cells are injected. All immunized animals develop tumors when the tumor dose per animal is higher than  $4 \times 10^5$  cells. In an attempt to define the best strategy of vaccination, we explored different priming and boosting strategies. All immunizations were performed with plasmid DNA cocktails that contained rat GM-CSF. All animals were challenged with  $2 \times 10^5$  cells. Rats were sacrificed on the day tumor

was detected. All tumor-free animals were sacrificed at day 16 following tumor cell inoculation. Ten rats were included per group. The best protection following immunization involved priming with hPSMA<sub>t</sub> and boosting with rPSMA<sub>t</sub> and all the animals were tumor free 16 days after tumor inoculation (figure 4; circles). In contrast, when priming was performed with hPSMA<sub>s</sub> and boosting with rPSMA<sub>t</sub>, three rats developed tumors on day 12, while the rest of the rats (66%) remained tumor free. The difference between these two groups was not statistically significant ( $\chi^2 = 3.52$  at  $p < 0.1$ ). Spleen cells from the hPSMA<sub>s</sub>-immunized animals, however, gave the strongest proliferative response after stimulation with recombinant PSMA (data not shown). Priming and boosting with rPSMA<sub>t</sub> resulted in only partial protection – three rats developed tumor on day 10, and 3 rats on day 11. Four of those rats (34%) remained tumor free until day 16 (figure 4, diamonds). The difference between this group and the control group, though, was statistically significant ( $\chi^2 = 5$  at  $p < 0.05$ ) since all control rats developed tumors by day 12 after tumor inoculation.

#### *Antibodies against PSMA in animals following immunization with different gene-based vaccines*

Rats immunized with “truncated” constructs (hPSMA<sub>t</sub> or rPSMA<sub>t</sub>) did not develop antibodies to PSMA as detected by ELISA (figure 5) or Western blot (data not shown), no matter whether GM-CSF was present or not. No antibodies against PSMA (detection by Western blot or ELISA) developed following immunization with plasmid, encoding the secreted PSMA (hPSMA<sub>s</sub>), or with GCP II, when GM-CSF was absent from the immunization cocktail (figure 5). Immunization with hPSMA<sub>s</sub> in the presence of GM-CSF led to formation of antibodies against the native protein (detected by ELISA; figure 5). The highest titer of antibodies against PSMA was observed when priming was performed with GCPII or tVacs, and boosting was performed with the recombinant protein and GM-CSF was added as an adjuvant (figures 5 and 6). Antibodies, developing following immunization with hPSMA<sub>s</sub> or GCPII, were of mixed Th2 and Th1 type, since both IgG1 and IgG2a subtypes were detected. On the contrary, priming with tVacs (hPSMA<sub>t</sub> or rPSMA<sub>t</sub>) and boosting with hPSMA<sub>s</sub> or GCP II led to formation of Th1, cytotoxic antibodies (IgG2a and IgG2b) with relatively high titer.

## **DISCUSSION**

Immunotherapy holds a great promise for controlling and treatment of cancer, but effective immunization against tissue-specific, self-antigens is necessary in order to increase the likelihood of success. In order to increase the effectiveness of immunization, we decided to use plasmid vectors that will provide:

- a) effective generation of antigen derived peptide in the transfected cell cytoplasm for MHC class I presentation
- b) presence of antigenic determinants that would more-likely break existing tolerance to autologous antigens targeted for cancer immunotherapy.

For this reason we compared vectors that encode for:

a) proteins that do not undergo posttranslational translocation to the endoplasmic reticulum but are degraded in the proteasome<sup>9,28</sup> – i.e. tVacs

b) xenogeneic rather than autologous (syngeneic) proteins.<sup>21,22,24</sup>

Similarly to the *in vitro* experiments, performed by us,<sup>9</sup> *in vivo* administration of tVacs or sVacs stimulates cytotoxic T cell responses (figure 3). The stronger cytotoxicity seen after immunization with hPSMA<sub>t</sub> may reflect stimulation of T cell clones that have high affinity receptors for the antigen. Alternatively, the stronger spleen cell cytotoxicity to the YM2 cells may be due to priming to sub-dominant epitopes which, unlike results *in vitro*, is maintained after boosting.<sup>9</sup> In such a case the stronger cytotoxicity may originate in the greater number of T cell clones recognizing the target cells. Some of this may have receptors with low affinity for the antigen and greater numbers of responders (respectively responder to target ratio) may be required for induction of cell death in target cells. Whether this is the case, however, is impossible to conclude from the current animal model. Additional experimentation, for example with a transgenic animal model (HLA-A\*0201/H-2 Kb mice) to evaluate responses to individual PSMA-derived epitopes, is required before any hypothesis can be put forward.

An important issue for effective vaccines is the development of potent adjuvants that can facilitate induction or augmentation of immunity. Despite the presence of non-methylated CpG motifs in plasmid DNA vaccines,<sup>29-35</sup> naked DNA immunization of rats obviously requires an additional adjuvant, and GM-CSF could act as such (figure 2; table 2). Granulocyte-macrophage colony-stimulating factor (GM-CSF) enhances immune responses by inducing the proliferation, maturation, and migration of dendritic cells.<sup>36</sup> Animals, immunized with a GM-CSF-plasmid DNA cocktail, are fully protected against transplantable tumors. This result is in agreement with earlier observations both in animals and in humans.<sup>4,37-39</sup> The expansion and differentiation of both T and B lymphocytes is affected<sup>36</sup> since the adjuvant effect of GM-CSF influences both cytotoxic T cell generation (figure 2) and antibody development (figure 5).

It has been shown that immunization with xenogeneic rather than autologous proteins is more likely to break existing tolerance<sup>21</sup> and we, as have others,<sup>24</sup> speculated that immunizations with plasmid DNA that encodes xenogeneic proteins will have the same advantage. This clearly is the case as far as priming is concerned (figure 4). Addition of GM-CSF to the immunization cocktail, however, leads to easier break of immune tolerance to PSMA since good protection against tumor development is observed even in those animals that were immunized with rPSMA<sub>t</sub> (table 2), a construct encoding the autologous protein.

Rats, primed with a construct encoding a human, but boosted with a construct encoding a rat protein, demonstrate the best protection against a transplantable tumor (figure 5). This may be due to the fact that priming with the construct encoding the xenogeneic protein successfully breaks the tolerant state. Boosting with an autologous protein or construct that encodes it, however, is necessary for fine tuning of the immune response.

Finally, release of the expressed product at the immunization site after gene-based vaccination seems to be necessary for development of antibody (figure 5). Priming and boosting with tVacs, whose product is not released extracellularly, but degraded in the proteasome, leads to no antibody production no matter whether an adjuvant is administered with the immunization cocktail. Others have reported similar findings following immunization with cytosolically retained or secreted viral proteins.<sup>40-47</sup> This is an important observation because, in cases where a tumor marker serves as a target antigen, development of high titer antibodies in patients' sera may interfere with marker testing and the use of the marker in disease monitoring. This is especially true when targeting secreted proteins such as PSA or PAP, where development of antibodies is of no therapeutic value. In those cases using tVacs instead of sVacs is clearly advantageous.

Production of antibodies after immunization for immunotherapy of cancer, however, can be beneficial in those instances where the target antigen is a membrane protein, such as PSMA.<sup>48-51</sup> Recently, Haas et al<sup>52</sup> showed, that immunization with a plasmid vector encoding a PSMA residue that is expressed on the membrane following expression, also leads to development anti-PSMA antibodies that bind to the native antigen. One would expect that the type of the immune response would impact the outcome of the treatment with Th1-type, cytotoxic antibodies that bind complement having the advantage of causing direct damage to the tumor cells. Were such the case, priming with tVacs and boosting with the native protein seem to represent the best immunization strategy (figure 6).

In conclusion we have shown that in vivo immunization with both tVacs and sVacs leads to generation of cytotoxic spleen cells. This shows that both tVacs and sVacs are immunogenic in rodents, and spleen cells from animals immunized with tVacs demonstrate stronger cytotoxicity. Co-administration of GM-CSF during immunization enhances the immune response and immune animals are protected against development of tumor when tumor cells that express the target antigen are inoculated. Priming with a construct encoding a xenogeneic protein and boosting with a construct encoding a syngeneic (autologous) protein confer the best protection. Immunization with gene-based vaccines whose products are retained in the cytoplasm and proteasomally degraded (tVacs) does not lead to antibody formation to the target antigen. Priming with tVacs and boosting with the native protein antigen results in production of cytotoxic antibodies.

#### References:

1. Epstein JE, Charoenvit Y, Kester KE, et al. Safety, tolerability, and antibody responses in humans after sequential immunization with a PfCSP DNA vaccine followed by

- the recombinant protein vaccine RTS,S/AS02A. *Vaccine*. 2004;22(13-14):1592-1603.
2. Boyd D, Hung CF, Wu TC. DNA vaccines for cancer. *IDrugs*. 2003;6(12):1155-1164.
  3. Srinivasan R, Wolchok JD. Tumor antigens for cancer immunotherapy: therapeutic potential of xenogeneic DNA vaccines. *J Transl Med*. 2004;2(1):12.
  4. Mincheff M, Tchakarov S, Zoubak S, et al. Naked DNA and adenoviral immunizations for immunotherapy of prostate cancer: a phase I/II clinical trial. *Eur Urol*. 2000;38(2):208-217.
  5. Gulley J, Chen AP, Dahut W, et al. Phase I study of a vaccine using recombinant vaccinia virus expressing PSA (rV-PSA) in patients with metastatic androgen-independent prostate cancer. *Prostate*. 2002;53(2):109-117.
  6. Pavlenko M, Roos AK, Lundqvist A, et al. A phase I trial of DNA vaccination with a plasmid expressing prostate-specific antigen in patients with hormone-refractory prostate cancer. *Br J Cancer*. 2004;91(4):688-694.
  7. Pardoll D. T cells and tumours. *Nature*. 2001;411(6841):1010-1012.
  8. Rodriguez A, Regnault A, Kleijmeer M, et al. Selective transport of internalized antigens to the cytosol for MHC class I presentation in dendritic cells. *Nat Cell Biol*. 1999;1(6):362-368.
  9. Mincheff M, Zoubak S, Altankova I, et al. Human dendritic cells genetically engineered to express cytosolically retained fragment of prostate-specific membrane antigen prime cytotoxic T-cell responses to multiple epitopes. *Cancer Gene Ther*. 2003;10(12):907-917.
  10. Silver DA, Pellicer I, Fair WR, et al. Prostate-specific membrane antigen expression in normal and malignant human tissues. *Clin Cancer Res*. 1997;3(1):81-85.
  11. Shneider BL, Thevananther S, Moyer MS, et al. Cloning and characterization of a novel peptidase from rat and human ileum. *J Biol Chem*. 1997;272(49):31006-31015.
  12. Halsted CH, Ling E, Luthi-Carter R, et al. Folylpoly-gamma -glutamate carboxypeptidase from pig jejunum. Molecular characterization and relation to glutamate carboxypeptidase II. *J Biol Chem*. 2000;275(39):30746.
  13. Chang SS, Bander NH, Heston WD. Monoclonal antibodies: will they become an integral part of the evaluation and treatment of prostate cancer--focus on prostate-specific membrane antigen? *Curr Opin Urol*. 1999;9(5):391-395.
  14. Chang SS, Gaudin PB, Reuter VE, et al. Prostate-Specific Membrane Antigen: Much More Than a Prostate Cancer Marker. *Mol Urol*. 1999;3(3):313-320.
  15. Beckett ML, Cazares LH, Vlahou A, et al. Prostate-specific membrane antigen levels in sera from healthy men and patients with benign prostate hyperplasia or prostate cancer. *Clin Cancer Res*. 1999;5(12):4034-4040.
  16. Murphy GP, Kenny GM, Ragde H, et al. Measurement of serum prostate-specific membrane antigen, a new prognostic marker for prostate cancer. *Urology*. 1998;51(5A Suppl):89-97.

17. Uria JA, Velasco G, Santamaria I, et al. Prostate-specific membrane antigen in breast carcinoma. *Lancet*. 1997;349(9065):1601.
18. Israeli RS, Powell CT, Corr JG, et al. Expression of the prostate-specific membrane antigen. *Cancer Res*. 1994;54(7):1807-1811.
19. Pinto JT, Suffoletto BP, Berzin TM, et al. Prostate-specific membrane antigen: a novel folate hydrolase in human prostatic carcinoma cells. *Clin Cancer Res*. 1996;2(9):1445-1451.
20. Mincheff M, Zoubak S. Gene-based vaccines for immunotherapy of prostate cancer - lessons from the past. *Gene Therapy and Molecular Biology*. 2004;8:465-474.
21. Fong L, Ruegg CL, Brockstedt D, et al. Induction of tissue-specific autoimmune prostatitis with prostatic acid phosphatase immunization: implications for immunotherapy of prostate cancer. *J Immunol*. 1997;159(7):3113-3117.
22. Fong L, Brockstedt D, Benike C, et al. Dendritic cell-based xenoantigen vaccination for prostate cancer immunotherapy. *J Immunol*. 2001;167(12):7150-7156.
23. Wolchok JD, Gregor PD, Nordquist LT, et al. DNA vaccines: an active immunization strategy for prostate cancer. *Semin Oncol*. 2003;30(5):659-666.
24. Gregor PD, Wolchok JD, Ferrone CR, et al. CTLA-4 blockade in combination with xenogeneic DNA vaccines enhances T-cell responses, tumor immunity and autoimmunity to self antigens in animal and cellular model systems. *Vaccine*. 2004;22(13-14):1700-1708.
25. Barinka C, Rinnova M, Sacha P, et al. Substrate specificity, inhibition and enzymological analysis of recombinant human glutamate carboxypeptidase II. *J Neurochem*. 2002;80(3):477-487.
26. Matzinger P. The JAM test. A simple assay for DNA fragmentation and cell death. *J Immunol Methods*. 1991;145(1-2):185-192.
27. Taglietti M. Vaccine adjuvancy: a new potential area of development for GM-CSF. *Adv Exp Med Biol*. 1995;378:565-569.
28. Mincheff M, Zoubak S, Altankova I, et al. Depletion of CD25+ cells from human T-cell enriched fraction eliminates immunodominance during priming with dendritic cells genetically modified to express a secreted protein. *Cancer Gene Ther*. 2005;12(2):185-197.
29. Klinman DM, Yi AK, Beaucage SL, et al. CpG motifs present in bacteria DNA rapidly induce lymphocytes to secrete interleukin 6, interleukin 12, and interferon gamma. *Proc Natl Acad Sci U S A*. 1996;93(7):2879-2883.
30. Klinman DM, Yamshchikov G, Ishigatsubo Y. Contribution of CpG motifs to the immunogenicity of DNA vaccines. *J Immunol*. 1997;158(8):3635-3639.
31. Klinman DM, Barnhart KM, Conover J. CpG motifs as immune adjuvants. *Vaccine*. 1999;17(1):19-25.
32. Krieg AM, Love-Homan L, Yi AK, et al. CpG DNA induces sustained IL-12 expression in vivo and resistance to *Listeria monocytogenes* challenge. *J Immunol*. 1998;161(5):2428-2434.

33. Krieg AM, Yi AK, Schorr J, et al. The role of CpG dinucleotides in DNA vaccines. *Trends Microbiol.* 1998;6(1):23-27.
34. Krieg AM. Mechanisms and applications of immune stimulatory CpG oligodeoxynucleotides. *Biochim Biophys Acta.* 1999;1489(1):107-116.
35. Krieg AM, Yi AK, Hartmann G. Mechanisms and therapeutic applications of immune stimulatory cpG DNA. *Pharmacol Ther.* 1999;84(2):113-120.
36. Perales MA, Fantuzzi G, Goldberg SM, et al. GM-CSF DNA induces specific patterns of cytokines and chemokines in the skin: implications for DNA vaccines. *Cytokines Cell Mol Ther.* 2002;7(3):125-133.
37. Tsuchiya Y, Igarashi M, Suzuki R, et al. Production of colony-stimulating factor by tumor cells and the factor-mediated induction of suppressor cells. *J Immunol.* 1988;141(2):699-708.
38. Pasquini S, Xiang Z, Wang Y, et al. Cytokines and costimulatory molecules as genetic adjuvants. *Immunol Cell Biol.* 1997;75(4):397-401.
39. Simmons SJ, Tjoa BA, Rogers M, et al. GM-CSF as a systemic adjuvant in a phase II prostate cancer vaccine trial. *Prostate.* 1999;39(4):291-297.
40. Drew DR, Lightowlers M, Strugnell RA. Humoral immune responses to DNA vaccines expressing secreted, membrane bound and non-secreted forms of the *Tania ovis* 45W antigen. *Vaccine.* 2000;18(23):2522-2532.
41. Haddad D, Liljeqvist S, Stahl S, et al. Comparative study of DNA-based immunization vectors: effect of secretion signals on the antibody responses in mice. *FEMS Immunol Med Microbiol.* 1997;18(3):193-202.
42. Ma X, Fornis X, Gutierrez R, et al. DNA-based vaccination against hepatitis C virus (HCV): effect of expressing different forms of HCV E2 protein and use of CpG-optimized vectors in mice. *Vaccine.* 2002;20(27-28):3263-3271.
43. Strasser JE, Arnold RL, Pachuk C, et al. Herpes simplex virus DNA vaccine efficacy: effect of glycoprotein D plasmid constructs. *J Infect Dis.* 2000;182(5):1304-1310.
44. Wang L, Menting JG, Black CG, et al. Differences in epitope recognition, isotype and titer of antisera to *Plasmodium falciparum* merozoite surface protein 4 raised by different modes of DNA or protein immunization. *Vaccine.* 2000;19(7-8):816-824.
45. Haddad D, Liljeqvist S, Stahl S, et al. Characterization of antibody responses to a *Plasmodium falciparum* blood-stage antigen induced by a DNA prime/protein boost immunization protocol. *Scand J Immunol.* 1999;49(5):506-514.
46. Haddad D, Liljeqvist S, Stahl S, et al. Differential induction of immunoglobulin G subclasses by immunization with DNA vectors containing or lacking a signal sequence. *Immunol Lett.* 1998;61(2-3):201-204.
47. Higgins TJ, Herold KM, Arnold RL, et al. Plasmid DNA-expressed secreted and nonsecreted forms of herpes simplex virus glycoprotein D2 induce different types of immune responses. *J Infect Dis.* 2000;182(5):1311-1320.



48. McDevitt MR, Barendswaard E, Ma D, et al. An alpha-particle emitting antibody ([<sup>213</sup>Bi]J591) for radioimmunotherapy of prostate cancer. *Cancer Res.* 2000;60(21):6095-6100.
49. Nanus DM, Milowsky MI, Kostakoglu L, et al. Clinical use of monoclonal antibody HuJ591 therapy: targeting prostate specific membrane antigen. *J Urol.* 2003;170(6 Pt 2):S84-88; discussion S88-89.
50. Bander NH, Nanus DM, Milowsky MI, et al. Targeted systemic therapy of prostate cancer with a monoclonal antibody to prostate-specific membrane antigen. *Semin Oncol.* 2003;30(5):667-676.
51. Huang X, Bennett M, Thorpe PE. Anti-tumor effects and lack of side effects in mice of an immunotoxin directed against human and mouse prostate-specific membrane antigen. *Prostate.* 2004;61(1):1-11.
52. Kuratsukuri K, Wang CY, Sone T, et al. Induction of antibodies against prostate-specific membrane antigen (PSMA) by vaccination with a PSMA DNA vector. *Eur Urol.* 2002;42(1):67-73.

**Table 1.** Gene-based vaccination induced tumor protection

<i>Injection with:</i>	<i>Number of animals with tumor (Total number of animals per group)</i>	<i>Mean size of tumor</i>
A. 3 x hPSMA <sub>t</sub>	2(10)	0.5 +/- 0.1 cm <sup>3</sup>
B. 3 x rPSMA <sub>t</sub>	5(10)	1.2 +/- 0.23 cm <sup>3</sup>
C. 3 x hPSMA <sub>s</sub>	7(10)	1.6 +/- 0.48 cm <sup>3</sup>
D. 3 x Empty vector	8(10)	>2.3 +/- 0.3 cm <sup>3</sup>
E. 3 x Saline	9(10)	>2.44 +/- 0.48 cm <sup>3</sup>

Differences in the number of tumor developing rats. Difference between groups A and B is not statistically significant ( $\chi^2 = 1.978$ ;  $p < 0.2$ ). Differences between groups A and E is statistically significant ( $\chi^2 = 5.05$ ;  $p < 0.025$ ), Differences between groups D and E (combined) and A or B are statistically significant ( $\chi^2 = 3.8$ ;  $p < 0.01$  for A and  $\chi^2 = 4.17$ ;  $p < 0.05$  for B).

**Table 2.** Addition of GM-CSF to the immunization cocktail enhances gene-based vaccination induced tumor protection

<i>Injection with:</i>	<i>Number of animals with tumor (Total number of animals per group)</i>	<i>Mean size of tumor</i>
F. 3 x hPSMA <sub>t</sub> + rGM-CSF	0(10)	-
G. 3 x rPSMA <sub>t</sub> + rGM-CSF	0(10)	-
H. 3 x hPSMA <sub>s</sub> + rGM-CSF	2(10)	1.1 +/- 0.1 cm <sup>3</sup>
I. 3 x Empty vector + rGM-CSF	9(10)	>1.89 +/- 0.34 cm <sup>3</sup>

Differences in the number of tumor developing rats. Difference between A and F is not statistically significant ( $\chi^2 = 2.22$ ;  $p < 0.2$ ). Difference between B and G is statistically significant ( $\chi^2 = 6.66$ ;  $p < 0.01$ ). Difference between C and H is statistically significant ( $\chi^2 = 5.05$ ;  $p < 0.2$ ).

## FIGURES AND LEGENDS TO FIGURES




Figure 1. Detection of PSMA in AT3-B1 cells transfected with hPSMAs

+/-L – presence or absence of lactacystin

1 and 2 – AT3B-1 cell clone YM1, stably transfected with hPSMAs

3 and 4 – AT3B-1 cell clone YM2, stably transfected with hPSMAs

5 – supernatant from YM2

6 – supernatant from YM1

7 – recombinant GCPII




Fig. 2. Cytotoxicity of rat spleen cell towards the YM2 clone. Copenhagen rats were immunized three times at 10 day-intervals, euthanized 14 days later and their spleen cells tested for cytotoxicity against the YM2 clone. A – rats were injected with either rPSMA (circles), empty vector (squares) or saline (diamonds). B – rats were injected with rPSMA with (filled in circles) or without rat recombinant GM-CSF. C – rats were injected with hPSMA with (filled in triangles) or without rat recombinant GM-CSF. D – rats were primed and boosted with cocktails containing rat GM-CSF and either hPSMA (triangles) or rPSMA (circles), or primed with hPSMA and boosted twice with rPSMA (squares).




Figure 3. Cytotoxicity of spleen cells from Copenhagen rats that were immunized against the YM2 clone. Rats were immunized three times at 10 day-intervals with a cocktail that contained rat GM-CSF and either hPSMA (diamonds) or hPSMA (squares).




Figure 4. Tumor development in rats following different prime-boosting regimens. All rats were immunized with a cocktail containing 100  $\mu\text{g}$  plasmid DNA and  $9\mu\text{g}/\text{m}^2$  body surface area rat GM-CSF. Circles: Rats were primed with hPSMA and boosted with rPSMA; triangles – primed with hPSMA and boosted with rPSMA; diamonds – primed and boosted with rPSMA; squares – primed and boosted with empty vector.




Figure 5. Immunization with the native protein leads to development of antibodies against its native conformation (data from ELISA, 1: 30 serum dilution) **A** – priming with hPSMA<sub>t</sub>, boosting with rPSMA<sub>t</sub> (with GM-CSF); **B** – 3 immunizations with hPSMA<sub>s</sub> (no GM-CSF); **C** – 3 immunizations with hPSMA<sub>s</sub> (with GM-CSF); **D** – 3 immunizations with GCPII (with GM-CSF); **E** – prime with GCPII, boost with hPSMA<sub>t</sub> and rPSMA<sub>t</sub> (with GM-CSF); **F** – prime with GCPII, boost with rPSMA<sub>t</sub> (with GM-CSF); **G** – prime with hPSMA<sub>t</sub>, boost with GCPII (with GM-CSF); **H** – prime with rPSMA<sub>t</sub>, boost with GCPII (with GM-CSF).




Figure 6. Immunization with tVacs primes for development of cytotoxic antibodies but boosting with the native protein is necessary for development of antibodies (data from ELISA, 1:30 dilution). Rats were primed and boosted twice with either GCPII or hPSMA<sub>s</sub>, or primed with GCPII and boosted with hPSMA<sub>t</sub> or rPSMA<sub>t</sub>; or primed with hPSMA<sub>t</sub> and boosted with GCPII.