

Award Number: W81XWH-04-1-0887

TITLE: Pim-1: A Molecular Target to Modulate Cellular Resistance to Therapy in Prostate Cancer

PRINCIPAL INVESTIGATOR: Michael Lilly, M.D.

CONTRACTING ORGANIZATION: Loma Linda University
Loma Linda, CA 92354

REPORT DATE: October 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE 01-10-2006			2. REPORT TYPE Annual		3. DATES COVERED 1 Oct 2005 – 30 Sep 2006	
4. TITLE AND SUBTITLE Pim-1: A Molecular Target to Modulate Cellular Resistance to Therapy in Prostate Cancer					5a. CONTRACT NUMBER	
					5b. GRANT NUMBER W81XWH-04-1-0887	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Michael Lilly, M.D. Email: mlilly@uci.edu					5d. PROJECT NUMBER	
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Loma Linda University Loma Linda, CA 92354					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited						
13. SUPPLEMENTARY NOTES Original contains colored plates: ALL DTIC reproductions will be in black and white.						
14. ABSTRACT The contract supports studies to define the role of the PIM1 kinase in acquired resistance to chemotherapy by prostate cancer cells. Data to date for specific aim #1 define a signaling pathway induced by docetaxel, involving sequential steps of JAK1/2 activation, STAT3 phosphorylation, expression of PIM1, and activation of NFkB signaling. Blockade of this pathway by expression of dominant negative PIM1 proteins blocks drug-induced upregulation of NFkB activity, and sensitizes cells to docetaxel. Other studies (specific aim #2) focus on identifying a mechanism through which PIM1 activates NFkB. We have unambiguously identified S937 as the major PIM1 phosphorylation site on the NFkB1/p105 precursor protein, through use of LCM/MS/MS analysis. Other kinases that can phosphorylate this site include AKT and PKA. Additional data (specific aim #3) have been published to describe a small molecule inhibitor of PIM1. This molecule can sensitize prostate cancer cells to the cytotoxic effects of docetaxel in an additive or synergistic manner.						
15. SUBJECT TERMS PIM1 kinase chemotherapy resistance prostate cancer						
16. SECURITY CLASSIFICATION OF:				17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U	19b. TELEPHONE NUMBER (include area code)			
				UU	29	

Table of Contents

Cover.....	1
SF 298.....	2
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusions.....	7
References.....	8
Appendices.....	9

INTRODUCTION

Studies under this funded activity are focused on characterizing the role of the PIM1 gene in acquired resistance to chemotherapy drugs, by prostate cancer cells. The proposal included three specific aims: 1) to define a novel signal transduction pathway activated by docetaxel, 2) to characterize the mechanism through which PIM1 activates and regulates NFkB signaling, and 3) to explore genetic and pharmacologic means of inhibiting PIM1 activity or expression to enhance the sensitivity of prostate cancer cells to docetaxel and other chemotherapy drugs. Substantial progress has been made in each of these areas during the 01 year of support.

BODY

We will outline our progress through reference to the specific aims described above. The first




Fig. 1. Time course of PIM1, pSTAT3 expression


specific aim was to outline a signal transduction pathway activated by docetaxel and involving upregulation of PIM1 expression. This pathway has been substantially defined. Using RWPE2 prostate cells, we noted that docetaxel treatment rapidly leads to an increase in expression of the PIM1 serine/threonine kinase. Expression becomes apparent at 3hrs after drug addition, peaks at 9-12hrs, and returns to baseline by 24hrs (Fig. 1). This increase in expression is accompanied by an increase in *pim-1* mRNA, as shown by real time-PCR analysis (Fig. 2). Thus the effects of docetaxel are primarily transcriptional or post-transcriptional.

We next wanted to define mechanisms through which *pim-1* could be transcriptionally upregulated. Transcription of *pim-1* is known to be activated by STAT transcription factors and by NFkB transcription factors. We examined


the time course of STAT3 activation after docetaxel treatment (Fig. 1), and noted that it paralleled the course of *pim-1* expression. We therefore suspected that docetaxel increased *pim-1* expression in a STAT3-dependent manner. This was directly demonstrated by use of decoy oligonucleotides (Fig. 3). Double-stranded DNA oligonucleotides matching a known STAT3 binding site blocked the drug-induced upregulation of *pim-1* expression, while a decoy based on a mutated (non-binding) STAT3 site did not. These data therefore establish a linear relationship among the following events: docetaxel treatment → STAT3 activation → *pim-1* expression.

We hypothesized that NFkB transcriptional activation would be a downstream event in this signal transduction pathway, because many chemotherapy drugs and other stressors are known to activate NFkB. We engineered RWPE2 cells to constitutively express a NFkB-dependent promoter/luciferase plasmid, and


Fig. 2. Expression of *pim-1* mRNA after docetaxel treatment




found that docetaxel treatment increased NFκB transcriptional activity. We then transiently infected these cells with a *pim-1*-encoding retrovirus. *Pim-1* expression also consistently increased NFκB transcriptional activity (Fig. 4). To determine if the drug-induced increase in NFκB activity occurred in a *pim-1*-dependent manner, we then infected the reporter cell line with a retrovirus encoding a dominant-negative form of *pim-1*, pimNT81. The



dominant negative *pim-1* cDNA completely blocked the drug-induced upregulation of NFκB activity, demonstrating that *pim-1* expression is a necessary upstream step in the drug-induced activation of NFκB (Fig. 5). In aggregate these studies establish a signal transduction pathway triggered by docetaxel treatment of RWPE2 prostate cancer cells.




dominant negative *pim-1* cDNA completely blocked the drug-induced upregulation of NFκB activity, demonstrating that *pim-1* expression is a necessary upstream step in the drug-induced activation of NFκB (Fig. 5). In aggregate these studies establish a signal transduction pathway triggered by docetaxel treatment of RWPE2 prostate cancer cells.




To determine if this pathway modified drug toxicity, we examined the effects of enforced expression of wild-type or NT81 *pim-1* cDNAs of docetaxel cell kill (Fig. 6). Docetaxel produced dose-dependent cell kill in RWPE1, 2 cells. Enforced expression of wild-type *pim-1*

cDNA markedly reduced cell death. In contrast, expression of the dominant negative NT81 cDNA enhanced cell death after docetaxel treatment. These data demonstrate that *pim-1* expression can modulate drug-induced cell death, and demonstrate that the survival pathway described above is a legitimate target for pharmacologic intervention. These data will be presented at the 2006 AACR meeting in poster form (1).




The goal of specific aim #2 was to define pathways through which the PIM1 kinase could activate NFκB transcriptional activity. We had hypothesized that PIM1 would phosphorylate the NFκB1/p105 precursor protein on serine-937, leading to proteolytic cleavage of the protein with release of active p50 protein as well as other sequestered NFκB components and the TPL2 kinase. To demonstrate the site of phosphorylation we used mass spectroscopy of tyrosin-digested fragments of p105 that had been phosphorylated *in vitro*. We had previously


demonstrated that PIM1-dependent phosphorylation happens exclusively on serine. Fragments were separated by LC/MS/MS analysis and mass/charge ratios were determined. The predicted peptide fragment that would result from phosphorylation at serine-937 was recovered, with a mass of 1016 (Fig. 7). Since there are several potential phosphorylation sites within this peptide, we proceeded to sequence the peptide with mass spectroscopy. Only the fragment corresponding to phosphoserine-937 was not recovered. These data unambiguously demonstrate that the major phosphorylation site of PIM1 on p105 is serine-937.



To further characterize phosphorylation on this site we prepared an antibody specific for phosphoNFκB1/p105(S937). This antibody reacts quite specifically with its antigen in whole cell lysates and immunoprecipitates. Fig. 8 demonstrates that recombinant GST-PIM1 protein phosphorylates NFκB1/p105 on serine 937 *in vitro*. Furthermore wild-type PIM1 phosphorylates p105 *in vivo*, while a dominant-negative PIM1 (NT81) does not.



The third specific aim proposed to use small molecule inhibitors of the PIM1 kinase as molecular probes to determine their effect on docetaxel sensitivity. A report describing one such molecule, the flavonol quercetagenin, has been accepted for publication and will appear in



the January, 2007 issue of *Molecular Cancer Therapeutics* (see appendix A and Fig. 9). We have demonstrated that quercetagenin in a moderately potent ($\text{IC}_{50} = 340\text{nM}$, specific, and cell-permeable inhibitor of PIM1 activity in prostate cancer cells. Key data include the demonstration that quercetagenin in competitive with ATP. A crystal structure of PIM1 in complex with quercetagenin, or with three other flavonoids, has been determined. We have also shown that quercetagenin is able to inhibit the activity of the PIM1 kinase in

prostate cancer cells at an IC_{50} of about $5.5\mu\text{M}$. Interestingly the activity of the AKT kinase is not inhibited at all under these conditions (Fig. 9).

We have recently obtained, and begun characterizing, novel small molecule inhibitors of PIM1 from Excelix Corporation. These molecules show additive, or at some concentrations synergistic, cell growth inhibition in combination with docetaxel. These studies confirm the central hypothesis of this overall project, that PIM1 kinase acts to inhibit cell death caused by the cytotoxic drug docetaxel, and that blocking the activity of PIM can potentiate cell kill and overcome cytotoxic drug resistance.

KEY RESEARCH ACCOMPLISHMENTS

- Definition of a novel survival pathway activated by docetaxel treatment, and involving sequential activation or expression of JAK2, STAT3, PIM1, and NF κ B components.
- Identification of serine-937 as the major phosphorylation site for PIM1 on the p105/NF κ B1 precursor protein
- Identification of quercetagenin as a moderately potent and specific, cell-permeable PIM1 kinase inhibitor
- Demonstration that XL-1075 and XL-1154 can show additive or synergistic cell kill in prostate cancer cells treated with docetaxel
- Abstract presented at the annual AACR meeting, Washington DC, April, 2006
- Manuscript accepted for publication in *Molecular Cancer Therapeutics*

REPORTABLE OUTCOMES

None in 02 year

CONCLUSIONS

Our data demonstrate that PIM1 is a critical component of a survival/stress pathway activated by docetaxel treatment of prostate cancer cells. This pathway leads to activation of NF κ B-dependent transcription, possibly by phosphorylation of p105/NF κ B1 by PIM1 at serine-937.

Targeting PIM1 kinase activity with quercetagenin, or other PIM1 kinase inhibitors, leads to additive or synergistic cell kill following docetaxel treatment.

REFERENCES


1. Zemskova M, Sahakian E, Lilly M: The PIM1 kinase is a critical component of a survival pathway that protects prostate cancer cells from docetaxel-induced death (abstract #2777), presented at 97th Annual Meeting of AACR, Washington, DC, April 2006.
2. Holder S, Zemskova M, Bremer R, Neidigh JW, Lilly MB: Characterization of a potent and selective small-molecule inhibitor of the PIM1 kinase (in press *Molecular Cancer Therapeutics*, 2006).

APPENDIX

Research data are presented throughout the body of this report.
The appendix contains three items:

1. AACR abstract #2777, approved for presentation at the 97th Annual Meeting, April, 2006, entitled "The PIM1 kinase is a critical component of a survival pathway that protects prostate cancer cells from docetaxel-induced death" by M. Zemskova, E. Sahakian, M. Lilly.
2. Galley proofs of manuscript accepted for *Molecular Cancer Therapeutics*
2. Curriculum vitae for Michael Lilly, MD

is, The Online Abstract Submission System <http://www.abstractsonline.com/submit/SubmitPrinterFriendlyVersion.a...>

 American Association for Cancer Research | SAVING LIVES THROUGH RESEARCH

97th Annual Meeting 2006
ABSTRACT SUBMITTER

[Print this Page for Your Records](#) [Close Window](#)

Control/Tracking Number: 06-AB-2777-AACR
Activity: Abstract Submission
Current Date/Time: 11/11/2005 1:35:09 AM

The PIM1 kinase is a critical component of a survival pathway that protects prostate cancer cells from docetaxel-induced death.

Short Title:
PIM1 and docetaxel-induced death

Author Block: *Marina Zemskova, Eva Sahakian, Michael B. Lilly*. Loma Linda University, Loma Linda, CA

The PIM1 serine-threonine kinase is a true oncogene that mediates survival and proliferation signals in human neoplasms, including leukemias and prostate adenocarcinomas. Enforced expression of PIM1 has been shown to enhance resistance of cells to cytotoxic agents and ionizing radiation. To examine potential pathways through which PIM1 produces drug resistance, we examined RWPE1 and RWPE2 prostate epithelial cells treated with docetaxel, a taxane used for treatment of prostate cancer. Cells treated with docetaxel demonstrated an increase in PIM1 protein and pim-1 mRNA 3-12 hrs after drug exposure. We then sought to establish upstream and downstream effectors of PIM1 expression. Docetaxel also induced expression of phosphoSTAT3 with similar kinetics. Drug-induced upregulation of PIM1 expression was abolished when cells were transfected with STAT3 decoy oligonucleotides, demonstrating that STAT3 activation by docetaxel is required for drug-induced upregulation of PIM1. Docetaxel treatment, and infection with a PIM1 expressing retrovirus, both induced activation of NFkB transcriptional activity in RWPE2 stably transfected with an NFkB/luciferase reporter plasmid. However, when a dominant negative PIM1 protein (NT81) was introduced by retroviral transduction, drug induced activation of NFkB activity was abolished. Retroviral transduction of wild-type PIM1 or dominant-negative (NT81) PIM1 increased or decreased survival of RWPE1 and RWPE2 cells treated with docetaxel. These data establish a survival pathway (drug → STAT3 → PIM1 → NFkB) induced by docetaxel and capable of impairing drug cytotoxicity. Targeting the PIM1 kinase, along with STAT3 and NFkB, may be a viable approach to enhancing cell kill by cytotoxic drugs such as docetaxel.

Author Disclosure Block: M. Zemskova, None; E. Sahakian, None; M.B. Lilly, None.

Additional Disclosures (Complete):
The presenter of this abstract will discuss commercial products, devices, or technology in this presentation, as outlined below. : No - [Type "none" in the first box below]
1. Generic Name: : none
I anticipate discussing an OFF-LABEL use of a commercial product/device in this educational activity. : No - [Type "none" in the box below]
If you selected "Yes" above, you must indicate the product/device and describe its intended use in the box below. If you selected "No" above, you must type "none" in the box below: : none

Investigational Use (Not approved by the FDA for any purpose):
I anticipate discussing an INVESTIGATIONAL use of a commercial product/device in this educational activity. : No - [Type "none" in the box below]
If you selected "Yes" above, you must indicate the product/device and describe its intended use in the box below. If you selected "No" above, you must type "none" in the box below: : none
I agree with the declaration statement above. : True
Name: : Marina Zemskova
Date [mm/dd/yyyy]: : 11/10/2005

Category and Subclass (Complete): CB02-07 Survival factors
Keywords/ Indexing (Complete): Kinases ; Resistance ; Docetaxel ; Prostate
Sponsor (Complete):
2006 Travel Awards (Complete):
Payment (Complete): Your credit card order has been processed on Friday 11 November 2005 at 1:34 AM.
Status: Complete

Characterization of a potent and selective small-molecule inhibitor of the PIM1 kinase

Sheldon Holder,^{1,2,3} Marina Zemskova,²
Chao Zhang,⁴ Maryam Tabrizizad,⁴
Ryan Bremer,⁴ Jonathan W. Neidigh,²
and Michael B. Lilly^{1,2,3}

¹Center for Molecular Biology and Gene Therapy, Departments of ²Biochemistry and Microbiology and ³Medicine, Loma Linda University School of Medicine, Loma Linda, California; and ⁴Novartis, Inc., Berkeley, California

Abstract

Q2 The *pim-1* kinase is a true oncogene that has been implicated in the development of leukemias, lymphomas, and prostate cancer, and is the target of drug development programs. We have used experimental approaches to identify a selective, cell-permeable, small-molecule inhibitor of the *pim-1* kinase to foster basic and translational studies of the enzyme. We used an ELISA-based kinase assay to screen a diversity library of potential kinase inhibitors. The flavonol quercetagetin (3,3',4',5,8,7-hydroxyflavone) was identified as a moderately potent, ATP-competitive inhibitor (IC₅₀, 0.34 μmol/L). Resolution of the crystal structure of PIM1 in complex with quercetagetin or two other flavonoids revealed a spectrum of binding poses and hydrogen-bonding patterns in spite of strong similarity of the ligands. Quercetagetin was a highly selective inhibitor of PIM1 compared with PIM2 and seven other serine-threonine kinases. Quercetagetin was able to inhibit PIM1 activity in intact RWPE2 prostate cancer cells in a dose-dependent manner (ED₅₀, 5.5 μmol/L). RWPE2 cells treated with quercetagetin showed pronounced growth inhibition at inhibitor concentrations that blocked *pim-1* kinase activity. Furthermore, the ability of quercetagetin to inhibit the growth of other prostate epithelial cell lines varied in proportion to their levels of PIM1 protein. Quercetagetin can function as a moderately potent and selective, cell-permeable inhibitor of the *pim-1* kinase, and may be useful for proof-of-concept studies to support the development of clinically useful PIM1 inhibitors. [Mol Cancer Ther 2007;6(1):1-10]

Received 7/1/06; revised 10/1/06; accepted 11/1/06.

Grant support: Department of Defense, Congressionally Directed Medical Research Program, Award W81XWH-04-1-0887.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Michael B. Lilly, Center for Molecular Biology and Gene Therapy, Loma Linda University School of Medicine, 11234 Anderson Street, Loma Linda, CA 92354. Phone: 909-659-8777; Fax: 909-659-0177. E-mail: mlilly@llu.edu

Copyright © 2007 American Association for Cancer Research.
doi:10.1158/1535-7183.MCT-06-0397

Introduction

The *pim* family of serine-threonine kinases is composed of three highly homologous genes, *pim-1*, *pim-2*, and *pim-3*. These enzymes are increasingly being recognized as important mediators of survival signals in cancers, stress responses, and neural development (1-6). In addition, these kinases are constitutively expressed in some tumors and function as true oncogenes. Thus, they are of significant interest as targets for therapeutic intervention.

Small-molecule inhibitors are important molecular probes for studying protein kinases. In addition, they may serve as prototype therapeutic agents for treating diseases resulting from unregulated kinase activity. Three prior reports have shown that known, miscellaneous kinase inhibitors can inhibit PIM1 function *in vitro*. Jacobs et al. (7) showed that several staurosporine and bisindolylmaleimide analogues, as well as the morpholino-substituted diazomycin LY294002, were able to inhibit PIM1 activity *in vitro*. Subsequently, Fabian et al. (8) presented an interaction map involving 113 kinases and 20 small-molecule kinase inhibitors now under clinical study. Only three inhibitors had detectable binding to (and presumably inhibitory activity against) PIM1—two staurosporine analogues and flavopiridol, a flavonoid undergoing evaluation as an inhibitor of cyclin-dependent kinases. A recent report (9) confirmed the activity of bisindolylmaleimide derivatives as well as some flavonoids *in vitro*. All of the identified inhibitors either lacked specificity for PIM1 or were only modestly active at low micromolar concentrations, or both. Furthermore, none of these reports showed that the test agents could selectively inhibit PIM1 activity in intact cells.

To further our basic and translational studies of the *pim* kinases, we have sought to identify small-molecule inhibitors of PIM1. We here report that the flavonol quercetagetin is a selective PIM1 inhibitor with nanomolar potency and can differentially inhibit the kinase in cell-based assays.

Materials and Methods

Cell Lines and Culture Methods

The prostate epithelial cell lines RWPE1, RWPE2, LNCaP, and PC3 were obtained from the American Type Culture Collection (Manassas, VA) and cultured in the recommended **Q3** medium. We produced additional pools of RWPE2 prostate cells that overexpressed *pim-1* through retroviral transduction. The coding region for the human *pim-1* gene was cloned into the pLNCX retroviral vector (Clontech). **Q4** Infectious viruses were produced in the GP-293 packaging cell line by cotransfection with retroviral backbone plasmids (pLNCX or pLNCX/*pim-1*) and with pVSV-G, a plasmid that expresses the envelope glycoprotein from

2 PIM1 Inhibitor

Q1

vesicular stomatitis virus. Forty-eight hours after transfection, the medium was collected and the virus particles were concentrated as described in the manufacturer's protocol (Clontech). RWPE2 cells were plated at 1×10^5 per 60-mm plate 16 to 18 h before infection. Cells were infected with 5×10^5 viral particles in the presence of 8 $\mu\text{g}/\text{ml}$ polybrene. After 6 h of incubation, the virus-containing medium was replaced with fresh medium, and on the next day G418 (400 $\mu\text{g}/\text{ml}$) was added to select infected cells. After 10 days of selection, stable cell pools were established and PIM1 expression was verified by immunoblotting.

For growth-inhibition experiments, cells were plated onto 24-well plates and fixed with formaldehyde at intervals. Cell number was quantified by crystal violet staining (10).

Recombinant p1m Kinases and Kinase Assays

We prepared recombinant PIM1 and PIM2 as glutathione S-transferase (GST) fusions in *Escherichia coli*, as described (11). For the inhibitor screening assays, a solid-phase kinase assay was developed based on our demonstration that PIM1 is a potent kinase for phosphorylating BAD on Ser¹¹² (11, 12). Ninety-six-well flat-bottomed plates (Corning) were coated overnight at 4°C with recombinant GST-BAD [1 $\mu\text{g}/\text{well}$ in HEPES buffer: 136 mmol/L NaCl, 2.6 mmol/L KCl, and 20 mmol/L HEPES (pH 7.5)]. The plates were then blocked for 1 h at room temperature with 10 mg/ml bovine serum albumin in HEPES buffer. The blocking solution was then removed and 5 μl of each inhibitor, dissolved in 50% DMSO, were added to each well. Then, 100 μl of kinase buffer [20 mmol/L MOPS (pH 7.0), 12.5 mmol/L MgCl₂, 1 mmol/L MnCl₂, 1 mmol/L EGTA, 150 mmol/L NaCl, 10 $\mu\text{mol}/\text{l}$ ATP, 1 mmol/L DTT, and 5 mmol/L β -glycerophosphate] containing 25 ng recombinant GST-PIM1 kinase were added to each well. The final concentration of each inhibitor was ~ 10 $\mu\text{mol}/\text{l}$. The plate was placed on a gel slab dryer prewarmed to 30°C, and the kinase reaction was allowed to proceed. The reaction was stopped after 60 min by removal of the reaction buffer, followed by the addition of 100 μl of HEPES buffer containing 20 mmol/L EDTA to each well. Phosphorylated GST-BAD was detected by an ELISA reaction, using as first antibody a monoclonal anti-phospho-BAD(S112) antibody (Cell Signaling), a secondary goat anti-mouse IgG-peroxidase conjugated antibody (Pierce), and Turbo-TMB peroxidase substrate (Pierce). The level of phosphorylated GST-BAD present was proportional to the absorbance at 450 nm.

For quantitative and kinetic studies of inhibitors against various BAD(S112) kinases, a solution phase assay was used. A biotinylated peptide based on the PIM1 phosphorylation site of human BAD was synthesized (GGAGA-VEIRSRHSSYPAGTE) and used as the assay substrate. Recombinant GST-PIM1 (25 ng/reaction) was preincubated with various concentrations of inhibitors in the previous kinase buffer (final volume 100 μl). The reaction proceeded by addition of substrate peptide, followed by incubation for 5 min in a 30°C water bath. The reaction was terminated by transferring the mixture to a streptavidin-coated 96-well plate (Pierce) containing 100 $\mu\text{l}/\text{well}$ of 40 mmol/L EDTA. The biotinylated peptide substrate was allowed to bind to

the plate at room temperature for 10 min. The level of phosphorylation was then determined by ELISA as described above. Curve fitting and enzyme analyses were done using GraphPad Prism version 4.00 for Windows (GraphPad Software, San Diego, CA). For the additional BAD(S112) kinases [PIM2, PDK2 (ribosomal S6 kinase 2), and PKA (cyclic AMP-dependent protein kinase)], reaction components were as described above. As with the PIM1 assays, an ATP concentration of 10 $\mu\text{mol}/\text{l}$ was used. Furthermore, with each kinase, linear reaction velocities for the duration of the reaction were confirmed (data not shown).

To further assess the specificity of quercetin as a PIM1 inhibitor, its activity against a panel of serine-threonine kinases was also studied through a commercial kinase inhibitor profiling service (KinaseProfiler; Upstate Biotechnology). All KinaseProfiler assays were conducted using 10 $\mu\text{mol}/\text{l}$ ATP concentrations.

Small-Molecule Library Screening

We obtained a library of 1,200 compounds that had structural affinity to known kinase inhibitors (TunTec, Inc.). The entire library was screened once with our solid-phase ELISA kinase assay, with each compound at ~ 10 $\mu\text{mol}/\text{l}$ concentration. Positive hits were rescreened at the same concentration. Compounds that had reproducible activity at 10 $\mu\text{mol}/\text{l}$ were then screened at a range of concentrations from 0.001 to 300 $\mu\text{mol}/\text{l}$. Additional flavonoids were purchased from Indofine Chemicals and were tested in a similar protocol.

Measurement of PIM1 Kinase Activity in Cells

RWPE2 cell pools, stably infected with empty retrovirus or p1m-1-encoding retrovirus, were seeded in six-well plates at 5×10^5 cells per well. After 18 h, the normal supplemented keratinocyte medium was removed and replaced with supplement-free keratinocyte medium. Cells were then incubated for an additional 20 h. Quercetin, or an equivalent volume of DMSO, was added to the cells 3 h before the end of the starvation period. At the conclusion of the starvation period, the cells were washed twice with PBS and subsequently lysed in a denaturing buffer with protease, phosphatase inhibitors. The lysates were normalized by total protein content (BCA protein assay, Pierce), then analyzed by immunoblotting with the following antibodies: monoclonal anti-PIM1 (Santa Cruz Biotechnology, Santa Cruz, CA); monoclonal anti- β -actin (Sigma); monoclonal anti-BAD (Transduction Laboratories); and monoclonal anti-phospho-BAD(S112), polyclonal anti-phospho-AKT(S473), and anti-AKT (all from Cell Signaling).

Cloning, Expression, Purification, and Crystallization of PIM1

The production, purification, and characterization of recombinant 6His-tagged PIM1 proteins for crystallography have been described previously (13). To obtain cocystals of complexes of the protein with ligands, the protein solution was initially mixed with the compound (dissolved in DMSO) at a final compound concentration of 1 mmol/L and then set up for crystallization. The protein was

Q8

Q9

Q10

Q11
Q12

crystallized by a sitting-drop, vapor-diffusion experiment in which equal volumes of protein (10–15 mg/ml, concentration) and reservoir solution [0.4–0.9 mol/L sodium acetate, 0.1 mol/L imidazole (pH 6.5)] were mixed and allowed to equilibrate against the reservoir at 4 °C. The crystals routinely grew to a size of 200 × 200 × 800 μm in ~2 to 3 days.

Structure Determination

X-ray diffraction data were collected at Advanced Light Source (Berkeley, CA). All data were processed and reduced with MOSFLM and scaled with SCALA, of the CCP4 suite of programs using the software ELVES. The space group of all crystals was determined to be $P6_3$, with the cell axes being approximately 99, 99, and 80, and one protein monomer being present in the asymmetrical unit. All structures were determined by molecular replacement using the apo PIM1 structure (1YVW; ref. 13) as a model, and refined by CNX and REFMACS. Crystallographic statistics are reported in Supplementary Table S1.⁵ The coordinates and structure factors for the structures have been deposited with the RCSB Protein Data Bank (accession codes).

Q13

Results

Screening of a Chemical Library with Structural Affinity to Known Kinase Inhibitors

As an initial approach to the identification of PIM1 inhibitors, we screened a library of small molecules whose structures were similar to those of known kinase inhibitors. Of the seven compounds that had a reproducible inhibitory activity at 10 μmol/L, six were flavonoids [quercetin, luteolin, kaempferol, 7-hydroxyflavone, (5)-5,7-dihydroxy-8-(3-methylbut-2-enyl)flavone, and (R)-5,7-dihydroxyflavone]. These compounds exhibited a range of inhibitory potencies (as IC_{50}) from 1.1 to 60 μmol/L. Thirty-seven other flavonoids failed to show detectable inhibitory activity at 10 μmol/L. These inactive compounds were characterized in most cases by bulky (charged or uncharged) groups at the 3, 3', 4', or 7 positions; lack of at least two hydrogen bond donors on the A or C rings; presence of glycoside linkages; or failure of all rings to adopt a planar conformation.

The most active compound in the chemical library was the flavonoid quercetin (IC_{50} , 1.1 μmol/L), a known inhibitor of kinases and many other enzymes (14–19). Furthermore, six of the seven compounds with a reproducible activity at 10 μmol/L were flavonoids. Hence, we screened additional flavonoids to identify molecules with inhibitory activity against the PIM1 kinase (Fig. 1). The most active molecule was the flavonoid quercetageitin (IC_{50} , 0.34 μmol/L). The four flavonoids with the highest inhibitory activity were characterized by the presence of five to six -OH groups distributed between the A and B rings. In comparison, the hydroxyl groups on the B ring seemed to be more critical

for the activity of the compounds than those on the A ring, as compounds with an unsubstituted B ring showed greatly reduced activity. Finally, a hydrophobic substituent at the 8 position was tolerated.

Quercetageitin Is a Selective, Potent Inhibitor of PIM1 *In Vitro*

To assess the selectivity of quercetageitin for PIM1, we determined its IC_{50} value toward the alternative BAD5(112) kinases RSK2, PKA, and PIM2 (Table 1). The IC_{50} of quercetageitin for PIM1 kinase was 0.34 μmol/L, whereas the corresponding values for the other kinases were 9- to 20-fold higher.

To further characterize the specificity of quercetageitin, its inhibitory activity was examined at 1 or 10 μmol/L against additional serine-threonine kinases (c-Jun-NH₂-kinase 1, PKA, Aurora-A, c-RAF, and PKC δ ; Fig. 2). At the lower concentration, the selectivity of quercetageitin was most apparent. In the presence of 1 μmol/L inhibitor, PIM1 activity was inhibited by 92%. In contrast, the activity of the other kinases was inhibited by only 0% to 41%. In aggregate, these studies established that quercetageitin was a severalfold more potent inhibitor for p11-1 kinase than for seven other serine-threonine kinases. In addition, quercetageitin was completely inactive against the c-Kit tyrosine kinase when tested at the 200 μmol/L concentration (data not shown).

Crystallographic Analysis of Quercetageitin in Complex with PIM1

Recently, several crystal structures of the p11-1 kinase have been solved and presented, including apo forms and the enzyme in complex with a variety of ligands (7, 9, 13, 20, 21). Because the PIM1 protein has several unique structural features around its ATP-binding pocket, including the lack of the canonical hydrogen bond donor from the hinge region typically used by kinases to bind ATP-like ligands, we determined the crystal structure for the kinase in complex with three flavonoid inhibitors: quercetageitin, myricetin, and 5,7,3',4',5'-pentahydroxyflavone (Fig. 3).

The three flavonoid inhibitors show two distinct binding poses, denoted here as orientations I and II, respectively. Quercetageitin, the compound with two hydroxyl groups on the B ring, adopts orientation I, whereas the compounds with a trisubstituted B ring (myricetin and 5,7,3',4',5'-pentahydroxyflavone) adopt orientation II.

The binding pose of quercetageitin in PIM1 (Fig. 3A) closely resembles that of quercetin in phosphatidylinositol 3-kinase γ (1E8W; ref. 22) and that of fisetin in CDK6 (1XC2; ref. 23), designated here as orientation I. As seen in the two earlier structures (Fig. 3D and E), the 3-OH of the quercetageitin (Fig. 3A) makes a canonical hydrogen bond with backbone carbonyl oxygen of the hinge residue Glu⁹¹. In addition, the B ring of quercetageitin binds deep inside the PIM1 ATP-binding pocket, with the 4-hydroxyl group hydrogen-bonded to the side chains of two highly conserved residues, Lys⁸⁷ and Glu⁸⁸. However, significant difference was also observed between the current structure and the two reported structures. In both 1E8W and 1XC2, the 4-keto group of the chromenone core of the compound

⁵Supplementary data for this article are available at Molecular Cancer Therapeutics Online (<http://mct.aacrjournals.org/>).

4 PIM1 inhibitor

formed a hydrogen bond with the same hinge amide nitrogen [Val⁶² in phosphatidylinositol 3-kinase γ (Fig. 3D) and Val¹⁰⁸ in CDK6 (Fig. 3E)]. However, there is no direct interaction between the 4-keto group of quercetagenin and the amide nitrogen of the corresponding residue Pro¹²⁰ in PIM1 because proline is incapable of acting as a hydrogen bond donor. Instead, the 4-keto group of quercetagenin makes close contact with the backbone C α of Arg¹²⁰ (3.4 Å). It is not clear whether this interaction makes a positive contribution to the binding of quercetagenin to PIM1.

The B ring of quercetagenin binds deep inside the PIM1 ATP-binding pocket. The 4'-hydroxyl group forms hydrogen bonds with both Lys⁶⁷ and Glu⁶⁶, two of the most conserved residues in kinases. As has been noted, satisfying the hydrogen bonding requirements at this region is one of the determining features of binding of compounds to PIM1 (13).


When compared with quercetagenin, the chromone core of myricetin (Fig. 3B) and 5,7,3',4',5'-pentahydroxyflavone (Fig. 3C) has flipped 180° in PIM1 such that the B ring is now oriented toward the entrance of the ATP pocket. A possible explanation for adopting this orientation is that the interior of the ATP pocket cannot accommodate the B ring with these hydroxyl substitutions. Although they bind in the same orientation, there are important differences

between the binding poses of the two compounds, which can be attributed to the presence or absence of the 3-hydroxyl group. The 3-hydroxyl group in myricetin still makes a hydrogen bond with the carbonyl oxygen of Glu⁶¹, despite the difference in binding orientation. Because of the adjacent 4-keto group, the 3-hydroxyl is likely to be most acidic of all the hydroxyl groups in the compound, and, as a result, it dictates the overall positioning of the compound. Another interaction that may contribute to the observed binding pose is a hydrogen bond between the 3'-hydroxyl group of myricetin and the carbonyl oxygen of Pro⁶³ (Fig. 3B). The importance of the 3-hydroxyl group is evident. The second compound, 5,7,3',4',5'-pentahydroxyflavone, lacking such a group, makes no direct interaction with the hinge region.

Quercetagenin Inhibits PIM1 Kinase Activity in Intact Cells

To determine if quercetagenin could act as a cell-permeable PIM1 inhibitor, we examined the activity of the flavonoid in RWPE2 prostate cancer cells. We studied the phosphorylation of endogenous BAD on Ser¹¹², under conditions of growth factor starvation, as an indicator of intracellular PIM1 activity (Fig. 4).

RWPE2 cells infected with a *neo*-expressing retrovirus showed little phospho-BAD(S112) when cultured overnight



Flavonoid	3	5	6	7	8	2'	3'	4'	5'	6'	C ₅₀ (nM)
quercetagenin	OH	OH	OH	OH	H	H	OH	OH	H	H	0.24
quercetin	OH	OH	H	OH	OH	H	OH	OH	H	H	0.43
5,7,3',4',5'-pentahydroxyflavone	H	OH	H	OH	H	H	OH	OH	OH	H	0.85
myricetin	OH	OH	H	OH	H	H	OH	OH	OH	H	0.28
flavetin	OH	H	H	OH	H	H	OH	OH	H	H	0.45
apigenin	H	OH	H	OH	H	H	H	OH	H	H	0.24
3,7,4'-trihydroxyflavone	OH	H	H	OH	H	H	H	OH	H	H	0.88
quercetin	OH	OH	H	OH	H	H	OH	OH	H	H	1.1
kaempferol	OH	OH	H	OH	H	H	H	OH	H	H	1.3
luteolin	H	OH	H	OH	H	H	OH	OH	H	H	1.6
morin	OH	OH	H	OH	H	OH	H	OH	H	H	2.7
7,3'-dihydroxyflavone	H	H	H	OH	H	H	OH	H	H	H	4.52
3,7'-dihydroxyflavone	OH	H	H	OH	H	H	H	H	H	H	4.6
7,3',4',5'-tetrahydroxyflavone	H	H	H	OH	H	H	OH	OH	OH	H	7.6
3,8,2',4'-tetrahydroxyflavone	OH	H	OH	H	H	OH	H	OH	H	H	8.5
(S)-6,7-dihydroxy-8-(3-methylbut-2-enyl)flavone											12.2
7-hydroxyflavone	H	H	H	OH	H	H	H	H	H	H	14
6,7-dihydroxyflavone	H	OH	H	OH	H	H	H	H	H	H	16
7,8,4'-trihydroxyflavone	H	H	H	OH	OH	H	H	OH	H	H	22
(R)-6,7-dihydroxyflavone											60.2
(S)-6,7-dihydroxyflavone											107

Figure 1. Identification of flavonoids with PIM1 inhibitory activity. Structures of all studied flavonoids with detectable PIM1 inhibitory activity, given with their C₅₀ values.

Table 1. Quercetagenin is a selective inhibitor of the PIM1 kinase over other BAD(S112) kinases

Kinase	IC ₅₀ (μmol/L)	Log IC ₅₀ (μmol/L)	SE of log IC ₅₀	R ²
PIM1	0.34	-0.46	0.12	0.98
PIM2	3.45	0.53	0.22	0.94
PKA	21.2	1.33	0.23	0.94
RSK2	2.62	0.45	0.09	0.99

NOTE: All data were derived from nonlinear regression analyses using a three-parameter logistic that assumes a Hill coefficient of -1.

in basal serum-free medium. However, cells with enforced expression of *pim-1* kinase had a 4-fold higher amount of phospho-BAD, reflecting the ability of the PIM1 protein to phosphorylate the endogenous BAD protein. When *pim-1*-expressing cells were treated with quercetagenin, phospho-BAD(S112) levels were markedly reduced in proportion to the concentration of the inhibitor. Half-maximal inhibition occurred at 5.5 μmol/L extracellular concentration. Quercetagenin did not inhibit the activity of the AKT kinase under these conditions, as indicated by persistent phosphorylation of AKT on Ser⁴⁷³. These data indicate that quercetagenin was able to selectively block the ability of PIM1 to phosphorylate BAD in intact cells.

Quercetagenin Treatment Reproduces a Known *pim-1* Knockdown Phenotype

If quercetagenin acts as a true PIM1 inhibitor, then it should reproduce a *pim-1*-dependent phenotype in the target cells. We have shown that PIM1 inhibition by genetic means (small interfering RNA) inhibits the proliferation of RWPE1 and RWPE2 cells (Supplementary Fig. S1).⁵ We therefore determined if quercetagenin could reproduce this phenotype. RWPE2 cells were treated with quercetagenin for up to 72 h (Fig. 5A). Marked dose-dependent growth inhibition was apparent by 24 h, leading to persistent growth arrest thereafter. Quercetagenin reproduced this *pim-1*-dependent phenotype at a drug concentration that inhibited the enzyme in cells (ED₅₀, 3.8 μmol/L; Fig. 5B). Similar results were seen in RWPE1 cells (data not shown). Apoptotic cells, showing cytoplasmic blebbing and detachment, were rare, but dividing cells virtually disappeared in cultures treated with quercetagenin at 6.25 μmol/L or higher concentrations (data not shown). DNA histograms obtained at 24 h after the addition of quercetagenin (6.25 μmol/L) or DMSO vehicle were very similar (Fig. 5C). Neither showed a <2n population suggestive of apoptosis. There was a slight increase in the proportion of cycling cells (S + G₂-M) in the drug-treated samples.

A PIM1 inhibitor would be predicted to inhibit the growth of cells that express the molecular target, more than cells with little or no *pim-1* expression. We examined the effects of quercetagenin on the growth of prostate cell lines that express a spectrum of PIM1 levels. RWPE2 cells expressed the highest amount of PIM1 protein, PC3 had an intermediate level, and LNCaP cells showed the lowest amount of kinase protein (Fig. 6A). Treatment of the

cells with various concentrations of quercetagenin for 72 h resulted in inhibition of cell growth (Fig. 6B). At all concentrations, RWPE2 cells were inhibited the most, being significantly more sensitive to quercetagenin growth inhibition than the other prostate cancer cell lines. PC3 cells showed intermediate growth suppression and were also significantly more sensitive than were LNCaP cells at quercetagenin concentrations of <12.5 μmol/L. Thus, the ability of the flavonol to inhibit proliferation was proportional to the amount of PIM1 protein in the target cells, particularly at lower drug concentrations. Although other interpretations are possible, these data support our observation that quercetagenin can act as a PIM1 inhibitor.

Discussion

The development of clinically useful small-molecule kinase inhibitors has been a seminal event in the world of oncology. Flavonoids were among the early scaffold structures identified as potential kinase inhibitors. However, although many flavones, isoflavones, and flavonols have been shown to regulate the activity of kinases in cell-based assays, fewer data exist to show that these molecules can directly bind and inhibit kinase targets both *in vitro* and in cells. It is clear that some flavonoids are ATP-competitive ligands for both tyrosine and serine-threonine kinases, as well as other ATP-binding enzymes. The flavonol quercetin is one such ligand, and its ability to directly bind to ATP-binding enzymes has been well shown. At low-micromolar concentrations, it directly binds and inhibits such diverse enzymes as the phosphatidylinositol 3-kinase (14), the epidermal growth factor receptor tyrosine kinase (15), retroviral reverse transcriptases (16), DNA gyrase (17), phosphodiesterases (18), and thioesterase reductase (19). Other direct flavonoid inhibitors have been described for RSK2 kinase (24), mitogen-activated protein/extracellular signal-regulated kinase 1 (25), and several cyclin-dependent kinases (23, 26–28). One such ligand, flavopipidol, has already entered clinical trials for the treatment of cancer. Others, such as PD98059, are familiar laboratory reagents for inhibition of kinase pathways. We now show, by means of crystallography, that quercetagenin is a direct ligand for the ATP-binding pocket of PIM1 kinase (Fig. 3).




Figure 2. Quercetagenin is a selective inhibitor of PIM1 kinase. Inhibitor activity of quercetagenin at 1 and 10 μmol/L final concentration against a spectrum of serine-threonine kinases of a panel of Kinases, assayed by KinaseProfiler assay. The activity in the presence of vehicle only was taken to be 100% activity. Columns, mean of duplicate determinations.

6 PIM1 Inhibitor




Figure 3. X-ray crystal structures of flavonoids bound to the ATP-binding sites of PIM1 and comparison with structures of flavonoids in complex with other kinases. Of the three compounds cocrystallized with PIM1, quercetagenin (A) orientates the B ring inside the pocket (orientation I), whereas both myricetin (B) and 5,7,3',4',5'-pentahydroxyflavone (C) flip the B ring out toward solvent (orientation II). Both binding orientations have been observed in the crystal structures of flavonoids bound to other kinases. 1E9W (D), quercetin in complex with phosphatidylinositol 3-kinase (I) and 1J02 (E), flavin in complex with CDK6) represent orientation I, whereas 2HCK (F, quercetin in complex with HCK) exemplifies orientation II. All pictures show residues that form hydrogen bonds with the inhibitors (Lys⁶⁷, Glu⁸⁹, Glu¹²¹, and Asp¹⁸⁶). In the PIM1 structures, the three residues near the ATP-binding site that differentiate PIM1 from PIM2 (Ser⁶⁴, Glu¹²⁴, and Val¹²⁶) are also shown. The inhibitors are colored by atom type: red, oxygen atoms; yellow, carbon atoms. Dashed purple lines, hydrogen bonds.

Specificity is always a concern with ATP pocket ligands. There are probably no absolutely selective inhibitors for a kinase but rather ligands that show a spectrum of affinities for their various targets. We have shown that quercetagenin is severalfold more active against PIM1 than against eight other serine-threonine kinases and a tyrosine kinase, either with *in vitro* assays or in cell cultures. Interestingly, quercetagenin showed 10-fold more selectivity for PIM1 than for the homologous PIM2 kinase (sequence identity 56%). The ATP-binding pockets of these two kinases are identical with the exception of three residues along the edge of the PIM1 ATP-binding pocket—Ser⁶⁴ (Ala⁶⁶ in PIM2), Glu¹²⁴ (Leu¹²⁰ in PIM2), and Val¹²⁶ (Ala¹²² in PIM2). Val¹²⁶ of PIM1 makes direct van der Waal's contact with the A ring of quercetagenin (Fig. 3A). Loss of such a contact due

to the Val-to-Ala substitution is likely a contributing factor to the reduced activity of the compound in PIM2. The other residues are located close to the hinge Arg⁸² (Arg⁷⁸ in PIM2). The polar side chains of Ser⁶⁴ and Glu¹²⁴ can form hydrogen bonds with Arg⁸², thus affecting its conformation. Substitutions of these residues to hydrophobic amino acids in PIM2 will change the local environment (Fig. 3A).

The only large-scale examination of the specificity of flavonoid kinase inhibitors was reported recently by Fabian et al. (8). This investigation used a competitive binding assay to predict the inhibitor potency and specificity of the test agents. Flavopiridol was tested for binding affinity to 119 kinases. Twenty-three kinases bound flavopiridol under the test conditions, with binding constants ranging from 0.019 to 6.6 μmol/L. Interestingly, the tested cyclin-dependent

kinases bound flavopiridol less well than did calcium/calmodulin-dependent protein kinase kinase 1. These data suggest that cyclin-dependent kinases may not be the only kinases inhibited in cells by flavopiridol. Both PIM1 and PIM2 were among the bound kinases, with binding constants of 0.52 and 0.65 $\mu\text{mol/L}$, respectively. Although there is no absolute correlation between binding constants and enzymatic activity, flavopiridol could conceivably inhibit the activity of both PIM1 and PIM2 in test systems.

Because quercetin has not been tested against a large number of other kinases, we cannot predict what other enzymes would be perturbed by this flavonoid. It is likely, however, that its spectrum of selectivity will be substantially different from that of flavopiridol. Quercetin showed clear preference for inhibiting PIM1 over PIM2, whereas flavopiridol did not. Furthermore quercetin inhibited the activity of the Aurora-A kinase (IC_{50} , $\sim 4 \mu\text{mol/L}$), a kinase that did not bind flavopiridol (8). The substantial




Figure 4. Quercetin inhibits PIM1 kinase activity in intact cells. **A**, RWPE2/neo or RWPE2/pim1 cells were cultured in unsupplemented keratinocyte medium overnight, then treated with quercetin (0–50 $\mu\text{mol/L}$) for 3 h. Lysates were then prepared and examined by immunoblotting with the indicated antibodies. **B**, quantitation of the pBAD(S112)/actin ratio in immunoblots by using densitometry on the digital file. IC_{50} , 5.55 $\mu\text{mol/L}$.

8 PIM1 inhibitor

homology between Aurora-A kinase and PIM1 kinase likely contributed to the low-level inhibitory activity of quercetin for the former; Aurora-A and PIM1 are 29% identical over their entire kinase domains; and the ATP binding pockets have 68% conserved amino acids.

An earlier, smaller-scale study looked at the effect of the flavonol quercetin on the *in vitro* kinase activity of 25 kinases, none of which were pim family kinases (29). At the tested concentration (20 $\mu\text{mol/L}$), quercetin inhibited the enzymatic activity of eight of the kinases. The propensity of this flavonol to form aggregates in aqueous solution has been advanced as an explanation for its widespread enzyme-inhibitory activity *in vitro* (30). We have not detected quercetin aggregates at concentrations of <10 $\mu\text{mol/L}$ in aqueous solution, using a light-scattering assay (data not shown). Thus, we feel that this artifact does not account for the ability of this flavonol to inhibit PIM1 at nanomolar concentrations.

Because of the potential ambiguities that may accompany the use of small-molecule kinase inhibitors, a series of standards have been proposed for their use (29). To validate the results, it is desirable to show that the effects

of an inhibitor disappear when a drug-resistant mutant of the protein kinase is overexpressed. Although convincing, this standard often fails due to the lack of an identified mutant with the desired properties. No such mutant has been identified for any of the pim kinases. Another potential standard is to show that the cellular effect of the drug occurs at the same concentrations that prevents the phosphorylation of an authentic physiologic substrate of the protein kinase. We have seen in these studies that half-maximal growth inhibition of prostate cancer cells occurred at a drug concentration (3.8 $\mu\text{mol/L}$) that approximated the IC_{50} for PIM1 enzyme inhibition in cells (5.5 $\mu\text{mol/L}$). Furthermore, the selectivity for prostate cancer growth inhibition, in proportion to endogenous PIM1 levels, was greatest at 6.25 $\mu\text{mol/L}$. Higher concentrations suppressed growth more, but the relationship to endogenous PIM1 levels was obscured. These data suggest that, at relatively low concentrations (perhaps 5–10 $\mu\text{mol/L}$), the growth-inhibitory effects of quercetin likely involve PIM1 antagonism. A third standard is to observe the same effect with at least two structurally unrelated inhibitors of the protein kinase. Previously described inhibitors of pim




Figure 5. Quercetin inhibits growth of prostate cancer cells at concentrations that inhibit PIM1 kinase activity. **A**, growth curve of RWPE2 cells with different concentrations of quercetin. Cell number is measured by crystal violet staining. Points, mean of triplicate determinations; from one of four similar experiments. **B**, calculation of ED_{50} at 24, 48, and 72 h of drug exposure. Average ED_{50} from all curves is 3.8 $\mu\text{mol/L}$. **C**, DNA histograms from RWPE2 cells treated with vehicle or quercetin (6.25 $\mu\text{mol/L}$, 24 h). Proportion of cells in G_0/G_1 or $\text{S} + \text{G}_2\text{-M}$ fractions. Columns, mean of triplicate determinations; from three independent experiments; bars, SD. P values show the probability of no difference by t test.




Figure 8. Quercetin inhibits the growth of prostate cancer cells in proportion to their content of PIM1. **A**, measurement of intracellular PIM1 in PC3, LNCaP, and RWPE2 prostate cancer cells by immunoblotting. **B**, growth inhibition by treatment of prostate cancer cells with quercetin for 72 h. Columns, mean of triplicate determinations from one of two similar experiments; bars, SD. *P* values were calculated by paired *t* tests and represent the probability that there is no difference between the two compared populations.

kinases are either less active or less specific flavonoids (7, 9), the same structural class as quercetin, or staurosporine analogues (8, 9, 21). We therefore used small interfering RNA as a genetic means to identify a *pim-1*-dependent phenotype. Passivation of prostate cells was suppressed with both the genetic and chemical inhibitors of PIM1 activity. These data show that quercetin is an authentic small-molecule inhibitor of PIM1 kinase.

The crystal structures of PIM1 complexed with quercetin, myricetin, and 5,7,3',4',5'-pentahydroxyflavone show that flavonoids bind to PIM1 in two distinct orientations. Although interesting, this is not a surprising observation, as flavones have shown a variety of binding modes in kinases (9, 22, 23, 26–28). An examination of the intermolecular interactions of each flavonoid with PIM1 does not clearly reveal why one orientation was adopted over the other. However, it is possible that the presence of these hydroxyl groups on the B ring of myricetin and 5,7,3',4',5'-pentahydroxyflavone discourages these two flavonoids from adopting the binding orientation observed

for quercetin. The hydrophobic side chain of Leu⁶⁹, which extends into the ATP pocket in the same region occupied by the B ring of quercetin (Fig. 3A), may be incompatible with the 5 hydroxyl group of myricetin and 5,7,3',4',5'-pentahydroxyflavone.

Both *pim-1* and *pim-2* can phosphorylate 4EBP-1, a regulator of protein translation (31, 32). Rapamycin was unable to block this effect. These data suggest that *pim* kinases may function in a parallel pathway to the phosphatidylinositol 3-kinase/AKT/mammalian target of rapamycin cascade to regulate and support protein synthesis under stress conditions. Because AKT-1 and PIM2 function cooperatively to induce lymphoma formation in transgenic mice (6), it may be necessary to target both pathways for effective antitumor effects. Several prototype AKT inhibitors have been described (33, 34). Our identification of quercetin as a PIM1 inhibitor provides a tool for tissue culture studies to investigate this hypothesis. Under the tested conditions, we found no evidence that quercetin inhibited the phosphorylation of AKT on Ser⁴⁷³. Thus, it may be possible to combine inhibitors of these kinases to detect additive or synergistic effects resulting from the blockade of the two kinase pathways.

References

- Wang Z, Bhattacharya N, Weaver M, et al. PIM1: a serine/threonine kinase with a role in cell survival, proliferation, differentiation, and tumorigenesis. *J Vet Sci* 2001;2:167–79.
- Gan KC, Wang L, Holey ER, et al. Structural basis of constitutive activity and a unique nucleotide binding mode of human PIM1 kinase. *J Biol Chem* 2005;280:6150–7.
- Lilly M, Kraft A. Enhanced expression of the MY 33,000 PIM1 kinase enhances factor-independent survival and inhibits apoptosis in murine myeloid cells. *Cancer Res* 1997;57:5348–55.
- Pficher TJ, Zhao S, Gilger JN, Jonaja B, Wojchowski DM. PIM1 kinase protects hematopoietic FDC cells from genotoxin-induced death. *Oncogene* 2000;19:3884–92.
- Lilly M, Sandholm J, Cooper JJ, Koskinen PJ, Kraft A. The PIM1 serine kinase promotes survival and inhibits apoptosis-related mitochondrial dysfunction in part through a bcl-2 dependent pathway. *Oncogene* 1999;18:4022–31.
- Hammaman PS, Fox CJ, Barbaum MJ, Thompson CB. Pim and Akt oncogenes are independent regulators of hematopoietic cell growth and survival. *Blood* 2005;105:4477–83.
- Jacob AMD, Black J, Futer O, et al. PIM1 ligand bound structures reveal the mechanism of serine/threonine kinase inhibition by LY294002. *J Biol Chem* 2005;280:13728–34.
- Fabian MA, Eggs WH III, Treiber DK, et al. A small molecule-kinase interaction map for clinical kinase inhibitors. *Nat Biotechnol* 2005;23:329–39.
- Ballock AN, DeBreceni JE, Fedorov OY, Nelson A, Manden ND, Krupp S. Structural basis of inhibitor specificity of the human protooncogene proximal insertion site in Moloney murine leukemia virus (PIM1) kinase. *J Med Chem* 2005;48:7804–14.
- Kung W, Silber E, Eppenberger U. Quantification of cells cultured on 96-well plates. *Anal Biochem* 1989;182:118–9.
- Yan B, Zemskova M, Holder S, et al. The PIM-2 kinase phosphorylates BAD on serine 112 and averts BAD-induced cell death. *J Biol Chem* 2003;278:45358–67.
- Aho TL, Sandholm J, Petala KJ, Mankonen HP, Lilly M, Koskinen PJ. PIM1 kinase promotes inactivation of the pro-apoptotic bcl protein by phosphorylating it on the Ser¹¹² gatekeeper site. *FEBS Lett* 2004;571:43–9.

13. Kumar A, Mandiyan V, Suzuki Y, et al. Crystal structures of proto-oncogene kinase PKM1: a target of aberrant somatic hypermutations in diffuse large cell lymphoma. *J Mol Biol* 2005;348:183–93.
14. Mayr GW, Windhoner S, Hillemeier K. Antiproliferative plant and synthetic polyphenols as specific inhibitors of vertebrate inositol-1,4,5-trisphosphate 3-kinases and inositol polyphosphate multikinase. *J Biol Chem* 2005;280:13229–40.
15. Lee LT, Huang YT, Hwang JJ, et al. Blockade of the epidermal growth factor receptor tyrosine kinase activity by quercetin and luteolin leads to growth inhibition and apoptosis of pancreatic tumor cells. *Anticancer Res* 2002;22:1815–27.
16. Chu SC, Hsieh YS, Lin JY. Inhibitory effects of flavonoids on Moloney murine leukemia virus reverse transcriptase activity. *J Nat Prod* 1992;55:179–83.
17. Pflüger A, Golob M, Hafner I, Oblok M, Solmajer T, Jenita R. Characterization of quercetin binding site on DNA gyrase. *Biochem Biophys Res Commun* 2003;308:530–8.
18. Ko WC, Shin CM, Lee YH, Chen JH, Huang HL. Inhibitory effects of flavonoids on phosphodiesterase isozymes from guinea pig and their structure-activity relationships. *Biochem Pharmacol* 2004;68:2007–94.
19. Lu J, Papp LV, Fang J, Rodriguez-Nieto S, Zhitovskiy B, Holmgren A. Inhibition of mammalian thionin oxidase by some flavonoids: implications for myricetin and quercetin anticancer activity. *Cancer Res* 2005;65:4410–8.
20. Bullock AN, Debeczeni J, Amos AJ, Krupp S, Turk BE. Structure and substrate specificity of the PKM1 kinase. *J Biol Chem* 2005;280:41875–82.
21. Debeczeni JE, Bullock AN, Adile GE, et al. Ruthenium half-sandwich complexes bound to protein kinase pkm-1. *Angew Chem Int Ed Engl* 2005;45:1580–5.
22. Walter BH, Record ME, Perisic O, et al. Structural determinants of phosphoinositide 3-kinase inhibition by wortmannin, LY294002, quercetin, myricetin, and staurosporine. *Mol Cell* 2000;6:909–19.
23. Lu H, Chang DJ, Banette B, Meijer L, Schube-Gahmen U. Crystal structure of a human cyclin-dependent kinase B complex with a flavonol inhibitor, fisetin. *J Med Chem* 2005;48:737–43.
24. Clark DE, Erlington TM, Smith JA, Frieson HF, Jr., Weber MJ, Lannigan DA. The serine/threonine protein kinase, p90 ribosomal S6 kinase, is an important regulator of prostate cancer cell proliferation. *Cancer Res* 2005;65:3108–16.
25. Alessi DR, Cende A, Cohen P, Dudley DT, Sakai AR. PD 098059 is a specific inhibitor of the activation of mitogen-activated protein kinase kinase *in vitro* and *in vivo*. *J Biol Chem* 1995;270:27489–94.
26. De AW, Jr., Mueller-Gleditsch HU, Schube-Gahmen U, Wortland PJ, Sauville E, Kim SH. Structural basis for specificity and potency of a flavonoid inhibitor of human CDK2, a cell cycle kinase. *Proc Natl Acad Sci U S A* 1998;95:2735–40.
27. Slicher F, Moretti J, Kuriyan J. Crystal structure of the Src family tyrosine kinase Hck. *Nature* 1997;385:602–9.
28. De Azevedo WF, Jr., Mueller-Gleditsch HU, Schube-Gahmen U, Wortland P, Sauville E, Kim SH. Structural basis for specificity and potency of a flavonoid inhibitor of human CDK2, a cell cycle kinase. *Proc Natl Acad Sci U S A* 1998;95:2735–40.
29. Davies SP, Reddy H, Cavano M, Cohen P. Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem J* 2000;351:95–105.
30. McGovern SL, Sheldahl BK. Kinase inhibitors: not just for kinases anymore. *J Med Chem* 2003;46:1470–83.
31. Fox CJ, Hammerman PS, Thompson CB. The Fim kinases control rapamycin-resistant T cell survival and activation. *J Exp Med* 2005;201:259–68.
32. Chen WW, Chan DC, Donald C, Lilly MB, Kraft AS. Fim family kinases enhance tumor growth of prostate cancer cells. *Mol Cancer Res* 2005;3:443–51.
33. Yang L, Dan HC, Sun M, et al. Akt/protein kinase B signaling inhibitor-2, a selective small molecule inhibitor of Akt signaling with antitumor activity in cancer cells overexpressing Akt. *Cancer Res* 2004;64:4094–9.
34. Banett SF, DeLeo-Jones D, Fu S, et al. Identification and characterization of pleckstrin-homology-domain-independent and isozyme-specific Akt inhibitors. *Biochem J* 2005;395:399–408.

MICHAEL BRIAN LILLY, MD

Updated 1/2006

Personal Data: Birth: 26 December, 1950, Atlanta, GA
Citizenship: USA

Office: Center for Molecular Biology and Gene Therapy
Loma Linda University School of Medicine
Loma Linda, CA 92354
Phone (909) 558-8777
Fax (909) 558-0177
e-mail: mlilly@llu.edu

Home: 511 West Olive Avenue
Redlands, CA 92373
(909) 792-5573

Education: Southern Adventist University, Collegedale, TN
1967-1971, B.A. (biology, chemistry)

Loma Linda University, Loma Linda, CA
1971-1975, M.D.

Postgraduate Training Internal Medicine residency
University of Alabama at Birmingham
Birmingham, AL 1975-1978

Hematology-Oncology Fellowship
University of Alabama at Birmingham
1978-1981

Faculty Positions: 6/81-6/82 Instructor in Medicine
UAB School of Medicine

6/82-10/88 Assistant Professor of Medicine,
UAB School of Medicine

6/82-10/88 Associate Scientist, Lurleen Wallace
Tumor Institute, Birmingham, AL

**Faculty Postions
(cont'd):**

6/89-9/98 Associate Professor of
Medicine, University of Washington School
of Medicine, Seattle, WA

4/96-10/96 Visiting Scientist, The Walter and Eliza
Hall Institute for Medical Research, Melbourne,
Victoria, AUSTRALIA

9/98 – present Professor of Medicine & Microbiology
Director, Center for Molecular Biology & Gene
Therapy Loma Linda University School of Medicine,
Loma Linda, CA

**Hospital Positions:
Alabama**

1981-1988 Attending Physician, University of
Hospitals and Clinic

1981-1988 Staff Oncologist, Birmingham VA Medical
Center, Birmingham, AL

1989-1998 Staff Oncologist, Seattle
VA Medical Center, Seattle, WA

1998-present Attending Physician, Loma Linda
University Medical Center, Loma Linda, CA

Honors:

1974 Alpha Omega Alpha

1980 National Research Service Fellow

1981 Fellow, American College of Physicians

Board Certification:

1979 American Board of Internal Medicine

1980 ABIM Subspecialty Exam, Hematology

1981 ABIM Subspecialty Exam, Med. Oncology

Licensure:

Alabama Medical License #7730 (3/77-12/91)

Washington State License #27864 (12/91 – 12/00)

California Medical License #G84932 (12/98 – present)

Organizations:

Fellow, American College of Physicians

Member, American Society of Hematology

Member, American Society for Bone Marrow
Transplantation

Member, American Society for Gene Therapy

**National
Professional
Responsibilities**

Member, ad hoc study sections for NIH:

1987 Diagnostic Radiology

1988 Experimental Therapeutics

Member, site visit team for program project

**Special Local
Responsibilities**

Dr. George Hahn, PI; Stanford University
1988, 1989
Member, site visit team for program project
Dr. Bayard Clarkson, PI, Memorial-Sloan
Kettering Inst., 1997
Member, Scientific Review Subcommittee
SVAMC, 1993, 1994, 1997
Member, Research & Development Committee
SVAMC, 1994, 1995
Member, Hospice Advisory Committee
SVAMC, 1994, 1995
Board Development Committee, Leukemia &
Lymphoma Society (Southern California Chapter),
2003

Consultant

Cetus Corporation (1986)
EncorePharma (2001-present)
Myriad Genetics (2002-present)
Exelixis Pharmaceuticals (2005-present)

GRANTS & CONTRACTS (PRINCIPAL INVESTIGATOR) *Note: This listing does not
include multicenter clinical trials in which Dr. Lilly was the local principal investigator.*

National Institutes of Health F32CA27980 *Hyperthermia of animal and human tumors;*
7/80-6/82

National Institutes of Health R01CA18138-11 *Prediction of thermal tolerance by in vivo*
NMR spectroscopy; 7/82-6/83

National Institutes of Health R01CA36790 *Assessment of hyperthermia by in vivo ³¹P-*
NMR spectroscopy; 9/84-9/87

Cetus Corporation *Characterization of a human granulocyte CSF; 7/85-6/86*

National Institutes of Health R01CA45672 *Cytokine signaling in myeloid leukemia;*
9/87-10/98

VA Merit Review Award *Non-protein hematopoietic agents; 10/90-4/97*

March of Dimes Birth Defects Foundation *Characterization of a 28kd protein related to*
G-CSF; 7/93-6/96

Lymphoma Research Foundation of America *Mechanism of action of the pim-1*
oncogene; 7/95-7/96

Roche Pharmaceuticals *Preclinical study of Roferon and bryostatin 1 in a melanoma model*; 1/98-12/99

Department of Defense, National Medical Technology Testbed #76-FY99: *Cell-permeable proteins for cell regulation*. 12/99 – 7/02

Leukemia Society of American Translational Award *Propionic Acid Analogues for CLL*. 9/1/01 – 8/31/05

Celgene Corporation, *Phase I-II trial of combined GM-CSF (sargramostim) and thalidomide for hormone-refractory prostate cancer* (5/02-5/04).

National Institutes of Health R03CA107820 *Molecular Targets of NSAIDs in Prostate Cancer*; (5/1/04 – 4/30/07)

Department of Defense, CDMRP Prostate Cancer Program PC040635 *Pim-1: A Molecular Target to Modulate Cellular Resistance to Therapy in Prostate Cancer* (10/04 – 10/07)

Pharmion Corporation, *Use of azacytidine to reverse silencing of GST-p1 in early prostate cancer*. (10/05 – 10/07)

GRANTS and CONTRACTS (Co-investigator)

National Institutes of Health R01CA097043 *Molecular pathology of 2-deoxy-5-azacytidine*; L. Sowers, PI; Michael Lilly, co-investigator (10% FTE). 7/1/03 – 6/30/08

PUBLICATIONS IN PEER-REVIEWED JOURNALS

1. Brezovich I, **Lilly M**, Durant J, Richards D: A practical system for clinical radiofrequency hyperthermia. *Int J Rad Onc Biol Phys* 7:423-430, 1981
2. Ng T, Evanochko W, Hiramoto R, Ghanta V, **Lilly M**, Lawson A Corbett T, Durant J, Glickson J: ³¹P-topical NMR spectroscopy of *in vivo* tumors. *Mag Res* 49:271-286, 1982.
3. **Lilly M**, Brezovich I, Chakraborty D, Atkinson W, Durant J, Ingram J, McElvein R: Hyperthermia with implanted electrodes: *in vitro* and *in vivo* correlations. *Int J Rad Onc Biol Phys* 9:373-382, 1983.
4. Evanochko W, Ng T, **Lilly M**, Kumar N, Durant J, Glickson J: *In vivo* ³¹P-NMR studies of the effect of cancer therapy on a murine mammary carcinoma. *Proc Natl Acad Sci USA* 80:334-338, 1983.

5. **Lilly M**, Ng T, Evanochko W, Kumar N, Elgavish G, Durant J, Hiramoto R, Ghanta V, Glickson J: *in vivo* ³¹P-NMR study of hyperthermia tumor treatment. *Cancer Res* 44:663-638, 1984.
6. Hiramoto R, Ghanta V, **Lilly M**: Reduction in tumor burden in murine osteosarcoma by hyperthermia and cyclophosphamide. *Cancer Res* 44:1405-1408, 1984.
7. Brezovich I, Atkinson W, **Lilly M**: Local hyperthermia with interstitial techniques. *Cancer Res* 44:46752s-4756s, 1984.
8. **Lilly M**, Brezovich I, Atkinson W: Hyperthermia with thermally self-regulating ferromagnetic implants. *Radiology* 154:243-244, 1985.
9. **Lilly M**, Katholi C, Ng T: Direct relationship between high-energy phosphate content and blood flow in thermally treated tumors. *JNCI* 75:885-889, 1985.
10. **Lilly M**, Omura G: Clinical pharmacology of oral intermediate dose methotrexate with or without probenecid. *Cancer Chemo Pharm* 15:220-222, 1985.
11. **Lilly M**, Carroll A, Prchal J: Lack of association between glutathione content and development of thermal tolerance in human fibroblasts. *Radiation Res* 106:41-46, 1986.
12. Tucker K, **Lilly M**, Heck L, Rado T: Characterization of a new human diploid myeloid leukemia cell line (PLB985) with granulocytic and monocytic differentiating capacity. *Blood* 70:372-378, 1987.
13. Devlin J, Devlin P, Myambo K, **Lilly M**, Rado T, Warren K: Isolation and expression of a cDNA encoding a human granulocyte colony-stimulating factor. *J Leukocyte Biol* 41:302-306, 1987.
14. **Lilly M**, Devlin J, Devlin P, Rado T: Production of granulocyte colony-stimulating factor by a human melanoma line. *Exp Hematol* 15:966-971, 1987.
15. Barton J, Parmley R, Butler T, Williamson S, **Lilly M**, Gualtieri R, Heck L: Differential staining of neutrophils and monocytes: surface and cytoplasmic iron-binding proteins. *Histochem J* 210:147-155, 1988.
16. Csepregy M, Yielding A, **Lilly M**, Scott C, Prchal J: Characterization of a new G6PD variant: G6PD Central City. *Am J Hematol* 28:61-62, 1988.
17. **Lilly M**, Kraft A: Leukemia-differentiating activity expressed by the human melanoma cell line LD1. *Leukemia Res* 12:213-218, 1988
18. Prchal J, Hauk M, Csepregy M, **Lilly M**, Berkow R, Scott C: Two apparent G6PD variants in normal XY man: G6PD Alabama. *Am J Med* 84:517-523, 1988.

19. Bailey A, **Lilly M**, Bertoli L, Ball E: An antibody which inhibits in vitro bone marrow proliferation in a patient with system lupus erythematosus and aplastic anemia. *Arthritis and Rheumatism* 32:901-905, 1989.
20. Kraft A, Williams F, Pettit R, **Lilly M**: Variable response of human myeloid leukemia lines and fresh cells to differentiating activity of bryostatin 1. *Cancer Res* 49:1287-1293, 1989.
21. Everson M, Brown C, **Lilly M**: IL6 and GM-CSF are candidate growth factors for chronic myelomonocytic leukemia cells. *Blood* 74:1472-1476, 1989.
22. Nemunaitis J, Andrews F, Mochizuki D, **Lilly M**, Singer J: Human marrow stromal cells: response to IL6 and control of IL6 expression. *Blood* 74:1693-1699, 1989.
23. Brezovich I, **Lilly M**, Meredith R, Weppleman B, Brawner W, Henderson R, Salter M: Hyperthermia of pet animal tumors with self-regulating ferromagnetic thermoseeds. *Intl J Hyperthermia* 6:117-130, 1990.
24. **Lilly M**, Tompkins C, Brown C, Pettit R, Kraft A: Differentiation and growth modulation of chronic myelogenous leukemia cells by bryostatin 1. *Cancer Res* 50:5520-5525, 1990.
25. **Lilly M**, Brown C, Pettit R, Kraft A: Bryostatin 1: a potential cytotoxic agent for chronic myelomonocytic leukemia cells. *Leukemia* 5:282-287, 1991.
26. Andrews D, **Lilly M**, Tompkins C, Singer J: Sodium vanadate, a tyrosine phosphatase inhibitor, affects expression of hematopoietic growth factors and extracellular matrix RNAs in SV40-transformed human marrow stromal cells. *Exp Hematol* 20:391-400, 1992.
27. **Lilly M**, Le T, Holland P, Hendrickson S: Expression of the pim-1 kinase is specifically induced in myeloid cells by growth factors whose receptors are structurally related. *Oncogene* 7:727-732, 1992.
28. Takahashi G, Andrews D, Tompkins C, Montgomery R, **Lilly M**, Singer J, Alderson M: Effect of granulocyte macrophage colony-stimulating factor (GM-CSF) and interleukin 3(IL3) on interleukin 8 (IL8) production in neutrophils and monocytes. *Blood* 81:357-364, 1993.
29. Polostkya A, Zhao C, **Lilly M**, Kraft A: A critical role for the intracellular domain of the alpha chain of the GM-CSF receptor in cell cycle transition. *Cell Growth & Diff* 4:5250531, 1993

30. Polostkya A, Zhao C, **Lilly M**, Kraft A: Mapping the intracytoplasmic regions of the alpha granulocyte-macrophage colony-stimulating factor receptor necessary for cell growth regulation. *J Biol Chem* 269:14607-14613, 1994
31. Takahashi G, Montgomery B, Stahl W, Crittenden C, Thorning D, **Lilly M**: Pentoxifylline inhibits tumor necrosis factor-alpha mediated cytotoxicity and activation of phospholipase A2 in L929 murine fibrosarcoma cells. *Intl J Immunopharm* 16:723-736, 1994
32. Sensebe L, Li J, **Lilly M**, Crittenden C, Herve P, Charbord P, Singer J: Non-transformed colony-derived stromal cell lines from normal human marrows. I. Growth requirements, characterization, and myelopoiesis-supportive ability. *Exp Hematol* 23:507-513, 1995
33. Asiedu C, Biggs J, **Lilly M**, Kraft A: Inhibition of leukemic cell growth by the protein kinase C activator Bryostatin 1 correlates with the dephosphorylation of cyclin-dependent kinase 2. *Cancer Res* 55:3716-3720, 1995
34. **Lilly M**, Vo K, Lee T, Takahashi G: Bryostatin 1 acts synergistically with interleukin 1 to promote the release of G-CSF and other myeloid cytokines from marrow stromal cells. *Exp Hematol* 24:613-621, 1996.
35. Matsuguchi T, Zhao Y, **Lilly MB**, Kraft AS: The cytoplasmic domain of the granulocyte-macrophage colony-stimulating factor (GM-CSF) receptor α subunit is essential for both GM-CSF-mediated growth and differentiation. *J Biol Chem* 272:17450-17459, 1997.
36. **Lilly M**, Kraft A: Enforced expression of the 33kd pim-1 kinase enhances factor-independent growth and inhibits apoptosis in murine myeloid cells. *Cancer Res* 57:5348-5355, 1997.
37. Matsuguchi T, **Lilly MB**, Kraft AS: Cytoplasmic domains of the human granulocyte-macrophage colony-stimulating factor receptor β chain (h β c) responsible for human GM-CSF-induced myeloid cell differentiation. *J Biol Chem* 273:19411-19418, 1998.
38. Frankel A, **Lilly M**, Kreitman R, Hogge D, Beran M, Freedman MH, Emanuel PD, McLain C, Hall P, Tagge E, Berger M, Eaves C: Diphtheria toxin fused to granulocyte-macrophage colony-stimulating factor is toxic to blasts from patients with juvenile myelomonocytic leukemia and chronic myelomonocytic leukemia. *Blood* 92:4279-4286, 1998.
39. **Lilly M**, Kiskonen P, Sandholm J, Cooper JJ, Kraft AS: The Pim-1 kinase prevents apoptosis-associated mitochondrial dysfunction, and supports cytokine-independent survival of myeloid cells in part through regulation of *bcl-2* expression. *Oncogene* 18:4022-4031, 1999.

40. **Lilly M**, Zemskova M, Frankel AE, Salo J, Kraft AS: Distinct domains of the human GM-CSF receptor alpha subunit mediate activation of JAK/STAT signaling and differentiation. *Blood* 97:1662-1670, 2001.
41. Wu X, Daniels T, Molinaro C, **Lilly MB**, Casiano C: Caspase cleavage of the nuclear autoantigen LEDGF/p75 abrogates its prosurvival function: implications for autoimmunity in atopic disorders. *Cell Death Differentiation* 9:915-924 (2002).
42. Lombano F, Kidder MY, **Lilly M**, Gollin YG, Block BS: Recurrence of microangiopathic hemolytic anemia after apparent recovery from the HELLP syndrome: A case report. *J Reprod Med* 47:875-877 (2002)
43. Frankel AE, Powell BL, **Lilly MB**. Diphtheria toxin conjugate therapy of cancer. *Cancer Chemother Biol Response Modif.* 2002;20:301-13. Review
44. Ionov Y, Le X, Tunquist BJ, Sweetenham J, Sachs T, Ryder J, Johnson T, **Lilly MB**, Kraft AS: Nuclear localization of the pim-1 protein kinase is necessary for its biologic effects. *Anticancer Res* 23(1A):167-78 (2003).
45. Yan B, Zemskova M, Holder S, Chin V, Kraft AS, Koskinen PJ, **Lilly MB**: The PIM-2 kinase phosphorylates BAD on serine-112 and reverses BAD induced cell death. *J Biol Chem* 278:45358-45367 (2003)
46. Aho TL, Sandholm J, Peltola KJ, Mankonen H, **Lilly M**, Koskinen PJ: Pim-1 kinase promotes inactivation of the pro-apoptotic Bad protein by phosphorylating it on the Ser¹¹² gatekeeper site. *FEBS Letters* 571:43-49 (2004).
47. Fodor I, Timiryasova T, Denes B, Yoshida J, Ruckle H, **Lilly M**: Vaccinia virus-mediated p53 gene therapy of bladder cancer in an orthotopic murine model. *J Urology* 173, 604-609 (2005).
48. Kim K-T, Baird K, Ahn J-Y, Meltzer P, **Lilly M**, Small D: Pim-1 is upregulated in constitutively activating FLT3 mutants and plays a role in FLT3-mediated cell survival. *Blood* 105(4), 1759-1767 (2005).
49. Chen WW, Chan DC, Donald C, **Lilly MB**, Kraft AS. Pim family kinases enhance tumor growth of prostate cancer cells. *Mol Cancer Res.* 2005 Aug;3(8):443-51.
50. Zemskova M, Wechter W, Yoshida J, Ruckle H, Reiter RE, **Lilly MB**: Gene expression profiling in R-flurbiprofen-treated prostate cancer: Identification of prostate stem cell antigen as a flurbiprofen-regulated gene. *Biochem Pharmacol* 72:1257-67 (2006)
51. Holder SL, Zemskova M, Bremner R, Neidigh J, **Lilly MB**: Identification of specific, cell-permeable small molecule inhibitor of the PIM1 kinase. (submitted, 2006)

BOOKS AND CHAPTERS:

Singer J, Slack J, **Lilly M**, Andrews D: Marrow stromal cells: response to cytokines and control of gene expression (in) *The Hematopoietic Microenvironment*. M. Wicha and M. Long, eds. Johns Hopkins Press, Baltimore, (1993).

RECENT ABSTRACTS:

Hromas R, Collins S, Bavisotto L, Hagen F, Raskind W, **Lilly M**, Kaushansky K: HEM-1, a potential transmembrane protein, is restricted to, yet ubiquitous in, hematopoietic cells. *Blood* 75:98a, 1990

Bianco J, Nemunaitis J, Andrews D, **Lilly M**, Shields A, Singer J: Combined therapy with pentoxifylline, ciprofloxacin, and prednisone reduces regimen related toxicity and accelerates engraftment in patients undergoing bone marrow transplantation. *Blood* 78:237a, 1991.

Lilly M, Sensebe L, Singer J: Characterization of cell-associated granulocyte colony-stimulating factor in human marrow stromal cells. *Blood* 78:261a, 1991 (oral presentation)

Takahasi G, **Lilly M**, Bianco J, Crittenden C, Singer J: Pentoxifylline inhibits tumor necrosis factor-alpha cytotoxicity and activation of phospholipase A2 in murine fibrosarcoma cells. *Blood* 78:323a, 1991.

Kirshbaum M, **Lilly M**: Multiple growth factors induce expression of the Bcl-2 protein in 32D murine hematopoietic cells, but differ in their ability to inhibit apoptosis. *Blood* 84:423a, 1994.

Lilly M, Pettit G: Identification of the cephalostatins as potent cytotoxic agents for myeloid leukemia cells. *Blood* 86:517a, 1995 (poster presentation)

Lilly M, Kraft A, Rotman E: Enforced expression of the human 33kd Pim-1 kinase enhances autonomous proliferation and tumorigenicity in factor-dependent murine FDCP1 cells. *Blood* 86:588a, 1995 (oral presentation).

Lilly M, Cooper JJ: Enforced expression of the human 33kd Pim-1 kinase prevents apoptosis-associated mitochondrial dysfunction and upregulates *bcl-2* mRNA expression in murine myeloid cells. (oral presentation, ASH 12/97)

Wu X, Molinaro C, **Lilly M**, Casiano C: Caspase-mediated cleavage of the transcription co-activator p75 during apoptosis (abstract #993). *Proc AACR* 41:155 (2000).

Quiggle DD, **Lilly M**, Murray ED, Gibson K, Leipold D, Gutierrez I, Loughman B, Wechter W: PK guided multi-dose, tolerance, and safety of E-7869 in prostate cancer patients (abstract #3874). *Proc AACR* 41:609 (2000)

Lilly M, Frankel AE, Salo J, Kraft AS: Distinct domains of the human GM-CSF receptor alpha subunit mediate activation of Jak/Stat signaling and differentiation (abstract #2455). *Blood* 96:572a (oral presentation, ASH 12/00)

Chen CS, **Lilly MB**, Wang FS, Howard FD, Houwen B: Rapid monitoring of peripheral blood stem cells (PBSC) mobilization by using cell membrane phospholipid content correlates well with CD34+ measurements, successful harvest and engraftment (abstract #1642). *Blood* 96:380a (poster presentation, ASH 12/00)

Lilly M, Ruckle H, Quiggle D, Gutierrez I, Murray D, Gibson K, Leipold D, Wechter W, Loughman B: Multi-dose phase I-II trial of E-7869 in prostate cancer patients: safety and time to PSA progression (TPSAP). *Proc AACR* 42:142 (2001)

Kastaros EP, Casiano C, Colburn KK, **Lilly M**, Weisbart RH, Kim J, Green LM: Lupus associated anti-guanosine antibodies: potential pathogenic effects. *Arthritis & Rheumatism* 44:S99 (2001)

Yan B, Zemskova M, Kraft AS, Koskinen PJ, **Lilly MB**: The pim-2 kinase phosphorylates Bad on serine-112 and reverses Bad induced cell death. (abstract #2919). *Blood* (poster presentation, ASH 12/02).

Lilly MB, Thorn S, Oberg K, Bashirova S, Zemskova M: The pim-1 serine/threonine kinase is primarily expressed in granulocytes and macrophages in inflamed tissues (abstract #981). *Blood* 102:276a (poster presentation, ASH 12/2003)

Neidigh J, Holder S, **Lilly MB**: Using docking to improve comparative modeling predictions: applications to pim-1 kinase. (poster presentation, *Structure-Based Drug Design 2004*, Boston, MA, April 26-28, 2004)

Chen CS, Zemskova M, Reiter R, **Lilly MB**: Gene expression profiling in R-flurbiprofen-treated prostate cancer: Identification of prostate stem cell antigen as a flurbiprofen-regulated gene. (poster presentation, *AACR 3rd Annual Conference on Frontiers in Cancer Prevention*, Seattle, WA; October 2004).

Lilly MB, Wechter W, Puuvula L, Henry H: R-Flurbiprofen (RFB) a non-steroidal anti-inflammatory drug (NSAID) with anti-tumor activity, inhibits the expression of CYP24 in murine prostate carcinomas. (poster presentation at *Biennial Vitamin D Conference "Vitamin D and Cancer Chemoprevention"*, NIH, Bethesda, MD, November 2004)