

REPORT DOCUMENTATION PAGE

*Form Approved
OMB No. 0704-0188*

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.

1. REPORT DATE (DD-MM-YYYY) 26-09-2007	2. REPORT TYPE Conference Proceedings	3. DATES COVERED (From - To)
---	--	------------------------------

4. TITLE AND SUBTITLE Multi-model Super-Ensemble Ocean Prediction: An Operational Example Using a Kalman Filter Approach in the Adriatic Sea	5a. CONTRACT NUMBER
	5b. GRANT NUMBER
	5c. PROGRAM ELEMENT NUMBER 0602435N

6. AUTHOR(S) Michel Rixen, Jeffery W. Book, Paul J. Martin, Nadia Pinardi, Paolo Oddo, Jacopo Chiggiato, Nello Russo	5d. PROJECT NUMBER
	5e. TASK NUMBER
	5f. WORK UNIT NUMBER 73-6648-07-5

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004	8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7330-06-7033
---	---

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5660	10. SPONSOR/MONITOR'S ACRONYM(S) ONR
	11. SPONSOR/MONITOR'S REPORT NUMBER(S)

12. DISTRIBUTION/AVAILABILITY STATEMENT
Approved for public release, distribution is unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT
Multi-model Super-Ensembles (SE) aim at combining optimally different models. A dynamic Kalman Filter version of this technique has been applied on a unique set of in-situ data and operational ocean models during the Dynamic of the Adriatic in Real-Time (DART) field experiment and is shown to significantly improve forecast skills.

20071119069

15. SUBJECT TERMS
Adriatic Sea, Models, Monitoring, Acoustics, Mesoscale Phenomena

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UL	18. NUMBER OF PAGES 1	19a. NAME OF RESPONSIBLE PERSON Jeffery Book
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 228-688-5251

38th CIESM Congress
Apr 9-13, 2009 in Turkey

MULTI-MODEL SUPER-ENSEMBLE OCEAN PREDICTION: AN OPERATIONAL EXAMPLE USING A KALMAN FILTER IN THE ADRIATIC SEA

M. Rixen^{1*}, J. Book², P. Martin², N. Pinardi³, P. Oddo³, J. Chiggiato⁴, N. Russo⁵

¹ NATO Undersea Research Center, La Spezia, Italy - rixen@nurc.nato.int

² Naval Research Laboratory, Stennis Space Center, MS39529, US

³ National Institute of Geophysics and Volcanology, Bologna, Italy

⁴ Servizio IdroMeteorologico, ARPA Emilia Romagna, Bologna, Italy

⁵ Universita Politecnica delle Marche, Ancona, Italy

Abstract

Multi-model Super-Ensembles (SE) aim at combining optimally different models. A dynamic Kalman Filter version of this technique has been applied on a unique set of in-situ data and operational ocean models during the Dynamics of the Adriatic in Real-Time (DART) field experiment and is shown to significantly improve forecast skills.

Keywords : Adriatic Sea, Models, Monitoring, Acoustics, Mesoscale Phenomena.

An increasing number of models are routinely providing operational weather forecasts and climate predictions. The SE technique [1], which uses an optimised combination of an ensemble of models, has previously been demonstrated to improve forecast skills in atmospheric models. Applications in the ocean are promising [2, 3]. However, they suffer from the lack of *in-situ* time series available in real-time.




Fig. 1. Time series of sound velocity (m/s) profiles, from top to bottom and left to right: SEPTR data, ADRICOSM, HOPS, NCOM and ROMS models.

The trawl-safe bottom mounted SEPTR platform developed at NURC (NATO Undersea Research Center), which is equipped with an ADCP and a CTD profiling unit transmitting data in real-time through a dedicated Globalstar link, now overcomes this limitation. During the DART (Dynamics of the Adriatic in Real-Time) field experiment in March 2006, 6 SEPTR were moored in the Gulf of Manfredonia in the Adriatic. In parallel, a suite of atmospheric, wave and ocean models were run operationally for the same period. This unique combination of data and models (Fig. 1) was used to derive SE products using a Kalman Filter-like Dynamic Linear Model [4] to account for the rapid changes in individual model skills and derive associated uncertainties. Our results suggest that the combination of SEPTR and operational models may decrease the errors on sound velocity profiles from 3-5 m/s down to 1-2 m/s (Fig. 2) at a marginal cost.




Fig. 2. Error on sound velocity for a 4-day forecast: (top) Standard Super-ensemble (bottom) Kalman Filter Dynamic Linear Model.

Acknowledgments

This work was performed at NURC and funded by the North Atlantic Treaty Organization.

References

- 1 - Krishnamurti T., Kishtawal C., LaRow T.E., Bachiocchi D.R., Zhang Z., Williford C.E., Gadgil S. and Surendran S.S., 1999. Improved weather and seasonal climate forecasts from multimodel superensemble. *Science*, 285 (5433): 1548-1550. DOI: 10.1126/science.285.5433.1548.
- 2 - Rixen M. and Ferreira-Coelho E., 2004. Operational prediction of acoustic properties in the ocean using multi-model statistics. *Oc. Mod.*, 11 (3-4): 428-440. DOI:10.1016/j.ocemod.2005.02.002.
- 3 - Rixen M. and Ferreira-Coelho E., 2006. Operational surface drift prediction using linear and non-linear hyper-ensemble statistics on atmospheric and ocean models. *J. Marine Syst.*, in press.
- 4 - Shin D. and Krishnamurti T., 2003. Short- to medium-range superensemble precipitation forecasts using satellite products: 1. Deterministic forecasting. *J. Geophys. Res.*, 108 (8): 8383. DOI 10.1029/2001JD001510.