




MSMR

A publication of the Armed Forces Health Surveillance Center



MEDICAL SURVEILLANCE MONTHLY REPORT

INSIDE THIS ISSUE:

| | |
|--|----|
| Pneumonia-influenza and severe acute respiratory illnesses, U.S. Armed Forces, January 1997-April 2008 | 2 |
| Allergic rhinitis, U.S. Armed Forces, 1998-2007 | 8 |
| Update: Deployment health assessments, U.S. Armed Forces, May 2008 | 12 |
| <i>Summary tables and figures</i> | |
| Sentinel reportable medical events, active components, U.S. Armed Forces, cumulative numbers through May 2007 and May 2008 | 16 |
| Acute respiratory disease, basic training centers, U.S. Army, June 2006-June 2008 | 23 |
| Deployment-related conditions of special surveillance interest | 24 |

Read the MSMR online at: <http://www.afhsc.army.mil>

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

| | | | | |
|--|------------------------------------|---|-----------------------------|----------------------------------|
| 1. REPORT DATE JUN 2008 | 2. REPORT TYPE | 3. DATES COVERED 00-00-2008 to 00-00-2008 | | |
| 4. TITLE AND SUBTITLE Medical Surveillance Monthly Report (MSMR). Volume 15, Number 5, June 2008 | | 5a. CONTRACT NUMBER | | |
| | | 5b. GRANT NUMBER | | |
| | | 5c. PROGRAM ELEMENT NUMBER | | |
| 6. AUTHOR(S) | | 5d. PROJECT NUMBER | | |
| | | 5e. TASK NUMBER | | |
| | | 5f. WORK UNIT NUMBER | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Center for Health Promotion and Preventive Medicine, Armed Forces Health Surveillance Center (AFHSC), 2900 Linden Lane, Suite 200, Silver Spring, MD, 20910 | | 8. PERFORMING ORGANIZATION REPORT NUMBER | | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | 10. SPONSOR/MONITOR'S ACRONYM(S) | | |
| | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | |
| 13. SUPPLEMENTARY NOTES | | | | |
| 14. ABSTRACT | | | | |
| 15. SUBJECT TERMS | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT | |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | Same as Report (SAR) | 18. NUMBER OF PAGES 28 |
| | | | | 19a. NAME OF RESPONSIBLE PERSON |

Update: Pneumonia-Influenza and Severe Acute Respiratory Illnesses, U.S. Armed Forces, January 1997-April 2008

Throughout history, military populations have been at high risk of acute respiratory illnesses.¹⁻⁴ In the U.S. military, acute respiratory illnesses are leading causes of hospitalizations and ambulatory visits of service members.⁵⁻⁷

Upper respiratory infectious illnesses are extremely common among U.S. service members, particularly among recruits and during fall-winter “cold and influenza” seasons. Upper respiratory illnesses are among the leading causes of medical encounters and limited duty dispositions of service members.^{6,7} Acute infections of the lower respiratory tract (e.g., pneumonias) are less frequent but more debilitating than upper respiratory illnesses. Consistently, pneumonias and influenza are among the leading causes of hospitalizations of service members.^{5,7} Finally, there are sporadic cases and rare outbreaks of severe, life threatening acute respiratory illnesses, usually in recruit camps and during training and operational deployments.⁸⁻¹⁰ In 2003, there were 19 cases of severe acute respiratory illnesses among U.S. service members deployed in the Middle East and south-central Asia. Two of the cases

were fatal – despite extensive investigation, the etiologies of the cases were not identified.¹⁰


This report summarizes frequencies, seasonal variability, and general trends of hospitalizations of U.S. service members for “pneumonia and influenza” and severe acute respiratory illnesses (e.g., acute respiratory failure, acute respiratory distress) from January 1997 to April 2008.

Methods:

The surveillance period was 1 January 1997 to 30 April 2008. The surveillance population included all individuals who served in any component of the U.S. Armed Forces any time during the surveillance period.

For surveillance purposes, a case of “pneumonia-influenza (P&I)” was defined as a hospitalization with a primary (first-listed) diagnosis of “pneumonia and influenza” (ICD-9: 480-487); or a primary diagnosis of “acute respiratory infection” (ICD-9 460-466) plus a secondary diagnosis (in diagnostic

Figure 1. Hospitalizations for pneumonia or influenza, by month, active component, U.S. Armed Forces, January 1997-April 2008



position 2 through 8) of “pneumonia and influenza” (ICD-9: 480-487).

A case of “severe acute respiratory illness (SARI)” was defined as a hospitalization with a primary diagnosis of “acute respiratory failure” (ICD-9-CM: 518.81) or “other pulmonary insufficiency” (which includes “acute respiratory distress, acute respiratory insufficiency, adult respiratory distress syndrome [ARDS]”) (ICD-9-CM: 518.82); or a secondary diagnosis (in diagnostic position 2 through 8) of “acute respiratory failure” (ICD-9-CM: 518.81) or “other pulmonary insufficiency” (ICD-9-CM: 518.82) plus a primary diagnosis of one of the following: “respiratory distress/insufficiency” (ICD-9-CM: 786.09), “acute respiratory infection” (ICD-9 460-466) or “pneumonia and influenza” (ICD-9: 480-487).

For surveillance purposes, a “respiratory illness year” was considered 1 July through 30 June of the following year. For each service member during each respiratory illness year, only one incident episode each of pneumonia-influenza and severe acute respiratory illness were included in analyses.

Results:

Pneumonia-influenza:

During the period, there were 11,038 incident hospitalizations for pneumonia and influenza (Table 1). The most and fewest cases per respiratory illness year were in 2002-3 (n=1,363) and 1998-9 (n=760), respectively (Table 1).

Over the entire period, the fewest cases were in the “spring” and “summer” (mean cases per month, April-June: 65.1; July-September: 72.5) and the most in the “fall” and “winter” (mean cases per month, October-December: 98.9; January-March: 87.9) (Table 1). In most years, the distinct seasonality in P&I case occurrence was apparent: in general, P&I-related hospitalizations sharply increased from late summer through

early fall and sharply decreased through the spring (Figure 1).

On average, the most cases per month each year were in October (mean: 99.8) and November (mean: 99.3) and the fewest in June (mean: 61.8). During the period, the most cases in any month were in December 2002 (n=200) and the fewest were in June 1999 and April 2008 (n=30) (Figure 1).

In general, there were relatively fewer P&I-related hospitalizations during the beginning and end of the surveillance period relative to the middle. For example, during 35 (80%) of the 44 months from January 1997 through August 2000 and during 28 (90%) of the 31 months from October 2005 through April 2008, there were “deficits” of P&I-related hospitalizations (relative to the means for the respective months overall) (Figure 2). In contrast, during 43 (70%) of the 61 months from September 2000 to September 2005, there were “excess” P&I-related hospitalizations (relative to the means for the respective months overall) (Figure 2). Of particular note, during the 10-month period from March-December 2003, there were more than 35 “excess” P&I-related hospitalizations each month and 507 excess hospitalizations overall (Figure 2).

Severe acute respiratory illness:

During the period, there were 626 incident hospitalizations for severe acute respiratory illnesses (SARI) (Table 1). The most and fewest cases per year were in 2006-7 (n=99) and 1998-9 (n=18), respectively (Table 1). Based on dispositions reported on hospital discharge records, case fatality overall was 9.4% (with no clear trend of increasing or decreasing case fatality over the period [data not shown]).


Numbers of SARI-related hospitalizations generally increased during the period (Table 1, Figures 3). For example, during 29 (81%) of the 36 months from January 1997 through December 1999, there were “deficits” of SARI-related

Table 1. Incident hospitalized cases of pneumonia-influenza and severe acute respiratory illness, by calendar quarter, U.S. Armed Forces, January 1997-April 2008

| <i>Pneumonia-influenza</i> | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | Total | Mean cases per month | |
|----------------------------|------|------|------|------|-------|-------|-------|------|-------|------|------|------|--------|----------------------|--------------|
| | | | | | | | | | | | | | | No. | Relative no. |
| Jan-Mar | 267 | 259 | 268 | 238 | 329 | 288 | 276 | 241 | 284 | 216 | 241 | 259 | 3,166 | 87.9 | 1.35 |
| Apr-Jun | 153 | 130 | 129 | 170 | 210 | 197 | 331 | 243 | 243 | 190 | 188 | 30 | 2,214 | 65.1 | ref |
| Jul-Sep | 165 | 169 | 185 | 199 | 232 | 306 | 352 | 203 | 235 | 168 | 180 | | 2,394 | 72.5 | 1.11 |
| Oct-Dec | 231 | 194 | 283 | 314 | 271 | 450 | 494 | 290 | 266 | 218 | 253 | | 3,264 | 98.9 | 1.52 |
| <i>Total</i> | 816 | 752 | 865 | 921 | 1,042 | 1,241 | 1,453 | 977 | 1,028 | 792 | 862 | 289 | 11,038 | 81.2 | |

| <i>Severe acute respiratory illness</i> | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | Total | Mean cases per month | |
|---|------|------|------|------|------|------|------|------|------|------|------|------|-------|----------------------|--------------|
| | | | | | | | | | | | | | | No. | Relative no. |
| Jan-Mar | 13 | 13 | 5 | 10 | 15 | 15 | 15 | 12 | 24 | 14 | 25 | 15 | 176 | 4.9 | 1.20 |
| Apr-Jun | 12 | 8 | 7 | 7 | 11 | 9 | 16 | 13 | 24 | 17 | 19 | 0 | 143 | 4.2 | 1.04 |
| Jul-Sep | 5 | 3 | 9 | 5 | 9 | 12 | 17 | 13 | 16 | 31 | 14 | | 134 | 4.1 | ref |
| Oct-Dec | 7 | 3 | 13 | 18 | 10 | 15 | 19 | 17 | 26 | 24 | 21 | | 173 | 5.2 | 1.29 |
| <i>Total</i> | 37 | 27 | 34 | 40 | 45 | 51 | 67 | 55 | 90 | 86 | 79 | 15 | 626 | 4.6 | |

Figure 2. Number (bars) and percent (line) of excess/deficit of hospitalizations for pneumonia-influenza, relative to the mean number for the respective month for the entire period, U.S. Armed Forces, January 1997-April 2008



hospitalizations (relative to the means for the respective months overall) (Figure 4). In contrast, during 33 (92%) of the 36 months from January 2005 to December 2007, there were “excess” SARI-related hospitalizations (relative to the means for the respective months overall) (Figure 4). Of note, during the 4-months from July through September 2006, there were approximately 27 “excess” (of 44 overall) SARI-related hospitalizations of U.S. service members (Figure 4).

As with pneumonia and influenza, the fewest SARI-related cases were in the “spring” and “summer” (mean cases per month, April-June: 4.2; July-September: 4.1) and the most in the “fall” and “winter” (mean cases per month, October-December: 5.2; January-March: 4.9) (Table 1). However, in contrast to the distinct seasonality of pneumonia-influenza incidence, there was no consistent month-to-month or seasonal variability in SARI-related hospitalizations (Table 1, Figure 3). For the entire period, the most cases in any month ($n=13$) were in June 2005, July 2006, and October 2006; while in eight months overall – including six (14%) of 43 months from January 1997 to July 2000, there were no SARI-related hospitalizations (Figure 3).

Finally, there was no apparent relationship between the number of pneumonia-influenza-related hospitalizations

each month and the number of SARI cases in the same month (data not shown).

Data analysis by Vicki Jeffries and Tannya Martin.

Editorial comment:

Compared to the general experience since 1997, in the past few years – particularly since the fall of 2005, there have been relatively few (accounting for the month) hospitalizations for pneumonia-influenza. U.S. military members receive annual influenza immunizations. The recent experience regarding pneumonia-influenza reflects, at least in part, the effectiveness of influenza vaccines against predominant circulating strains.¹¹

In contrast, since the fall of 2002 – particularly since January 2005, there were more hospitalizations for severe acute respiratory illness than in the preceding 5-years. Because the case definition used for this analysis required a respiratory illness-specific primary diagnosis (indicative of the primary reason for hospitalization), it is unlikely that the increase reflects complications of combat-related trauma or associated care. Also, in response to the cluster of “severe acute pneumonitis” cases among deployed service members in

Figure 3. Hospitalizations for severe acute respiratory illness, by month, U.S. Armed Forces, January 1997-April 2008





Figure 4. Number (bars) and percent (line) of excess/deficit of hospitalizations for severe acute respiratory illness, relative to the mean number for the respective month for the entire period, U.S. Armed Forces, January 1997-April 2008



the spring-summer of 2003⁹, the MSMR has tracked incident episodes of “severe acute pneumonia” among service members deployed to or within 30 days of returning from service in Afghanistan or Iraq (using a similar but not identical case definition as for this report) (see page 26). Since 2004, there have been consistently few cases per month of severe acute pneumonia temporally related to deployment (range, cases per month: 0-4; annual mean cases per month: 0.6-1.2) (see page 26). Thus, deployment-related cases may explain some but not all of the increase in severe acute respiratory illness cases since 2002. Another contributory cause is likely adenovirus type 14 which has recently emerged as a significant pathogen among U.S. military trainees.¹²⁻¹⁴


This analysis clearly documents the seasonality of pneumonia-influenza incidence among U.S. service members. From 1997 through 2007, the fewest pneumonia-influenza hospitalizations were in June and the most in October. In most years of the period, there were sharp increases in pneumonia-influenza hospitalizations beginning in the late summer (July-August). The findings suggest that annual activities to counter seasonal pneumonia-influenza risk should be conducted as soon as possible after the current year’s vaccines are available.

References:

1. Sartin JS. Infectious diseases during the Civil War: the triumph of the “Third Army”. *Clin Infect Dis*. 1993 Apr;16(4):580-4.


2. Oxford JS, Sefton A, Jackson R, et al. World War I may have allowed the emergence of “Spanish” influenza. *Lancet Infect Dis*. 2002 Feb;2(2):111-4.
3. Gray GC, Callahan JD, Hawksworth AW, Fisher CA, Gaydos JC. Respiratory diseases among U.S. military personnel: countering emerging threats. *Emerg Infect Dis*. 1999 May-Jun;5(3):379-85.
4. Gray GC, Mitchell BS, Tueller JE, Cross ER, Amundson DE. Pneumonia hospitalizations in the US Navy and Marine Corps: rates and risk factors for 6,522 admissions, 1981-1991. *Am J Epidemiol*. 1994 Apr 15;139(8):793-802.
5. Hospitalizations among active duty members, US Armed Forces, 2007. *MSMR* 2008 Apr;15(3); 2-8.
6. Ambulatory visits among active duty members, US Armed Forces, 2007. *MSMR* 2008 Apr;15(3); 9-14.
7. Absolute and relative morbidity burdens attributable to various illnesses and injuries, U.S. Armed Forces, 2007. *MSMR* 2008 Apr;15(3); 15-20.
8. Sanchez JL, Craig SC, Kolavic S, et al. An outbreak of pneumococcal pneumonia among military personnel at high risk: control by low-dose azithromycin postexposure chemoprophylaxis. *Mil Med*. 2003 Jan;168(1):1-6.
9. Outbreak of group A streptococcal pneumonia among Marine Corps recruits—California, November 1-December 20, 2002. *MMWR*. 2003 Feb 14;52(6):106-9.
10. Severe acute pneumonitis among deployed U.S. military personnel—Southwest Asia, March-August 2003. *MMWR*. 2003 Sep 12;52(36):857-9
11. Update: Influenza activity --- United States, September 30, 2007-April 5, 2008, and composition of the 2008--09 influenza vaccine. *MMWR*. 2003 Apr 18;57(15):404-9.
12. Metzgar D, Osuna M, Kajon AE, et al. Abrupt emergence of diverse species B adenoviruses at US military recruit training centers. *J Infect Dis*. 2007 Nov 15;196(10):1465-73. Epub 2007 Oct 31.
13. Acute respiratory disease associated with adenovirus serotype 14--four states, 2006-2007. *MMWR*. 2007 Nov 16;56(45):1181-4.
14. Binn LN, Sanchez JL, Gaydos JC. Emergence of adenovirus type 14 in US military recruits--a new challenge. *J Infect Dis*. 2007 Nov 15;196(10):1436-7. Epub 2007 Oct 31.

Figure 5. Relationship between pneumonia-influenza hospitalizations each month and severe acute respiratory illness cases during the same month, U.S. Armed Forces, January 1997-April 2008



SURVEILLANCE SNAPSHOT

Medical encounters for occupational exposures to blood/other body fluids,
by military status, U.S. Armed Forces, 1998-2007



Allergic Rhinitis, U.S. Armed Forces, 1998-2007

Allergic rhinitis (“hay fever”) affects an estimated 10-30% of the general population of the United States.¹ Between 2000 and 2005, expenditures for ambulatory care and treatment of allergic rhinitis in the U.S. increased 73% to \$11.2 billion.²

In 2007, approximately 48,000 U.S. service members had at least one medical encounter for allergic rhinitis. Allergic rhinitis was the primary (first-listed) diagnosis for nearly 3% of all medical encounters of service members in 2007.³ It is a significant cause of morbidity and lost duty time, especially during the spring and fall allergy seasons.

This report summarizes frequencies, rates, seasonal trends and correlates of risk of allergic rhinitis among U.S. service members during the past 10 years.

Methods:

The surveillance period was 1 January 1998 to 31 December 2007. The surveillance population included all service members who served in an active component of the U.S. Armed Forces any time during the surveillance period. A case of allergic rhinitis was defined as an outpatient encounter with a diagnoses of allergic rhinitis (ICD-9-CM:

477) in any diagnostic position. For this analysis, incident ambulatory visits were limited to one per service member per 30 consecutive days of follow-up.

Results:

During the 10-year surveillance period, 438,330 service members had approximately 1.2 million ambulatory medical encounters related to allergic rhinitis. Approximately 79% of all affected individuals had more than one allergic rhinitis-related clinic visit during the period. Among service members with multiple visits, the mean number was 2.7 visits per year.

There were 833,772 incident (one per person per 30 days) ambulatory visits for allergic rhinitis during the surveillance period (Table 1). Incidence rates increased with age and were higher among females than males and in the Air Force than the other services (Table 1). Numbers of affected individuals and incidence rates generally increased during the first seven years and were stable during the last three years of the surveillance period (Figure 1).


Throughout the period, there were distinct seasonal trends in allergic rhinitis incidence. Each year, there were large spring and smaller fall outbreaks (Figure 2). During the ten spring outbreaks, numbers of incident ambulatory visits

Table 1. Number and rates (per 1,000 person-years) of incident* ambulatory visits for allergic rhinitis, active components, U.S. Armed Forces, 1998 - 2007

| | 1998 | | 1999 | | 2000 | | 2001 | | 2002 | | 2003 | |
|----------------------------|-----------------|------|-----------------|------|-----------------|-------|-----------------|-------|-----------------|-------|-----------------|-------|
| | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate |
| Total | 57,414 | 41.3 | 66,679 | 49.0 | 76,307 | 56.0 | 84,059 | 61.7 | 83,986 | 60.4 | 88,683 | 62.9 |
| Service | | | | | | | | | | | | |
| Air Force | 26,657 | 73.1 | 32,626 | 91.5 | 35,427 | 100.7 | 38,123 | 109.5 | 37,414 | 103.9 | 40,232 | 109.6 |
| Army | 18,065 | 37.9 | 19,389 | 41.4 | 24,298 | 51.3 | 28,043 | 59.2 | 28,289 | 58.9 | 28,677 | 58.5 |
| Navy | 9,993 | 26.5 | 11,618 | 31.9 | 13,320 | 36.4 | 13,955 | 37.9 | 14,415 | 38.2 | 15,685 | 41.6 |
| Marine Corps | 2,699 | 15.8 | 3,046 | 17.8 | 3,262 | 19.0 | 3,938 | 23.0 | 3,868 | 22.5 | 4,089 | 23.1 |
| Gender | | | | | | | | | | | | |
| Female | 15,607 | 80.6 | 18,216 | 94.5 | 21,233 | 107.5 | 23,661 | 117.4 | 23,702 | 114.0 | 25,546 | 120.7 |
| Male | 41,807 | 35.0 | 48,463 | 41.5 | 55,074 | 47.3 | 60,398 | 52.1 | 60,284 | 51.0 | 63,137 | 52.7 |
| Race/ethnicity | | | | | | | | | | | | |
| Black non-Hispanic | 12,329 | 45.0 | 13,952 | 51.8 | 16,315 | 60.0 | 18,274 | 67.0 | 17,814 | 65.5 | 19,135 | 71.4 |
| Asian | 2,260 | 44.7 | 2,766 | 53.0 | 3,277 | 60.2 | 3,591 | 64.5 | 3,629 | 61.6 | 4,100 | 66.5 |
| White non-Hispanic | 32,066 | 38.8 | 39,578 | 47.1 | 47,002 | 54.9 | 51,963 | 60.9 | 52,247 | 60.1 | 54,164 | 61.3 |
| Hispanic | 8,624 | 46.0 | 7,970 | 54.3 | 6,885 | 54.4 | 6,933 | 55.3 | 7,028 | 53.4 | 7,536 | 54.8 |
| Native American | 528 | 37.5 | 649 | 42.8 | 771 | 45.9 | 891 | 48.6 | 941 | 47.3 | 1,056 | 49.1 |
| Other | 1,607 | 43.9 | 1,764 | 48.3 | 2,057 | 56.7 | 2,407 | 67.3 | 2,327 | 62.4 | 2,692 | 69.3 |
| Age group | | | | | | | | | | | | |
| <20 | 2,722 | 20.6 | 3,134 | 23.0 | 4,145 | 28.1 | 5,299 | 33.3 | 5,342 | 33.5 | 5,116 | 33.7 |
| 20-24 | 12,532 | 29.8 | 14,688 | 35.3 | 17,009 | 39.6 | 19,827 | 45.1 | 20,106 | 43.4 | 21,865 | 45.0 |
| 25-29 | 12,144 | 41.9 | 14,102 | 50.9 | 15,612 | 58.0 | 16,271 | 62.9 | 16,402 | 62.9 | 17,819 | 65.9 |
| 30-34 | 10,841 | 46.8 | 11,847 | 54.6 | 13,339 | 64.3 | 14,343 | 71.7 | 14,028 | 70.0 | 14,567 | 72.6 |
| 35-39 | 11,115 | 56.5 | 13,260 | 67.9 | 14,645 | 76.1 | 15,638 | 83.3 | 14,824 | 80.2 | 14,894 | 82.9 |
| >39 | 8,060 | 68.0 | 9,648 | 81.7 | 11,557 | 99.3 | 12,681 | 108.7 | 13,284 | 109.9 | 14,422 | 117.5 |
| Military occupation | | | | | | | | | | | | |
| Combat | 6,776 | 24.1 | 7,700 | 28.0 | 9,149 | 33.3 | 10,077 | 36.9 | 9,961 | 36.2 | 10,207 | 36.6 |
| Health care | 8,976 | 73.2 | 9,988 | 85.7 | 10,591 | 92.6 | 11,606 | 100.8 | 11,487 | 99.0 | 12,189 | 104.4 |
| Other | 41,662 | 42.2 | 48,991 | 50.6 | 56,567 | 58.1 | 62,376 | 64.1 | 62,538 | 62.6 | 66,287 | 65.3 |

*One ambulatory visit per service member per 30-day period

Figure 1. Number and rates of incident ambulatory visits for allergic rhinitis, active components, U.S. Armed Services, 1998-2007



sharply increased in March and peaked in April (n=5) or May (n=5). The most incident visits per day during any months of the period were in April 2004 (incident visits per day: 464) and April 2005 (incident visits per day: 467).

During each fall season except one, the peaks of incidence were in September (Figure 2). Invariably, the magnitudes of the peaks of incidence in the fall were much smaller than those in the spring — in general, there were approximately 70% more incident visits during the peak months of each spring season compared to the respective fall seasons (ratio, incident visits during peak months of spring and fall, 1998-2007: 1.44-2.03) (Figure 2). Seasonal trends in gender- and age-defined subgroups reflected the overall trend (data not shown).

Data analysis by Vicki Jeffries.


Editorial comment:

In the past 10 years, approximately 70,000 U.S. service members per year have received medical care for allergic rhinitis. Between 1998 and 2005, the number of service members treated each year for allergic rhinitis increased by approximately two-thirds. The trend may reflect a true increase in incidence of symptomatic allergic rhinitis, changes in health care seeking for treatment of allergic rhinitis symptoms, and/or changes in case ascertainment (e.g., diagnosis, reporting).

Table 1 continued. Number and rates (per 1,000 person-years) of incident* ambulatory visits for allergic rhinitis, active components, U.S. Armed Forces, 1998 - 2007

| | 2004 | | 2005 | | 2006 | | 2007 | | Total | | | |
|----------------------------|-----------------|-------|-----------------|-------|-----------------|-------|-----------------|-------|-----------------|-------|------------|--|
| | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate | Incident Visits | Rate | Rate ratio | |
| Total | 93,233 | 65.9 | 94,408 | 68.6 | 94,766 | 69.6 | 94,237 | 69.3 | 833,772 | 60.5 | | |
| Service | | | | | | | | | | | | |
| Air Force | 41,807 | 111.7 | 41,646 | 117.8 | 41,543 | 120.5 | 41,506 | 124.5 | 376,981 | 106.1 | ref | |
| Army | 31,128 | 63.2 | 31,884 | 65.6 | 32,453 | 65.9 | 32,448 | 63.6 | 274,674 | 56.7 | 0.53 | |
| Navy | 16,123 | 43.4 | 16,355 | 45.6 | 16,119 | 46.6 | 15,582 | 46.7 | 143,165 | 39.4 | 0.37 | |
| Marine Corps | 4,175 | 23.6 | 4,523 | 25.4 | 4,651 | 26.1 | 4,701 | 25.8 | 38,952 | 22.3 | 0.21 | |
| Gender | | | | | | | | | | | | |
| Female | 26,851 | 127.5 | 26,707 | 132.8 | 26,315 | 133.2 | 25,473 | 130.3 | 233,311 | 116.1 | ref | |
| Male | 66,382 | 55.1 | 67,701 | 57.6 | 68,451 | 58.8 | 68,764 | 59.1 | 600,461 | 51.0 | 0.44 | |
| Race/ethnicity | | | | | | | | | | | | |
| Black non-Hispanic | 19,540 | 75.5 | 19,928 | 81.8 | 19,407 | 83.0 | 18,790 | 82.4 | 175,484 | 67.7 | ref | |
| Asian | 4,391 | 67.8 | 4,847 | 73.9 | 4,811 | 72.6 | 4,836 | 72.3 | 38,508 | 64.5 | 0.95 | |
| White non-Hispanic | 57,315 | 64.5 | 57,280 | 66.2 | 57,795 | 67.3 | 56,802 | 65.9 | 506,212 | 58.8 | 0.87 | |
| Hispanic | 7,952 | 56.7 | 8,253 | 58.9 | 8,454 | 60.1 | 8,448 | 59.9 | 78,083 | 55.1 | 0.81 | |
| Native American | 1,163 | 50.1 | 1,254 | 52.5 | 1,276 | 52.2 | 1,209 | 49.4 | 9,738 | 48.3 | 0.71 | |
| Other | 2,872 | 73.0 | 2,846 | 75.3 | 3,023 | 80.9 | 4,152 | 111.9 | 25,747 | 69.1 | 1.02 | |
| Age group | | | | | | | | | | | | |
| <20 | 4,527 | 31.3 | 4,307 | 33.6 | 4,626 | 37.9 | 4,236 | 34.9 | 43,454 | 31.0 | ref | |
| 20-24 | 23,303 | 47.3 | 22,447 | 46.7 | 21,291 | 44.8 | 20,476 | 43.7 | 193,544 | 42.3 | 1.37 | |
| 25-29 | 19,245 | 69.1 | 20,148 | 71.6 | 20,171 | 70.7 | 20,220 | 69.2 | 172,134 | 62.3 | 2.01 | |
| 30-34 | 15,462 | 77.2 | 15,724 | 80.1 | 15,728 | 81.8 | 15,580 | 82.0 | 141,459 | 69.5 | 2.24 | |
| 35-39 | 15,032 | 87.1 | 15,173 | 92.5 | 15,743 | 97.5 | 16,132 | 99.4 | 146,456 | 81.5 | 2.63 | |
| >39 | 15,664 | 124.5 | 16,609 | 132.1 | 17,207 | 137.9 | 17,593 | 139.8 | 136,725 | 112.5 | 3.63 | |
| Military occupation | | | | | | | | | | | | |
| Combat | 11,530 | 40.6 | 12,019 | 41.7 | 12,396 | 41.0 | 12,503 | 43.6 | 102,318 | 36.3 | ref | |
| Health care | 12,691 | 108.4 | 12,928 | 114.1 | 12,868 | 115.3 | 13,004 | 118.2 | 116,328 | 100.9 | 2.78 | |
| Other | 69,012 | 68.1 | 69,461 | 71.3 | 69,502 | 73.3 | 68,730 | 71.3 | 615,126 | 62.7 | 1.73 | |

Figure 2. Average number per day of incident visits for allergic rhinitis, by calendar month, active components, U.S. Armed Forces, 1998-2007



Of interest, studies in civilian populations suggest that reports of allergic rhinitis significantly increased between 2000 and 2005.²

There are limitations to this summary that should be considered when interpreting the results. For example, for this report, incident ambulatory visits were limited to one per person per 30 consecutive days of follow-up. In turn, incident episodes of seasonal allergic rhinitis – e.g., from exposures to pollens of specific trees, grasses, weeds, etc. – are appropriately attributed to the seasons when the instigating allergens are present. However, individuals with perennial allergic rhinitis – e.g., from exposures to dust, dander, molds, etc. that may persistently contaminate indoor environments – may be overrepresented because they can count as an incident case once every 30 days of follow-up. Also, it is likely that many service members with allergic rhinitis seek care from unit medical personnel and/or treat themselves with over-the-counter medications. Because such episodes are not documented with electronic medical records, the results reported here likely underestimate the actual numbers of individuals affected and the health care costs associated with treatment. Finally, many individuals with “allergic

rhinitis” may receive non-specific symptomatic diagnoses (e.g., headache, cough, malaise) during routine medical visits – the result would be underascertainment of the number of affected individuals and associated health care costs.

Allergic rhinitis does not cause significant disability; however, its symptoms (e.g., nasal obstruction, headache, watery eyes, sneezing) and potential side effects of treatment (e.g., drowsiness, lethargy) can degrade the performance of physically and mentally demanding tasks.⁴ Thus, allergic rhinitis is militarily relevant because it affects so many service members, decreases military operational capabilities, and causes significant lost duty time and high health care costs (e.g., clinic visits, medication).

The management of allergic rhinitis in military populations should aim to minimize the frequency of attacks, the severity of symptoms during attacks, the impacts on military operational performance, and health care costs. However, the management of allergic rhinitis in military populations and settings is difficult. Service members are stationed in or deployed to many regions of the U.S. and overseas; and among susceptible individuals, exposures that can precipitate symptoms often vary across geographic locations, indoor

environments, and activities – and are unpredictable. Also, military operational considerations often limit or eliminate options for avoiding significant indoor (e.g., military billets) and outdoor (e.g., field exercises, deployment operations) allergen exposures. Still, there are many preventive interventions, countermeasures, and treatments that can reduce the very large occupational and health care burdens associated with allergic rhinitis in military members.^{1,4}

References:

1. Berger WE. Overview of allergic rhinitis. *Ann Allergy Asthma Immunol.* 2003 Jun;90(6 Suppl 3):7-12.
2. Agency for Healthcare Research and Quality. Allergic rhinitis: trends in use and expenditures, 2000 and 2005. Accessed 24 July 2008: http://www.meps.ahrq.gov/mepsweb/data_files/publications/st204/stat204.pdf
3. Armed Forces Health Surveillance Center. Absolute and relative morbidity burdens attributable to various illness and injuries, U.S. Armed Forces, 2007. *Medical Surveillance Monthly Report (MSMR)*. 2008 Apr;15(3):15-20.
4. Spector SL. Overview of comorbid associations of allergic rhinitis. *J Allergy Clin Immunol.* 1997 Feb;99(2):S773-80.

Update: Deployment Health Assessments, U.S. Armed Forces, May 2008

The health protection strategy of the U.S. Armed Forces is designed to deploy healthy, fit, and medically ready forces, to minimize illnesses and injuries during deployments, and to evaluate and treat physical and psychological problems (and deployment-related health concerns) following deployment.

In 1998, the Department of Defense initiated health assessments of all deployers prior to and after serving in major operations outside of the United States.¹ In March 2005, the Post-Deployment Health Reassessment (PDHRA) program was begun to identify and respond to health concerns that persisted for or emerged within three to six months after return from deployment.²

This report summarizes responses to selected questions on deployment health assessments completed since 2003. In addition, it documents the natures and frequencies of changes in responses from before to after deployments.

Methods:

Completed deployment health assessment forms are transmitted to the Armed Forces Health Surveillance Center (AFHSC) where they are incorporated into the Defense Medical Surveillance System (DMSS).³ In the DMSS, data recorded on health assessment forms are integrated with data that document demographic and military characteristics and medical encounters (e.g., hospitalizations, ambulatory visits) at fixed military and other (contracted care) medical facilities of the Military Health System. For this analysis, DMSS was searched to identify all pre (DD2795) and post (DD2796)

deployment health assessment forms completed since 1 January 2003 and all post-deployment health reassessment (DD2900) forms completed since 1 August 2005.

Results:

Since January 2003, 1,894,503 pre-deployment health assessment forms, 1,905,239 post-deployment health assessment forms, and 487,129 post-deployment health reassessment forms were completed at field sites, transmitted to the AFHSC, and integrated into the DMSS (Figure 1). Throughout the period, there were intervals of approximately 2-4 months between peaks of pre-deployment and post-deployment health assessments (that were completed by different cohorts of deployers) (Figure 1). Post-deployment health reassessments rapidly increased between February and May 2006 (Figure 1). Since then, numbers of reassessment forms per month have been relatively stable (reassessment forms per month, June 2007-May 2008: mean: 21,274; range: 10,558-33,392) (Figure 1, Table 1).

Between June 2007 and May 2008, nearly three-fourths (73.3%) of deployers rated their "health in general" as "excellent" or "very good" during pre-deployment health assessments (Figure 2). During the same period, only 58.1% and 51.9% of redeployers rated their general health as "excellent" or "very good" during post-deployment assessments and post-deployment reassessments, respectively (Figure 2).

From pre-deployment to post-deployment to post-deployment reassessments, there were sharp increases in the proportions of deployers who rated their health as "fair" or "poor"

Figure 1. Total deployment health assessment and reassessment forms, by month, U.S. Armed Forces, January 2003-May 2008




Table 1. Deployment-related health assessment forms, by month, U.S. Armed Forces, June 2007-May 2008

| | Pre-deployment assessment DD2795 | | Post-deployment assessment DD2796 | | Post-deployment reassessment DD2900 | |
|--------------|----------------------------------|------------|-----------------------------------|------------|-------------------------------------|------------|
| | No. | % | No. | % | No. | % |
| Total | 347,733 | 100 | 345,360 | 100 | 255,292 | 100 |
| 2007 | | | | | | |
| June | 23,784 | 6.8 | 18,605 | 5.4 | 17,512 | 6.9 |
| July | 23,812 | 6.8 | 20,404 | 5.9 | 16,745 | 6.6 |
| August | 35,124 | 10.1 | 32,729 | 9.5 | 18,743 | 7.3 |
| September | 32,679 | 9.4 | 43,105 | 12.5 | 18,744 | 7.3 |
| October | 36,635 | 10.5 | 34,936 | 10.1 | 17,107 | 6.7 |
| November | 19,374 | 5.6 | 31,004 | 9.0 | 16,825 | 6.6 |
| December | 24,519 | 7.1 | 37,523 | 10.9 | 22,432 | 8.8 |
| 2008 | | | | | | |
| January | 40,103 | 11.5 | 32,709 | 9.5 | 33,392 | 13.1 |
| February | 34,034 | 9.8 | 20,622 | 6.0 | 32,339 | 12.7 |
| March | 24,859 | 7.1 | 21,455 | 6.2 | 25,109 | 9.8 |
| April | 29,540 | 8.5 | 24,950 | 7.2 | 25,786 | 10.1 |
| May | 23,270 | 6.7 | 27,318 | 7.9 | 10,558 | 4.1 |

(Figure 2). For example, prior to deployment, approximately one of 40 (2.5%) deployers rated their health as “fair” or “poor”; however, 3-6 months after returning from deployment (during post-deployment reassessments), approximately one of seven (14.1%) respondents rated their health as “fair” or “poor” (Figure 2).

During the past 12 months, the proportion of deployers who assessed their general health as “fair” or “poor” before deploying remained consistently low (% “fair” or “poor” “health in general,” pre-deployment health assessments, June 2007-May

2008, by month: mean: 2.5% [range: 1.8-3.3%]) (Figure 3). The proportion of redeployers who assessed their general health as “fair” or “poor” around times of return from deployment was consistently and clearly higher than before deploying (% “fair” or “poor” “health in general,” post-deployment health assessments, May 2007-April 2008, by month: mean: 6.9% [range: 5.7-8.1%]) (Figure 3). Finally, the proportion of redeployers who assessed their general health as “fair” or “poor” 3-6 months after redeploying was sharply higher than at redeployment (% “fair” or “poor” “health in general,” post-deployment health reassessments, June 2007-May 2008, by month: mean: 14.1% [range: 11.3-20.3%]) (Figure 3).

More than half of service members who rated their overall health before deployment chose a different descriptor after deploying, but usually by a single category (on a five-category scale). The proportions of deployers whose self-rated health improved by more than one category from pre-deployment to reassessment remained relatively stable between June 2007 and May 2008 (mean: 1.3%, range:1.0-1.7%) (Figure 4). The proportions of service members whose self-assessed health declined by more than one category was relatively stable between May and September 2007 and has generally increased since September 2007 (mean: 16.6, range 13.6-21.9%) (Figure 4).

In general, on post-deployment assessments and reassessments, members of Reserve components and members of the Army were much more likely than their respective counterparts to report mental health-related symptoms and health and exposure-related concerns – and in turn, to have indications for medical and mental health follow-ups (“referrals”) (Table 2).

Among Reserve versus active component members, relative excesses of health-related concerns and provider-indicated

Figure 2. Percent distributions of self-assessed health status as reported on deployment health assessment forms, U.S. Armed Forces, June 2007-May 2008





Figure 3. Proportion of deployment health assessment forms with self-assessed health status as “fair” or “poor”, U.S. Armed Forces, June 2007-May 2008



referrals were much greater 3-6 months after redeployment (DD2900) than either before deploying (DD2795) or at redeployment (DD2796) (Table 2, Figures 5,6). For example, among both active and Reserve component members of all Services, mental or behavioral health referrals were more common after deployment than before (Figure 5). However, from the time of redeployment to 3-6 months later, mental health referrals sharply increased among Reserve component members of the Army, Navy, and Marine Corps (but not Air Force) (Table 2, Figure 5). Of note in this regard, the largest absolute increase in mental health referrals from redeployment to 3-6 months later was for Reserve component members of the Army (post-deployment: 4.9%; reassessment: 11.6%) (Table 2, Figure 5).

Finally, over the past three years, Reserve component members have been approximately twice as likely as active to report “exposure concerns” on post-deployment health assessments (DD2796) (“exposure concerns,” post-deployment assessments, by month, June 2007-May 2008: Reserve: mean: 27.0%, range: 22.2-32.6%; active: mean: 16.5%; range: 13.2-22.7%) (Table 2, Figures 6,7). Sharply higher proportions of both Reserve and active component members endorsed exposure concerns 3-6 months after (DD2900) compared to around times (DD2796) of redeployment (“exposure concerns,” post-deployment reassessments, by month, June 2007-May 2008: Reserve: mean: 36.2%, range: 31.0-41.7%; active: mean: 20.7%; range: 18.3-28.3%) (Figure 7).

Editorial comment:

In general, since 2003, proportions of U.S. deployers to Iraq and Afghanistan who report medical or mental health-related symptoms (or have indications for medical or mental health referrals) on deployment-related health assessments increased from pre-deployment to post-deployment to 3-6 months post-deployment, are higher among members of the Army than the other Services, and are higher among Reserve than the active component members.

Regardless of the Service or component, deployers often rate their general health worse when they return compared to before deploying. This is not surprising because deployments are inherently physically and psychologically demanding. Clearly, there are many more – and more significant – threats to the physical and mental health of service members when they are conducting or supporting combat operations away from their families in hostile environments compared to when serving at their permanent duty stations (active component) or when living in their civilian communities (Reserve component).

However, many redeployed service members rate their general health worse 3-6 months after returning from deployment compared to earlier. This finding may be less intuitively understandable. Symptoms of post-traumatic stress disorder (PTSD) may emerge or worsen within several months after a life threatening experience (such as military service in a war zone). PTSD among U.S. veterans of combat duty in Iraq has been associated with higher rates of physical health problems after redeployment.⁴ The post-deployment health reassessment at 3-6 months post-deployment is designed to detect service members with symptoms not only of PTSD but also persistent or emerging deployment-related medical and mental health problems.

Among British veterans of the Iraq war, Reservists reported more “ill health” than their active counterparts.⁵ Roles, traumatic experiences, and unit cohesion while deployed were associated with medical outcomes after redeployment; however, PTSD symptoms were more associated with problems at home (e.g., reintegration into family, work, and other aspects of civilian life) than with events in Iraq.⁵ The finding may explain, at least in part, the differences in prevalences of mental health symptoms, medical complaints, and provider-indicated mental health referrals among Reserve compared to active members — 3-6 months after returning from deployment compared to earlier.

Post-deployment health assessments may be more reliable several months after redeployment compared to earlier. Commanders, supervisors, family members, peers, and providers of health care to redeployed service members should be alert to emerging or worsening symptoms of physical and psychological problems for several months, at least, after returning from deployment.

Figure 4. Proportion of service members whose self-assessed health status improved (“better”) or declined (“worse”) (by 2 or more categories on 5-category scale) from pre-deployment to reassessment, by month, U.S. Armed Forces, June 2007-May 2008




Figure 5. Percent of deployers with mental or behavioral health referrals, by Service and component, by timing of health assessment, U.S. Armed Forces, June 2007-May 2008




Table 2. Percentage of service members who endorsed selected questions/received referrals on health assessment forms, U.S. Armed Forces, June 2007–May 2008

| | Army | | | Navy | | | Air Force | | | Marine Corps | | | All service members | | |
|---------------------------------------|-------------------|--------------------|-------------------|-------------------|--------------------|-------------------|-------------------|--------------------|-------------------|-------------------|--------------------|-------------------|---------------------|--------------------|-------------------|
| | Pre-deploy DD2795 | Post-deploy DD2796 | Reassessmt DD2900 | Pre-deploy DD2795 | Post-deploy DD2796 | Reassessmt DD2900 | Pre-deploy DD2795 | Post-deploy DD2796 | Reassessmt DD2900 | Pre-deploy DD2795 | Post-deploy DD2796 | Reassessmt DD2900 | Pre-deploy DD2795 | Post-deploy DD2796 | Reassessmt DD2900 |
| Active component | n=136,993 | n=163,974 | n=74,976 | n=17,858 | n=9,310 | n=6,144 | n=61,417 | n=49,303 | n=44,342 | n=36,993 | n=25,467 | n=35,355 | n=253,261 | n=248,054 | n=160,817 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| General health "fair" or "poor" | 4.2 | 8.2 | 17.8 | 1.6 | 4.2 | 7.6 | 0.5 | 2.2 | 4.9 | 2.0 | 3.1 | 10.8 | 2.8 | 6.3 | 12.3 |
| Health concerns, not wound or injury | 13.5 | 28.5 | 41.4 | 5.6 | 11.6 | 26.3 | 4.1 | 15.1 | 17.5 | 4.4 | 8.5 | 31.1 | 9.3 | 23.2 | 31.9 |
| Health worse now than before deployed | 0.0 | 23.1 | 28.8 | 0.0 | 9.9 | 16.5 | 0.0 | 8.4 | 10.3 | 0.0 | 11.0 | 21.1 | 0.0 | 18.4 | 21.5 |
| Exposure concerns | 0.0 | 21.3 | 26.1 | 0.0 | 12.3 | 16.7 | 0.0 | 7.1 | 14.8 | 0.0 | 7.7 | 22.7 | 0.0 | 16.7 | 21.9 |
| PTSD symptoms (2 or more) | 0.0 | 17.7 | 19.5 | 0.0 | 5.1 | 11.2 | 0.0 | 2.9 | 3.5 | 0.0 | 5.4 | 13.3 | 0.0 | 13.0 | 13.4 |
| Depression symptoms (any) | 0.0 | 33.8 | 37.3 | 0.0 | 18.5 | 29.3 | 0.0 | 9.3 | 15.2 | 0.0 | 23.8 | 34.6 | 0.0 | 27.3 | 30.3 |
| Referral indicated by provider (any) | 6.2 | 32.7 | 24.5 | 6.8 | 21.8 | 26.0 | 1.7 | 11.6 | 10.5 | 4.8 | 14.9 | 30.2 | 4.9 | 26.3 | 22.0 |
| Mental health referral indicated* | 1.0 | 8.6 | 7.7 | 1.1 | 4.0 | 6.9 | 0.4 | 1.8 | 2.8 | 0.4 | 2.9 | 7.4 | 0.8 | 6.5 | 6.3 |
| Medical visit following referral† | 96.1 | 99.4 | 98.1 | 85.6 | 89.0 | 95.8 | 76.3 | 93.7 | 95.4 | 72.9 | 67.2 | 87.8 | 91.2 | 96.0 | 94.7 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| Reserve component | n=69,531 | n=80,050 | n=73,252 | n=4,651 | n=2,965 | n=4,878 | n=16,257 | n=13,025 | n=12,984 | n=4,033 | n=1,266 | n=3,361 | n=94,472 | n=97,306 | n=94,475 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |
| General health "fair" or "poor" | 2.1 | 9.9 | 19.6 | 0.6 | 4.4 | 8.3 | 0.3 | 2.0 | 4.7 | 1.9 | 5.1 | 11.9 | 1.7 | 8.6 | 16.7 |
| Health concerns, not wound or injury | 16.2 | 40.0 | 56.9 | 3.3 | 23.1 | 38.0 | 1.9 | 22.9 | 19.2 | 4.9 | 17.6 | 44.5 | 12.6 | 36.9 | 50.3 |
| Health worse now than before deployed | 0.0 | 29.6 | 38.3 | 0.0 | 18.5 | 23.3 | 0.0 | 11.9 | 11.1 | 0.0 | 20.0 | 26.8 | 0.0 | 26.8 | 33.4 |
| Exposure concerns | 0.0 | 31.1 | 39.7 | 0.0 | 31.7 | 28.7 | 0.0 | 9.2 | 19.9 | 0.0 | 16.8 | 28.2 | 0.0 | 28.0 | 36.0 |
| PTSD symptoms (2 or more) | 0.0 | 14.3 | 24.8 | 0.0 | 4.5 | 12.9 | 0.0 | 1.9 | 3.6 | 0.0 | 4.6 | 20.0 | 0.0 | 12.2 | 21.1 |
| Depression symptoms (any) | 0.0 | 29.5 | 40.0 | 0.0 | 17.0 | 26.9 | 0.0 | 8.0 | 14.9 | 0.0 | 30.7 | 35.9 | 0.0 | 26.2 | 35.7 |
| Referral indicated by provider (any) | 8.6 | 31.2 | 38.1 | 5.1 | 19.2 | 25.4 | 0.4 | 11.5 | 11.3 | 5.3 | 33.6 | 40.0 | 6.9 | 28.3 | 33.8 |
| Mental health referral indicated* | 0.7 | 4.9 | 11.6 | 0.4 | 2.3 | 5.6 | 0.0 | 1.2 | 1.0 | 0.1 | 4.4 | 9.5 | 0.5 | 4.3 | 9.7 |
| Medical visit following referral† | 95.4 | 96.0 | 26.3 | 90.1 | 78.8 | 30.6 | 51.9 | 56.0 | 26.1 | 48.9 | 80.7 | 23.1 | 93.0 | 93.6 | 26.4 |
| | % | % | % | % | % | % | % | % | % | % | % | % | % | % | % |

*Includes behavioral health, combat stress and substance abuse referrals

†Record of inpatient or outpatient visit within 6 months after referral

References:

1. Undersecretary of Defense for Personnel and Readiness. Department of Defense Instruction (DODI) Number 6490.3. Subject: Deployment health, dated 11 August 2006. Accessed on 19 March 2007 at: <http://www.dtic.mil/whs/directives/corres/pdf/649003p.pdf>.
2. Assistant Secretary of Defense (Health Affairs). Memorandum for the Assistant Secretaries of the Army (M&RA), Navy (M&RA), and Air Force (M&RA), subject: Post-deployment health reassessment (HA policy: 05-011), dated 10 March 2005. Washington, DC. <http://www.ha.osd.mil/policies/2005/05-011.pdf>. Accessed 18 October 2006.

3. Rubertone MV, Brundage JG. The Defense Medical Surveillance System and the Department of Defense Serum Repository: Glimpses of the Future of Public Health Surveillance. *Am J Public Health* 2002 Dec;92, (12):1900-04.
4. Hoge CW, Terhakopian A, Castro CA, Messer SC, Engel CC. Association of posttraumatic stress disorder with somatic symptoms, health care visits, and absenteeism among Iraq war veterans. *Am J Psychiatry*. 2007 Jan;164(1):150-3.
5. Browne T, Hull L, Horn O, et al. Explanations for the increase in mental health problems in UK reserve forces who have served in Iraq. *Br J Psychiatry*. 2007 Jun;190:484-489.

Figure 6. Ratio of percents of deployers who endorse selected questions, Reserve versus active component, on pre-deployment health assessments (DD2795) and post-deployment health reassessments (DD2900), U.S. Armed Forces, June 2007-May 2008





Figure 7. Proportion of service members who endorse exposure concerns on post-deployment health assessments, U.S. Armed Forces, January 2004-May 2008



Sentinel reportable events for service members and beneficiaries at U.S. Army medical facilities, cumulative numbers* through May 2007 and May 2008



Army

| Reporting locations | Number of reports all events† | | Food-borne | | | | | | | | Vaccine preventable | | | | | |
|------------------------|-------------------------------|--------------|----------------|-----------|-----------|----------|------------|-----------|-----------|-----------|---------------------|----------|-------------|-----------|-----------|-----------|
| | | | Campylo-bacter | | Giardia | | Salmonella | | Shigella | | Hepatitis A | | Hepatitis B | | Varicella | |
| | | | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 |
| NORTH ATLANTIC | | | | | | | | | | | | | | | | |
| Washington, DC Area | 114 | 159 | . | 1 | 3 | 2 | 1 | 1 | . | . | . | . | 2 | 1 | 1 | 6 |
| Aberdeen, MD | 17 | 0 | . | . | 1 | . | . | . | . | . | . | . | . | . | . | . |
| FT Belvoir, VA | 105 | 125 | 6 | 2 | . | . | 1 | 1 | 1 | . | . | . | . | . | . | . |
| FT Bragg, NC | 523 | 645 | 1 | . | . | . | 7 | 7 | 2 | . | . | . | . | . | . | . |
| FT Drum, NY | 108 | 199 | . | . | . | . | . | . | . | . | . | . | 2 | . | . | . |
| FT Eustis, VA | 83 | 400 | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| FT Knox, KY | 124 | 345 | . | 2 | . | . | 1 | . | . | . | . | . | 1 | . | . | . |
| FT Lee, VA | 191 | 153 | . | . | . | . | . | . | . | . | . | . | 2 | 1 | . | 1 |
| FT Meade, MD | 23 | 190 | . | . | . | 1 | . | . | . | . | . | . | . | . | . | . |
| West Point, NY | 14 | 44 | . | . | . | . | . | . | . | . | . | . | 3 | . | . | . |
| GREAT PLAINS | | | | | | | | | | | | | | | | |
| FT Sam Houston, TX | 312 | 394 | . | . | 1 | . | 2 | 4 | . | 8 | . | . | 2 | . | 6 | . |
| FT Bliss, TX | 53 | 229 | . | . | . | . | . | 1 | . | . | . | . | . | . | . | . |
| FT Carson, CO | 328 | 412 | . | 1 | 2 | . | . | 2 | . | . | . | . | . | . | . | . |
| FT Hood, TX | 891 | 966 | 3 | 2 | 2 | . | 3 | 9 | 6 | 5 | . | . | . | . | 1 | 1 |
| FT Huachuca, AZ | 49 | 43 | . | . | . | . | 5 | 1 | . | 1 | . | . | . | . | . | . |
| FT Leavenworth, KS | 22 | 19 | 1 | . | . | . | . | . | . | . | . | . | . | . | . | . |
| FT Leonard Wood, MO | 182 | 322 | . | . | . | 2 | . | 1 | . | . | . | . | . | 1 | 5 | . |
| FT Polk, LA | 76 | 76 | . | . | 3 | . | 2 | . | . | 1 | . | . | . | . | 1 | 1 |
| FT Riley, KS | 130 | 285 | 1 | . | . | . | 3 | 1 | . | . | . | . | . | 1 | . | . |
| FT Sill, OK | 77 | 148 | . | . | . | . | 1 | . | . | . | . | . | . | . | 1 | . |
| SOUTHEAST | | | | | | | | | | | | | | | | |
| FT Gordon, GA | 314 | 507 | . | . | . | . | . | 7 | . | 9 | . | . | 1 | . | . | 1 |
| FT Benning, GA | 180 | 181 | 1 | . | 1 | . | 2 | . | . | . | . | . | 1 | . | 1 | . |
| FT Campbell, KY | 0 | 144 | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| FT Jackson, SC | 85 | 111 | . | . | . | . | . | . | . | . | . | . | 1 | 1 | . | . |
| FT Rucker, AL | 25 | 37 | . | . | . | . | . | 1 | 5 | . | . | . | . | . | . | . |
| FT Stewart, GA | 459 | 329 | 1 | 1 | . | 1 | 4 | 6 | 9 | . | . | . | 1 | 5 | 1 | . |
| WESTERN | | | | | | | | | | | | | | | | |
| FT Lewis, WA | 268 | 506 | 1 | 2 | 1 | . | 1 | . | 1 | 1 | . | . | . | . | 1 | . |
| FT Irwin, CA | 40 | 27 | 1 | . | . | . | 1 | 1 | . | 1 | . | . | . | . | . | . |
| FT Wainwright, AK | 95 | 186 | . | 3 | . | . | . | . | . | . | . | . | . | . | . | . |
| OTHER LOCATIONS | | | | | | | | | | | | | | | | |
| Hawaii | 332 | 380 | 12 | 16 | . | 1 | 8 | 7 | . | 3 | . | . | 1 | 3 | . | . |
| Germany | 327 | 639 | 4 | 7 | 1 | 2 | 2 | 2 | 1 | 1 | . | . | . | 4 | 1 | . |
| Korea | 260 | 310 | . | . | . | . | . | . | . | . | . | . | . | . | 2 | 1 |
| Total | 5,807 | 8,511 | 32 | 37 | 15 | 9 | 44 | 52 | 25 | 30 | 0 | 0 | 17 | 17 | 21 | 11 |

*Events reported by June 7, 2007 and 2008

†Seventy medical events/conditions specified by Tri-Service Reportable Events Guidelines and Case Definitions, May 2004.

Note: Completeness and timeliness of reporting vary by facility.

Sentinel reportable events for service members and beneficiaries at U.S. Army medical facilities, cumulative numbers* through May 2007 and May 2008



Army

| Reporting location | Arthropod-borne | | | | Sexually transmitted | | | | | | | | Environmental | | | |
|------------------------|-----------------|-----------|-----------|-----------|----------------------|--------------|------------|------------|-----------------------|-----------|-------------------------|-----------|---------------|-----------|-----------|-----------|
| | Lyme disease | | Malaria | | Chlamydia | | Gonorrhea | | Syphilis [‡] | | Urethritis [§] | | Cold | | Heat | |
| | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 |
| NORTH ATLANTIC | | | | | | | | | | | | | | | | |
| Washington, DC Area | 1 | 2 | . | . | 67 | 48 | 9 | 15 | 2 | 4 | . | . | . | . | . | . |
| Aberdeen, MD | . | . | . | . | 8 | . | 3 | . | . | . | . | . | . | . | . | . |
| FT Belvoir, VA | . | . | . | . | 70 | 67 | 10 | 1 | 2 | . | . | . | . | . | . | . |
| FT Bragg, NC | . | . | 1 | 3 | 387 | 424 | 67 | 82 | . | 1 | 41 | 28 | 1 | . | 15 | 8 |
| FT Drum, NY | . | 3 | . | . | 62 | 141 | 13 | 14 | . | . | . | . | . | . | . | . |
| FT Eustis, VA | . | . | . | . | 71 | 100 | 2 | 13 | . | 3 | . | . | . | . | 1 | . |
| FT Knox, KY | . | . | . | . | 101 | 69 | 17 | 18 | . | 1 | . | . | . | . | . | . |
| FT Lee, VA | 1 | 1 | . | . | 151 | 101 | 23 | 39 | . | 1 | . | . | 1 | . | . | . |
| FT Meade, MD | . | 1 | . | . | 16 | 28 | 5 | 1 | 1 | . | . | . | 1 | . | . | . |
| West Point, NY | 2 | 5 | . | . | 8 | 11 | . | . | . | . | . | . | . | . | . | . |
| GREAT PLAINS | | | | | | | | | | | | | | | | |
| FT Sam Houston, TX | 1 | . | . | . | 144 | 123 | 25 | 31 | 2 | 11 | . | . | . | 1 | 1 | . |
| FT Bliss, TX | . | . | . | . | 42 | 147 | 7 | 31 | 1 | 3 | . | . | . | . | . | . |
| FT Carson, CO | . | . | . | . | 202 | 260 | 32 | 25 | 1 | . | 5 | 12 | 1 | . | . | . |
| FT Hood, TX | . | . | . | . | 659 | 676 | 98 | 122 | 1 | . | 51 | 36 | . | . | . | . |
| FT Huachuca, AZ | . | . | . | . | 38 | 35 | 6 | 5 | . | . | . | . | . | . | . | . |
| FT Leavenworth, KS | . | . | . | . | 19 | 19 | 2 | . | . | . | . | . | . | . | . | . |
| FT Leonard Wood, MO | . | . | . | . | 119 | 83 | 20 | 11 | 1 | . | . | . | 2 | 3 | 2 | . |
| FT Polk, LA | . | . | 14 | . | 44 | 49 | 8 | 18 | 1 | . | . | . | . | . | 2 | 6 |
| FT Riley, KS | . | 2 | . | 1 | 87 | 150 | 5 | 12 | . | . | . | . | . | 1 | 2 | . |
| FT Sill, OK | . | . | . | . | 53 | 41 | 14 | 10 | . | . | . | . | 1 | . | 2 | 4 |
| SOUTHEAST | | | | | | | | | | | | | | | | |
| FT Gordon, GA | . | . | . | . | 221 | 223 | 32 | 60 | 1 | . | . | . | . | . | . | . |
| FT Benning, GA | . | . | 1 | . | 115 | 114 | 39 | 32 | . | 1 | . | . | 1 | . | 3 | 8 |
| FT Campbell, KY | . | . | . | . | . | 51 | . | 2 | . | 1 | . | . | . | . | . | . |
| FT Jackson, SC | . | . | . | . | 66 | 94 | 16 | 16 | 2 | . | . | . | . | . | . | . |
| FT Rucker, AL | . | 1 | . | . | 18 | 26 | 1 | 7 | . | 1 | . | . | . | . | 1 | . |
| FT Stewart, GA | . | . | . | 1 | 327 | 257 | 73 | 45 | 2 | 2 | . | . | . | . | 1 | . |
| WESTERN | | | | | | | | | | | | | | | | |
| FT Lewis, WA | . | . | 1 | . | 232 | 389 | 19 | 39 | . | 1 | 6 | 8 | . | . | . | . |
| FT Irwin, CA | . | . | . | . | 29 | 21 | 3 | 3 | . | . | . | . | . | . | 5 | 1 |
| FT Wainwright, AK | . | 1 | . | . | 69 | 119 | 3 | 14 | . | . | . | . | 10 | 11 | . | . |
| OTHER LOCATIONS | | | | | | | | | | | | | | | | |
| Hawaii | . | . | . | 1 | 230 | 263 | 27 | 35 | . | . | . | . | . | . | 2 | . |
| Germany | 7 | 16 | 3 | 5 | 201 | 356 | 56 | 81 | 1 | 4 | 3 | . | . | 8 | 7 | 4 |
| Korea | . | . | 1 | . | 214 | 261 | 22 | 31 | . | 4 | 1 | . | 20 | . | . | 3 |
| Total | 12 | 32 | 21 | 11 | 4,070 | 4,746 | 657 | 813 | 18 | 38 | 107 | 84 | 38 | 24 | 44 | 34 |

‡Primary and secondary.

§Urethritis, non-gonococcal (NGU).

Sentinel reportable events for service members and beneficiaries at U.S. Navy medical facilities, cumulative numbers* through May 2007 and May 2008



Navy

| Reporting location | Number of reports all events† | | Food-borne | | | | | | | | Vaccine preventable | | | | | | |
|------------------------------|-------------------------------|------------|----------------|----------|----------|----------|------------|-----------|----------|----------|---------------------|----------|-------------|----------|-----------|----------|---|
| | | | Campylo-bacter | | Giardia | | Salmonella | | Shigella | | Hepatitis A | | Hepatitis B | | Varicella | | |
| | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | |
| NATIONAL CAPITOL AREA | | | | | | | | | | | | | | | | | |
| Annapolis, MD | 0 | 10 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Bethesda, MD | 17 | 19 | 1 | . | . | 1 | . | 1 | . | . | . | . | . | 1 | . | . | . |
| Patuxent River, MD | 0 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| NAVY MEDICINE EAST | | | | | | | | | | | | | | | | | |
| Albany, GA | 0 | 3 | . | . | . | . | . | . | . | . | . | . | . | 2 | . | . | . |
| Atlanta, GA | 3 | 1 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Beaufort, SC | 100 | 4 | . | . | . | . | . | . | 1 | . | . | . | . | . | . | . | . |
| Camp Lejeune, NC | 141 | 27 | . | . | . | . | . | 1 | . | . | . | . | . | . | . | . | . |
| Cherry Point, NC | 60 | 42 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Great Lakes, IL | 134 | 10 | . | . | 1 | . | 1 | . | . | . | . | . | . | . | . | . | . |
| Jacksonville, FL | 108 | 27 | 1 | . | . | . | 2 | 4 | 3 | . | . | . | . | . | . | . | 1 |
| Mayport, FL | 23 | 21 | 1 | . | . | . | 4 | 3 | . | 1 | . | . | . | . | . | . | . |
| NABLC Norfolk, VA | 32 | 4 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| NBMC Norfolk, VA | 198 | 88 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| NEHC Norfolk, VA | 4 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | 2 | . |
| North Charleston, SC | 3 | 14 | . | . | . | . | . | 1 | . | 1 | . | . | . | . | . | . | . |
| Pensacola, FL | 42 | 21 | . | . | . | . | 2 | . | 1 | 1 | . | . | . | . | . | 3 | . |
| Portsmouth, VA | 0 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Washington, DC | 0 | 3 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Guantanamo Bay, Cuba | 1 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Europe | 16 | 9 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| NAVY MEDICINE WEST | | | | | | | | | | | | | | | | | |
| Camp Pendleton, CA | 13 | 14 | . | . | . | 1 | 1 | . | . | . | . | . | . | . | . | . | . |
| Corpus Christi, TX | 3 | 1 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Fallon, NV | 0 | 1 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Ingleside, TX | 0 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Lemoore, CA | 1 | 10 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Pearl Harbor, HI | 0 | 3 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| San Diego, CA | 268 | 22 | 1 | . | 2 | . | 2 | . | 2 | 1 | . | . | 28 | 1 | . | . | . |
| Guam | 27 | 3 | . | . | . | . | 1 | . | . | . | . | . | . | . | . | . | 2 |
| Japan | 25 | 22 | . | . | . | . | . | . | . | . | . | . | . | . | . | 1 | . |
| NAVAL SHIPS | | | | | | | | | | | | | | | | | |
| COMNAVAIRLANT/CINCLANTFLEET | 5 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| COMNAVSURFPAC/CINCPACFLEET | 20 | 2 | . | . | . | . | . | . | . | . | . | . | . | . | . | 1 | . |
| Total | 1,244 | 381 | 4 | 0 | 3 | 2 | 13 | 10 | 7 | 4 | 0 | 0 | 28 | 4 | 7 | 3 | |

*Events reported by June 7, 2007 and 2008

†Seventy medical events/conditions specified by Tri-Service Reportable Events Guidelines and Case Definitions, May 2004.

Note: Completeness and timeliness of reporting vary by facility.

Sentinel reportable events for service members and beneficiaries at U.S. Navy medical facilities, cumulative numbers* through May 2007 and May 2008



Navy

| Reporting location | Arthropod-borne | | | | Sexually transmitted | | | | | | | | Environmental | | | | |
|------------------------------|-----------------|----------|----------|----------|----------------------|------------|------------|-----------|-----------|----------|-------------|----------|---------------|----------|-----------|----------|---|
| | Lyme disease | | Malaria | | Chlamydia | | Gonorrhea | | Syphilis‡ | | Urethritis§ | | Cold | | Heat | | |
| | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | |
| NATIONAL CAPITOL AREA | | | | | | | | | | | | | | | | | |
| Annapolis, MD | . | . | . | . | . | 5 | . | . | . | . | . | . | . | . | . | . | . |
| Bethesda, MD | . | . | . | 1 | 12 | 8 | . | 1 | 1 | . | . | . | . | . | . | . | . |
| Patuxent River, MD | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| NAVY MEDICINE EAST | | | | | | | | | | | | | | | | | |
| Albany, GA | . | . | . | . | . | 1 | . | . | . | . | . | . | . | . | . | . | . |
| Atlanta, GA | . | . | . | . | 1 | 1 | 1 | . | 1 | . | . | . | . | . | . | . | . |
| Beaufort, SC | . | . | . | . | 78 | 1 | 6 | . | 2 | . | . | . | . | . | 7 | 1 | . |
| Camp Lejeune, NC | 8 | . | 1 | . | 111 | 15 | 17 | 3 | . | . | . | 5 | . | . | 4 | 2 | . |
| Cherry Point, NC | . | . | . | . | 52 | 24 | 4 | 5 | 1 | . | . | . | . | . | . | . | . |
| Great Lakes, IL | . | . | . | . | 117 | 10 | 11 | . | . | . | . | . | . | . | . | . | . |
| Jacksonville, FL | . | . | . | . | 84 | 11 | 12 | . | 2 | . | . | . | . | . | . | . | . |
| Mayport, FL | . | . | . | . | 15 | 11 | . | 2 | 1 | . | . | . | . | . | . | . | . |
| NABLC Norfolk, VA | . | . | . | . | 30 | 4 | 2 | . | . | . | . | . | . | . | . | . | . |
| NBMC Norfolk, VA | . | . | . | . | 165 | 73 | 32 | 10 | . | 1 | . | . | . | . | . | . | . |
| NEHC Norfolk, VA | . | . | . | . | 2 | . | . | . | . | . | . | . | . | . | . | . | . |
| North Charleston, SC | . | . | . | . | 3 | 7 | . | 1 | . | 1 | . | . | . | . | . | . | . |
| Pensacola, FL | . | . | . | . | 25 | 14 | 3 | . | . | . | . | . | . | . | 4 | . | . |
| Portsmouth, VA | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Washington, DC | . | 1 | . | . | . | 2 | . | . | . | . | . | . | . | . | . | . | . |
| Guantanamo Bay, Cuba | . | . | . | . | 1 | . | . | . | . | . | . | . | . | . | . | . | . |
| Europe | . | . | . | 1 | 15 | 6 | 1 | 2 | . | . | . | . | . | . | . | . | . |
| NAVY MEDICINE WEST | | | | | | | | | | | | | | | | | |
| Camp Pendleton, CA | . | . | . | . | 10 | 11 | 1 | . | 1 | . | . | . | . | . | . | . | . |
| Corpus Christi, TX | . | . | . | . | 2 | . | 1 | 1 | . | . | . | . | . | . | . | . | . |
| Fallon, NV | . | . | . | . | . | 1 | . | . | . | . | . | . | . | . | . | . | . |
| Ingleside, TX | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Lemoore, CA | . | . | . | . | . | 3 | . | . | . | . | . | . | . | . | . | . | . |
| Pearl Harbor, HI | . | . | . | . | . | 2 | . | . | . | 1 | . | . | . | . | . | . | . |
| San Diego, CA | 1 | . | . | . | 165 | 18 | 29 | 1 | 3 | 1 | . | . | . | . | . | . | . |
| Guam | . | . | . | . | 23 | 1 | 3 | . | . | . | . | . | . | . | . | . | . |
| Japan | . | . | . | . | 18 | 16 | 4 | 3 | . | . | . | . | . | . | 1 | . | . |
| NAVAL SHIPS | | | | | | | | | | | | | | | | | |
| COMNAVAIRLANT/CINCLANTFLEET | . | . | . | . | 5 | . | . | . | . | . | . | . | . | . | . | . | . |
| COMNAVSURFPAC/CINCPACFLEET | . | . | . | . | 14 | 2 | 5 | . | . | . | . | . | . | . | . | . | . |
| Total | 9 | 1 | 1 | 2 | 948 | 247 | 132 | 29 | 12 | 4 | 0 | 5 | 0 | 0 | 16 | 3 | |

‡Primary and secondary.

§Urethritis, non-gonococcal (NGU).

Sentinel reportable events for service members and beneficiaries at U.S. Air Force medical facilities, cumulative numbers* through May 2007 and May 2008



Air Force

| Reporting location | Number of reports all events† | | Food-borne | | | | | | | | Vaccine preventable | | | | | |
|-------------------------------|-------------------------------|--------------|---------------|----------|----------|----------|------------|----------|----------|----------|---------------------|----------|-------------|-----------|-----------|-----------|
| | | | Campylobacter | | Giardia | | Salmonella | | Shigella | | Hepatitis A | | Hepatitis B | | Varicella | |
| | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 |
| Air Combat Cmd | 753 | 831 | 1 | 2 | 1 | 2 | . | 1 | . | 3 | . | . | 4 | 16 | 6 | 3 |
| Air Education & Training Cmd | 277 | 409 | . | . | . | 2 | 3 | 1 | . | . | . | . | 3 | 1 | 4 | 3 |
| Lackland, TX | 0 | 0 | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| USAF Academy, CO | 22 | 9 | . | . | . | . | 2 | . | . | . | . | . | . | . | . | . |
| Air Force Dist. of Washington | 9 | 8 | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| Air Force Materiel Cmd | 213 | 324 | . | 1 | . | . | 4 | 2 | . | 2 | . | . | . | . | 1 | . |
| Air Force Special Ops Cmd | 59 | 103 | . | . | . | . | . | . | 1 | . | . | . | . | 2 | . | . |
| Air Force Space Cmd | 143 | 214 | 1 | . | 2 | . | 5 | 2 | . | . | . | . | 2 | 1 | 1 | 1 |
| Air Mobility Cmd | 331 | 486 | . | 1 | . | 2 | 1 | 2 | 2 | . | . | . | 3 | 2 | 2 | 4 |
| Pacific Air Forces | 228 | 219 | . | 4 | 1 | 2 | 3 | 1 | . | . | . | . | 2 | 4 | 8 | 2 |
| PACAF Korea | 77 | 95 | . | . | . | . | . | . | . | . | . | . | 4 | . | 1 | . |
| U.S. Air Forces in Europe | 141 | 191 | 2 | 1 | . | . | . | . | 1 | . | . | . | . | . | . | 1 |
| Total | 2,253 | 2,889 | 4 | 9 | 4 | 8 | 18 | 9 | 4 | 5 | 0 | 0 | 18 | 26 | 23 | 14 |

*Events reported by June 7, 2007 and 2008

†Seventy medical events/conditions specified by Tri-Service Reportable Events Guidelines and Case Definitions, May 2004.


Note: Completeness and timeliness of reporting vary by facility.

| Reporting location | Arthropod-borne | | | | Sexually transmitted | | | | | | | | Environmental | | | |
|-------------------------------|-----------------|-----------|----------|----------|----------------------|--------------|------------|------------|-----------|----------|-------------|----------|---------------|----------|----------|----------|
| | Lyme disease | | Malaria | | Chlamydia | | Gonorrhea | | Syphilis‡ | | Urethritis§ | | Cold | | Heat | |
| | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 | 2007 | 2008 |
| Air Combat Cmd | 2 | 3 | . | . | 409 | 414 | 33 | 38 | . | . | . | 1 | . | 3 | 2 | . |
| Air Education & Training Cmd | . | 2 | . | . | 220 | 192 | 19 | 15 | . | . | . | . | . | . | . | . |
| Lackland, TX | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . | . |
| USAF Academy, CO | . | . | . | . | 18 | 9 | . | . | . | . | . | . | . | . | . | . |
| Air Force Dist. of Washington | . | . | . | . | 9 | 3 | . | . | . | . | . | . | . | . | . | . |
| Air Force Materiel Cmd | . | 2 | . | 1 | 176 | 174 | 20 | 14 | . | 2 | . | . | . | . | . | . |
| Air Force Special Ops Cmd | . | 1 | . | 1 | 52 | 75 | 6 | 16 | . | . | . | . | . | . | . | . |
| Air Force Space Cmd | 1 | 1 | . | . | 116 | 132 | 10 | 6 | . | . | . | . | . | . | . | . |
| Air Mobility Cmd | 3 | 3 | . | . | 280 | 303 | 17 | 29 | 2 | 1 | . | . | 2 | 2 | 2 | 2 |
| Pacific Air Forces | . | . | . | . | 185 | 174 | 8 | 8 | . | 1 | . | . | 1 | . | . | . |
| PACAF Korea | . | . | . | . | 57 | 70 | 1 | 3 | 2 | . | . | . | 2 | . | . | . |
| U.S. Air Forces in Europe | 2 | . | . | 1 | 99 | 146 | 9 | 13 | . | . | . | . | . | . | . | . |
| Total | 8 | 12 | 0 | 3 | 1,621 | 1,692 | 123 | 142 | 4 | 4 | 0 | 1 | 3 | 5 | 4 | 2 |

‡Primary and secondary.

§Urethritis, non-gonococcal (NGU).


Acute respiratory disease (ARD) and streptococcal pharyngitis rates (SASI*), basic combat training centers, U.S. Army, by week, June 2006-June 2008




*Streptococcal-ARD surveillance index (SASI) = ARD rate x % positive culture for group A streptococcus
 ARD rate = cases per 100 trainees per week
 ARD rate \geq 1.5 or SASI \geq 25.0 for 2 consecutive weeks are surveillance indicators of epidemics

Deployment-related conditions of special surveillance interest, U.S. Armed Forces, by month and service, January 2003-May 2008

Traumatic brain injury, hospitalizations (ICD-9: 800-804, 850-854, 959.01)*



Traumatic brain injury, multiple ambulatory visits (without hospitalization), (ICD-9: 800-804, 850-854, 959.01)†




Reference: Army Medical Surveillance Activity. Traumatic brain injury among members of active components, U.S. Armed Forces, 2002-2007. *MSMR*. Aug 2007; 14(5):2-6.

*Indicator diagnosis (one per individual) during a hospitalization while deployed to/within 30 days of returning from OEF/OIF.

†Two or more ambulatory visits at least 7 days apart while deployed to/within 365 days of returning from OEF/OIF.

Deployment-related conditions of special surveillance interest, U.S. Armed Forces, by month and service, January 2003-May 2008


Amputations (ICD-9: 887, 896, 897, V49.6 to V49.7, PR 84.0 to PR 84.1)*



Reference: Army Medical Surveillance Activity. Deployment-related condition of special surveillance interest: amputations. Amputations of lower and upper extremities, U.S. Armed Forces, 1990-2004. *MSMR*. Jan 2005;11(1):2-6.

*Indicator diagnosis (one per individual) during a hospitalization or ambulatory visit while deployed to/within 365 days of returning from OEF/OIF.

Heterotopic ossification (ICD-9: 728.12, 728.13, 728.19)†




Reference: Army Medical Surveillance Activity. Heterotopic ossification, active components, U.S. Armed Forces, 2002-2007. *MSMR*. Aug 2007; 14(5):7-9.

†One diagnosis during a hospitalization or two or more ambulatory visits at least 7 days apart while deployed to/within 365 days of returning from OEF/OIF.

Deployment-related conditions of special surveillance interest, U.S. Armed Forces, by month and service, January 2003-May 2008


Deep vein thrombophlebitis/pulmonary embolus (ICD-9: 415.1, 451.1, 451.81, 451.83, 451.89, 453.2, 453.40 to 453.42 and 453.8)*



Reference: Isenbarger DW, Atwood JE, Scott PT, et al. Venous thromboembolism among United States soldiers deployed to Southwest Asia. *Thromb Res.*2006;117(4):379-83.

*Indicator diagnosis (one per individual) during a hospitalization while deployed to/within 90 days of returning from OEF/OIF.

Severe acute pneumonia (ICD-9: 518.81, 518.82, 518.3, 480-487, 786.09)†




Reference: Army Medical Surveillance Activity. Deployment-related condition of special surveillance interest: severe acute pneumonia. Hospitalizations for acute respiratory failure (ARF)/acute respiratory distress syndrome (ARDS) among participants in Operation Enduring Freedom/Operation Iraqi Freedom, active components, U.S. Armed Forces, January 2003-November 2004. *MSMR.* Nov/Dec 2004;10(6):6-7.

†Indicator diagnosis (one per individual) during a hospitalization or ambulatory visit while deployed to/within 30 days of returning from OEF/OIF.

Deployment-related conditions of special surveillance interest, U.S. Armed Forces, by month and service, January 2003-May 2008

Leishmaniasis (ICD-9: 085.0 to 085.9)*



Reference: Army Medical Surveillance Activity. Deployment-related condition of special surveillance interest: leishmaniasis. Leishmaniasis among U.S. Armed Forces, January 2003-November 2004. *MSMR*. Nov/Dec 2004;10(6):2-4.

*Indicator diagnosis (one per individual) during a hospitalization, ambulatory visit, and/or from a notifiable medical event during/after service in OEF/OIF.

Commander
U.S. Army Center for Health Promotion
and Preventive Medicine
ATTN: MCHB-TS-EDM
5158 Blackhawk Road
Aberdeen Proving Ground, MD 21010-5422

STANDARD
U.S. POSTAGE
PAID
APG, MD
PERMIT NO. 1

OFFICIAL BUSINESS

Executive Editor

COL Robert F. DeFraitas, MD, MPH (USA)

Senior Editors

Mark V. Rubertone, MD, MPH
LTC Steven Tobler, MD, MPH (USA)

Editor

John F. Brundage, MD, MPH

Technical Writer-Editor

Ellen Wertheimer, MHS

Webmaster/Graphic Artist

Rick McNerney

Service Liaison

Lt Col Sean I. Moore, MD, MS (USAF)

Lead Analyst

Toan Le, ScD

The *Medical Surveillance Monthly Report* (MSMR) is prepared by the Armed Forces Health Surveillance Center (AFHSC), US Army Center for Health Promotion and Preventive Medicine (USACHPPM).

Data in the MSMR are provisional, based on reports and other sources of data available to AFHSC.

Inquiries regarding content or material to be considered for publication should be directed to: Editor, Armed Forces Health Surveillance Center, 2900 Linden Lane, Suite 200 (Attn: MCHB-TS-EDM), Silver Spring, MD 20910. E-mail: mshr.afhsc@amedd.army.mil

To be added to the mailing list, contact the Armed Forces Health Surveillance Center at (301) 319-3240. E-mail: mshr.afhsc@amedd.army.mil

Views and opinions expressed are not necessarily those
of the Department of Defense.