



*Negative First-Term Outcomes Associated
with Lower Extremity Injury
during Recruit Training among
Female Marine Corps Graduates*

*D. W. Trone
A. Villasenor
C. A. Macera*



Naval Health Research Center

Report No. 05-19

Approved for Public Release; Distribution Unlimited.

*Naval Health Research Center
140 Sylvester Road
San Diego, California 92106*

Negative First-Term Outcomes Associated with Lower Extremity Injury during Recruit Training among Female Marine Corps Graduates

Guarantor: Daniel W. Trone, MA

Contributors: Daniel W. Trone, MA; Adriana Villaseñor, MPH; Caroline A. Macera, PhD

This study assessed the impact of lower extremity injuries and stress fractures during recruit training on first-term outcomes among female Marine Corps graduates. Injury data were collected from women recruits at Parris Island, South Carolina (1995–1999) and negative first-term outcomes were obtained from the Career History Archival Medical and Personnel System. The three negative outcomes included (1) failure to complete first-term of service, (2) failure to achieve rank of corporal, and (3) failure to reenlist. Overall, 22% did not complete their first-term enlistment and 12% of those who did were not promoted to corporal. After adjustment for demographic characteristics, not completing first term and not being promoted to corporal were both associated with injuries or stress fracture during training. Reenlistment was not associated with training injuries. Our findings indicate lower extremity injuries among women undergoing Marine Corps recruit training are associated with poor first-term outcomes even among those who graduate.

Introduction

Lower extremity musculoskeletal injuries are a significant problem in military populations and have a major impact on operational readiness. In 1999, musculoskeletal conditions, including stress fractures, were responsible for at least 14,000 hospital admissions and almost 4.3 million outpatient visits among active duty personnel in the four U.S. Armed Services.¹ These conditions also account for limited duty rates of 40 to 120 days per 100 soldiers per month.² In training populations, the incidence of both musculoskeletal injury and stress fracture is higher among women than among men and accounts for considerable attrition.³ For male Marine Corps recruit training graduates, the occurrence of a training injury, particularly a stress fracture, is associated with poor long-term military outcomes,^{1,3} but little is known about female Marine Corps graduates. This research augments a Center for Naval Analyses report⁴ by including injuries sustained during recruit training as a possible determinant of first-term success. There are no articles published in scientific or medical literature on the associ-

ation between lower extremity injury sustained during recruit training among female Marine Corps recruit graduates and first-term outcomes.

The Chief of Naval Operations' Top Five Readiness Priorities for the Navy⁵ emphasize efforts to reduce first-term enlistment attrition. First-term attrition and low first-term reenlistment directly affect manpower, and current and future readiness. First-term attrition and injury prevention are critical to military operational readiness. The General Accounting Office estimated that in fiscal year (FY) 1996 alone, the services lost \$390 million due to enlistees who separated before they completed 6 months of service.⁶

This study was designed to examine the impact of musculoskeletal injuries and stress fractures during recruit training on postrecruit training attrition in female Marines, using multiple career outcomes including failure to complete the first-term enlistment, failure to achieve the rank of corporal during the first term, and failure to reenlist after completion of the first term.

Subjects

Subjects were women who attended the U.S. Marine Corps Recruit Depot (MCRD) at Parris Island in Beaufort, South Carolina. The Fourth Battalion at MCRD Parris Island is the only entry-level training site for Marine Corps female recruits; therefore, this sample has the potential to represent the entire population at risk from a single location. After providing informed consent, recruits were enrolled into the study and completed a questionnaire. These volunteers were obtained in two cohorts, the first between June 1995 and September 1996 and the second between June and September 1999. Since data were collected for more than a year, the investigators believe the seasonal variations in recruit accessions are limited.⁷

Methods

Subjects, Recruitment, and Follow-Up

The study design was historical cohort. This design uses an exposure, illness, or injury, about a population, as it was at some time in the past and determines the subsequent status of the population with regard to the outcome of interest. In 1999, the Naval Health Research Center, San Diego, California, completed a study of 3,786 female Marine Corps recruits which formed the historical cohort for the current study. One thousand seventy-one female recruits who did not complete basic training (28%) were not included in the prospective analysis, resulting in a sample size of 2,715 female Marines. The institu-

Behavioral Science and Epidemiology Program, Musculoskeletal Injury Epidemiology, Naval Health Research Center, PO Box 85122, San Diego, CA 92186-5122.

Preliminary findings were presented in a Poster Session at the American College of Sports Medicine Annual Meeting, June 1–4, 2005, Nashville, TN.

The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of the Army, Department of Defense, or the U.S. government. Approved for public release; distribution is unlimited. This research has been conducted in compliance with all applicable federal regulations governing the protection of human subjects in research (Protocol NHRC.2002.0003).

This manuscript was received for review in October 2005. The revised manuscript was accepted for publication in May 2006.

tional review board approved the original studies before they were conducted and, at a later date, approved the prospective follow-up of these recruits.

The historical recruits had been followed through the 12 weeks of training or to graduation to ascertain the incidence of stress fracture and other musculoskeletal injury, discovered by self-referral during sick call to the branch medical clinic or battalion aid station. Recruits who reported symptoms of stress fracture or other musculoskeletal injury were examined and diagnosed by a medical doctor; some recruits entered the medical rehabilitation platoon (MRP) for injury rehabilitation, healed, and were reassigned to a new training platoon, eventually graduating longer than 12 weeks on board.

Outcome Data

The Career History Archival Medical and Personnel System (CHAMPS) was used to obtain information on first-term outcomes.⁸ Using information codes from the Defense Enrollment Eligibility Reporting System, CHAMPS is able to provide a detailed chronological record of changes in Military Occupational Specialty (MOS), duty station assignments, and medical and career events.⁸ First-term was defined as the period after MCRD graduation, and included School of Infantry training and any specialized MOS training before matriculating into the Fleet Marine Force, up to 48 months of service. Failure to complete the first-term enlistment was defined as less than 48 months of completed service in the Marine Corps. Two additional negative first-term outcomes were assessed: failure to achieve the rank of corporal within the first-term enlistment period and failure to reenlist (defined as less than 52 months of active service in the Marine Corps) among those who completed their first-term enlistment.

Injury Data

For analysis purposes, two injury measures were created: "Any musculoskeletal injury" included one or more lower extremity musculoskeletal injury; "Stress fracture" included one or more stress fracture at any body site. The recruits were followed through training to ascertain musculoskeletal injury and stress fracture incidence, discovered by self-referral during sick call to the branch medical clinic or battalion aid station. Recruits who reported symptoms of a musculoskeletal injury or stress fracture were examined and diagnosed by a medical doctor. Stress fracture case diagnoses were based on (1) the clinical presentation of localized pain of insidious onset, without prior acute trauma, aggravated by repetitive weight-bearing activities and relieved with rest; and (2) a confirmatory (+) radiograph or bone scan or both at a site consistent with the clinical presentation. Cases were coded according to the criteria of the International Classification of Diseases (Ninth Revision, Clinical Modification).

Demographic Data

From the survey questionnaire, self-reported age, weight, height, and race/ethnicity were obtained from the trainees. A measure of body mass (BMI) was calculated as weight (in kilograms) divided by the square of height (in meters).

All Marine Corps recruits must have the equivalent of a high school education. In addition, an aptitude test is administered

upon entry into the service. The Armed Forces Qualification Test (AFQT) is designed to measure the potential for military success and to screen out individuals who are not likely to complete the initial training program.^{4,6,9,10} Furthermore, the higher the AFQT score, the more opportunities for advancement there are within the military.⁴ The AFQT score is based on raw scores for 4 out of 10 sections of the Armed Services Vocational Aptitude Battery: word knowledge, paragraph comprehension, arithmetic reasoning, and mathematics knowledge.¹⁰ The raw scores are combined, weighted, and converted to percentiles and categorized into six categories: The highest category (I) consists of those in the 93rd to 99th percentile, category II (65–92%), category IIIA (50–64%), category IIIB (31–49%), category IV (10–30%) and category V (1–9%). No recruit can be in the lowest 10th percentile (category V). Since only a small proportion (6%) of recruits can be within the 10th to 30th percentiles, we combined categories IIIB and IV; therefore, four AFQT groups were used in the analysis.

Statistical Analysis

Descriptive statistics were used to characterize participating recruits. Descriptive data included means, medians, percentages, and ranges. Initially, univariate logistic models were used to identify statistically significant differences between each injury exposure (lower extremity injury and/or stress fracture) and the demographic variables for each of the three attrition outcomes. Adjusted logistic regression models were developed to identify statistically significant differences between injury exposure sustained during recruit training (either any lower extremity overuse injury or stress fracture) and each of the three attrition outcomes while adjusting for all of the demographic variables. In all cases, statistical significance was determined by a $p < 0.05$ or confidence intervals (CIs) that did not include 1.0.

Results

Recruit Characteristics

A total of 2,715 women from the historical sample graduated from Marine Corps basic training. The final sample size was 2,420 women, after omitting subjects with missing follow-up information. The prospective sample included 1,071 women (44%) who incurred a lower extremity injury during training and 87 women (3.6%) who had a confirmed stress fracture during training (Table I). The majority of women were non-Hispanic Caucasian (62%), followed by non-Hispanic African American (18%), and Hispanic (14%). The "non-Hispanic, other" category (6%) consisted of Asian and Pacific Islander and Native American/Native Alaskan. The women had completed a median of 12 years of education, and 96.6% of the women fell within AFQT categories II, IIIA, or IIIB (a score range of 31–92%). (The DoD has established guidelines for recruiting, accession attrition, retention, and reenlistment.^{11,12} Recruiters rely on AFQT scores as a primary measure of recruit potential. The distribution of AFQT scores for our study sample (Table I) are comparable to the AFQT scores for fiscal year 1998 non-prior service accessions for Marine Corps women,⁹ indicating that this study sample is a strong representation of the Marine Corps women,⁹ indicating that this study sample is a strong representation of the Marine Corps-wide female recruit population: category I,

TABLE I

SELECTED CHARACTERISTICS OF 2,240 FEMALE MARINE CORPS GRADUATES, PARRIS ISLAND, 1995-1996, 1999

Characteristic	Mean (SD) or %	n
Lower extremity injury during training	44.3	1,071
Lower extremity stress fracture during training	3.6	87
Race/ethnicity		
Caucasian, non-Hispanic	61.7	1,494
African American, non-Hispanic	18.3	444
Hispanic	14.4	349
Other, non-Hispanic ^a	5.5	133
AFQT categories (percentiles)		
I (93-99)	3.3	79
II (65-92)	37.2	901
IIIA (50-64)	31.7	767
IIIB (31-49)	27.7	671
IV (10-30)	0.1	2
Age (years) entering basic training	19.0 (1.96)	
Weight (pounds) entering basic training	127.7 (14.58)	
Height (inches) entering basic training	64.4 (2.56)	
Calculated BMI (kg/m ²)	21.6 (1.92)	

^a Includes Asian/Native American/Native Alaskan/Native Hawaiian/Pacific Islander.

3.3% vs. 4.7%; category II, 37.2% vs. 33.2%; category IIIA, 31.7% vs. 30.1%; and category IIIB, 27.7% vs. 31.8% for our sample compared with the DoD population report, respectively.⁹) Study participants ranged in age from 17 to 32 years, with an average age of 19 years. At entry to basic training, these women had a mean height of 64.4 inches, mean weight of 127.7 pounds, and mean BMI of 21.6 kg/m².

First-Term Outcome Measures

Overall, 22% of the 2,420 women who graduated from basic training were discharged before the end of their first-term service. Among the 1,879 women who completed their first term, 228 (12%) were not promoted to the rank of corporal during the first term and about half (51%) did not reenlist (Table II).

TABLE II

FIRST-TERM OUTCOMES FOR FEMALE MARINE CORPS GRADUATES, PARRIS ISLAND, 1995-1996, 1999

Outcomes	n	%	Total Sample
Failure to complete first term ^a	541	22.4	2,420
Not promoted to Corporal rank within first term	228	12.1	1,879
No reenlistment beyond first term ^b	949	50.5	1,879

^a Defined as less than 48 months of completed service in the Marine Corps according to CHAMPS.

^b Defined as less than 52 months of active service in the Marine Corps according to CHAMPS.

Failure to Complete First Term

As shown in Table III, after adjusting for race/ethnicity, AFQT categories, age, and BMI, women who had a lower extremity injury during training were less likely to complete their first-term enlistment than women who did not have an injury (adjusted odds ratio (AOR) = 1.7; 95% CI, 1.4-2.1). All race/ethnicity groups were more likely than non-Hispanic Caucasians to complete their first-term enlistment. After adjusting for race/ethnicity, AFQT categories, age, and BMI, women who had a stress fracture during training were less likely than those who did not to complete their first-term enlistment (AOR = 2.4; 95% CI, 1.5-3.8).

Failure to be Promoted to Corporal

After adjusting for race/ethnicity, AFQT categories, age, and BMI (Table IV), women who had a lower extremity injury or a stress fracture during recruit training were less likely to be promoted to corporal during their first-term enlistment than women who did not have an injury (AOR = 1.6; 95% CI, 1.3-2.1; and AOR = 2.1; 95% CI, 1.1-3.9, respectively). In both injury models, Hispanics were more likely than non-Hispanic Caucasians to be promoted to corporal. In addition, women scoring between 10 and 64% on the AFQT (categories IIIA and B) were less likely to be promoted when compared with the highest category (category I). Age had no bearing on the relationship, while BMI was associated with failing to be promoted (as BMI increased, the likelihood of nonpromotion also increased).

Failure to Reenlist

The AORs shown in Table V showed no association between either sustaining a lower extremity injury or stress fracture during recruit training and subsequent reenlistment (Table V). However, women scoring below 65% on the AFQT (categories IIIA, IIIB, and IV) were less likely to reenlist than women scoring 93% or higher (AOR = 2.7; 95% CI, 1.5-4.9 for category IIIA and AOR = 2.9; 95% CI, 1.6-5.2 for categories IIIB and IV).

Figure 1 summarizes the findings for all three attrition outcomes and lower extremity injuries and stress fractures during training. The strongest relation for both lower extremity injury and stress fracture is seen for failure to complete the first term enlistment (AOR = 1.7 and 2.4, respectively). A moderate but still statistically significant association for lower extremity injury and stress fracture is found for not being promoted to corporal during the first term (AOR = 1.6 and 2.1, respectively). The relation for both lower extremity injury and stress fracture and failure to reenlist is no different than the reference group (AOR = 1.1 and 1.2, respectively).

Discussion

Stress fracture rates during Marine Corps recruit training are well documented.¹³⁻¹⁶ The stress fracture incidence among Marine Corps women during recruit training has been reported to be 5.7%¹³; in this study, the stress fracture incidence during training was 5.1% (95% CI, 4.4%-5.8%). Medical treatment can also affect recruit training graduation rates. Stress fracture standard of care was followed but taken on a case-by-case basis; metatarsal stress fractures usually heal in 2 to 4 weeks and

TABLE III

AORS FOR FAILURE TO COMPLETE FIRST-TERM ENLISTMENT AND LOWER EXTREMITY INJURY OR STRESS FRACTURE DURING TRAINING, 2,420 FEMALE MARINE CORPS GRADUATES, PARRIS ISLAND, 1995-1996 AND 1999

	Lower Extremity Injury		Stress Fracture	
	AOR ^a	95% CI	AOR ^a	95% CI
Lower extremity injury during training				
No	1.00 ^b		1.00 ^a	
Yes	1.71	(1.41, 2.08)	2.40	(1.53, 3.75)
Race/ethnicity				
Caucasian, non-Hispanic	1.00 ^b		1.00 ^b	
African American, non-Hispanic	0.73	(0.56, 0.95)	0.76	(0.58, 0.99)
Hispanic	0.51	(0.37, 0.70)	0.49	(0.36, 0.70)
Other, non-Hispanic ^c	0.52	(0.32, 0.85)	0.53	(0.33, 0.88)
AFQT categories (percentiles)				
I (93-99)	1.00 ^b		1.00 ^b	
II (65-92)	0.66	(0.39, 1.12)	0.71	(0.42, 1.19)
IIIA (50-64)	0.85	(0.50, 1.44)	0.92	(0.54, 1.56)
IIIB and IV (10-49)	0.85	(0.50, 1.45)	0.90	(0.53, 1.54)
Age (years) entering basic training	1.02	(0.97, 1.07)	1.01	(0.96, 1.06)
Calculated BMI (kg/m ²)	1.04	(0.99, 1.10)	1.04	(0.99, 1.10)

^a Adjusted for all other variables in Table III.

^b Indicates reference category.

^c Includes Asian/Native American/Native Alaskan/Native Hawaiian/Pacific Islander.

femoral shaft or femoral neck stress fractures typically heal in 4 months.¹⁷ A 1-year follow-up of the long-term impact of stress fractures sustained in men undergoing recruit training describes the impact of stress fractures after graduation.¹⁸ In that analysis, 13.6% of male recruits who sustained a stress fracture had a protracted recovery period, experiencing pain at the stress fracture site associated with prolonged activity months after returning to duty,¹⁸ suggesting that some recruits who sustain a stress fracture during recruit training do not fully recover before returning to duty. That study also found that 10.6% had a recurrent stress fracture and only 47% had an uneventful recovery and were asymptomatic after the first period of recovery.

Furthermore, recruits who did not sustain a stress fracture during basic training were at a low risk (1.7%) of stress fracture during the 1-year follow-up.¹⁸

The Naval Health Research Center completed a first-term enlistment follow-up of male Marine Corps recruits using a historical cohort design similar to this one. Among a historical cohort of 1,131 male recruits from MCRD San Diego studied in 1993, those who suffered from a stress fracture during recruit training and still graduated were over two times more likely to be discharged before the end of their first-term enlistment for any reason, and more than six times more likely to be discharged before the end of their first-term enlistment due to a physical

TABLE IV

AORS FOR PERSONNEL NOT PROMOTED TO CORPORAL WITHIN THE FIRST-TERM ENLISTMENT AND LOWER EXTREMITY INJURY AND STRESS FRACTURE DURING TRAINING, 1,879 FEMALE MARINE CORPS GRADUATES, PARRIS ISLAND, 1995-1996, 1999

	Lower Extremity Injury		Stress Fracture	
	AOR ^a	95% CI	AOR ^a	95% CI
Lower extremity injury during training				
No	1.00 ^b		1.00 ^b	
Yes	1.64	1.28, 2.10	2.06	1.09, 3.90
Race/ethnicity				
Caucasian, non-Hispanic	1.00 ^b		1.00 ^b	
African American, non-Hispanic	0.90	0.65, 1.24	0.93	0.68, 1.28
Hispanic	0.52	0.35, 0.78	0.51	0.34, 0.76
Other, non-Hispanic ^c	0.87	0.52, 1.46	0.89	0.53, 1.49
AFQT categories (percentiles)				
I (93-99)	1.00 ^b		1.00 ^b	
II (65-92)	2.64	0.81, 8.66	2.74	0.84, 8.96
IIIA (50-64)	4.27	1.30, 14.03	4.55	1.39, 14.92
IIIB and IV (10-49)	5.14	1.56, 16.93	5.33	1.62, 17.51
Age (years) entering basic training	0.98	0.92, 1.05	0.97	0.91, 1.04
Calculated BMI (kg/m ²)	1.09	1.02, 1.16	1.08	1.02, 1.16

^a Adjusted for all other variables in Table IV.

^b Indicates reference category.

^c Includes Asian/Native American/Native Alaskan/Native Hawaiian/Pacific Islander.

TABLE V

AORS FOR FAILURE TO REENLIST AND LOWER EXTREMITY INJURY OR STRESS FRACTURE DURING TRAINING, 1,879 FEMALE MARINE CORPS GRADUATES, PARRIS ISLAND, 1995-1996, 1999

	Lower Extremity Injury		Stress Fracture	
	AOR ^a	95% CI	AOR ^a	95% CI
Lower extremity injury during training				
No	1.00 ^b		1.00 ^b	
Yes	1.14	(0.94, 1.37)	1.20	(0.69, 2.09)
Race/ethnicity				
Caucasian, non-Hispanic	1.00 ^b		1.00 ^b	
African American, non-Hispanic	0.70	(0.54, 0.90)	0.70	(0.55, 0.90)
Hispanic	0.89	(0.69, 1.16)	0.89	(0.68, 1.15)
Other, non-Hispanic ^c	0.88	(0.60, 1.31)	0.89	(0.60, 1.32)
AFQT (percentiles)				
I (93-99)	1.00 ^b		1.00 ^b	
II (65-92)	1.76	(0.98, 3.14)	1.78	(0.99, 3.18)
IIIA (50-64)	2.71	(1.49, 4.87)	2.76	(1.53, 4.96)
IIIB and IV (10-49)	2.87	(1.58, 5.20)	2.91	(1.60, 5.27)
Age (years) entering basic training	0.97	(0.92, 1.01)	0.96	(0.92, 1.01)
Calculated BMI (kg/m ²)	1.03	(0.98, 1.08)	1.03	(0.64, 2.09)

^a Adjusted for all other variables in Table V.^b Reference category.^c Includes Asian/Native American/Native Alaskan/Native Hawaiian/Pacific Islander.

disability.¹⁹ However, these results were not published in the scientific or medical literature and the ORs were not adjusted for demographics and other covariates.

In the last 10 years, recruits who suffered from a stress fracture at MCRD San Diego (males only) or Parris Island (males and females), could have been sent to MRP, healed, then immediately discharged, or rolled back into training. There is no command policy or medical policy regarding stress fractures and retention, and no sex-specific practices. Current medical practice at MCRD Parris Island is to recommend separation for

any recruit who experiences two or more stress fractures during recruit training. As a general rule the medical officer will separate recruits with severe stress fractures (pelvis, femoral neck, bilateral tibia) and sacral stress fractures (mostly women) because they take a long time to heal (T. May, unpublished observation).

Current medical practice at MCRD San Diego is to treat all stress fracture recruits on an individual basis. In general, if a recruit is in the MRP for any injury for an extended period of time, he is +less likely to resume training. Multiple metatarsal stress fractures (three or more), bilateral tibia stress fractures, or one femoral neck stress fracture usually means separation (T. May, unpublished observation).

Once a recruit graduates from basic training, efforts to retain qualified personnel focus on job satisfaction, health care, housing, and quality-of-life issues.^{12,20,21} The Marine Corps career force requires 25% of the eligible first-term Marines to reenlist.¹² Marines have to be eligible for reenlistment and must request reenlistment within their expiration-of-active service time frame. First-term reenlistment requirements indicate that an eligible candidate must be corporal or higher, have a first class physical fitness test, no history or assignment to a weight-control program, no nonjudicial punishment, court-martial, or civilian convictions, and there must be "boat space" for their MOS,²² which means that fast-filling MOSs could require command approval for a lateral move to a new MOS to reenlist. In contrast, a qualified Marine in a highly technical MOS may be eligible for a Selective Reenlistment Bonus. The Marine Corps can offer incentives to high-quality personnel to retain, educate, and train the first-term Marine based on career potential. Unplanned attrition in the MOS Corps-wide means the Marine has to be replaced.²²

Musculoskeletal injuries are a significant problem in military recruit populations where unfit individuals are exposed to sudden increases in the volume and intensity of physical activity. Musculoskeletal injuries during military training are one of the




Fig. 1. AORs for first-term outcome indicators and lower extremity injury or stress fracture during training for 2,240 female Marine Corps graduates and 1,879 women who completed their first-term enlistment. AORs adjusted for race/ethnicity, AFQT category, age, and BMI. +, Statistically significant difference ($p < 0.05$) from the reference category.

major causes of medical attrition.³ Outpatient visits due to injuries are generally the result of musculoskeletal disorders associated with physical training and vigorous operational activities. These injuries have a significant impact on readiness. A fracture can account for >100 lost duty days and a simple sprain can result in several weeks of limited duty, either would restrict deployment. In the Department of the Navy, 58% of medical separations are for musculoskeletal/degenerative disorders.¹ According to Department of Defense (DoD) reports, it costs an average of \$11,350 to recruit and train an active duty enlistee through basic training and initial skills training.²³ If in fiscal year 1998, 4,000 sailors and Marines were medically separated (based on trends described elsewhere), the cost could be \$45.4 million in recruiting and training costs alone. Not computed in this total are total costs related to disability, hospitalization, prescription medication, and lost man-hours. All told, DoD pays \$1.5 billion per year in orthopedic disability costs.² Furthermore, in relation to the cost of injuries, a relatively small amount of resources are devoted to prevention, surveillance, best practices program implementation, and research of injury causal factors in the Department of the Navy. The current status of injury surveillance and prevention efforts in the DoD is fragmented.¹

The Marine Corps Recruiting Commands, in collaboration with the MCRD Recruit Training Regiments, are putting together a physical training program designed for incoming recruits. This program, soon to be administered by recruiters, is a primary prevention strategy whose goal is to change the risk of injury during recruit training and increase the graduation rate by increasing the incoming recruit's physical fitness. The "Poolee Physical Training Manual" and video are in postproduction. However, the recruiter's involvement is limited because there are potential legal ramifications if a recruiter "breaks" a recruit before they ship.

It is clear that training injuries are very costly in terms of health care resources and lost training days during recruit training.^{1,2} Injuries can result in temporary and permanent impairments that can interfere with a service member's ability to perform. For example, the estimated fiscal impact of physical disability payments for the U.S. Army was \$500 million for 1994 alone.² The cost of physical disability in the Marine Corps increased from \$10.1 million in 1989 to \$28.5 million in 1995.² A musculoskeletal injury can result in immediate separation from recruit training; however, for those recruits who suffered a musculoskeletal injury and completed training, the impact of such injuries has not been assessed beyond the training environment, along the accession pathway, and into the Fleet Marine Force.

Musculoskeletal injuries are costly not only because of health care costs but also the financial loss from attrition. The potential long-term impact of musculoskeletal injuries and stress fracture training-related injuries emphasizes the importance of early identification and prevention in training environments. In light of the increased escalation to fight the Global War on Terrorism, emphasis should be made to prevent training-related injuries, not on separating those who sustain an injury and therefore increase the recruit graduation rates for all the military services.

This study has limitations. We did not collect specific sepa-

ration or reenlistment codes and therefore only analyzed broad-level associations between recruit training injury and the three outcomes of interest. Obstacles to analyzing more specific associations are stated in a General Accounting Office report²⁴:

DoD's current data on attrition is inconsistent and incomplete for two reasons. First, the services interpret DoD's definitions of separation codes differently and therefore place enlistees with identical situations in different discharge categories. Second, DoD's separation codes capture only the officially assigned reason for discharge, when many other factors may result in an enlistee's separation. DoD has not issued guidance for applying these separation codes.

However, it is known that first-term attrition before expiration-of-active service may be due to failure to meet medical, physical fitness or weight-control standards, pregnancy, failure to successfully complete alcohol or drug abuse rehabilitation, or unsatisfactory conduct or job performance.²⁵ Also not known is how many Marines who sustained an injury during recruit training were affected by first-term enlistment Stop-Loss orders, effectively requiring them to stay beyond the end of their active service, or if Stop-Loss orders affect only career Marines and not first-term enlistees. Not knowing Stop-Loss information may understate the effect of recruit injury on first-term enlistment outcomes.

The recruits who are diagnosed with severe stress fractures (pelvis, femoral neck, bilateral tibia) and sacral stress fractures (mostly women) are usually separated because they take a long time to heal. Therefore, this study does not represent the first-term outcomes of the recruits diagnosed with the most severe stress fractures and cannot infer outcomes to that group. Furthermore, first-term separation, failure to achieve the rank of corporal, or reenlisting might be due to some other trait and not due to the basic training injury, especially stress fracture, and therefore does not confirm or deny a causal relationship.

There were 295 (10.9%) of the 2,715 women from the historical sample who graduated from Marine Corps basic training with missing follow-up information who were excluded from the prospective sample. It is possible that the 295 omitted women could be significantly different than the final sample of 2,420 women.

Comments

These data suggest that a lower extremity injury during recruit training impedes future military success, even among the women who graduate. Several differences by injury status during training were found in three measures of military success (attrition, failure to complete first-term enlistment, and failure to achieve the rank of corporal during the first term). Women who graduated after incurring training injuries and especially stress fractures were less likely to complete their first-term enlistment. Because 44% of the women who graduated had incurred a lower extremity injury during training, this could significantly affect military readiness. This effect is even stronger among female graduates who had a stress fracture during training, but only approximately 4% of women are in this category. In addition, women who graduated after incurring recruit training injuries, especially stress fractures, and completed

their first-term enlistment were less likely to be promoted to corporal during the first-term enlistment than uninjured women. Among women who completed the first-term, there was no effect of injury or stress fracture during training on reenlistment status. Future work needs to be done to compare these rates with rates for male recruits and to explore variables other than age, race, and education that may affect this relation. In addition, follow-up on the efficacy of the Marine Corps Recruiting Commands' Poolee Physical Training Program a few years after its implementation by recruiters would help understand the association between the magnitude of change in fitness before recruit training and injury risk and graduation rates. Too many separations can affect the operational readiness of the Fleet Marine Force. The need for all of the services to write instructions and policy requiring injury prevention during the accession pathway may be warranted.

Acknowledgments

We thank the U.S. Marines at MCRD Parris Island, particularly the Fourth Battalion Commanding Officer, for their interest and support of this research. We thank the medical staff at MCRD Parris Island Sports Medicine Clinic (SMART), particularly CAPT Scott Flinn, MC USN. We also acknowledge the contributions of Mrs. Kelli Betsinger, our on-site project coordinator, and Mr. Timothy Bockelman, the Physical Fitness Advisor to the Recruit Training Regiment. Finally, we thank the research staff at Naval Health Research Center, especially Ms. Michelle Stoia for her technical writing expertise.

This research was supported by the Army Reimbursable, U.S. Army Medical Research Acquisition Activity (USAMRAA), Bone BAA Award DAMD 17-02-IA-0001, was conducted at the Naval Health Research Center, San Diego, California, under research Work Unit Number 60206.

References

- Jones BH, Amoroso PJ, Canham ML, Weyandt MB, Schmitt JB: Atlas of injuries in the U.S. Armed Forces. *Milit Med* 1999; 164(Suppl 8): 1-1-9-26.
- U.S. Army: Department of the Army Inspector General's Report, Disability Cost Estimate, 1995. Washington, DC, Physical Disability Agency, 1995.
- Kaufman KR, Brodine S, Shaffer RA: Military training-related injuries: surveillance, research, and prevention. *Am J Prev Med* 2000; 18: 54-63.
- Quester AO, North JH, Kimble TH: Identifying successful Marine Corps recruits. CRM 89-314. Alexandria, VA, Center for Naval Analyses, April 1990.
- Chief of Naval Operations: Top Five Readiness Priorities for the Navy: Manpower; Current Readiness; Future Readiness; Quality of Service; Alignment. Statement, written, to the Senate Armed Services Committee on Status of the Navy, September 27, 2000. Washington, DC, Department of the Navy, 2000.
- Booth-Kewley S, Larson GE, Ryan MAK: Predictors of Navy attrition. I. Analysis of 1-year attrition. *Milit Med* 2002; 167: 760-9.
- North JH: Seasonal patterns in the senior and graduate markets for Marine Corps recruits. CRM 91-230. Alexandria, VA, Center for Naval Analyses, March 1992.
- Gunderson EK, Miller MR, Garland CF: Career History Archival Medical and Personnel System (CHAMPS): Data Resource for Cancer, Chronic Disease, and Other Epidemiological Research. NHRC Technical Report No. 02-06. San Diego, CA, Naval Health Research Center, 2002.
- Population Representation in the Military Services, Fiscal Year 1998. Service data from OASD (FMP) (MPP)/Accession Policy, submitted in accordance with DoD Instruction 7730.56, 25th Annual Department of Defense Report. Washington, DC, Department of Defense, November 1999.
- Kilburn MR, Hanser LM, Klerman JA: The AFQT and its role in enlistment. In: Estimating AFQT Scores for National Education Longitudinal Study (NELS) Respondents, Ch 2. Rand Document MR-818-OSD/A. Santa Monica, CA, The Rand Corporation, Arroyo Center, 1998.
- Fiscal 2006 Defense Budget: Recruiting Retention and Personnel, FDCH Congressional Testimony, March 16, 2005. Statement of Lieutenant General HP Osman, Deputy Commandant for Manpower and Reserve Affairs, US Marine Corps. Washington, DC, Department of Defense, 2006.
- Military Personnel Overview, FDCH Congressional Testimony, March 11, 2003. Statement of Lieutenant General GL Parks, US Marine Corps. Washington, DC, Department of Defense, 2003.
- Shaffer RA, Brodine SK, Ito SI, Le AT: Epidemiology of illness and injury among U.S. Navy and Marine Corps female training populations. *Milit Med* 1999; 164: 17-21.
- Almeida SA, Maxwell-Williams K, Shaffer RA, Brodine SK: Epidemiologic patterns of musculoskeletal injuries and physical training. *Med Sci Sports Exerc* 1999; 31: 1176-82.
- Shaffer RA, Brodine SK, Almeida SA, Williams KM, Ronaghy S: Use of simple measures of physical activity to predict stress fractures in young men undergoing rigorous physical training program. *Am J Epidemiol* 1999; 149: 236-42.
- Linenger JM, West LA: Epidemiology of soft-tissue/musculoskeletal injury among U.S. Marine recruits undergoing basic training. *Milit Med* 1992; 157: 491-3.
- Orava S: Stress fractures. *Br J Sports Med* 1980; 14: 40-4.
- Milgrom C, Giladi M, Chisin R, Dizian R: The long-term followup of soldiers with stress fracture. *Am J Sports Med* 1985; 13: 398-400.
- Bosworth AM, Hurtado SL, Shaffer RA: Long-term outcomes associated with lower extremity stress fractures in recruit training. In: Program and Abstracts of the San Diego Biostatistics and Epidemiology Research Exchange. La Jolla, CA, Biostatistics and Epidemiology Research Exchange, May 5, 1998.
- Hindelang RL, Schwerin MJ, Farmer WL: Quality of life (QOL) in the U.S. Marine Corps: the validation of a QOL model for predicting reenlistment intentions. *Mil Psychol* 2004; 16: 115-34.
- Sanchez RP, Bray RM, Vincus AA, Bann CM: Predictors of job satisfaction among active duty and reserve/guard personnel in the U.S. military. *Mil Psychol* 2004; 16: 19-35.
- Maradmin 255/05, signed June 6, 2005, Fiscal Year 2006 Enlisted Retention Guidelines. Available at www.usmc.mil/maradmins/maradmins2000.nsf/maradmins; accessed July 12, 2006.
- Military attrition: better screening of enlisted personnel could save DoD millions of dollars. GAO/NSIAD-97-102. Washington, DC, General Accounting Office, March 1997.
- Military attrition: DoD could save millions by better screening enlisted personnel. GAO/NSIAD-97-39. Washington, DC, General Accounting Office, January 1997.
- Armed Services and Government News: Keeping soldiers in service. *Army Times*, June 13, 2005.

Copyright of *Military Medicine* is the property of Association of Military Surgeons of the United States and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. Report Date (DD MM YY) 05 20 2007	2. Report Type Final	3. DATES COVERED (from - to) November 2001 to November 2004
--	--------------------------------	---

4. TITLE AND SUBTITLE Negative First-Term Outcomes Association With Lower Extremity Injury During Recruit Training Among Female Marine Corps Graduates	5a. Contract Number: Army Reimbursable 5b. Grant Number: 5c. Program Element: 5d. Project Number: 5e. Task Number: 5f. Work Unit Number: 60206
--	---

6. AUTHORS Daniel W. Trone, MA; Adriana Villaseñor, BS; Caroline A. Macera, PhD	
---	--

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center P.O. Box 85122 San Diego, CA 92186-5122	9. PERFORMING ORGANIZATION REPORT NUMBER Report No. 05-19
---	---

8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES) Chief, Bureau of Medicine and Surgery Code M53 2300 E St NW Washington DC 20372-5300	10. Sponsor/Monitor's Acronyms(s) 11. Sponsor/Monitor's Report Number(s)
--	---

12 DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.
--

13. SUPPLEMENTARY NOTES Published in <u>Military Medicine</u> , 2007, 172(1), 83-89

14. ABSTRACT (maximum 200 words)

The objective of this study was to assess the impact of lower extremity injuries and stress fractures during recruit training on first-term outcomes among female Marine Corps graduates. Information on injuries was collected from women undergoing training in Parris Island, SC (1995–1999) and negative first-term outcomes were obtained from the Career History Archival Medical and Personnel System. First-term was defined as the period after MCRD graduation, and included School of Infantry training and any specialized military occupational (MOS) training before matriculating into the Fleet Marine Force, up to 48 months of service. The three negative outcomes included (1) failure to complete the first-term of service, (2) failure to achieve the rank of Corporal, and (3) failure to reenlist. Overall, 22% did not complete the first-term enlistment, and 12% of those who did were not promoted to Corporal. After adjustment for demographic characteristics, not completing the first term and not being promoted to Corporal were both associated with injuries or stress fracture during training. Reenlistment status was not associated with training injuries. Our findings indicate that lower extremity injuries among women undergoing Marine Corps recruit training are associated with poor first-term military outcomes even among those who graduate.

15. SUBJECT TERMS first-term attrition, injuries, military, stress fracture, women
--

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UNCL	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON Commanding Officer	
a. REPORT UNCL	b. ABSTRACT UNCL	c. THIS PAGE UNCL			19b. TELEPHONE NUMBER (INCLUDING AREA CODE) COMM/DSN: (619) 553-8429	