AD

Award Number: W81XWH-06-1-0541

TITLE: International Conference on Natural Orifice Transluminal Endoscopic Surgery (NOTES)

PRINCIPAL INVESTIGATOR: Michael Marohn

JHU (410) 502-0723

CONTRACTING ORGANIZATION: American Society for Gastrointestinal Endoscopy Oak Brook, IL 60523-2141

REPORT DATE: June 2006

TYPE OF REPORT: Final Proceedings

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT	DOCUMENTATION	I PAĜE	Form Approved OMB No. 0704-0188
Public raporting burdan for this collection of inforr data naedad, and complating and raviewing this t this burdan to Department of Dafanse, Washingto 4302. Raspondants should be awara that notwit	nation is astimated to avarage 1 hour per raspor collection of information. Sand commants ragarc on Haadquarters Sarvices, Diractorate for Inform standing any other provision of law. no person i	nsa, including tha tima for rav ding this burdan astimata or a ation Operations and Raport shall be subject to any penalt	iviawing instructions, searching axisting data sources, gathering and maintaining the any othar aspect of this collection of information, including suggastions for reducing rts (0704-0188), 1215 Jafferson Davis Highway, Suite 1204, Artington, VA 22202- ity for failing to comply with a collection of information if it does not display a currantly and the comply with a collection of information.
valid OMB control number. PLEASE DO NOT R	ETURN YOUR FORM TO THE ABOVE ADDRE	SS.	
1. REPORT DATE	Einal Proceedings		9 Mar 2006 - 30 Jun 2006
4. TITLE AND SUBTITLE	That Toceedings		5a, CONTRACT NUMBER
International Conference on N (NOTES)	Vatural Orifice Transluminal E	Endoscopic Surge	ery 5b. GRANT NUMBER W81XWH-06-1-0541
(			5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)			5d. PROJECT NUMBER
Michael Marohn			5e. TASK NUMBER
			5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION	NAME(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT
American Society for Gastroir Oak Brook, IL 60523-2141	itestinal Endoscopy		NUMBER
9. SPONSORING / MONITORING A U.S. Army Medical Research	GENCY NAME(S) AND ADDRESS	(ES)	10. SPONSOR/MONITOR'S ACRONYM(S)
Fort Detrick, Maryland 21702	2-5012		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)
Approved for Public Release;	Distribution Unlimited		
13. SUPPLEMENTARY NOTES Original contains colored pla	ates: ALL DTIC reproductions	s will be in black a	and white.
14. ABSTRACT			-
	2	0091	029219
15. SUBJECT TERMS NOT PROVIDED			
16. SECURITY CLASSIFICATION C	)F:		18. NUMBER 19a. NAME OF RESPONSIBLE PERSON OF PAGES USAMRMC
a. REPORT b. ABSTRA U U	CT C. THIS PAGE U	UU	384 <b>19b. TELEPHONE NUMBER</b> (include area
		A	Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std. 239.18 Q F10-01-00095

Volume 20 • Number 5 May 2006

and Other Interventional Techniques

• LAPAROSCOPY • FLEXIBLE ENDOSCOPY • INTERVENTIONAL RADIOLOGY • ULTRASOUND •

Official Journal of the Society of American Gastrointestinal and Endoscopic Surgeons European Association for Endoscopic Surgery

464 Surg Endosc ISSN 0930-2794 20(5) 711-844 (2006)

RGIGA

MII SCS



Online First Immediately Online springerlink.com

## I BUILD RELATIONSHIPS THROUGH BARIATRICEDGE<sup>SM</sup>

DEVELOPED BY YOUR TRUSTED PROVIDER OF STATE-OF-THE-ART BARIATRIC SURGICAL TOOLS AND DEVICES. ETHICON ENDO-SURGERY BARIATRICEDGE<sup>5M</sup> DELIVERS PREMIER BARIATRIC SERVICE SOLUTIONS WITH A VISION OF DRAMATICALLY IMPROVED HEALTH FOR BARIATRIC PATIENTS AND AN UNMATCHED FOCUS ON PATIENT OUTCOMES AND CARE.

THROUGH BARIATRICEDGE<sup>™</sup>, IOFFER PASSION AND EXPERTISE IN BARIATRIC SURGERY. ONGOING TREATMENT AND CARE, PRACTICE SUPPORT, AND PATIENT EDUCATION.

TODD DAWSON Bariatric Account Manager

> STHICON ENDO-SURGERY, INC Submen-Submen concurv BARIATRIC EDGE

1011



Now Cleared for Laparoscopic Use!

SurgiWrap is the only bioresorbable sheet to create a physical barrier while minimizing soft tissue attachment in both open and laparoscopic procedures.

## SurgiWrap is...

- Made of 70:30 Poly(L-lactide-co-D,L-lactide)
- Transparent and offers significant mechanical strength
- Easily handled in a wet environment

## And...

Can be secured for precise placement and complete surgeon control



Visit us at SAGES Booth #311





# SAGES gratefully acknowledges the 2006 meeting levels donors for their generous support.

## PLATINUM LEVEL DONORS

Autosuture and Valleylab, divisions of Tyco Healthcare

Ethicon Endo-Surgery, Inc.

Karl Storz Endoscopy

Olympus America

## GOLD LEVEL DONORS

Inamed Health Stryker Endoscopy

## SILVER LEVEL DONORS

Boston Scientific Davol, Inc. General Surgery News Gore & Associates, Inc.

## **BRONZE LEVEL DONORS**

Adolor Corporation and GlaxoSmithKline Aesculap B-K Medical Systems Cook Surgical Medtronic Synovis Surgical Innovations SurgRX Taut, Inc. Tissue Science Laboratories



and Other Interventional Techniques

B. Millat, Montpellier

J. H. Peters, Los Angeles

J. L. Ponsky, Cleveland

D. Meijer, Rotterdam

Sir Alfred Cuschieri, Dundee





#### Editors-in-Chief

#### **Associate Editors**

## Technical Editors Multimedia Editor

#### **Editorial Board**

- D. Arvidsson, Stockholm
  M. E. Bailey, Guildford
  K. N. M. A. Bax, Utrecht
  H. J. Bonjer, Rotterdam
  L. M. Brunt, St. Louis
  G. Buess, Tübingen
  G. B. Cadiere, Brussels
  M. P. Callery, Boston
  J. Cervantes, Mexico City
  S. C. Chung, Hong Kong
- R. Clayman, St. Louis
- E. Croce, Monza
- M. A. Cuesta, Amsterdam
- B. Dallemagne, Liege

**B. V. MacFadyen. Jr.**, Augusta (Coordinating)

- D. Birkett, Burlington
- J. Jakimowicz, Eindhoven
- A. M. Lacy, Barcelona
- G. Berci, Los Angeles
- S. D. Schwaitzberg, Boston
- B. Schirmer, Charlottesville A. Darzi, London E. Lezoche, Rome M. Edye, New York T. Lobe, Memphis W. Schwesinger, San Antonio J. Marescaux, Strasbourg A. Fingerhut, Poissy N. Soper, Chicago D. Fletcher, Melbourne J. Marks, Wynnewood G. V. Stiegmann, Denver K. H. Fuchs, Frankfurt J. M. Marks, Mayfield Heights C. Sugawa, Detroit J. F. Gigot, Brussels A. Montori, Rome C. Sutton, Guildford D. Gossot, Paris J. Mouiel, Nice L. Swanstrom, Portland F. L. Greene, Charlotte L. Nathanson, Herston G. Vitale, Louisville J. K. Gupta, Birmingham E. Neugebauer, Cologne W. Wayand, Linz G. W. Holcombe, Kansas City S. Ohashi, Osaka S. Wexner, Weston A. Park, Baltimore J. Hunter. Portland R. L. Whelan, New York J. Perissat, Bordeaux N. Katkhouda, Los Angeles B. Wolfe. Sacramento E. Phillips, Los Angeles F. Köckerling, Hannover A. Pietrabissa, Pisa S. Wren, Palo Alto R. Kolvenbach, Dusseldorf J. Sandor, Budapest E. Laporte Rosella, Barcelona C. Z. Zeng, Shanghai R. Satava, Seattle G. Larson, Louisville

Editor Emeritus: K. A. Forde, New York Founding Editor: B. C. Manegold, Mannheim

Official Journal of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) and European Association for Endoscopic Surgery (E.A.E.S.)

Association of Endoscopic Surgeons of Great Britain and Ireland Société Française de Chirurgie Endoscopique (S.F.C.E.) Società Italiana di Chirurgia Endoscopica e Chirurgia Mini-invasiva (S.I.C.E.) Surgical Endoscopy Association, Branch of the Society of Polish Surgeons

# SURGICAL-ENDOSCOPY

and Other Interventional Techniques

The journal represents the surgical aspects of interventional endoscopy, ultrasound, and other techniques in the fields of gastroenterology, obstetrics, gynecology, and urology, as well as gastroenterologic, thoracic, traumatic, orthopedic, and pediatric surgery. It is a worldwide forum for the description and discussion of various aspects of interventional endoscopy and ultrasound as integral elements of surgical practice. Interventional endoscopy and ultrasound have permitted decisive advances in surgery. The journal serves the international surgical community for the transfer of information relating to practice, theory, and research in the various medical and surgical disciplines.

#### Submit manuscripts online at http://mc.manuscriptcentral.com/

surgendosc. If you are unable to submit via the internet, contact Bernadine Richey, Surgical Endoscopy Coordinating Editorial Office, publishingservices@fcc.net, 845-353-3106 (U.S. phone), 845-348-3948 (U.S. fax) for submission instructions.

#### Subscription Information

Surgical Endoscopy is published monthly by Springer Science + Business Media, Inc. (Springer), Volume 20 (12 issues) will be published in 2006.

ISSN: 0930-2794 (print version)

ISSN: 1432-2218 (electronic version)

#### Subscription Rates, Orders, Inquiries:

Please contact our Customer Service department for the latest rates and pricing information: *The Americas (North, South, Central America,* and the Caribbean): MAIL: Journals Customer Service P.O. Box 2485 Secaucus, NJ 07096, USA TEL: 800-777-4643; 212-460-1500 (outside North America)

FAX: 201-348-4505

E-MAIL: journals-ny@springer-sbm.com

Outside of the Americas: MAIL: Journals Customer Service Springer Distribution Center (SDC) Haberstr. 7 69126 Heidelberg, GERMANY TEL: 49-6221-345-4303 FAX: 49-6221-345-4229 E-MAIL: sdc-journals@springer-sbm.com

#### Society Subscription Information:

Membership in the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) and the European Association for Endoscopic Surgery (E.A.E.S.) includes a special subscription rate to *Surgical Endoscopy*. For information regarding membership please contact one of the following:

Membership Administrator, SAGES 11300 Olympic Boulevard, Suite 600 Los Angeles, CA 90064, USA TEL: (310) 437-0544 FAX: (310) 437-0585

Secretariat, E.A.E.S. Office c/o Mrs. Ria Palmen Luchthavenweg 81, Unit 139A 5657 EA Eindhoven, The Netherlands TEL: +31 40 252 52 88 FAX: +31 40 252 31 02 E-MAIL: ria.palmen@eaes-eur.org or info@eaes-eur.org website: www.eaes-eur.org

Change of Address: Allow six weeks for all changes to become effective. All communica-

tions should include both old and new addresses (with zip codes) and should be accompanied by a mailing label from a recent issue.

Back Volumes: Prices for back volumes are available on request.

*Microform Editions:* Available from: University Microfilms International, 300 N. Zeeb Road, Ann Arbor, MI 48106, USA.

SpringerAlerts Service: The SpringerAlert service is an innovative, free-of charge service that notifies users via e-mail whenever new SpringerLink articles and journals become available, and automatically sends the table of contents and direct links to the abstracts of a new issue of a journal in SpringerLink. Register for the SpringerAlert service at http://www.springerlink. com/alerting

#### Copyright

Submission of a manuscript implies that the work described has not been published before (except in the form of an abstract or as part of a published lecture, review or thesis): that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as-tacitly or explicitly-by the responsible authorities at the institution where the work was carried out. Transfer of copyright to Springer becomes effective if and when the article is accepted for publication. The copyright covers the exclusive right (for U.S. government employees: to the extent transferable) to reproduce and distribute the article, including reprints, translations, photographic reproductions, microform, electronic form (offline, online) or other reproductions of similar nature

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e.g., as reprints), as well as all translation rights. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from Springer. The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

An author may self-archive an author-created version of his/her article on his/her own website and his/her institution's repository, including his/ her final version; however he/she may not use the publisher's PDF version which is posted on www.springerlink.com. Furthermore, the author may only post his/her version provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The original publication is available at www.springerlink.

com." Please use the appropriate DOI for the article (go to Linking Options in the article, then to OpenURL and use the link with the DOI). Articles disseminated via www.springerlink.com are indexed, abstracted, and referenced by many abstracting and information services, bibliographic networks, subscription agencies, library networks and consortia.

While the advice and information in this journal is believed to be true and accurate at the date of its publication, neither the authors, the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein. Springer publishes advertisements in this journal in reliance upon the responsibility of the advertiser to comply with the all legal requirements relating to the marketing and sale of products or services advertised. Springer and the editors are not responsible for claims made in the advertisements published in the journal. The appearance of advertisements in Springer publications does not constitute endorsement, implied or intended, or the product advertised or the claims made for it by the advertiser.

Special regulations for photocopies in the USA. Photocopies may be made for personal or inhouse use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the *Copyright Clearance Center, Inc.*, 222 Rosewood Drive, Danvers, MA 01923, USA (WEB: http://www.copyright.com; FAX: 978-646-8600; TEL: 978-750-8400), stating the journal's ISSN, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works or resale. In these cases, specific written permission must first be obtained from the publisher.

The Canada Institute for Scientific and Technical Information (CISTI) provides a comprehensive, world-wide document delivery service for all Springer journals. For more information, or to place an order for a copyright-cleared Springer document, please contact Client Assistant, Document Delivery, CISTI, Ottawa K1A 0S2, Canada. TEL: 613-993-9251; FAX: 613-952-8243; E-MAIL: cisti.docdel@nrc.ca.

You may also contact *INFOTRIEVE* (WEB: http://www.infotrieve.com/; E-MAIL: orders@infotrieve.com; FAX: 310-208-5971; TEL: 800-422-4633),

UnCover: http://www.ingenta.com; UMI: http://www.umi.com/; ISI: http://www.isinet.com

#### Responsible for Advertisements

M.J. Mrvica Associates, Inc. 2 West Taunton Avenue Berlin, NJ 08009, USA TEL: (856) 768-9360 FAX: (856) 7653-0064 E-MAIL: mjmrvica@mrvica.com

#### Office of Publication

Springer, 233 Spring Street, New York, NY 10013, USA. Telephone: 212-460-1500; e-mail: journals-ny@springer-sbm.com website: springeronline.com Production Editor: Ray Ramonas Printed in the United States on acid-free paper. ©2006 Springer Science + Business Media, Inc.

Periodicals postage paid at New York, New York and additional mailing offices.

**Postmaster:** Send address changes to: Surgical Endoscopy, Springer, 233 Spring Street, New York, NY 10013, USA.

Universal Seal Accommodates Multiple Instrument Sizes

**Reduced Drag Force** 

**Enhanced Seal System** 

**Improved Retention** 

**Better Tactile Sensation** 

**Decreased Tipping and Rolling** 

More Ergonomic Housing

**Increased Seal Durability** 

## **Reduced Instrument Hang-ups**

Lower Profile Gas Stopcock

**Clear Cannula** 



For more information, contact your local Ethicon Endo-Surgery representative or call 1-800-USE-ENDO.



Compared to current ENDOPATH® trocars and/or other leading trocars. Test data on file. ©2004 Ethicon Endo-Surgery, Inc. All rights reserved. ENDOPATH is a registered trademark and XCEL is a trademark of Ethicon Endo-Surgery, Inc. DSL 04-0715



genzyme

# DON'T COMPROMISE

#### With Sepramesh"IP, you don't have to choose between safety, strength and ease of use in hernia repair.

On one side, you get the safety of a unique hydrogel coating. On the other, the trusted strength of polypropylene mesh. All incorporated into one easy-to-use, integrated design. Why compromise? Call 1-800-261-1570 or visit www.SeprameshiP.com.

Sepramesh<sup>ai</sup>IP is indicated for use in the reconstruction of soft tissue deticiencies, such as for the repair of hernias. In infants or children with tuture growth potential, Sepramesh IP may not stretch significantly as the patient grows. Do not use Sepramesh IP for the reconstruction of cardiovascular detects. Please see full product information for more details.



Copyright © 2005 by Generate Corporation. September 15 intrademark and September (is) a register of bratemark of Generate Corporation. "No Compromises" is a service mark of Generation Code No. 68724 a. 02/2005

Valent Provent PL AND AND ADD

and Other Interventional Techniques

and Ot

#### CONTENTS

#### Editorial

711 NOTES: Gathering Momentum David Rattner, Robert H. Hawes

#### Original articles

713	Laparoscopic management of gastrointestinal stromal tumors
	S. Q. Nguyen, C. M. Divino, JL. Wang, S. H. Dikman

717 Morbidity of laparoscopic surgery for complicated appendicitis: an international study

J. Cueto, B. D'Allemagne, J. A. Vázquez-Frías, S. Gomez, F. Delgado, L. Trullenque, R. Fajardo, S. Valencia, L. Poggí, J. Ballí, J. Diaz, R. González, J. H. Mansur, M. E. Franklin

- Utility of staging laparoscopy in subsets of biliary cancers: laparoscopy is a powerful diagnostic tool in patients with intrahepatic and gallbladder carcinoma D. Goere, G. D. Wagholikar, P. Pessaux, N. Carrère, A. Sibert, V. Vilgrain, A. Sauvanet, J. Belghiti
- 726 Laparoscopic ultrasound navigation in liver surgery: technical aspects and accuracy

M. Kleemann, P. Hildebrand, M. Birth, H. P. Bruch

- 730 Advantages of advanced laparoscopic systems J. Heemskerk, R. Zandbergen, J. G. Maessen, J. W. M. Greve, N. D. Bouvy
- 734 The anatomical significance and techniques of laparoscopic rectal surgery L. Chengyu, J. Xiaoxin, Z. Jian, G. Chen, Y. Qi
- 139 Impaired esophageal function in morbidly obese patients with gastrocsophageal reflux disease: evaluation with multichannel intraluminal impedance
   E. Quiroga, F. Cuenca-Abente, D. Flum, E. P. Dellinger, B. K. Oelschlager
- The MISTELS program to measure technical skill in laparoscopic surgery: evidence for reliability
   M. C. Vassiliou, G. A. Ghitulescu, L. S. Feldman, D. Stanbridge,
   K. Leffondré, H. H. Sigman, G. M. Fried
- 748 Laparoscopic subtotal splenectomy in hereditary spherocytosis: to prescrve the upper or the lower pole of the spleen?
  C. Vasilescu, O. Stanciulea, S. Tudor, D. Stanescu, A. Colita, R. Stoia, D. Coriu, A. Colita, C. Arion
- 753 Laparoscopic cholecystectomy using a newly developed laparoscope manipulator for 10 patients with cholclithiasis

K. Tanoue, T. Yasunaga, E. Kobayashi, S. Miyamoto, I. Sakuma, T. Dohi, K. Konishi, S. Yamaguchi, N. Kinjo, K. Takenaka, Y. Maehara, M. Hashizume

- 757 Modified needlescopic video-assisted thoracic surgery for primary spontaneous pneumothorax: the long-term effects of apical pleurectomy versus pleural abrasion Y.-C. Chang, C.-W. Chen, S.-H. Huang, J.-S. Chen
- 763 Systematic evaluation of different approaches for minimizing hemodynamic changes during pneumoperitoneum
   T. Junghans, D. Modersohn, F. Dörner, J. Neudecker, O. Haase, W. Schwenk
- 770 Linear stapling of the short gastric vessels reduces blood loss and splenectomy at oesophageal and gastric surgery J. F. Dowdall, O. J. McAnena
- 773 Zoom endoscopic monitoring of small bowel allograft rejection
  T. Kato, J. J. Gaynor, S. Nishida, N. Mittal, G. Selvaggi, D. Levi, J. Moon, J. Thompson, P. Ruiz, J. Madariaga, A. G. Tzakis
- 783 Bravo capsule induction of csophageal hypercontractility and chest pain
   C. Tharavej, J. A. Hagen, G. Portale, C.-C. Hsieh,
  - T. A. K. Gandamihardja, J. C. Lipham, J. H. Peters,
  - S. R. DeMeester, P. F. Crookes, C. G. Bremner,
  - T. R. DeMeester
- 787 Laparoscopic hepatic resection
   G. Borzellino, A. Ruzzenente, A.-M. Minicozzi,
   F. Giovinazzo, C. Pedrazzani, A. Guglielmi
- 791 The "stamp method": a new treatment for perforated peptic ulcer?
  M. J. O. E. Bertleff, R. S. B. Liem, H. L. Bartels,
  P. H. Robinson, J. F. A. Van der Werff, H. J. Bonjer,
  J. F. Lange
- Flucidating the relationship between cardiac preload and renal perfusion under pneumoperitoneum
   Simon Bergman, Arni Nutting, Liane S. Feldman, Melina C. Vassiliou, Christopher G. Andrew,
   Sebastian Demyttenaere, Debbie Woo, Franco Carli, Luc Jutras, Jean Buthieu, Donna D. Stanbridge,
   Gerald M. Fried
- 801 Do all patients with abnormal intraoperative cholangiogram merit endoscopic retrograde cholangiopancreatography?
   S. Varadarajulu, M. A. Eloubeidi, C. M. Wilcox, R. H. Hawes, P. B. Cotton
- 806 Urgent laparoscopic cholecystectomy in the management of acute cholecystitis: timing does not influence conversion rate Y.-C. Wang, H.-R. Yang, P.-K. Chung, L.-B. Jeng, R.-J. Chen
- 809 It is not worthwhile to perform ilcoscopy on all patients K. K. Y. Yoong, T. Heymann
- **Rigid sigmoidoscopy: a potential hazard for** cross-contamination
   D. Z. Lubowski, G. L. Newstead

# Set your sights

#### Introducing the INSCOPE<sup>™</sup> Precision Balloon Dilator

Improved accuracy under direct visualization\*

#### INCREASING THE LEVEL OF CONTROL

- Improves centering accuracy with 3 distinct marker bands
- Maintains position throughout procedure
- 100° proximal taper facilitates endoscopic visualization

#### SMOOTH, CONSISTENT REMOVAL

- Designed for rapid balloon deflation
- Co-extruded stainless steel wire prevents catheter elongation upon retraction
- Asymmetrical design with 50° distal taper
- Seamless transition from catheter to low-stretch balloon

#### ENHANCING PROCEOURAL EFFICIENCY

- Assured 3-stage dilation with a single intubation
- Available in 6 sizes, ranging in diameter from 6 mm to 20 mm

\*Compared to other balloons indicated for esophageal dilation.

Stricture -

Three distinct marker bands for improved centering

## INSCOPE"

PRECISION BALLOON DILATOR

Insight. Innovation. In Your Hands.

For more information, call 1-866-INSCOPE or visit www.INSCOPE.com.

For complete product deteils, pleese see Instructions for Use.

©2005 InScope, a division of Ethicon Endo-Surgery, Inc., a Johnson & Johnson company. 12/05 DSL#05-1452

- 815 Preconditioning-like amelioration of erythropoietin against laparoscopy-induced oxidative injury
   E. Ates, S. Yilmaz, E. Ihtiyar, B. Yasar, E. Karahuseyinoglu
- 820 Laparoscopic ventral hernia repair using expanded polytetrafluoroethylene–polyester mesh compound M. Farrakha
- Robotic surgery and training: electromyographic correlates of robotic laparoscopic training
   T. N. Judkins, D. Oleynikov, K. Narazaki, N. Stergiou

#### Technique

 Retroperitoneal hand-assisted laparoscopic surgery for endoscopic adrenalectomy
 N. Sata, M. Shiozawa, A. Suzuki, K. Kurihara, J. Ohki, H. Nagai

#### Letters to the editor

- 834 An alternative gallbladder extraction technique in laparoscopic cholecystectomy D. C. T. Wong, C. N. Tang
- A correlation between the shape of the epiglottis and obstructive sleep apnea
   M. Gazayerli, W. Bleibel, A. Elhorr, E. Elakkary, D. Maxwell, R. Seifeldin

#### 838 News and Notices

Indexed and abstracted in *Index Medicus—MEDLINE*, *Excerpta Medica/EMBASE*, *Science Citation Index* Included in the *MEDLARS* system and in *Current Contents* 

Ind	lex to Advertisers
Davol	p. A14
Ethicon Endosurgery	pp. A5, A9, A11, Cover 2
Fleet	pp. A12, A13
Genzyme	p. A6
Karl Storz	Cover 3
MAST Biosurgery	p. A1
Medtronic	Cover 4

Universal Seal Accommodates Multiple Instrument Sizes

**Reduced Drag Force** 

**Enhanced Seal System** 

**Improved Retention** 

**Better Tactile Sensation** 

**Decreased Tipping and Rolling** 

More Ergonomic Housing

**Increased Seal Durability** 

## **Reduced Instrument Hang-ups**

Lower Profile Gas Stopcock

**Clear Cannula** 



For more information, contact your local Ethicon Endo-Surgery representative or call 1-800-USE-ENDO.



Compared to current ENOOPATH® trocars and/or other leading trocars. Test data on file. ©2004 Ethicon Endo-Surgery, Inc. All rights reserved. ENOOPATH is a registered trademark and XCEL is a trademark of Ethicon Endo-Surgery, Inc. 0SL 04-0715

## NEW—the optimal formula for adult medical hydration from Fleet Laboratories<sup>™</sup>

elect the board c sur Inter-Faith fini Bre 42nd counties in North Carolina. For 11 amo your and. pers Bowel Prep's Estate you omas WANTED you 1 chair Best Friend shol dep ourism and conf the Allen, as Must be an oral hydration NO Jsiness rep harma therapy, easy to use, res taste great, and contain eboard 14 le Software up the optimal amount of you ley of 42nd in Raleigh, Na+, K+, and Cl-. il of Mama Japel Hill 1 of Summit p were elect hoerleading aurant f directors hant



HydraLife<sup>™</sup> is now available through the following select wholesale suppliers: CARDINAL 3713393 McKesson 2184224 Ameri-Source East 4915955 Ameri-Source West 334603

Reference: 1. Barclay RL, Depew WT, Vanner SJ. Carbohydrate-electrolyte rehydration protects against intravascular volume contraction during colonic cleansing with orally administered sodium phosphate. Gastrointest Endosc, 2002;56:5:633-638.

## Introducing New HydraLife<sup>™</sup> Oral Rehydration Drink From Fleet Laboratories<sup>™</sup>

For your adult patients undergoing bowel prep, water may not be sufficient to maintain adequate hydration. Now you can give them new HydraLife<sup>™</sup> from Fleet<sub>®</sub>—a powerful formulation of electrolytes and carbohydrates to help your patients avoid the risk of dehydration.

## **Complements Bowel Preparation**

Because patients undergoing bowel prep often experience a transient electrolyte shift that may result in dehydration, HydraLife<sup>™</sup> was specially formulated to replenish essential electrolytes. In fact, a recent study suggests that using an electrolyte drink during the bowel prep process leads to improved efficacy in colonic cleansing.<sup>1</sup>

## HydraLife<sup>™</sup>—The Next Clear Liquid of Choice

- Specially formulated to maintain adequate hydration and supply electrolytes that may be lost during bowel prep
- Crisp orange-tangerine flavor suited for adult taste without the excessive sweetness of children's products
- Available in ready-to-mix powder for enhanced portability and convenience

## Give Your Patients the Optimal Hydration Formulation\*

	HydraLife™	<b>Pedialyte</b> <sup>®†</sup>	Gatorade®‡	WHO <sup>s</sup>	
Sodium	367 mg	367 mg	163 mg	612 mg	
Potassium	417 mg	278 mg	56 mg	278 mg	- 10-00-00
Chloride	828 mg	440 mg	NA	818 mg	
Dextrose	5.68 g	7.10 g	19.88 mg	4.80 g	
Calories	21.3	27	75	18	

\* Milligrams per 12 ounces.

<sup>†</sup> Pedialyte is a registered trademark of Ross Products Division, Abbott Laboratories.

<sup>4</sup> Gatorade is a registered trademark of PepsiCo, Inc.

<sup>8</sup> World Health Organization (low sodium version).



Contains recommended electrolytes for rehydration

NEW HydraLife

Maintain the electrolyte balance

Oral Rehydration Drink

## What does Davol mean to you?

Bard\* 3DMax\* mesh is designed specifically for the laparoscopic repair of inguinal hernias. Its unique anatomical design makes it easy to place. With built-in memory, the pre-formed mesh contours precisely to the inguinal anatomy and may require less operative time. Some surgeons place **3DMax** mesh without fixation, a technique that may reduce the chance of post-operative pain and the additional surgical costs associated with fixation. To learn more about the use of **3DMax** mesh in laparoscopic TEP and TAPP procedures, request our free video introduction kit at www.davol.com/3D or call Davol at 800-556-6275.



Laparoscopic inguinal hernia repair

Bard\* 3DMax\* Mesh

#### BARD

Trusted solutions from the hernia repair company.



\* Bard, and 3DMax are trademarks and/or registered trademarks of C.R. Bard, Inc. or an affiliate. Consult product labels and inserts for any indications, contraindications, hazards, warnings, cautions and instructions for use.

Editorial

Surg Endosc (2006) 20: 711–712 DOI: 10.1007/s00464-006-0263-5

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

### **NOTES: Gathering Momentum**

It has been several years since Kalloo in Baltimore and Reddy and Rao in Hyderabad, India began accessing the peritoneal cavity via a transgastrie route. Though initially greeted with skepticism—some even called it blasphemy- the idea that intraperitoneal surgery might be performed without an abdominal incision appears to be worth pursuing. In a relatively short time span, the Hopkins group demonstrated the feasibility of performing a gastrojejunostomy via a totally endoseopie/ transgastrie route. Even more stunning are the reports from Hyderabad of a series of 7 transgastrie human appendectomies with good results.

Acknowledging the potential of this novel approach, the leadership of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) and the American Society for Gastrointestinal Endoscopy (ASGE) appointed a group of 14 members to study and comment on this new technique. This group met in New York City in July 2005 and published its deliberations as the NOTES Working Group White Paper [1]. The White Paper delineated the anticipated technical barriers to further development of NOTES, emphasized the need for development to be carried out by interdisciplinary teams of surgeons and gastroenterologists and emphasized that any human procedures be performed only with IRB approval. Since the White Paper was authored by a small group, it also mandated that a next step should be a larger and more inclusive gathering of interested parties to challenge the tenets put forth in the White Paper and to create a roadmap for NOTES development.

The first international conference on NOTES was held in Scottsdale, Arizona March 9–11, 2006. One hundred forty physicians from 11 countries came as teams (each team had at least one gastroenterologist and one surgeon with access to animal lab facilities). After a morning of lectures describing accomplishments to date as well as the challenges anticipated to move the field forward, participants were assigned to eight separate working groups with the task of developing a detailed roadmap for overcoming the eight technical barriers that had been identified in the original White Paper. The roadmaps from the Working Groups were presented at the closing session the following day. These presentations as well as the White Paper and other related resources have been posted on a Web site, www.nosear.org, for interested parties to see. Equally important was the launch of NOSCAR-the Natural Orifice Surgery Consortium for Assessment and Research.

NOTES is an emerging transdisciplinary therapy based upon a disruptive technology. If NOTES is to reach a stage of widespread clinical applicability there will need to be further innovation and true collaboration on multiple fronts. Experimental work needs to be done to understand the physiologie disruption and infectious complications of NOTES. Better devices are needed for gastrie elosure, suturing, tissue grasping and manipulation, and anastomosis. Further research is needed to optimize procedure performance. Ultimately, there will be a need for collaborative elinical trials to test the value of NOTES. The establishment of NOSCAR represents a collaborative vision built on the strength and leadership of two strong innovative organizations- SAGES and ASGE. It is our hope that NOSCAR ean provide leadership as well as a collaborative common ground to prospectively shape this emerging therapeutic discipline. In short, The NOSCAR group is a rare attempt to prospectively manage a disruptive technology.

We see NOSCAR as the appropriate vehicle for the following tasks:

- 1) Produce White Papers that define the large challenges needing thought and research.
- 2) Track portfolios i.e. groups of similar research projects that address challenges laid out in the White Paper.
- Provide organization for research projects in such a way as to enhance collaboration and attract funding to key areas of study.
- 4) Collect (and, in fact, require) submission of data to build a robust outcomes database.
- 5) Foster collaborative elinical trials.

We anticipate that SAGES and ASGE will establish a joint committee that will guide NOSCAR to identify and foster needed research directions. NOSCAR, through its parent societies is likely to establish a request for proposal process, raise research funds, vet grant applications and oversee a consortium of labs and clinical study 712

groups. In this fashion, NOSCAR should become the repository for maintaining a portfolio of research projects from interested groups around the world.

As NOTES matures and enters clinical trials it is envisioned that NOSCAR will create and maintain a clinical case registry. Ad hoc NOTES meetings seem likely in the future and NOSCAR, under societal guidance, would organize such meetings. Down the road, if NOTES is shown to be a beneficial technology, NO-SCAR might help define scope of practice, competency measures, and work with regulatory agencies on reimbursement issues.

All this is heady talk given the paucity of data currently available. However, the overwhelming sense among the 140 physicians in Scottsdalc was that NOTES will develop into a mainstream clinical capability in the near future. Times have changed since the introduction of laparoscopic cholecystectomy nearly two decades ago. The public and profession are no longer willing to accept indiscriminate introduction of new technology, and physicians are focused on keeping quality patient care first. By creating NOSCAR, we hope to introduce NOTES in a safe and responsible way that will provide an even less invasive way of undergoing surgical procedures.

#### Reference

 Rattner DW, Kalloo A, SAGES/ASGE working group on natural orifice translumenal endoscopic surgery (2006) White paper. Surg Endosc 20: 329–333

#### **David Rattner**

Massachusetts General Hospital Boston, MA, 02114, USA

#### **Robert H. Hawes**

Medical University of South Carolina Charleston, SC, 29425, USA

Online publication: 28 April 2006

## Original articles

Surg Endosc (2006) 20: 713-716 DOI: 10.1007/s00464-005-0435-8

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

### Laparoscopic management of gastrointestinal stromal tumors

S. Q. Nguyen,<sup>1</sup> C. M. Divino,<sup>1</sup> J.-L. Wang,<sup>1</sup> S. H. Dikman<sup>2</sup>

<sup>1</sup> Department of Surgery, Division of General Surgery, Mount Sinai School of Medicine, 5 E. 98th Street, 15th Floor, Box 1259, New York, NY 10029-6574, USA <sup>2</sup> Department of Pathology, Mount Sinai School of Medicine, 5 E. 98th Street, 15th Floor, Box 1259, New York, NY 10029-6574, USA

Received: 17 June 2005/Accepted: 8 November 2005/Online publication: 21 February 2006

#### Abstract

Background: Surgery remains the standard for nonmetastatic gastrointestinal stromal tumors (GISTs). Laparoscopic surgery should be considered for these tumors as their biologic behavior lends them to curative resection without requiring large margins or extensive lymphadenectomies.

Methods: A retrospective review was performed of patients who underwent laparoscopic treatment of GISTs by surgeons at the Mount Sinai Medical Center from 2000-2005. Records were reviewed with respect to patient demographics, medical history, diagnostic workup, operative details, postoperative course, and pathologic characteristics.

Results: Laparoscopic surgery was attempted in 43 patients with GISTs. The average age was 65 years and 21 were women. Fifty-six percent of patients presented with anemia or gastrointestinal bleeding. The tumors were located in the stomach (65%) and in the small bowel (35%). The mean tumor sizes were 4.6 cm (stomach) and 3.7 cm (small bowel). Gastric operations included laparoscopic wedge (29%), sleeve (21%), and partial (29%) gastrectomies. The three gastric conversions were due to local invasion of tumor into adjacent organs or proximity to the gastroesophageal junction. Small bowel operations included laparoscopic resections with extracorporeal (47%) and intracorporeal anastamoses (33%). Conversion in small bowel operations was associated with coincidental pathology in addition to the GIST. This consisted of an associated bowel perforation and a synchronous colonic carcinoma. There was one mortality and a 9% morbidity rate, including an evisceration requiring reoperation. All tumors were pathologically confirmed with CD117 immunohistochemistry.

Conclusions: In light of their biologic behavior, GISTs should be considered for laparoscopic resection. This minimally invasive approach to these tumors can be performed safely and reliably.

Key words: Abdominal - Digestive - General - GI SAGES

Although gastrointestinal stromal tumor (GIST) has been recognized clinically for almost two decades, recent advances in understanding its molecular pathogenesis have brought about rapid improvements in its management. The observation that almost all GISTs express CD117 antigen has led to the use of Imatinib for targeted systemic therapy. However, surgery still remains the primary therapy for nonmetastatic GISTs. Total excision of the tumor is the most significant factor for outcome, with 5 year survival rates of 40% to 55% after complete resection [6, 9].

Surgery for GIST involves en bloc resection of the tumor along with any involved structures. Large margins are unnecessary because these tumors usually grow out of the primary organ instead of dilfusely infiltrating. Lymphadenectomy usually is unwarranted because nodal involvement is rare [6, 9]. Typically, wedge resection of the stomach or segmental resection of the intestine is adequate therapy. Laparoscopic surgery may be considered ideal for these tumors because their biologic behavior predisposes them to curative resection without the requirement of large margins or extensive lymphadenectomies. In addition, many of these tumors are diagnosed at pathologic analysis after surgery. Therefore, a diagnostic laparoscopic exploration with excisional biopsy also can be a curative resection. Nevertheless, there is a paucity of data in the literature on the use of minimally invasive surgery for these tumors. This study aimed to investigate the feasibility and safety of laparoscopic surgery for the resection of GIST tumors.

#### Patients and methods

A retrospective review of patients who underwent laparoscopic surgery for GISTs by surgeons at the Mount Sinai Medical Center from 2000 to 2005 was performed. Cases were identified through the use of a

Correspondence to: C. M. Divino

#### Table 1. Presenting symptoms

Anemia Gastrointestinal bleeding Dysphagia/gastrocsophagcal diseasc Abdominal pain Abdominal abscess/perforation Small bowel obstruction Incidental finding at surgery Incidental finding at workup for another illness Stomach (n = 28) Upper Endoscopy Computed Tomography Endoscopic Untrasound Magnetic Resonance Imaging Found at Surgery Push Enteroscopy Capsule Endoscopy Small Bowel (n = 15) Push Enteroscopy Capsule Endoscopy Small Bowel Series Found at Surgery T%

hospital pathologic specimen database. Records were reviewed with respect to patient demographics, medical history, presenting symptoms, diagnostic workup, operative details, postoperative course, and pathologic characteristics. Institutional review board approval was obtained before the study was begun.

#### Results

Laparoscopic surgery was attempted for 43 patients with GISTs during the study period. The average patient age was 65 years (range, 41–92 years), and 22 of the patients were men. Most of the patients presented with signs or symptoms of anemia such as syncope, fatigue, or light-headedness (Table 1). One patient had experienced cryptogenic recurrent minor bleeding for many years, but presented with an acute abdomen attributable to tumor perforation. One-fifth of the GISTs were found incidentally at the workup for other illnesses or at exploration for other causes.

The diagnostic methods used to find the GISTs in this study are outlined in Fig. 1. All the tumors were located in the stomach (67%) or the small bowel (33%). All GISTs originating from the stomach were found using esophagogastroduodenoscopy or computed tomography (CT) scan, except in two patients, whose GISTs were found incidentally at surgery. Patients with small bowel tumors typically underwent numerous diagnostic procedures such as upper and lower endoscopies, contrast studies, capsule studies, and endoscopic ultrasound before the tumors were ultimately found. In two patients presenting with acute bleeding, small bowel tumors were eventually found at urgent exploration after a multitude of negative diagnostic examinations.

Laparoscopic resection was attempted for all the patients (Table 2). The mean gastric tumor size was 4.6 cm (range, 0.4-11.5 cm). Most gastric tumors were removed via laparoscopic wedge resection, partial gastrectomy, or sleeve gastrectomy. These tumors were located at the fundus or along the greater curvature. The largest tumor (11.5 cm), located at the greater curvature, was removed successfully by laparoscopic sleeve resection. Three patients underwent laparoscopic subtotal gastrectomy with gastrojejunostomy for antral tumors. One patient with enlarged retroperitoneal lymph nodes found during exploration underwent laparoscopic lymphadenectomy after gastric sleeve resection of a 8.5 cm fundal tumor. A laparoscopic distal pancreatectomy also was performed in a patient undergoing partial gastrectomy for a tumor invading posteriorly into the pancreas. One cardiac lesion was intraluminally resected via a combined laparoscopic-endoscopic technique.



20%

40%

60%

80%

0%

Laparoscopic intragastric trocars were used to shell out the tumor, and the mucosal defect was endoluminally closed using suture with endoscopic assistance.

Three gastric operations were converted to open surgery (11%). One patient had a cardiac lesion, requiring conversion for an esophagogastrectomy. Another patient had a fundal tumor. However, laparoscopic wedge resection was difficult because of the tumor's proximity to the gastroesophageal junction. The last gastric conversion occurred in the case of a 10.5 cm tumor that had locally invaded the transverse colon and pancreas. After laparoscopic sleeve resection, the case was converted to a distal pancreatectomy, splenectomy, and transverse colectomy.

Laparoscopic surgery was attempted in 15 patients with small bowcl tumors (Table 2). The mean small bowel GIST was 3.7 cm (range, 0.4-8.5 cm) in size. Most of the patients underwent laparoscopic segmental bowel resection with intra- or extracorporeal anastamoses. The largest small bowel tumor (8.5 cm) was successfully removed laparoscopically. One tumor in the proximal ileum had a lengthy serosal stalk, and an excision at the base was performed without bowel resection. Two conversions occurred for the patients with small bowel tumors (13%). One patient underwent a converted ileocolic resection for a colonic carcinoma, and an adjacent GIST tumor was found at pathologic examination. The other conversion involved a patient who underwent a laparoscopic exploration for a spontaneous intraabdominal abscess. The source of the abscess was a perforated GIST, and an open small bowel resection was performed.

The operative and postoperative data are outlined in Table 3. All the tumors were pathologically confirmed to be GISTs by CD117 immunohistochemistry. There were no incidences of tumor rupture or spillage during the laparoscopic operations. However, the large gastric tumor involving the pancreas and colon could not be

Table 2. Operations performed

Stomach $(n = 28)$	n (%)	Small Bowel $(n = 15)$	n (%)
Laparoscopic	25 (89)	Laparoscopic	13 (87)
Wedge resection	8 (29)	SBR-extracorporeal anastamosis	7 (47)
Partial gastrectomy	8 (29)	SBR-intracorporeal anastamosis	5 (33)
Sleeve gastrectomy	6 (21)	Excision at tumor stalk	1 (7)
Subtotal gastrectomy	3 (11)		
Esophagogastrectomy	1 (4)		
Intraluminal excision at stalk	1 (4)		
Converted	3 (11)	Converted	2 (13)
Partial gastrectomy	1	Ileocolic resection	1
Sleeve gastrectomy	1	SBR	1
Esophagogastrectomy	1		

SBR = Small bowel resection

Table 3. Operative and postoperative characteristics

	Range
Median Blood loss	50 cc (20-1000 cc)
Median Operative time	143 min (46-336 min)
Median Length of Stay	4 days (1-50 days)
Morbidity	4 (9%)
Evisceration/Pancreatic stump leak	1
Myocardial Infarction	1
Pneumonia	1
Anastamotic bleed	1
Mortality	1 (2%)
Reoperation	1 (2%)

completely removed en bloc, even after conversion to open surgery.

One death occurred in our group, giving a mortality rate of 2%. The patient who died had undergone a partial gastreetomy for a 7 cm GIST. Early in the postoperative period, a colonic careinoma was diagnosed, and the patient underwent a laparoscopie left hemicolectomy. His recovery was complicated by renal failure and sepsis, leading to death.

Four postoperative complications occurred, giving a morbidity rate of 9%. An anastamotic bleed in a patient with a midjejunal lesion was controlled using push enteroscopy with cauterization. The patient who underwent conversion for sleeve resection, colectomy, pancreatectomy, and splenectomy had to return to the operating room 7 days postoperatively after a coughing spell causing evisceration. At exploration, a pancreatic stump leak also was noted. Drains were placed, and the patient was placed on total parenteral nutrition and bowel rest. The patient eventually recovered and was discharged on postoperative day 50.

#### Discussion

Every GIST is considered to have malignant potential, and complete surgical resection is the primary treatment method. The tumor should be removed en bloc along with any involved structures and organs. Even small tumors (<2 cm) should be approached with aggressive management rather than watchful waiting. Wide margins are not necessary, and lymphadenectomy usually is not required because lymph node involvement is rare [7, 11]. Gentle dissection and avoidance of tumor rupture with subsequent peritoneal seeding are imperative. Therefore, some believe that traditional open surgery should be the mainstay of operative therapy [3].

Our series comprised a group of patients successfully treated with laparoscopic surgery. We believe that laparoscopic techniques may be ideal for GISTs because wide resections and extensive lymphadenectomies usually are not needed. Limited resections with negative margins usually are adequate. Most gastric tumors are in locations accessible by laparoscopic wedge resection or partial gastrectomy, and nearly all small bowel lesions can be removed via laparoscopic small bowel resection. The no-touch technique is imperative to minimize the risk of tumor rupture. We had no cases of tumor rupture, and only one tumor could not be removed en bloe because of local invasion and size.

Few other reports of laparoscopic GIST management exist in the literature. Our study represents the largest series reported to date. Most other studies involve case reports or small series of gastrie submucosal tumors with a proportion of GISTs [2, 4, 5, 15, 17]. One group used laparoscopic wedge resection for 34 gastric submucosal tumors, 14 of which were GISTs [14]. In this series, tumor size usually was small, but limited resections were successfully performed laparoscopically, and there were no reports of tumor rupture. Matthews et al. [12], comparing laparoscopic and open resection of gastric GIST, found equivalent operating time and blood loss between the two groups, but shorter hospital stay in the former group.

Three cases (11%) of gastrie GISTs required conversion. One of these cases required conversion beeause the aforementioned tumor was invading adjacent organs. The remaining cases were converted because of the tumor's proximity to the gastroesophageal junction or because of local invasion of adjacent organs. This is consistent with reports in the literature, which suggest that laparoscopic resection without anastamosis should be avoided near the esophagogastric junction due to the risk of elinically significant deformity or stenosis of the area [1, 16]. Laparoscopic intraluminal resections have been described for tumors in this area, limiting the need for extensive resections. With this technique, transgastric trocars and instruments are used to enucleate submucosal lesions or excise stalked tumors at their mucosal base [16, 18]. One GIST in our series was successfully removed using this laparoscopic transgastric technique. For another patient who underwent conversion, a gastrostomy was made, and the stalked tumor was excised from its intraluminal base.

There is a paucity of literature regarding laparoscopic resection for small bowel GISTs. Furthermore, only case report describes laparoscopic management of small bowel tumors in general [10, 13]. In our series, small bowcl GISTs were resected laparoscopically in 15 cases, with only two conversions. These conversions were not because of difficulty with laparoscopic resection. The one patient had an intestinal perforation of an unknown cause, and the other GIST was found incidentally during an operation for another malignancy. In the remaining cases, laparoscopic small bowel resection was performed without difficulty, even for tumors up to 8.5 cm in diameter. No small bowel tumors ruptured intraoperatively. Despite the lack of data in the literature regarding laparoscopic surgery for small bowel GISTs, our series suggests that laparoscopic resection is feasible for these tumors.

There was acceptable morbidity and mortality in our series. The one death actually was unrelated to the operation for the GIST. It occurred after a secondary operation for another malignancy found in the early postoperative period. Most of the other morbidities were managed conservatively, except in the case of one patient requiring reoperation for evisceration.

Because GISTs usually are not confirmed until pathologic analysis, a preoperative diagnosis is rare. Therefore, clinicians must be suspicious of all gastrointestinal submucosal tumors. Recent consensus reports have stated that these tumors are of uncertain malignant potential, and that all tumors should be resected despite an uncertain diagnosis [3, 8]. Therefore, laparoscopic resection may be ideal in these cases because it prevents morbidity when large laparotomies are performed for lesions of unclear pathology. Limited resections can be performed in a minimally invasive manner to achieve both tissue diagnosis and curative resection. However, little information is known regarding the long-term safety of laparoscopic GIST resection. Longer follow-up evaluation is needed to assess the recurrence rate and the survival of patients undergoing laparoscopic surgery, as compared with traditional open operations.

#### Conclusion

In light of their biologic behavior, GISTs may be good candidates for minimally invasive surgery. Because large margins and extensive lymphadenectomies are rarely indicated, laparascopic resection usually is feasible and can be performed safely. Conversion is associated with proximity of the tumor to the gastroesophageal junction, local tumor invasion into adjacent organs, and coincidental intraabdominal pathology.

#### References

- Aogi K, Hirai T, Mukaida H, Toge T, Haruma K, Kajiyama G (1999) Laparoscopic resection of submucosal gastric tumors. Surg Today 29: 102–106
- Basso N, Rosato P, De Leo A, Picconi T, Trentino P, Fantini A, Silecchia G (2000) Laparoscopic treatment of gastric stromal tumors. Surg Endosc 14: 524–526
- 3. Blay JY, Bonvalot S, Casali P, Casali P, Choi H, Debiec-Richter M, Dei Tos AP, Emile JF, Gronchi A, Hogendoorn PC, Joensuu H, Le Cesne A, Mac Clure J, Maurel J, Nupponen N, Ray-Coquard I, Reichardt P, Sciot R, Stroobants S, van Glabeke M, van Oosterom A, Demetri GD, GIST consensus meeting panelists (2005) Consensus Meeting for the Management of Gastrointestinal Stromal Tumors. Ann Oncol 16: 566– 578
- Buyske J, McDonald M, Fernandez C, Munson JL, Sanders LE, Tsao J, Birkett DH (1997) Minimally invasive management of lowgrade and benign gastric tumors. Surg Endosc 11: 1084–1087
- Cheng HL, Lee WJ, Lai IR, Yuan RH, Yu SC (1999) Laparoscopic wedge resection of benign gastric tumor. Hepatogastroenterology 46: 2100–2104
- Dematteo RP, Lewis JJ, Leung D, Mudan SS, Woodruff JM, Brennan MF (2000) Two hundred gastrointestinal stromal tumors: recurrence patterns and prognostic factors for survival. Ann Surg 231: 51–58
- Eisenberg BL, Judson I (2004) Surgery and Imatinib in the management of GIST: emerging approaches to adjuvant and neoadjuvant therapy. Ann Surg One 11: 465–475
- Fletcher CDM, Berman JJ, Corless C, Gorstein F, Lasota J, Longley BJ, Miettinen M, O'Leary TJ, Remotti H, Rubin BP, Shmookler B, Sobin LH, Weiss SW (2002) Diagnosis of gastrointestinal stromal tumors: a consensus approach. Hum Pathol 33: 459–465
- Joensuu H, Fletcher C, Dimitrijevic S, Silberman S, Roberts P, Demetri G (2002) Management of malignant gastrointestinal stromal tumors. Lancet Oncol 3: 655–664
- Kok KY, Mathew VV, Yapp SK (1998) Laparoscopic-assisted bowel resection for a bleeding leiomyoma. Surg Endosc 12: 995– 996
- Lehnert T (1998) Gastrointestinal sarcoma (GIST): a review of surgical management. Ann Chir Gynaecol 87: 297–305
- Matthews BD, Walsh RM, Kercher KW, Sing RF, Pratt BL, Answini GA, Heniford BT (2002) Laparoscopic vs open resection of gastric stromal tumors. Surg Endosc 16: 803–807
- Napolitano L, Waku M, De Nicola P, Di Bartolomeo N, Aceto L, Innocenti P (2004) Surgical laparoscopic therapy of small bowel tumors: review of the literature and report of two cases. G Chir 25: 235-237
- Otani Y, Ohgami M, Igarashi N, Kimata M, Kubota T, Kumai K, Kitajima M, Mukai M (2000) Laparoscopic wedge resection of gastric submucosal tumors. Surg Laparosc Endosc Percutan Tech 10: 19–23
- Rothlin M, Schob O (2001) Laparoscopic wedge resection for benign gastric tumors. Surg Endosc 15: 893–895
- Tagaya N, Hikami H, Kogurc H, Kubota K, Hosoya Y, Nagai H (2001) Laparoscopic intragastric staple resection of gastric submucosal tumors located near the esophagogastric junction. Surg Endosc 16: 177–179
- Tagaya N, Mikami H, Igarashi A, Ishikawa K, Kogure H, Ohyama O (1997) Laparoscopic local resection for benign nonepithelial gastric tumors. J Laparoendosc Adv Surg Tech A 7: 53–58
- Walsh RM, Ponsky J, Brody F, Matthews BD, Heniford BT (2003) Combined endoscopic/laparoscopic intragastric resection of gastric stromal tumors. J Gastrointest Surg 7: 386–392

Surg Endosc (2006) 20: 717–720 DOI: 10.1007/s00464-005-0402-4

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

# Morbidity of laparoscopic surgery for complicated appendicitis: an international study

J. Cueto,<sup>1</sup> B. D'Allemagne,<sup>2</sup> J. A. Vázquez-Frias,<sup>1</sup> S. Gomez,<sup>3</sup> F. Delgado,<sup>3</sup> L. Trullenque,<sup>3</sup> R. Fajardo,<sup>4</sup> S. Valencia,<sup>5</sup> L. Poggi,<sup>6</sup> J. Ballí,<sup>7</sup> J. Diaz,<sup>7</sup> R. González,<sup>7</sup> J. H. Mansur,<sup>7</sup> M. E. Franklin<sup>8</sup>

<sup>1</sup> Department of Surgery, Hospitals ABC y Angeles Lomas, Mexico City, Mexico

<sup>2</sup> Department of Surgery, St. Joseph Clinic, Liege, Belgium

<sup>3</sup> Hospital Dr. Peset Aleixandre, Valencia, Spain

<sup>4</sup> Clinica de Merida, Merida, Yucatan, Mexico

<sup>5</sup> Department of General Surgery, University of California, Los Angeles, CA, USA

<sup>6</sup> Department of Surgery, Clinica Angloamericana, Lima, Peru

<sup>7</sup> Department of Surgery, Hospital San Jose, Monterrey, N.L., Mexico

<sup>8</sup> Texas Endoscopic Institute, Department of Surgery, University of Texas Health Science Center, San Antonio, TX, USA

Received: 3 June 2005/Accepted: 12 December 2005/Online publication: 16 March 2006

#### Abstract

*Background:* Although laparoscopic appendectomy has some advantages over open appendectomy, some reports do show more postoperative intraabdominal abscesses.

*Methods:* A retrospective review of complicated appendicitis managed surgically by eight surgical groups from six countries was undertaken. Among 3,433 patients with appendicitis, 1,017 (29.5%) had complicated appendicitis, which included perforated or gangrenous appendicitis with or without localized or disseminated peritonitis. There were 74 preoperative abseesses (7.4%) and 5 small bowel obstructions.

*Results:* One patient died. There were 29 postoperative intraabdominal abscesses (2.8%) and 112 mostly minor complications. Conversion to laparotomy was necessary for 28 patients (2.7%). The surgical time ranged from 32 to 132 min (mean, 62 min), and the hospital stay ranged from 1 to 18 days (mean, 3.5 days).

*Conclusions:* The morbidity rates, particularly for intraabdominal absesses, were less for laparoseopic appendectomy in complicated appendicitis than those reported in the literature for open appendectomy, whereas operating times and hospital stays were similar.

Key words: Laparoscopie appendeetomy — Complieated appendieitis — Postoperative intraabdominal abscesses

Correspondence to: J. Cueto

Although not considered the "gold-standard," laparoscopic appendectomy (LA) is widely used, and it is generally accepted that it has several advantages over the eonventional open appendectomy (OA). On the other hand, some studies report that LA results in increased costs, a longer operating time, and more intraabdominal abseesses (1AAs), but these issues remain controversial. In 1995, the European Association for Endoscopic Surgery endorsed the procedure and stated that there was no evidence of increased prevalence of postoperative septic complications, and that the results were directly proportional to the experience and skills of the surgieal group [11]. Several recent metaanalyses and other reports agree with those statements and document the efficacy and safety of LA [2, 6-9, 13, 14, 17-21, 23, 25-31, 33, 34, 36-38, 41, 42]. To doeument the morbidity of LA in complicated appendicitis (CA), an international retrospective study was undertaken.

#### Material and methods

Eight groups from six countries participated in a retrospective review of CA. Among 3,433 patients with appendicitis, 1,017 had CA (29.5% of all appendectomics). Complicated appendicitis rates varied in the series of the different contributors from 13% to 48% (Table 1). There were 511 male and 506 female patients. Complicated appendicitis was diagnosed at the time of operation, pathologic examination, or hoth as gangrenous appendicitis, perforation with local peritonitis (purulent material in the periappendicular area or in the sac of Douglas), or diffuse peritonitis with or without abscess. The complications reported were mortality, residual IAA, port-site cellulitis, localized collections (PI), and reoperations. Duration of the procedure and hospital stay also were analyzed.

Presented in part at the 12th European Association for Endoscopic Surgery (EAES) International Congress, 9–12 June 2004, in Barcelona, Spain

Table 1. Results of laparoscopic appendectomy for complicated appendicitis (CA)

	Total Appys	СА	G	LP	DP	Abscess (preop)	OR time (min) n (range)	Conversion	Hospital stay (days) n (range)	Complications 1AA
D'Allemagne	574	154	52	54	48	0			5.8	9
Fajardo	445	121	21	65	35	II	52 (31-114)	3	2.3 (2-9)	3
Franklin	342	46	22	15	9	0	48 (38-128)	0	1.8 (1-3)	I
Poggi	634	315	144	101	70	22	70 (40-130)	2	2 (2.5-13)	3
TEC's group	537	94	38	40	16	0	57 (35-140)	0	2.6 (1-4)	4
Delgado's group	620	209	107	92	10	30	74 (40-120)	11	5	7
Cucto	281	78	25	40	13	11	60 (30-132)	2	5 (3-18)	2
Total	3,433	1,017	409	407	201	74	57.4 (30-140)	18		29

Appys, appendectomies; G, gangrene; LP, local peritonitis; DP, diffuse peritonitis; OR, operating room; IAA, intraabdominal abscesses

#### Results

Of the 1,017 CA patients, 409 had gangrenous appendicitis, 407 had local peritonitis, and 201 had diffuse peritonitis. Preoperatively, 74 patients had an abscess (7.4%) and 5 had small bowel obstruction.

A 92-year-old woman with diffuse peritonitis died of multiple organ failure. Postoperatively, IAA developed in 29 patients (2.8%). A total of 17 underwent reoperation, with 8 undergoing laparoscopy, 8 requiring laparotomy, and 1 undergoing transrectal drainage. Nine abscesses were drained by CT-guided puncture, whereas three patients with localized small collections received only antibiotics. One fistula developed after percutaneous drainage in a diabetic patient, which healed with conservative treatment.

A total of 112 minor complications (11%) were found. In 78 patients, PI developed, including port-site cellulites and 11 small purulent collections, all of which were treated successfully with drainage, antibiotics, and local wound care on an outpatient basis. A prolonged hospital stay for 13 patients was necessitated by ileus (n = 9), pneumonitis (n = 3), and parenteral nutrition (n = 1). There was conversion to laparotomy for 28 patients (2.7%), and surgical time ranged from 32 to 132 min (mean, 62 min). The hospital stay ranged from 1 to 18 days (mean, 3.5 days), although it must be considered that the criteria for discharge varies in different countries. For noncomplicated appendicitis managed by LA (2,415 patients), the surgical groups reported 4 IAAs that required relaparoscopy. This occurred at the beginning of the laparoscopic experience in the surgical groups.

#### Discussion

Although appendectomy accounts for 6% of all surgical procedures [10] and is among the most common surgical emergency procedures, it has not yet become the "goldstandard" as has laparoscopic cholecystectomy. This might be attributable to the fact that laparoscopic operating rooms, instruments, and nursing staff may not be available or adequate at some hospitals during the night or holiday shifts. As a matter of fact, in many surgical teaching programs, appendectomy, traditionally considered an "easy operation" by many, still is practiced routinely by the conventional method. On the other hand, LA is performed by surgical groups with special interest in mini-invasivc surgery, as mentioned by Kazemier et al. [24]. Nevertheless, the procedure is being used with increasing frequency. In several metaanalysis and prospective studies [2, 6–9, 13, 14, 17–21, 23, 25–31, 33, 34, 36–38, 41], its advantages and benefits have been objectively documented. The two main objections cited for its acceptance are higher costs [5, 18] and increased IAA [12, 16, 35, 39, 43] in CA, as reported by the Cochrane review [40]. In that report, the prevalence of IAA was 2.7% in OA, as compared with 4.7% in LA. These data are in direct conflict with the aforementioned studies.

The results presented in this report support the conclusions of Kazemeier et al. [23, 30], Ball et al. [3], Guller et al. [17] and others [9, 19, 30, 42], showing that 1AA is less frequent after LA than after OA. It might be said that our results are biased because the participating surgical groups had special interest and expertise in laparoscopic surgery, but they nevertheless prove that if the group is properly equipped and trained, LA in CA can be used with extremely low morbidity, offering the advantages already recognized with this approach.

In these series, more abscesses (IAA) were found preoperatively (n = 74, 7.4%) than postoperatively (n = 29, 2.8%) (p < 0.05). In the noncomplicated appendicitis group, LA for 2,715 patients resulted in only 4 postoperative 1AAs, and these reportedly occurred at the beginning of the group's laparoscopic experience. There was one fecal fistula after a CT-guided drainage for an IAA in a 62 year-old diabetic patient with diffuse peritonitis and a pelvic abscess. This fistula closed spontaneously with parenteral nutrition and intravenous antibiotics.

It is precisely in CA that the well-known advantages of LA can benefit a patient: thorough inspection of the entire peritoneal cavity, debridement, irrigation and lavage under direct visualization, avoidance of large abdominal incisions, less immunologic compromise, and fewer pulmonary complications. In a recent review by Novitsky et al. [32], the advantages of laparoscopic surgery over laparotomy in terms of acute inflammatory reaction as well as cell-mediated and peritoneal immunity are emphasized. Whether these advantages result from avoidance of large abdominal incisions, minimal organ manipulation and/or exposure of the abdominal viscera to room air, or decreased postoperative pain and/or pulmonary morbidity remains to be elucidated. The reoperation rate of 1.7% is very low. With regard to postoperative 1AA, all the authors in this study prefer a CT-guided drainage as the first line of treatment. If this fails, a relaparoseopy is the procedure of ehoice.

The prevalence of CA in this group of patients is 28% (13–48% range between the different groups), which is in agreement with elassical and recent studies such as that of Al-Omran et al. [1]. The percentage of patients with CA presenting with gangrenous appendicitis, local peritonitis, and diffuse peritonitis also appears to be in line with the reports for this emergency procedure.

Some of the patients with CA may represent a formidable challenge for exposure, dissection, suturing, and thorough irrigation and lavage. Thus, the procedure cannot be ealled an "easy operation." In the past 15 years, since the introduction and popularization of laparoscopie eholeeysteetomy, we all have experienced a learning curve for complicated endoscopic procedures. Certainly, many eases of CA fit into this eategory. Ball et al. [3] strongly emphasized the faet, supported by observations made in teaching hospitals, that appendeetomies frequently are performed as an emergeney procedure during the night, and if the resident staff does not have the experience, skills, and guidance to perform such surgery, then residual sepsis will follow. Sometimes, when teehnieal difficulties appear, an inexperieneed surgeon will opt for an "early eonversion," and then infection of the wounds may complicate the postoperative period, with direct bearing on the expenses.

On the other hand, in the series described in this report, very few PIs (local wound infections) occurred, a fact consistently reported in all papers comparing OA and LA [2, 5, 15, 16–18, 23, 24, 27, 30, 34, 39, 41, 42]. It is important to retrieve the purulent/gangrenous appendiceal specimens in a sterile bag [44], and to perform trocar-site lavage to prevent septic complications.

In the Coehrane study [40], the reviewers eoneluded that "where surgical expertise and equipment is available and affordable, LA seems to have various advantages over OA." They also concluded that "laparoseopic surgery should routinely be employed at least in special cases, for instance, in young females and in obese patients." An increasing number of surgeons, including the authors of this report, already use LA routinely in all cases of acute appendicitis.

The duration of the operation was 14 min longer for LA in the Cochrane review [40], but many recent reports show similar operating room times, and this finding affects the direct costs of the procedure. Although these may be higher in some instances, the hospital stay, the use of analgesies, and the period of convalescence were less in the current series, all of which must also be considered.

The diagnostie advantage of laparoseopy needs to be stressed because the preoperative etiology eannot be established in 15% to 35% of patients with acute abdomen, as shown in the Coehrane review [40] and in a previous report from our group [7]. This is especially true for women of reproductive age, for individuals at both extremes of life, and for patients eurrently receiving antibioties, steroids, or ehemotherapy. A diagnostie laparoseopy not only establishes or confirms the diagnosis, but also is a very efficient therapeutic tool.

The hospital stay reported earlier must be seen in a special context because the social security systems of some European eountries are at variance with routine practice in the United States. For example, patients in those eountries are not required to eomply with a limited hospital stay, but ean have what would be considered a long hospital stay in our practices. In a recent report [4], LA in CA is followed by a very short hospital stay, sometimes less than 24 h, a faet that our group eonsiders risky because it is our policy to provide at least 48 to 72 h of intravenous antibioties. This is important because a substantial number of OAs, whether performed for noneomplicated appendicitis or CA, result in wound infections and abseesses, which may delay the hospital stay or require readmission and/or prolonged outpatient eare, which must be taken into account when costs are analyzed.

The conversion rate for CA in this study was very low, and this may reflect the experience of the surgeons involved. Before converting to laparotomy in CA, as with any other complicated endoscopic procedures, elinicians must consider the insertion of one or more trocars for additional traction, suction, suturing instruments, and the like. One example of this may be a retroceeal appendix, which frequently in OA requires enlargement of the incision, which subsequently may become infected. Ancedotally, for one patient in our group with an abscess and bowel obstruction, even after placement of an additional trocar, technical difficulties persisted, and Katkhouda et al.'s [22] recommendation of a "finger-assisted procedure" was carried out in a trocar site enlarged to 12 mm, avoiding a laparotomy.

Finally, the definitive proof of the statement that LA in CA has as good if not better results than OA, partieularly with regard to the development of IAA, ean be eonfirmed only in a prospective randomized study.

#### Conclusions

Laparoseopie appendeetomy has special advantages for treating patients with CA. In this study, it is shown that morbidity, particularly IAA, was less than that reported for both OA and LA in other studies. Operating times were similar to those for OA, but the hospital stay was shorter. These findings have led many surgeons to adopt LA because of its diagnostie and therapeutie advantages over OA.

#### References

- Al-Omran M, Mamdani M, McLeod RS (2003) Epidemiologic features of acute appendicitis in Ontario, Canada. Can J Surg 46: 263–268
- Atwood SE, Hill AD, Murphy PG, Thornton J, Stephens RB (1992) A prospective randomized trial of laparoscopic versus open appendectomy. Surgery 112: 497–501
- Ball CG, Kortbeck JB, Kirkpatrick AW, Mitchell P (2004) Laparoscopic appendectomy for complicated appendicitis. Surg Endosc 18: 969–973
- Brosseuk DT, Bathe OF (1999) Day-care laparoscopic appendectomies. Can J Surg 42: 138–142

- Chung RS, Rowland Dy, Li P, Diaz J (1999) A metaanalysis of randomized controlled trials of laparoscopic versus conventional appendectomy. Am J Surg 177: 250–256
- Cox MR, McCall JL, Toouli J, Padbury RT, Wilson TG, Wattchow DA, Langcake M (1996) Prospective randomized comparison of open versus laparoscopie appendicectomy in men. World J Surg 20: 263–266
- Cueto J, Rojas O, Garteiz D, Rodriguez M, Weber A (1997) The efficacy of laparoscopic surgery in the diagnosis and treatment of peritonitis: experience with 107 cases in Mèxico City. Surg Endosc 11: 366–370
- Cueto GJ, Valencia-Reyes MS, Vázquez-Frías JA, Castañeda P, Nevarez R, Weber A (2000) Technical modifications for laparoscopic appendeetomy and other pelvis procedures using microinstruments. Surg Laparose Endose Percutan Tech 10: 211–214
- Cueto J, Vazquez JA (2003) Appendectomy. In: Cueto J, Gagner M, Jacobs M (eds). Laparoscopic surgery. McGraw-Hill, New York, ehap 39, pp 287–294
- De Dombal FT, Clamp SE, Wardle KS (1987) Measuring surgical performance in acute abdominal pain; some reflections from international studies. Epidemiol Rcv 9: 1
- European Association for Endoscopic Surgery (EAES) (1995) Consensus development conferences on laparoscopic cholecysteetomy, appendeetomy, and hernia repair. Surg Endosc 9: 550– 563
- Eypasch E, Sauerland S, Lefering R, Neugebauer EA (2002) Laparoscopic versus open appendectomy: hetween evidence and eommon sense. Dig Surg 19: 518–522
- Fingerhut A, Millat B, Borrie F (1999) Laparoscopic versus open appendectomy: time to decide. World J Surg 23: 835–845
- Frazee RC, Roberts JW, Symmonds RE, Snyder SK, Hendricks JC, Smith RW, Custer MD III, Harrison JB (1994) A prospective randomized trial comparing open versus laparoscopic appendeetomy. Ann Surg 219: 725–731
- Garbutt JM, Soper NJ, Shannon WD, Botero A, Littemberg E (1999) Meta-analysis of randomized controlled trials comparing laparoscopic and open appendectomy. Surg Laparosc Endosc 1: 17–26
- Golub R, Siddiqui F, Pohl D (1998) Laparoscopic versus open appendectomy: a meta analysis. J Am Coll Surg 186: 545–553
- Guller U, Hervey SH, Purves H, Muhlbaier LH, Peterson ED, Eubanks S, Pietrobon R (2004) Laparoscopic versus open appendeetomy: outcomes comparison based on a large administrative database. Ann Surg 239: 43-52
- Hansen JB, Smithers BM, Schache D, Wall D, Miller BJ, Menzies BL (1996) Laparoscopic versus open appendectomy: prospective randomized trial. World J Surg 20: 17–21
- Hart R, Rajgopal C, Plewes A, Sweeney J, Davies W, Gray D, Taylor B (1996) Laparoseopie versus open appendectomy: a prospective randomized trial 81 patients. Can J Surg 39: 457–462
- Hehebrand D, Troidi H, Spangenberger W, Neugebauer E, Schwalm T, Gunther MW (1994) Laparoskopische oder Klassische appendektomie? Eine prospektiv random-isierte studie. Chirurg 65: 112–120
- Hellberg A, Rudberg C, Kullman E, Enochsson L, Fenyo G, Graffner H, Hallerback B, Johansson B, Anderberg B, Wenner J, Ringqvist I, Sorensen S (1999) Prospective randomized multicentre study of laparoscpic versus open appendicectomy. Br J. Surg. 86: 48
- 22. Katkhouda N, Friedlander MH, Grant SW, Achanta KK, Essani R, Paik P, Velmahos G, Campos G, Mason R, Mavor E (2000) Intraabdominal abseess rate after laparoscopic appendectomy. Am J Surg. 180: 456–459; discussion 460–461
- Kazemier G, de Zeeuw GR, Lange JF, Hop WCJ, Bonjer HJ (1997) Laparoscopie vs open appendectomy: a randomized clinical study. Surg Endosc 11: 336–340

- 24. Kazemier G, Steyerberg EA, Bonjer HJ (1997) Meta-analysis of randomized clinical trials comparing open and laparoscopic appendectomy. Scientific Session. The Soeiety of Gastrointestinal Endoscopie Surgeons (SAGES) Meeting of the Americas, San Diego California 19–22 March
- Khalili TM, Hiatt JR, Savar A, Lau C, Margulies DR (1999) Perforated appendicitis is not a contraindication to laparoscopy. Am Surg 65: 965–967
- Klingler A, Henle KP, Beller S, Reehner J, Zerz A, Wetseher GJ, Sziniez G (1998) Laparoseopie appendectomy does not ehange the incidence of postoperative infectious complications. Am J Surg 175: 232–235
- Kum CK, Ngoi SS, Goh PM, Tekant Y, Isaac JR (1993) Randomized controlled trial comparing laparoscopic and open appendectomy. Br J Surg 80: 1599–1600
- Laine S, Rantala A, Gullichsen R, Ovaska J (1997) Laparoscopic appendectomy: is it worthwhile? Surg Endosc 11: 95–97
- Maearulla E, Vallet J, Abad JM, Hussein H, Fernandez E, Nieto B (1997) Laparoscopic versus open appendectomy: a prospective randomized trial. Surg Laparosc Endosc 4: 335–339
- McCall JL, Sharples K, Jadallah F (1997) Systematic review of randomized controlled trials comparing laparoscopic with open appendectomy. Br J Surg 84: 1045–1050
- Moberg AC, Montegomery A (1997) Appendicitis: laparoscopic versus conventional operation: a study and review of the literature. Surg Laparosc Endosc 7: 459–463
- Novitsky YW, Litwin DEM, Callery MP (2004) The net immunologic advantage of laparoscopic surgery. Surg Endosc 18: 1411– 1419
- 33. Ortega AE, Hunter JG, Peters JH, Swanstrom LL, Schirmer B (1995) Prospective, randomized comparison of laparoscopic appendectomy with open appendectomy. Laparoscopic Appendectomy Study Group. Am J Surg 169: 208–213
- Ozmen MM, Zulfikaroglu B, Tanik A, Taner I (1999) Laparoscopic versus open appendectomy: prospective randomized trial. Surg Laparosc Endosc Percutan Tech 9: 187–189
- 35. Paik PS, Towson JA, Anthone GJ, Ortega AE, Simons AJ, Beart RW Jr (1997) Intraabdominal abscesses following laparoscopic and open appendectomies. J Gastrointest Surg 1: 188–193
- Paya K, Rauhofeer U, Rebhandl W, Deluggi St, Horcher E (2000) Perforating appendicitis: an indication for laparoscopy? Surg Endose 14: 182–184
- Reid RI, Dobbs BR, Frizelle FA, Risk factors for postappendectomy intraabdominal abseess. Aust N Z J Surg 199: 373–374
- Reiertsen O, Larsen S, Trondsen E, Edwin B, Faerden AE, Rosseland AR (1997) Randomized controlled trial with sequential design of laparoscopic versus conventional appendicectomy. Br J Surg 84: 842–847
- Sauerland S, Lefering R, Holthausen U, Neugebauer EA (1998) Laparoscopic vs conventional appendectomy: a meta-analysis of randomized controlled trials. Lagenbecks Arch Surg 383: 289–295
- Sauerland S, Lefering R, Neugebauer EA (2004) Laparoscopic versus open surgery for suspected appendicitis. Coehrane Database Syst Rev 18; CD 001546
- Slim K, Pezet D, Chipponi J (1998) Laparoscopic or open appendeetomy: critical review of randomized, controlled trials. Dis Colon Rectum 4: 398–403
- So JB, Chiong EC, Chiong E, Cheah WK, Lomanto D, Goh P, Kum CK (2002) Laparoscopic appendectomy for perforated appendicitis. World J Surg 26: 1485–1488
- Tang E, Ortega AE, Anthone GJ, Bcart RW Jr (1996) Intraahdominal abscesses following laparoscopic and. Appendectomies. Surg Endosc 10: 327–328
- 44. Weber A, Vazquez JA, Valencia S, Cueto J (1998) Retrieval of specimens in laparoscopy using reelosable zipper-type plastic bags: a simple, cheap and useful method. Surg Laparosc Endosc 8: 457–459

Surg Endosc (2006) 20: 721–725 DOI: 10.1007/s00464-005-0583-x

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

## Utility of staging laparoscopy in subsets of biliary cancers

Laparoscopy is a powerful diagnostic tool in patients with intrahepatic and gallbladder carcinoma

D. Goere,<sup>1</sup> G. D. Wagholikar,<sup>1</sup> P. Pessaux,<sup>1</sup> N. Carrère,<sup>1</sup> A. Sibert,<sup>2</sup> V. Vilgrain,<sup>2</sup> A. Sauvanet,<sup>1</sup> J. Belghiti<sup>1</sup>

<sup>1</sup> Department of Hepato-biliary Surgery, Hospital Beaujon, 100, bd du Gl Leclerc, 92110 Clicby, France <sup>2</sup> Department of Radiology, Hospital Beaujon, 100, bd du Gl Leclerc, 92110 Clicby, France

Received: 25 August 2005/Accepted: 26 October 2005/Online publication: 27 February 2006

#### Abstract

*Background:* The aim of this study was to evaluate the utility of staging laparoscopy in patients with biliary cancers in the era of modern diagnostic imaging.

*Methods:* From September 2002 through August 2004, 39 consecutive patients with potentially resectable cholangioearcinoma underwent preoperative staging laparoseopy before laparotomy. Preoperative imaging included ultrasonography and triphasic computed tomography for all patients and magnetic resonance cholangiography in 35 patients (90%). Final pathologieal diagnosis included 20 hilar cholangiocarcinomas (HC), 11 intrahepatic cholangiocarcinomas (IHC), and eight gallbladder carcinomas (GBC).

Results: During laparoscopy, unresectable disease was found in 14/39 patients (36%). The main causes of unreseetability were peritoneal eareinomatosis (11/14) and liver metastases (5/14). At laparotomy, nine patients (37%) were found to have advanced disease precluding resection. Vascular invasion and nodal metastases were the main causes of unresectability during laparotomy (eight out of nine). In detecting peritoneal metastases and liver metastases, laparoseopy had an accuracy of 92 and 71%, respectively. All patients with vascular or nodal involvement were missed by laparoscopy. For prediction of unresectability disease, the yield and accuracy of laparoscopy were highest for GBC (62% yield and 83% accuracy), followed by 1HC (36% yield and 67% aceuraey) and HC (25% yield and 45% accuracy)

*Conclusion:* Staging laparoseopy ensured that unneeessary laparotomy was not performed in 36% of patients with potentially resectable biliary careinoma after extensive preoperative imaging. In patients with biliary careinoma that appears resectable, staging laparoseopy allows detection of peritoneal and liver metastasis in one third of patients. Both vascular and lymph nodes invasions were not diagonsed by this procedure. Due to these limitations, laparoscopy is more useful in ruling out dissemination in GBC and 1HC than in HC.

Key words: Biliary earcinoma — Laparoscopy — Staging

Biliary eaneers are aggressive tumors with poor prognosis, resulting in formidable diagnostic and therapeutie challenges. Resection is the only potentially curative treatment. Complete tumor extirpation is paramount for long-term survival. The majority of patients have advaneed disease at presentation, precluding radical exeision. Despite improvements in preoperative imaging modalities, the resectability rate has not increased beyond 40-60% [6, 9, 13, 15, 18]. Laparoscopy has been proposed as a staging modality to identify patients with occult disseminated disease that is missed on preoperative evaluation and thereby avoid the drawbacks of performing unnecessary laparotomy [8, 14, 24]. Laparoseopy resulted in decreased hospital stay and expenses and reduces delay to referral for palliative eare [7]. The yield of laparoscopy could vary according to the type of tumor and the preoperative investigations performed for staging the disease. The aim of this study was to evaluate the utility of staging laparoscopy in patients with biliary eancers in the era of modern diagnostic imaging and to identify patients with oecult advanced disease.

#### Materials and methods

From September 2002 through August 2004, 39 consecutive patients with potentially resectable biliary cancers underwent preoperative staging laparoscopy at our institution. Patients with potentially

Correspondence to: J. Belghiti

resectable intrahepatic cholangiocarcinoma (IHC), hilar cholangiocarcinoma (HC), and gallhladder carcinoma (GBC) were included in the study. There were 25 mcn and 14 women, with a median age of 56 years (range, 26-76). Chest and abdominal triphasic computed tomography (CT) with vascular reconstruction was performed in all patients. A magnetic resonance cholangiopancreatography (MRCP) was obtained in most patients and in some cases allowed more precise determination of the degree of biliary tree involvement than could be obtained with CT. When a vascular invasion (vena porta, hepatic artery, and their proximal branches) was suspected on CT scan or MRI. an angiography was performed. Cholangiography with biliary drainage (either endoscopic or percutaneous transhepatic) was performed in patients with biliary obstruction to decrease the serum bilirubin level below 50 µmol/L. Preoperative portal embolization was performed in patients who needed extensive liver resection (remnant liver volume < 30% of total liver volume). Clinicopathological data and radiological investigations were examined in a weekly multidisciplinary conference, and only patients ascertained to have potentially resectable disease were included in the study. Pathological diagnosis was confirmed in all patients pre-, intra-, or postoperatively. Previous upper abdominal surgery was not a contraindication for inclusion in the study

Staging laparoscopy was performed immediately prior to laparotomy in eight patients; for the other patients, laparotomy was delayed until the biliribun level was <50 µmol/L. For patients in whom portal vein embolization was planned as a preparation for resection, laparoscopy was performed prior to embolization. Staging laparoscopy was performed according to the standard described technique. Pneumoperitoneum was performed with open technique through an umbilical incision. Insufflation was performed with CO<sub>2</sub> gas to a pressure of 15 mmHg. A 0° laparoscope was used. Additional ports were placed to facilitate liver retraction and perform adhesiolysis when necessary. A complete laparoscopic examination entailed satisfactory assessment of all the greater sac structures: the diaphragmatic and undersurface of the liver, the lesser and greater omentum, the scrosal surface of the hollow viscera, and parietal peritoneum of the pelvis and anterior abdominal wall. In the presence of ascites, peritoneal fluid was harvested for cytology. All doubtful lesions located on the peritoneum or on the liver far away from the main lesion were biopsied and sent for frozen section examination. If cytopathology confirmed malignancy, the procedure was terminated. Peritoned cytology was not performed due to its slight additional values [12].

Patients with no evidence of dissemination at laparoscopy underwent laparotomy and resectability was confirmed. An aggressive approach was adopted for resections to achieve a tumor-free resection margin. Our surgical philosophy and approach toward patients with IC and those undergoing liver resections have been previously reported [2, 22].

Clinicopathological details, investigation records, operative findings, and postoperative outcome in patients with unresectable diseases were entered prospectively in a database and analyzed. Yield of laparoscopy was calculated as the ratio of patients identified by laparoscopy harboring occult advanced disease to the number of patients undergoing the procedure. Accuracy of laparoscopy was the proportion of patients with unresectable disease who were diagnosed by laparoscopy.

#### Results

#### Preoperative imaging and pathological diagnosis

Concerning preoperative imaging, all patients underwent ultrasonography and chest and abdominal CTs. Thirty-five patients (90%) had a MRCP. Vascular examination was completed by arteriography in nine patients (23%). Endoscopic or transhepatic biliary opacification was performed in patients with HC and GBC and was completed by a preoperative biliary drainage (eight internal and 18 external) in 26 patients (66%). The mean number of preoperative investigations per patient was 3.9. Final diagnosis was confirmed by pathology after resection or by biopsy if palliative treatment was performed and included 20 HCs, 11 IHCs, and eight GBCs.

#### **Operative** findings

Laparoscopic examination was completed in all except one patient. This patient with HC, who had undergone previous upper abdominal surgery, had dense adhesions to the anterior abdominal wall hindering satisfactory examination, and exploration was performed by limited laparotomy. Findings at laparoscopy and laparotomy are outlined in Fig. 1. Fourteen patients (36%) had evidence of advanced disease on laparoscopy and 25 patients were eligible for laparotomy. One patient without advanced disease on laparoscopy underwent portal vein embolization before the planned laparotomy, but 4 weeks later ascites and an aspect of peritoneal carcinomatosis were diagnosed by CT scan and precluded laparotomy. The remaining 24 patients (63%) underwent laparotomy. At laparotomy, nine patients (37%) were found to have advanced disease precluding resection, and 15 (63%) underwent resection of the tumor. Finally, 23 patients had unresectable disease; laparoscopy thus correctly identified 14/23 patients (61%) who had unresectable disease.

The details of findings responsible for unresectability are summarized in Table 1. Laparoscopy correctly identified peritoneal metastases in all except one patient; thus, the accuracy of laparoscopy in detecting peritoneal metastases was 92%. The only patient in whom laparoscopy failed had localized tumor deposits in the pelvis. In identifying liver metastases, laparoscopy had an accuracy of 71% (five out of seven patients). One patient with a large IC in the right liver underwent right portal vein embolization to induce hypertrophy in the remnant liver after staging laparoscopy ruled out dissemination. However, at exploration 40 days later, he had metastases in the left lobe of the liver and regional lymph nodes. Although laparoscopy was not repeated prior to exploration, this was considered a failure of diagnostic laparoscopy (by intent-to-treat analysis). Laparoscopy also failed to detect liver metastases in the patient in whom dense adhesions required conversion to laparotomy for exploration. All patients with vascular invasion or metastases to distant lymph nodes were missed on laparoscopy. Vascular invasion was diagnosed during laparotomy in five patients. In two of these patients, vascular invasion was suspected during laparoscopy, but laparotomy was necessary to obtain a pathological diagnosis.

The yield and accuracy of laparoscopy werc highest for GBC (62% yield and 83% accuracy), followed by IHC (36% yield and 67% accuracy) and HC (25% yield and 45% accuracy), as listed in Table 2.

Before laparoscopy, portal vein embolization was scheduled to prepare the resection in 10 patients. Among these patients, unresectable disease was diagnosed by laparoscopy in five; five patients had a preoperative portal vein embolization and three underwent liver resection.



Table 1. Factors precluding resection identified by laparoscopy or by laparotomy<sup>a</sup>

Factors precluding resection	Identified at laparoscopy	Identified only at laparotomy	Accuracy of laparoscopy (%)
Cirrhosis	1	0	1/1 (100)
Peritoneal metastases	11	1	11/12 (92)
Liver metastases	5	2	5/7 (71)
Adjacent organ infiltration	1	f	1/2 (50)
Advanced nodal spread	0	3	0/3 (0)
Vascular invasion	0	5	0/5 (0)

<sup>a</sup> Some patients had multiple factors responsible for unresectability

#### Postoperative stay and complications

For the 15 patients undergoing only laparoscopy, the median postoperative stay was 4 days (range, 2–24). One patient with IC and advanced disease diagnosed by laparoscopy developed postoperative aspiration pneumonia resulting in a hospital stay of 24 days. For the nine patients with unresectable disease diagnosed at laparotomy, the median stay was 11 days (range, 7–23). There were no procedural complications attributable to laparoscopy.

#### Discussion

In the current study, the overall accuracy of laparoscopy in correctly identifying patients with unresectable discase was 61%. This result was obtained despite an extensive and homogeneous preoperative assessment including a mean of 3.9 imaging procedures per patient. Both yield and accuracy of laparoscopy varied markedly among the subgroups of biliary cancers. It was highest for GBC, followed by IHC and HC. Laparoscopy identified all but one case of peritoneal dissemination (92% accuracy), and it identified 71% of patients with liver metastases. The ability to identify unresectable disease due to lymph nodal and vascular invasion was more limited with laparoscopy. Fig. 1. Flowchart showing findings at laparoscopy and laparotomy. \*The delay between staging laparoscopy and reevaluation by imaging after a portal vein embolization was 4 weeks.

 Table 2. Yield and accuracy of laparoscopy in detecting unresectable disease in individual cancers

Type of tumor	п	Overall yield (%)	Accuracy (%)
GBC	8	62 (5/8)	83 (5/6)
IHC	11	36 (4/11)	67 (4/6)
HC	20	25 (5/20)	45 (5/11)
Entire series	39	38 (14/39)	61 (14/23)

GBC, gallbladder carcinoma; IHC, intrahepatic cholangiocarcinoma; HC, hilar cholangiocarcinoma

A large number of patients with biliary cancers have advanced disease at presentation. Survival in patients who have incomplete tumor extirpation is identical to that in patients who do not undergo surgery and receive only palliative therapies [5, 13, 15]. Hence, the importance of avoiding a nontherapeutic laparotomy cannot be overemphasized. Despite improvements in imaging modalities, the incidence of nontherapeutic laparotomy in patients with proximal biliary cancers remains high, from 25 to 46% in the most recent series, and results in increased hospital stay and potential morbidity in addition to greater treatment expenses [9, 13, 18]. Staging laparoscopy has been proposed as an important tool to identify occult dissemination that can be missed on preoperative imaging and avoid unnecessary explorations.

The yield and accuracy of staging laparoscopy depend on the type of cancer. Gallbladder carcinoma has a propensity for peritoneal spread and liver metastases that can be easily identified by laparoscopy. In an analysis of 100 patients with biliary cancers, almost half of the GBC patients had advanced disease on laparoscopy [25]. Vascular invasion and lymph node metastases of HC, which represent the major cause of unresectability, are often missed during staging laparoscopy. In the study of the Memorial Sloan-Kettering Cancer Center, the overall yield for detecting unresectable disease in HB was significantly lower than that obtained for GBC. Therefore these authors suggested performing laparoscopy only in patients with HC with a high risk of unresectable disease, such as T2 or T3 lesions [25]. There are few data in the literature on the role of laparoscopy in patients with IHC, a less common biliary cancer but with an increasing incidence. In our study, more than one-third of patients with IHC had advanced disease at laparoscopy and were spared an unnecessary laparotomy. Laparoscopy had an accuracy of 67% in identifying advanced disease. Staging laparoscopy should be integral in the management protocol of patients with IHC. Groups advocating aggressive surgery in cases of IHC have also changed their policy and started using laparoscopy to assess the resectability [18].

Laparoscopic ultrasound (LUS) has been proposed as a complementary investigation to obviate some of the limitations of laparoscopy and has added to the yield in patients with pancreatic and liver tumors [4, 6, 19, 21, 23]. In the current series, LUS was not routinely used and would not have been helpful in the diagnosis of liver metastases. Indeed, in two cases liver metastases were diagnosed by laparotomy performed because of adhesions in one case and 40 days after laparoscopy in the other. Also, it is difficult to identify vascular invasion and lymph nodal involvement on LUS. In the series of van Delden et al. [23], who were the first to study the use of LUS in proximal biliary cancers, nine of 31 patients (28%) had occult disease at laparoscopy and only one patient with unresectable disease was detected solely on the basis of LUS. Tumor adherence to major vascular structures and extensive biliary involvement by cancer are often difficult to determine radiographically. Accuracy may be further limited by the presence of biliary stents and secondary inflammation [25]. Furthermore, metastatic lymph nodes are difficult to diagnose based on ultrasound findings and mandate biopsy. Conversely, in a study from Edinburgh, UK, the yield of staging laparoscopy in HC was considerably increased by LUS, from 24 to 42%, by detecting local advanced disease, whereas accuracy in detecting liver metastases was not increased [6]. Also, the authors emphasized the long period (from 1992 to 2003) during which this study was conducted and advancements of preoperative imaging during this time, which could explain the considerable yield of LUS observed.

The strategy of preoperative external biliary drainage and portal vein embolization is increasingly being adopted in jaundiced patients who need major liver resection to reduce the risk of postoperative liver failure and improve surgical outcome [17]. The high incidence of occult advanced disease in potentially resectable patients highlights the need to perform laparoscopy before preoperative intervention for optimization is performed. Our study shows that early detection of dissemination by laparoscopy can avoid unnecessary invasive procedures and lead to immediate institution of palliative care.

Detection of advanced disease by laparoscopy not only helped in avoiding the pain and morbidity of an unnecessary laparotomy but also resulted in a shorter hospital stay. This is important for patients with unresectable biliary cancers, for whom the median survival is only a few months. Early discharge from the hospital improves quality of life and helps in an earlier institution of systemic chemotherapy [1, 11, 20]. Also, pathological malignancy is easier to confirm by harvesting an unsuspected liver metastases during laparoscopy than by obtaining tissue samples of HC by either transhepatic or endoscopic maneuvers [10, 16].

In summary, laparoscopy helped avoid unnecessary laparotomy in more than one-third of patients, resulting in significantly shorter hospital stay. Laparoscopy is accurate in detecting peritoneal and liver metastases but misses the majority of patients with vascular invasion and lymph nodal disease. Due to these limitations, laparoscopy is more useful in ruling out dissemination in GBC and IHC than in HC. We recommend the routine use of laparoscopy in all patients with potentially resectable GBC and IHC and selectively in patients with large HC, or when prior invasive procedures for optimization, such as portal vein embolization, are contemplated.

#### References

- Alberts SR, Al-Khatib H, Mahoney MR, Burgart L, Cera PJ, Flynn PJ, Finch TR, Levitt R, Windschitl HE, Knost JA, et al. (2005) Gemcitabine, 5-fluorouracil, and leucovorin in advanced biliary tract and gallbladder carcinoma: a North Central Cancer Treatment Group phase II trial. Cancer 103: 111–118
- Belghiti J, Hiramatsu K, Benoist S, Massauli P, Sauvanet A, Farges O (2000) Seven hundred forty-seven hepatectomies in the 1990s: an update to evaluate the actual risk of liver resection. J Am Coll Surg 191: 38–46
- Burke EC, Jarnagin WR, Hochwald SN, Pisters PW, Fong Y, Blumgart LH (1998) Cholangiocarcinoma: patterns of spread, the importance of hepatic resection for curative operation, and a presurgical clinical staging system. Ann Surg 228: 385–394
- Callery MP, Strasberg SM, Doherty GM, Soper NJ, Norton JA (1997) Staging laparoscopy with laparoscopic ultrasonography: optimizing resectability in hepatobiliary and pancreatic malignancy. J Am Coll Surg 185: 33–39
- 5. Chamberlain RS, Blumgart LH (2000) Hilar cholangiocarcinoma: a review and commentary. Ann Surg Oncol 7: 55-66
- Connor S, Barron E, Wigmore S, Madhavan K, Parks R, Garden J (2005) The utility of laparoscopic assessment in the preoperative staging of suspected hilar cholangiocarcinoma. J Gastrointest Surg 4: 476–480
- Cuschieri A (2001) Role of video-laparoscopy in the staging of intra-abdominal lymphomas and gastrointestinal cancer. Semin Surg Oncol 20: 167–172
- D'Angelica M, Fong Y, Weber S, Gonen M, DeMatteo RP, Conlon K, Blumgari H, Jarnagin WR (2003) The role of staging laparoscopy in hepatobiliary malignancy: prospective analysis of 401 cases. Ann Surg Oncol 10: 183–189
- Dixon E, Vollmer CM, Sahajpal MC, Grant D, Doig C, Hemming A, Taylor B, Langer B, Greig P, Gallinger S (2005) An aggressive surgical approach leads to improved survival in patients with gallbladder cancer. Ann Surg 241: 385–394
- Domagk D, Poremba C, Dietl KH, Senninger N, Heinecke A, Domschke W, Menzel J (2002) Endoscopic transpapillary biopsies and intraductal ultrasonography in the diagnostics of bile duct strictures: a prospective study. Gut 51: 240–244
- 11. Ducreux M, Van Cutsem E, Van Laethem JL, Gress TM, Jeziorski K, Rougier P, Wagener T, Anak O, Baron B, Nordlinger B;EO-RTC Gastro Intestinal Tract Cancer Group (2005) A randomised phase II trial of weekly high-dose 5-fluorouracil with and without folinic acid and cisplatin in patients with advanced biliary tract carcinoma: results of the 40955 EORTC trial. Eur J Cancer 41: 398–403
- Fernandez-del Castillo C, Rattner DW, Warshaw AL (1995) Further experience with laparoscopy and peritoneal cytology in the staging of pancreatic cancer. Br J Surg 82: 1127–1129

- Hemming AE, Reed AI, Fujita S, Foley D, Howard RJ (2005) Surgical management of hilar cholangiocarcínoma. Ann Surg 241: 693–702
- Jarnagin WR, Bodniewicz J, Dougherty E, Conlon K, Blumgart LH, Fong Y (2000) A prospective analysis of staging laparoscopy in patients with primary and secondary hepatobiliary malignancies. J Gastrointest Surg 4: 34–43
- Jarnagin WR, Fong Y, DeMatteo RP, Gonen M, Burke EC, Bodniewicz J, Youssel M, Kimstra D, Blumgart MH (2001) Staging, resectability, and outcome in 225 patients with hilar cholangiocarcinoma. Ann Surg 234: 507–517
- Jung GS, Huh JD, Lee SU, Han BH, Chang HK, Cho YD (2002) Bile duct: analysis of percutaneous transluminal forceps biopsy in 130 patients suspected of having malignant biliary obstruction. Radiology 224: 725–730
- Kawasaki S, Imamura H, Kobayashi A, Noike T, Miwa S, Miyagawa S (2003) Results of surgical resection for patients with hilar bile duct cancer: application of extended hepatectomy after biliary drainage and hemihepatic portal vein embolization. Ann Surg 238: 84–92
- Lang H. Sotiropoulos G, Fruhauf N, Domland M, Paul A, Kind EV, Malago M, Broelsch CE (2005) Extended hepatectomy for intrahepatic cholangiocellular carcinoma. When is it worthwhile? Single center experience with 27 resections in 50 patients over a 5-year period. Ann Surg 241: 134–143
- Lo CM, Lai EC, Liu CL, Fan ST, Wong J (1998) Laparoscopy and laparoscopic ultrasonography avoid exploratory laparotomy

in patients with hepatocellular carcinoma. Ann Surg 227: 527-532

- Malik IA, Aziz Z, Zaidi SH, Sethuraman G (2003) Gemeitabine and cisplatin is a highly effective combination chemotherapy in patients with advanced cancer of the gallbladder. Am J Clin Oncol 26: 174–177
- van Nieveen Dijkum EJ, de Wit LT, van Delden OM, Kruyt PM, van Lanschot JJ, Rauws EA, Obertop H, Gouma DJ (1999) Staging laparoscopy and laparoscopic ultrasonography in more than 400 patients with upper gastrointestinal carcinoma. J Am Coll Surg 189: 459–465
- Valverde A, Bonhomme N, Farges O, Sauvanet A, Flejou JF, Belghiti J (1996) Resection of intrahepatic cholangiocarcinoma: a Western experience. J Hepatobiliary Pancreat Surg 6: 122– 127
- van Delden OM, de Wit LT, van Nieveen Dijkum EJ, Smits NJ, Gouma DJ, Reeders JW (1997) Value of laparoscopic ultrasonography in staging of proximal bile duct tumors. J Ultrasound Mcd 16: 7–12
- Vollmer CM, Drebin JA, Middleton WD, Teefey SA, Linchan DC, Soper NJ, Eagon CJ, Strasberg SM (2002) Utility of staging laparoscopy in subsets of peripancreatic and biliary malignancies. Ann Surg 235: 1–7
- Weber SM, DeMatteo RP, Fong Y, Blumgart LH, Jarnagin WR (2002) Staging laparoscopy in patients with extrahepatic biliary carcinoma. Analysis of 100 patients. Ann Surg 235: 392– 399

Surg Endosc (2006) 20: 726–729 DOI: 10.1007/s00464-005-0196-4

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

## Laparoscopic ultrasound navigation in liver surgery: technical aspects and accuracy

M. Kleemann,<sup>1</sup> P. Hildebrand,<sup>1</sup> M. Birth,<sup>2</sup> H. P. Bruch<sup>1</sup>

<sup>1</sup> Department of Surgery, University Hospital of Schleswig-Holstein-Campus Lübeck, Ratzeburger Allee 160, 21358 Lübeck, Germany
<sup>2</sup> Department for General, Visceral, Thoracic, and Vascular Surgery, Hanse Klinikum Stralsund, Große Parower Straße 47-53, 18435 Stralsund, Germany

Received: 7 April 2005/Accepted: 25 October 2005/Online publication: 16 March 2006

#### Abstract

The functional-anatomic structure of the liver according to Couinaud classification based on the intrahepatical course of the vascular structures is the basis of all modern liver surgery. Consequently, the use of intraoperative ultrasound is an undisputed requirement for every liver resection. Exact following of the planned resection plane can be realized only with the application of permanent online navigation based on intraoperative ultrasound during the dissection of the hepatical tissue. Now that the authors have established ultrasound navigated resection in open liver surgery using a navigated parenchymal dissecting instrument, they intend to transfer this technique from open to laparoscopic liver surgery. A special adapter was developed to connect an ultrasound-based navigation system to laparoscopic instruments. The authors present the first results in terms of technical aspects and feasibility.

Key words: Laparoscopic surgery — Laparoscopic ultrasound — Liver — Navigation

Despite recent advances in laparoscopic techniques and instrumentation, laparoscopic liver surgery still is limited to a select patient population. One major reason may be the lack of tactile sensation and orientation during dissection of the liver parenchyma. Laparoscopic fenestration of solitary giant liver cysts has been reported [4]. Laparoscopic resections are feasible and safe for selected patients with left-sided or right peripheral lesions requiring limited resection [2] (Table 1).

Although laparoscopic resection of metastatic liver diseasc seems feasible, this approach still is debated [7]. The safe anterolateral segments according Couinaud classification are shown in Fig. 1. However, direct translation of the information received sonographically into the resection procedure can cause difficulties, especially with segment/sector ectomies. After the course of the vessels has been projected onto the liver capsule in accordance with the ultrasound picture, dissection of the hepatic tissue itself is performed currently without the support of pictures. As a result, there could be significant deviations in the planned resection plane (Fig. 2). Exact following of the planned resection plane can be realized only by application of a permanent online navigation based on intraoperative ultrasound during dissection of the hepatic tissue.

Now that we have established an ultrasound-navigated system for open liver surgery with online navigation of the dissection instrument, we will use this technique also in laparoscopic surgery to navigate under laparoscopic ultrasound control (e.g., during interventional ablation procedures or liver resections).

#### Materials and methods

The US-Guide 2000 is an independent navigation system compatible with all ultrasound machines. It is based on an electromagnetic tracking system with six degrees of freedom [5]. On the system monitor, the ultrasound B-picture, overlaid by the navigation data in real time received from the ultrasound and transferred into the navigation system, is displayed. The position finding necessary for the navigation is based on a calculation of distance and angle in accordance with the common satellite navigation. A newly developed adapter allows the navigation system to be combined with a laparoscopic ultrasound probe (B-K Medical 8566, Denmark, Fig. 3a and b). The computer is connected to a transmitter and two magnetic sensors. The weak magnetic field created by the transmitter has to cover the abdominal part in which the intervention is performed. At the same time, the sensors fixed to the head of the laparoscopic ultrasound probe, the interventional or dissection instrument, must remain within the magnetic field throughout the whole procedure. The navigation system

Correspondence to: M. Kleemann

Table 1. Literature overview of laparoscopic liver surgery

Autbor	Patients (n)	Duration (min)	Conversion rate (%)	Complication rate (%)	Hospital duration (days)
D 1009 (101	17	102 5	5.0	11.0	7.0
Rau 1998 [10]	1 /	183.5	5.9	11.8	7.8
Cherqui 2000 [2]	30	214	6.6	20	9.6
Mouicl 2000 [9]	30	79	10	20	
Fong 2000 [6]	11	248	55	40	9.2
Berends 2001 [1]	10	180	20	0	6
Tang 2002 [11]	11	190	9	9	11.3
Gigot 2002 [7]	37		13.5	22	7
Descottes 2003 [3]	87		10	5	5



Fig. 1. Anterolateral segments 11, 111, 1Vb, V, and VI (sbaded grey) according to Couinaud classification. Adapted from Gigot et al. [8].

recognizes the position and anterograde orientation of the needle in relation to the laparoscopic ultrasound head and target structure, then projects this virtually onto the ultrasound picture. Additional functions of the ultrasound machine (e.g., duplex sonography) are available during the intervention and can be used to recognize and avoid vessels.

First, the adapter is placed at the bead of the laparoscopic ultrasound probe to connect the electromagnetic tracker to the adapter. The nearer the adapter can he placed to the tip of the instrument, the bigher the accuracy of the system. For calibration with an ultrasound phantom, the distance between the adapter and the ultrasound probe must be determined and calibrated with the software of the navigation system. Then the other tracker is placed at a laparoscopic dissection instrument huilt for laser dissection and calibrated as mentioned earlier. In phantom testing and in a liver organ model, the virtual resection line then is overlaid to the laparoscopic ultrasound B-mode picture, offering the possibility of navigated ablation or resection. Second, the system is integrated in a liver organ model to detect disturbances attrihutable to trocar and camera instruments.

#### Results

For transmission of the ultrasound picture data, the ultrasound machine and the navigation system must be connected by an interface. For correct calculation and display of the course followed by the resection line, it is necessary to define the length of the hepatic dissection instrument (i.e., distance from the instrument tip to the sensor position). For the navigated resection, the US-Guide 2000 is positioned close to the area of intervention. The two magnetic sensors, connected by cable to the navigation system, are then attached to the laparoscopic ultrasound head with special adapters. The transformer adapter that takes on the transformer sensor is a small plastic mount connected to the ultrasound head. The transformer sensor is attached to it in the same manner as a connecting clamp. The ultrasound Bmode picture is then presented with the overlaid navigation data (Fig. 4a). We used metal troears up to 25 mm in size to bring the system into the abdominal cavity (Fig. 4b).

Laparoscopic navigation under ultrasound guidance is technically feasible in this model. Even when the tip of the ultrasound probe was angulated, no disturbances of the navigation system were obvious, due to the closc approximation of the laparoscopic ultrasound head and electromagnetic sensor. Anatomic landmarks in liver tissue could be safely reached. No interaction of the electromagnetic tracking system and the laparoscopic equipment (e.g., trocar and laparoscopic camera) could be seen.

#### Conclusion

In this report, we describe the use of a laparoscopic navigation system with permanent sonographically guidance. Therefore, the exact following of a planned resection level can be transferred online to the liver organ. This improves the precision of laparoscopic liver dissection and may lead to an improvement in the quality of the operation. Laparoscopic navigation under ultrasound guidance offers a new technique and tool for the visceral surgeon. Especially in laparoscopic surgery, this method may improve orientation in interventions or resections in liver surgery. Our preliminary results show the feasibility of this technique in the field of laparoscopic surgery. To date, the size of the electromagnetic sensors are limiting the minimally invasive use of navigation because the sensors in our studies still measure 8  $\times 8 \times 6$  mm. Further studies investigating accuracy and reproducibility in the laparoscopic operation field are necessary for evaluation of this new technique.





Fig. 2. Deviation of planned (red line) vs real (yellow lines) resection line without navigation.



Fig. 3. a Adapter with mounted tracking sensor placed as near as possible to the tip of the laparoscopic ultrasound probe. b Adapter with mounted tracking sensor at the head of the laparoscopic ultrasound probe. The total diameter of the system is 17.5 mm.

#### References

- 1. Berends FJ, Meijer S, Prevoo W, Bonjer HJ, Cuesta MA (2001) Technical considerations in laparoscopic liver surgery. Surg Endose 15:794-798
- 2. Cherqui D, Husson E, Hammoud R, Malassagne B, Stephan F, Bensaid S, Rotman N, Fagniez PL (2000) Laparoscopic liver resections: a feasibility study in 30 patients. Ann Surg 232: 753-762
- 3. Descottes B, Glineur D, Lachachi F, Valleix D, Paineau J, Hamy A, Morino M, Bismuth H, Castaing D, Savier E, Honore P, Detry O, Legrand M, Azagra JS, Goergen M, Ceuterick M, Mareseaux J, Mutter D, de Hemptinne B, Troisi R, Weerts J, Dallemagne B, Jehaes C, Gelin M, Donckier V, Aerts R, Topal B, Bertrand C, Mansvelt B, Van Krunekelsven L, Herman D, Kint M, Totte E, Schockmel R, Gigot JF (2003) Laparoscopic liver resection of benign liver tumors. Surg Endose 17:23-30


Fig. 4. a Ultrasound B-mode picture of the liver with the navigation line (*trajectory*) of the dissection instrument. The yellow circlc indicates the point at which the dissection line intersects with the B-mode picture plane (out-of-plane function). b Laparoscopic ultrasound prohe (B-K Medical 8566) connected to the navigation system through a 22 mm trocar in an experimental environment. Problems of sterilization are not yet solved.

- Diez J, Decaud J, Gutierrez L. Suhl A, Merello J (1998) Laparoscopic treatment of symptomatic cysts of the liver. Br J Surg 85: 25
- Flock of Birds. Technical description of DC Magnetic Trackers. Ascension Technology Corporation, Ascension Technology Corporation POB 527, Burlington, VT 05402, 2000
- Fong Y, Jarnagin W, Conlon KC, De Matteo R, Dougherty E, Blumgart LH (2000) Hand-assisted laparoscopic liver resection: lcssons from an initial experience. Arch Surg 135:854–859
- Gigot JF, Glincur D, Santiago Azagra J, Goergen M, Ceuterick M, Morino M, Etienne J, Marescaux J, Mutter D, van Krunckelsven L, Descottes B, Valleix D, Lachachi F, Bertrand C, Mansvelt B, Huhens G, Saey JP, Schockmel R (2002) Laparoscopic liver resection

for malignant liver tumors: preliminary results of a multicenter European study. Ann Surg 236: 90-97

- 8. Köckerling F, Schwartz SI (2001) Liver Surgery. Operative Techniques and Avoidance of Complications J.A. Barth, Heidelberg
- Mouiel J, Katkhouda N, Gugenheim J, Fahiani P (2000) Possibilities of laparoscopic liver resection. J Hepatobiliary Panereat Surg 7: 1–8
- Rau HG, Buttler E, Meyer G, Schardey HM, Schildherg FW (1998) Laparoscopic liver resection compared with conventional partial hepatectomy—a prospective analysis. Hepatogastroenterology 45: 2333–2338
- Tang CN, Li MK (2002) Laparoscopic-assisted liver resection. J Hepatobiliary Pancreat Surg 9: 105–110

Surg Endosc (2006) 20: 730–733 DOI: 10.1007/s00464-005-0456-3

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

## Advantages of advanced laparoscopic systems

J. Heemskerk,<sup>1</sup> R. Zandbergen,<sup>2</sup> J. G. Maessen,<sup>2</sup> J. W. M. Greve,<sup>1</sup> N. D. Bouvy<sup>1</sup>

<sup>1</sup> Department of Surgery, Maastricht University Hospital, P. DeByelaan 25, PO Box 5800, 6202 AZ Maastricht, The Netherlands
<sup>2</sup> Department of Cardio-Thoracic Surgery, Maastricht University Hospital, P. DeByelaan 25, PO Box 5800, 6202 AZ Maastricht, The Netherlands

Received: 25 June 2005/Accepted: 6 November 2005/Online publication: 9 March 2006

#### Abstract

*Background:* Conventional laparoscopy offers great benefits to our patients, but suffers from major technical drawbacks. Advanced laparoscopic systems are being developed adressing some of these drawbacks.

*Methods:* We performed a training-box based study, performing laparoscopic tasks using conventional laparoscopy and advanced laparoscopic systems in order to assess the influence of these technical drawbacks in order to predict where the biggest advantages of newly developed surgical systems can be expected.

*Results:* The most significant technical drawbacks were two-dimensional vision, disturbed cye-hand target axis and (possibly to a lesser extent) the rigid instruments with a limited five degrees of freedom.

*Conclusion:* Major advances in advanced laparoscopy might only be expected using console-based robot-arm manipulated systems like the daVinci surgical system, or a combination of a high-quality 3-dimensional vision system, restoration of the eye-hand-target axis and the use of an advanced handheld instrument offering seven degrees of freedom such as the Radius surgical system.

Key words: Advanced laparoscopy — Degrees of freedom — Robot — Da Vinci — Fulcrum — Radius — Stereoscopy

Minimal invasive surgery is one of the great advances in medicine in recent decades, aiming at maximal reduction of surgical trauma. However, laparoscopic surgeons sacrifice dexterity to provide patients with less invasive surgery. The following are major drawbacks of laparoscopy:

- 1. Two-dimensional (2D) vision using a conventional monitor reduces perception of depth.
- 2. A disturbed eye-hand-target axis decreases ergonomics and dexterity.

- 3. The long, inflexible instruments used in laparoscopic surgery magnify the surgeon's natural hand tremor.
- 4. The rigid instruments with five degrees of freedom limit the surgeon's natural range of motion, decreasing dexterity.
- 5. Fixed abdominal entry points result in limited freedom of motion and movement of the tip of the instrument to the opposite direction of the outer part of the instrument, a technical drawback known as the fulcrum effect.
- 6. Camera instability increases fatigue.
- 7. Limited tactile feedback decreases dexterity.

These factors probably all contribute to the relatively long learning curve in laparoscopic surgery [15].

Advanced stereoscopic and instrument manipulating surgical systems are being developed in order to address some of the shortcomings related to conventional laparoscopy, potentially leading to faster and more accurate laparoscopy [6, 7]:

- 1. A variety of stereoscopic systems are being developed. Although stereoscopy rarely offers convincing depth perception [9], its use might improve laparoscopic performance [2, 8, 12, 16, 18].
- 2. The disturbed eye-hand-target axis is difficult to restore using conventional laparoscopic equipment. Although ergonomic monitor placement is crucial, the ideal situation of projecting the image exactly where the operation takes place is difficult to achieve without a console-based surgical system [1, 4].
- 3. Tremor can be diminished using a robot arm manipulated system with tremor filtration [11].
- 4. Both handheld and console-based surgical systems offer the full seven degrees of freedom, increasing dexterity [11, 17].
- 5. The fulcrum effect is difficult to address using conventional laparoscopic instruments. Although extensive training leads to faster automation to the fulcrum effect [3, 10], only robot arm manipulated

Correspondence to: J. Heemskerk

systems can restore intuitive movement of the instrument's tip in the direction of the surgeon's hand, increasing dexterity.

- 6. Camera instability due to exhausting camera holding can be restored using a variety of mechanical or robot arm manipulated systems.
- 7. No commercially available surgical systems have been able to restore normal sensitivity in tactile feedback.

We performed a training box-based study, describing time consumption and accuracy in both inexperienced users and expert laparoscopic surgeons performing laparoscopic tasks using conventional laparoscopy, the Radius Surgical System (Tuebingen Scientific, Tuebingen, Germany), and the da Vinci Surgical System (Intuitive Surgical, Mountain View, CA, USA) in a variety of settings. The aim of the study was to assess the significance of the previously described technical drawbacks for laparoscopic surgery in order to predict where the major advantages of newly developed surgical systems can be expected.

#### Materials and methods

#### Participants

Ten inexperienced and 10 experienced volunteers were selected to perform laparoscopic tasks using various laparoscopic systems. The inexperienced group consisted of 10 volunteers without any previous laparoscopic experience. The experienced group consisted of 10 expert laparoscopic and thoracoscopic surgeons from the Departments of Surgery and Cardio-Thoracic Surgery of Maastricht University Hospital. All of them had extensive experience in laparoscopy or thoracoscopy, having performed more than 100 laparoscopic or thoracoscopic procedures.

#### Conventional laparoscopy

All conventional laparoscopic tasks were performed using a pelvic trainer with one 12 mm video port and two 12 mm trocar ports (Versaport, US Surgical Corporation, Norwalk, CT, USA). A 10 mm, 0° digital video camera (Endoeye, Olympus, Hamhurg, Germany) was used, and the image was displayed on a 14-in. high-resolution 100 Hz monitor. Camera handling was done using a simple rigid standard. Manual laparoscopic drills were performed using disposable 5 mm laparoscopic instruments (Endo Clinch II, AutoStutre, Norwalk, CT, USA) and a 5 mm laparoscopic needle driver (Storz 26173SC laparoscopic needle driver, Karl Storz Endoskope, Tuttlingen, Germany).

#### Radius surgical system

All Radius-assisted laparoscopic tasks were performed in the same pelvic trainer using the same trocars described previously. The Radius handheld manipulator (Radius Surgical System) was used instead of conventional laparoscopic instruments. This laparoscopic instrument enables the surgeon to perform laparoscopic tasks offering the full seven degrees of freedom instead of the five degrees of freedom in conventional laparoscopy, potentially increasing dexterity and improving performance. Unfortunately, the tip of the Radius instrument was not suitable to grasp the heads used in task 1. Therefore, this task was not done using the Radius Surgical System.

#### da vinci surgical system

All da Vinci-assisted laparoscopic tasks were performed in the same pelvic trainer as described previously using a 12 mm video port and two 7 mm trocars. We used three arms of the four-armed da Vinci Surgical System. This robotic surgical system consists of a surgeon's console, patient side cart, Endowrist instruments, and InSite Vision System. The surgeon's console offers an ergonomic position to the surgeon, translating the surgeon's intuitive movements into precise, real-time movements of the instruments. The patient side cart offers four robot arms, executing the surgeon's commands while offering tremor filtration and movement downscaling if desired. The Endowrist instruments attached to the patient side cart offer the full seven degrees of freedom. The InSite Vision System provides high-quality stereoscopic stable vision, projecting the tips of the instruments where the fingertips of the surgeon are located.

Three arms of the four-armed da Vinci system were used. One arm handled the camera and the other two arms manipulated two da Vinci laparoscopic Dehakey forceps. The da Vinci tasks were performed using the da Vinci Surgical System in stereoscopic 3D InSite vision, in 2D InSite vision, and in a conventional monitor-viewed modus.

#### Tasks

Three laparoscopic tasks were devised to test dexterity, two-handed coordination, and suturing. Each participant was instructed about the main features of the endoscopic tasks to be performed and on how to use the surgical systems. The participants were allowed to manipulate each surgical system for 5 min to become familiar with the controls and setup. Questions were allowed before and during the tests, hut no assistance was provided. The same order of tasks was performed for every participant, hut the sequence of the use of the different surgical systems changed in order to prevent a learning curve from interfering with the results.

#### Task 1: pick up and drop

A comparable laparoscopic drill was used in other studies [5, 13, 14], in which a receptacle (40 mm opening and 10 mm high) containing five beads was used. The task was to pick up a bead from the receptacle with the right-handed instrument and transfer it halfway to a second receptacle. The bead had to be taken over with the left-handed instrument and dropped inside the second receptacle. Time was recorded from starting position with the instrument in focus hut outside the initial receptacle to the fifth bead dropped into the final receptacle. Inaccuracy was defined as 10 points for every head accidentally dropped outside the receptacle. The task was performed eight times —twice per suitable instrument. The tip of the Radius instrument was unfit for this task.

#### Task 2: cap the needle

This task was performed as described previously [5] using a 19-gauge  $\times$  1.5-in. aspiration needle with Luer Lock (Terumo Europe NV, Leuven, Belgium) and its cap. The task was to cap the needle after grasping both pieces from the floor of the training hox, keeping both cap and needle above the box floor. Time was recorded from starting position with the instruments in focus but 5 cm from the needle and its cap to the moment when the needle and cap were securely coupled and held by one instrument. Inaccuracy was defined as 10 points for every cap or needle accidentally dropped or touching the box floor. The task was performed 10 times—twice per instrument.

#### Task 3: suturing and knot tying

This task has previously heen described [5, 14, 19], and it consists of using a size 8 latex glove and a Vicryl 3–0 polyglactin suture with FS-1 24 mm 3/8 circular needle (Johnson & Johnson, New Brunswick, NJ, USA). The task was to pass the needle through two separated 5 mm dots on the glove and then tie a double knot. Time was recorded from

	Task	Conventional	Radius	daVinci 3-D	daVinei 2-D	daVinci-M
Time	Task 1, first trial	171 (233,108)		85 (108,62)	96 (111,82)	140 (189,90)
	Task 1, second trial	121 (167,74)		65 (74.56)	92 (107.76)	118 (143,94)
	Task 2, first trial	138 (220,55)	100 (156,43)	40 (44,35)	64 (74,55)	105 (121,89)
	Task 2, second trial	103 (158,49)	62 (80,44)	55 (40,70)	64 (52,77)	109 (137,80)
	Task 3, first trial	332 (500,164)	304 (477,132)	122 (165,79)	158 (192,124)	182 (238,127)
	Task 3, second trial	323 (515,132)	240 (335,148)	87 (105,69)	155 (200,110)	186 (251,120)
Inaceuracy	Task 1, first trial	18 (28,7)		9 (13,4)	10 (16,3)	24 (44,4)
	Task 1, second trial	11 (17,5)		6 (6,5)	9 (16,2)	14 (26,2)
	Task 2, first trial	23 (37,9)	14 (20,8)	15 (10,20)	23 (21,24)	30 (26,34)
	Task 2, second trial	16 (27,4)	9 (9,8)	17 (12,22)	24 (19,28)	37 (37,36)
	Task 3, first trial	23 (44,2)	27 (41,13)	0 (0,0)	4 (7, 1)	7 (11, 2)
	Task 3, second trial	27 (46,7)	24 (45,2)	2 (3,0)	5 (9,0)	12 (24,0)

Table 1. Mean time consumption and inaccuracy scores in the total group (inexperienced, experienced subgroups)

starting position with the instruments in focus but 5 cm away from the needle to the moment when the suture was securely tied. Inaccuracy was defined as 10 points per millimeter distance between the black dot and the needle entry through the glove. Twenty points was added if the knot was too loose or the suture broke. The task was performed 10 times—twice per setup.

#### Statistical analysis

Data were stored in an Excel XP database (Microsoft, Redmond, WA, USA) and analyzed using SPSS version 11.0.1 (SPSS, Chicago, IL, USA). Comparison of groups was done using Pearson's chi-square test. Comparison of two related samples was done using a nonparametric Wilcoxon signed rank test. A p value  $\leq 0.05$  was defined as statistically significant.

#### Results

#### Study population

The median age of the study population was 32 years (range, 21–52), 36 years in the experienced group (range, 32–52) and 23 years in the inexperienced group (range, 21–35). In total, 20 participants performed two tasks twice using five different setups. One task was performed twice using four setups, leading to a total of 560 tasks. Performing every task, time consumption, and accuracy were registered, leading to a total of 1,120 analyzable data points.

#### Time and accuracy

Time consumption was compared by performing different tasks using various instruments and setups. Inexperienced participants took substantially more time to complete a task than the experienced surgeons. Conventional surgery was most time-consuming, whereas the da Vinci system with stereoscopy was the fastest. Task 3 was far more time-consuming than tasks 1 and 2. The benefit of using advanced surgical systems seemed less for experienced users compared to inexperienced users.

Accuracy was compared using the different surgical systems. Higher numbers of failures and mistakes resulted in higher inaccuracy scores. Inexperienced participants had higher inaccuracy scores than expert surgeons. Conventional laparoscopy and the use of the da Vinci system with monitor-viewed vision resulted in the highest inaccuracy scores, whereas use of the da Vinci system with stereoscopic InSite vision resulted in the lowest inaccuracy scores and thus the best results. Table 1 shows mean time consumption and inaccuracy scores for the total group and for the inexperienced and experienced subgroups separately.

#### Comparing instruments

Using the Radius Surgical System, two tasks were performed twice each, describing time consumption, accuracy, and score. This resulted in 12 data samples. The other instruments were used to perform three tasks twice each, describing time, accuracy and score. This resulted in 18 data samples. Using a nonparametric Wilcoxon signed rank test, related samples could be compared in order to assess significant superiority of one setup or the other. Results are depicted in Table 2.

#### Discussion

Conventional laparoscopy suffers from seven technical drawbacks as described previously. This study was conducted in order to assess these drawbacks.

The role of 2D vision was assessed by comparing results of da Vinci in 2D InSite vision mode with da Vinci in stereoscopic InSite vision mode. The da Vinci system in 3D mode seemed faster and more accurate in all 12 data samples (Table 1). This difference was significant in five of 12 data samples (Table 2), suggesting high-definition stereoscopic vision does indeed lead to faster and better performance of laparoscopic tasks.

The role of the disturbed eye-hand-target axis was assessed by comparing the results of da Vinci in the 2D InSite vision mode with da Vinci in the monitor-vicwed mode. Two-dimensional InSite vision seemed faster and more accurate in all 12 data samples (Table 1). This difference was significant in five of 12 data samples (Table 2), suggesting restoration of the disturbed eychand-target axis does improve performance of laparoscopic tasks. Table 2. Number of significantly better results comparing first instrument versus second instrument (number of comparing data samples)

	Total group	Inexperienced	Experienced
Conventional vs Radius	0 vs 1 (8)	0 vs 2 (8)	0 vs 0 (8)
Conventional vs daVinci 3-D	0 vs 9 (12)	0 vs 10 (12)	0 vs 1 (12)
Conventional vs daVinci 2-D	0 vs 5 (12)	0 vs 8 (12)	1 vs 0 (12)
Conventional vs daVinci-M	1 vs 0 (12)	0 vs 2 (12)	2 vs 0 (12)
Radius vs daVinei 3-D	0 vs 5 (8)	0 vs 7 (8)	0 vs 2 (8)
Radius vs daVinci 2-D	1 vs 4 (8)	0 vs 5 (8)	1 vs 0 (8)
Radius vs daVinci-M	1 vs 1 (8)	1 vs 1 (8)	3 vs 0 (8)
daVinci 3-D vs daVinci 2-D	5 vs 0 (12)	3 vs 0 (12)	4 vs 0 (12)
daVinci 3-D vs daVinci-M	12 vs 0 (12)	10 vs 0 (12)	6 vs 0 (12)
daVinci 2-D vs daVinci-M	5 vs 0 (12)	4 vs 0 (12)	0 vs 0 (12)

The role of limited degrees of freedom in conventional laparoseopy was assessed by comparing conventional laparoseopy with the Radius Surgical System. Although the tip of the Radius instrument was not fit for task 1, the other two tasks seemed to be performed faster and more accurate using the Radius instrument in seven of eight data samples (Table 1). This suggests that offering seven degrees of freedom might improve laparoseopie performance.

The roles of tremor enhancement, fulerum effect, and limited tactile feedback could not be assessed separately in this study. However, comparing the Radius system, which suffers from tremor enhancement, fulcrum effect, and limited tactile feedback, with the da Vinci system in monitor-viewed mode, no significant difference was present (Tables 1 and 2). This suggests these three drawbacks do not play an important role and might not require a technical solution.

The role of eamera instability was not assessed in this study since camera holding was done by a rigid standard or robot arm.

Analysis of the experienced and inexperienced subgroups showed a more significant advantage of the use of advanced laparoscopic systems in the inexperienced group. This suggests that extensive training in laparoscopy may reduce the need for advanced stereoscopic or manipulating laparoscopic systems.

We conclude that the most significant improvements with regard to the previously mentioned technical drawbacks in conventional laparoscopy are high-definition 3D vision, restoration of the disturbed eye-hand-target axis, and (possibly to a lesser extent) the use of instruments offering the full seven degrees of freedom. Major improvements in laparoscopic surgery may only be expected from either a console-based surgical system, such as the da Vinei Surgical System, or a combination of a high-definition 3D vision system with ergonomic monitor placement (or a head-mounted display), with a handheld, seven degrees of freedom instrument, such as the Radius Surgical System.

Advanced laparoscopic surgery is in its infaney, and major improvements in the availability of specifically designed surgical systems are expected soon, offering great opportunities for the future. However, more research is needed in order to develop affordable and feasible instruments offering high-quality 3D vision, a restoration of the eye-hand-target axis, and seven degrees of freedom.

#### References

- Berguer R, Forkey DL. Smith WD (2001) The effect of laparoscopic instrument working angle on surgeons' upper extremity workload. Surg Endosc 15: 1027–1029
- Cheah WK, Lenzi JE, So J, et al. (2001) Evaluation of a headmounted display (HMD) in the performance of a simulated laparoscopic task. Surg Endosc 15: 990–991
- 3. Crothers IR, Gallagher AG, McClure N, et al. (1999) Experienced laparoscopic surgeons are automated to the "fulcrum effect": an ergonomic demonstration. Endoscopy 31: 365–369
- 4. Emam TA, Hanna G, Cushieri A (2002) Ergonomic principles of task alignment, visual display, and direction of execution of laparoscopic bowel suturing. Surg Endose 16: 267–271
- Garcia-Ruiz A, Gagner M, Miller JH, et al. (1998) Manual versus robotically assisted laparoscopic surgery in the performance of basic manipulation and suturing tasks. Arch Surg 133: 957–961
- Hanly EJ, Talamini MA (2004) Robotic abdominal surgery. Am J Surg 188(Suppl): 19S-26S
- Hazey JW, Melvin WS (2004) Robot-assisted general surgery. Semin Laparose Surg 11: 107–112
- Herron DM, Lantis JC 2nd, Maykel J, et al. (1999) The 3-D monitor and head-mounted display. A quantitative evaluation of advanced laparoscopic viewing technologies. Surg Endose 13: 751–755
- Hofmeister J, Frank TG, Cushieri A, et al. (2001) Perceptual aspects of two-dimensional and stereoscopic display techniques in endoscopic surgery: review and current problems. Semin Laparosc Surg 8: 12–24
- Jordan JA, Gallagher AG, McGuinan J, et al. (2000) Randomly alternating image presentation during laparoscopic training leads to a faster automation to the "fulcrum" effect. Endoscopy 32: 317–321
- Moorthy K, Munz Y, Dosis A, et al. (2004) Dexterity enhancement with robotic surgery. Surg Endosc 18: 790–795
- Mueller MD, Camartin C, Dreher E, et al. (1999) Three-dimensional laparoscopy. Gadget or progress? A randomized trial on the efficacy of three-dimensional laparoscopy. Surg Endosc 13: 469–472
- Prasad SM, Maniar HS, Soper NJ, et al. (2002) The effect of robotic assistance on learning curves for basic laparoscopic skills. Am J Surg 183: 702–707
- Scott DJ, Young WN, Tesfay ST, et al. (2001) Laparoscopic skills training. Am J Surg 182: 137–142
- Subramonian K, DeSylva S, Bishai P, et al. (2004) Acquiring surgical skills: a comparative study of open versus laparoscopic surgery. Eur Urol 45: 346–351
- Taffinder N, Smith SG, Huber J, et al. (1999) The effect of a second-generation 3D endoscope on the laparoscopic precision of novices and experienced surgeons. Surg Endose 13: 1087–1092
- Usui S, Inoue H, Yoshida T, et al. (2004) Preliminary report of multi degrees of freedom forceps for endoscopic surgery. Surg Laparosc Endosc Percut Techn 14: 66–72
- van Bergen P, Kunert W, Buess GF (1999) Three-dimensional (3-D) video systems: bi-channel or single-channel optics? Endoscopy 31: 732-737
- Yohannes P, Rotariu P, Pinto P, et al. (2002) Comparison of robotic versus laparoscopic skills: is there a difference in learning curve? Urology 60: 39–45

Surg Endosc (2006) 20: 734–738 DOI: 10.1007/s00464-004-2247-7

© Springer Science+Business Media, Inc. 2006

SUNU/GA/

and Other Interventional Techniques

# The anatomical significance and techniques of laparoscopic rectal surgery

L. Chengyu, J. Xiaoxin, Z. Jian, G. Chen, Y. Qi

Beijing Fuxing Hospital, Capital University of Medical Science, Beijing 100038, China

Received: 28 October 2004/Accepted: 10 June 2005/Online publication: 16 March 2006

#### Abstract

*Background:* Because there are difficulties associated with the technique of laparoscopic colorectal surgery, thorough knowledge of the anatomy is particularly important. We pay close attention to anatomical features during laparoscopic rectal surgery. In this study, we analyze the association of the anatomy with the operative procedure.

*Methods:* Laparoscopic rectal surgery was performed on 117 patients (66 men) with benign and malignant diseases in the rectum by the complete laparoscopy or hand-assisted technique. All operations were mainly performed by the first author. The association between anatomy and the operation was analyzed.

*Results:* The mean operative time was 144 min (range, 87–235). The hand-assisted technique was performed in two patients. Four patients required conversion to laparotomy due to the amount of fat in three patients and disruption of the Endo-stapler in one patient, for a conversion rate of 1.7%. Operative blood loss was small, averaging 126 ml (range, 50–350). No injury of the ureters, major bleeding in front of the sacrum, or other operation-related severe complications occurred during or after operation. In one case, dissecting disrupted the anterior left wall of the rectum.

*Conclusion:* By mastering the anatomical features of laparoscopic rectum surgery, operative mistakes and complications can be reduced. Particular attention must be paid to the anatomy of the obese patient undergoing laparoscopy. It is very convenient that the corresponding skills can be applied in the course of dissection and exposure.

Key words: Rectum — Surgical procedure — Laparoscopy — Anatomy Laparoscopic colon surgery was successfully applied by Fowler and Jacobs in 1990. Since then, the technique has developed rapidly, and the feasibility of laparoscopic colorectal surgery has been demonstrated.

Prospective and retrospective studies have shown the safety and feasibility of the operation. However, the operative range for the colorectum is extensive, not just cholystectomy, which is rather limited in some regions. The blood supply and adjacent tissues of the colorectum are complicated and often variable. The operation of the colorectum involves not only simple resection but also reconstruction of the intestinal tract. Moreover, it involves the radical problem of malignancy. These factors result in technical difficulties of laparoscopic colorectum surgery [3, 6, 10]. Therefore, a detailed understanding of the anatomy is very important. A total of 117 patients with benign and malignant diseases of the rectum were laparoscopically operated on by us since 2001. We gained detailed knowledge of the anatomical features and their association with the operative procedure.

#### Materials and methods

One hundred and seventeen patients with rectal diseases underwent laparoscopic surgery. All operations were mainly performed by the first author. There were 66 men and 51 women, with a median age of 50.4 years (range, 17–76). There were four rectal tubulovillous adenomas with severe atypical hyperplasia and three rectal tubulovillous adenomas with severe atypical hyperplasia and focal cancer certified by histologic examination. We used the GEN 300 ultrasonic knife (Johnson & Johnson) and Ligasure vessel sealing system (Valleylab). Hand-assisted kits were the hand-port (Smith & Nephew)and Lap Dise (Johnson & Johnson).

#### Results

The operative period ranged from 87 to 235 min, with an average of 144 min. The hand-assisted kits were used in two operations. Four patients were converted to open operation; thus, the conversion rate was 1.7%. Obesity

Correspondence to: L. Chengyu



Fig. 1. The left ureter lies lateral to the ovarian vessel above the pelvic entrance.

resulted in three operative conversions, and one conversion was due to the breakdown of the laparoseopic Endo-stapler so that the lower reetum could not be transected.

Blood loss during the operation was minimal, with an average of 126 ml (range, 50–350). No injury of the ureter or major bleeding in front of the sacrum and other operation-related complications occurred during or after the operations. Only in one case did dissection disrupt the anterior left wall of the rectum.

An anastomatic leak occurred in one patient 5 days after operation, and it was treated by intermittent washing and continuous suction through a drainage tube. This patient with reetal cancer and liver metastasis (a 64-year-old male) had been treated by interventional therapy of the liver and general chemotherapy for more than 3 months. Then, he was transferred to surgery for operation.

One patient (a 76-year-old male) died due to major bleeding of a stress ulcer. The bleeding continued even though multiple treatments were performed. His family deeided to stop treatment on day 8 after the operation. This patient had had blood hypertension disease for more than 20 years. He took enterie-coated aspirin for approximately 3 years, but it was stopped 1 week before the operation.

There were also two cases of pneumonia, one case of deep vein thrombosis in the lower limb, and one case of heart failure after operation. However, they were all cured by corresponding treatments.

#### Discussion

Laparoseopic colorectal surgery has become more standardized in the course of continuous praetice. Ten of thousands of procedures have been performed and followed up for more than 10 years. The standardized operative procedure is sufficient to reach the surgical aim. Moreover, its minimally invasive advantages cannot be obtained by open operation [1, 2, 5, 8].



Fig. 2. The left ureter runs across the end of the common iliac artery at the pelvic entrance.

Laparoscopic colorectal surgery was first performed in 1993 in China [10, 11]. To date, it has been used in approximately 50 large hospitals, mainly on malignant tumors. Due to continuous improvements in the operative technique and the application of the ultrasonic knife and Ligasure vessel scaling system, inexperienced surgeons can make full use of the experience of others and skip the "learning curve" of laparoscopic colorectal surgery. The fact that there were no operative techniquerelated complications in our 117 patients further proves the clinical feasibility of this kind of operation.

Identifying the urcter is the first important step in laparoseopie reetal surgery. We conventionally expose the left ureter. First, the white filmy adhesion between the lateral leaf of the mesosigmoid and the peritoneum in the left iliae gutter is snipped. Then, the lower loosely eonnective tissuc is further dissected. The testicular (ovarian) vessel passes lateral to the ureter. The left ureter is often near the root part of the mesosigmoid. The left and right ureters go aeross the end of the common iliac artery and the origin of lateral iliae artery, respectively, at the pelvie entrance. Its peristalsis can be seen while the ureter is poked by dissecting forceps (Figs. 1-3). After entering the pelvis, it first desecnds along the lateral wall and then turns anteromedially at the level of the seiatie thorn. In the male, the ureter reaches the fundus of the bladder posteroinferior to the defercnt duct. In the female, the ureter runs anteriomedially in the base of the broad ligament, crossing the uterine artery beneath the artery 2 em lateral to the uterine cervix and then reaches the bladder wall. Because the right ureter is more lateral, it does not need to be exposed. In most eireumstanees, the right urcter can be seen going down at the pelvie entrance through the peritoneum, especially in nonobese patients (Figs. 4 and 5).

The inferior mesenteric artery originates from the anterior wall of the lower part of the aorta at approximately the level of L3, 3 or 4 cm below the aortie bifureation, and then running in a lower left direction. It divides into the left colie artery superiolateral, the sigmoid artery lateral, and superior rectal artery

Fig. 3. The left external iliac artery which is variable presses over the ureter.

patient.

Fig. 5. The right ureter can be seen through the peritoneum in another

Fig. 4. The right ureter can be seen through the peritoneum.

downward. When the assistant or the left hand of the operator elevates the upper part of the mesosigmoid using the grasper in the left superior direction, the inferior mcsenteric artery will be drawn straight. Then, the pedicle of the inferior mesenteric artery can be generally seen. The root of the inferior mesenteric artery can easily be dissected at the base of the pedicle and then cut (Fig. 6).

The superior rectal artery enters the pelvic cavity through the mesosigmoid and descends left and anterior to the sacral promontory. It divides the left and right branches in the upper part of the rectum at the level of S3, and then runs down to the rectum posteriorly to laterally (Fig. 7). Although the mesorectum is cut with the ultrasonic knife, the branches in the mesorectum bleed easily, especially in patients who are older and those with blood hypertension, coronary heart disease, and arteriosclerosis whose vessels are hard and fragile. The cut end of the bleeding vessels can be clamped with the dissecting forceps to stop the bleeding by electric coagulation. If Ligasure is used, the chance that major bleeding will occur is largely reduced.

Fig. 6. The inferior mesenteric artery is drawn straight. The pedicle of the inferior mesenteric artery can be seen.

The middle rectal artcry originates from the anterior branch of the internal iliac artery and enters into the lower part of the rectum through the rectal lateral ligament. It can be cut near the rectal wall with the ultrasonic knife or Ligasure.

In obese women, especially those who are short, the fat accumulation makes the mesorectum thick and the peritoneum at the anterior and lateral walls hangs loosely into the pelvic cavity. Therefore, the pelvic cavity in these patients becomes smaller and narrow so that exposure to the pclvic cavity is more difficult to obtain (Fig. 8). Performing the anus-preserving procedure in low rectal cancer is very difficult. Use of the hand-assisted procedure for keeping off the fat blockage or the immediate conversion to open operation must often be performed, and these do not represent failure of the operation.

After the peritoneal reflection before the rectum is incised, there are the bladder, the ampulla of the deferent duct, and the seminal vesicle in the upper part and the prostate in the lower part in this rectovesical septum in the male. In the female, the plane called the









Fig. 7. The inferior mesenteric artery runs down into the superior rectal artery.



Fig. 9. The posterior wall of the vagina can be seen by the guiding method through the vagina.



**Fig. 8.** In a short, obese woman, the mesorectum is thick, and the peritoneum of the pelvic cavity at the anterior and lateral walls hangs loosely into the pelvic cavity.

rectovaginal septum is looser than the male's, so it is easily dissected. The guiding method through the anus or vagina can often achieve twice the results with half the effort (Fig. 9). The assistant should place one linger into the anus or vagina or both (binanual examination). The primary operator touches the wall of the reetum or vagina with the head of the dissecting forceps, meeting the assistant's fingers in the anus or vagina, to seek the dissecting plane indirectly. The disruption of the left anterior wall of the reetum in one case by dissecting occurred due to the faulty assistant's guide at the perineum in the early stage of the operation.

While dissecting the presacral space, the spacious and bright field of vision under laparoscopy significantly improves the situation of the narrow and difficult to expose postreetal space in the conventional open operation. The blunt dissection by the operator's fingers is often needed in the conventional open operation, and it is sometimes performed in a semiblinded condition. The



Fig. 10. The spacious and bright field of vision in front of the sacrum under laparoscopy.

elear plane for dissecting under laparoscopy largely reduces the dangerous occurrence of major bleeding of the presacrum (Fig. 10). On the other hand, the two bundles of the hypogastric nerve plexus can be much more easily seen in laparoscopy than in the open operation, which is very helpful for preserving the autonomic nerve and the bladder, and thus preserving sexual function after the operation.

The superior hypogastric nerve plexus from the aortie plexus sends out a couple of hypogastric nerves at approximately the level of the saeral promontory, 3 cm below the aortie bifurcation, one on each side so that it is shaped like a "moustachio" (Fig. 11). The hypogastric nerve descends on either side between the internal iliae artery and the rectal inherent faseia and converges with the anterosuperior part of the pelvic nerve plexus lateral to the rectum beneath the peritoneal reflection. The hypogastrie nerves often seem to be "put" in front of the sacrum after dissection under laparoscopy. If the

distribution and course of the hypogastric nerve are clear, it usually is not injured, unless the tumor has invaded the rectal inherent fascia [4, 7]. In this condition, preservation of the autonomic nerve is no longer considered.

seem to be "put" in front of the sacrum.

The tiny pelvic splanchnic plexus comes from the anterior foramen of S2–S4, crossing with the middle rectal artery at approximately the central part of the rectal lateral ligament. This plexus is difficult to dissect in any kind of operation (open or laparoscopic). In order to preserve this plexus, dissecting and cutting should be done near the rectal wall (Fig. 12) [7, 9].

Anatomy is the cornerstone of any surgical operation. A thorough knowledge of the anatomic features of laparoscopic rectal surgery can reduce operative mistakes and complications.

#### References

- Bacal D (2003) Experience with laparoseopic medial and lateral dissection of the rectosigmoid for cancer. World J Surg 27: 1337–1338
- 2. Bretagnol F, Rullier E, Couderc P, et al. (2003) Technical and oncological feasibility of laparoscopic total mesoreetal excision

with pouch coloanal anastomosis for rectal eaneer. Colorectal Dis 5: 451–453

and pelvic splanchnic plexus is clearly exposed.

- Feliciotti F, Guerrieri M, Paganini AM, et al. (2003) Long-term results of laparoseopic versus open resections for rectal cancer for 124 unselected patients, Surg Endosc 17: 1530–1535
- Kiran RP, Delaney CD, Senagore AJ, et al. (2004) Operative blood loss and use of blood products after laparoscopic and conventional open colorectal operations. Arch Surg 139: 39–42
- Leroy J, Jamali F, Forbes L, et al. (2004) Laparoscopic total mesorectal excision (TME) for rectal cancer surgery: long-term outcomes. Surg Endose 18: 281–289
- Patankar SK, Larach SW, Ferrara A, et al. (2003) Prospective comparison of laparoscopic vs open resections for colorectal adenocarcinoma over a ten-year period. Dis Colon Rectum 46: 601–611
- Quah HM, Jayne DG, Eu KW, et al. (2002) Bladder and sexual dysfunction following laparoscopically assisted and conventional open mesorectal resection for cancer. Br J Surg 89: 1551–1556
- Senagore AJ, Duepree HJ, Delaney CP, et al. (2003) Results of a standardized technique and postoperative care plan for laparoscopic sigmoid colectomy; a 30-month experience. Dis Colon Rectum 46: 503–509
- Uyama I, Sugioka A, Matsui H, et al. (2001) Laparoscopic lateral node dissection with autonomic nerve preservation for advanced lower reetal eaneer. J Am Coll Surg 193: 579–584
- Zheng Minhua (2002) The eurrent state and evaluation of laparoscopic colorectal surgery. Chin Min Inv Surg 2(Suppl): 25–27
- Zhou Zongguang, Li Li, Shu Ye, et al. (2002) Laparoscopic total mesoreetal excision with anus-preservation for low reetal cancer. Chin J Surg 40: 899–901





Surg Endosc (2006) 20: 739–743 DOI: 10.1007/s00464-005-0268-5

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

# Impaired esophageal function in morbidly obese patients with gastroesophageal reflux disease: evaluation with multichannel intraluminal impedance

E. Quiroga, F. Cuenca-Abente, D. Flum, E. P. Dellinger, B. K. Oelschlager

The Swallowing Center, Department of Surgery, University of Washington, 1959 NE Pacific Street, Box 356410, Seattle, WA 98195-6410, USA

Received: 16 April 2005/Accepted: 26 November 2005/Online publication: 16 March 2006

#### Abstract

*Background:* Morbid obesity is associated with gastroesophagcal reflux disease (GERD), and both have an independent association with motility disorders. Impaired esophageal function is thought to play a role in the development of dysphagia after fundoplication and bariatric procedures (especially restrictive procedures). The authors aimed to define both the physiology and the underlying pathophysiology of swallowing using a novel technique, multichannel intraluminal impedance (MII), which ean accurately determine the elearance of a swallowed bolus through the esophagus, in combination with traditional manometry, which ean measure peristalsis.

*Methods:* Simultaneous MII, manometry, and pH monitoring were performed for 10 asymptomatie subjects, 22 consecutive nonobese patients with GERD (GERD), and 22 consecutive morbidly obese patients with GERD (MO-GERD) who were under evaluation for antireflux and bariatrie surgery at the University of Washington. In this study, MII was defined as abnormal if less than 80% of swallowed liquid boluses cleared the esophagus completely.

*Results:* All GERD and MO-GERD patients had abnormal pH monitoring. The manometric findings were similar for the GERD and MO-GERD patients. All the asymptomatic subjects had normal manometry and impedance test results. Abnormal manometry would have predieted that approximately 23% of GERD and MO-GERD patients had defective emptying. However, when measured with impedance, esophageal clearance was found to be defective in two times as many GERD and nearly three times as many MO-GERD patients.

*Conclusions:* In patients with GERD, impedance often detcets impairments in esophageal motility not identified by manometry. Morbidly obese patients with GERD

have a higher ineidence of impaired esophageal motility than nonobese patients with GERD. This may have implications for bariatric procedures, especially those that are restrictive.

**Key words:** GERD — Esophageal impedance — Esophageal manometry — Esophageal motility — Morbid obesity

Gastroesophageal reflux disease (GERD) and morbid obesity are among the most common diseases in the United States. More than 30% of the U.S. population experiences regular symptoms or manifestations of GERD [4, 15]. More than 50% of U.S. adults are obese, and more than 5% are severely obese [13]. Surgieal intervention plays prominently in the management of each disease. Fundoplication is an excellent alternative to antisecretory therapy for GERD. Currently, the only reliable and effective method of long-term weight loss for morbidly obese patients is bariatrie surgery [7]. Moreover, GERD is commonly associated with obesity [16], oeeurring in as many as 50% to 70% of patients undergoing bariatric surgery [6, 14]. Thus, there are common eonsiderations between antireflux and bariatrie surgery.

Esophageal motility may be important in the etiology and pathophysiology of GERD. It potentially determines the type of antireflux procedure a patient may be able to tolerate without dysphagia. This is one reason why esophageal manometry is recommended before a restrictive antireflux procedure is performed [8]. Although most bariatric procedures also increase resistance to flow through the gastroesophageal junction, manometry is rarely performed preoperatively. Thus, our knowledge concerning the prevalence and nature of esophageal motility disorders in obese patients with GERD is limited.

Correspondence to: B. K. Oelschlager

Manometry measures the presence and strength of peristalsis, and thus provides information about clearance only indirectly. Multichannel intraluminal impedance (MII) is a new technology that allows the transit and clearance of swallowed material within the esophageal lumen to be evaluated. It can therefore directly measure the effectiveness of esophageal transit and clearance.

In this study, we sought to determine the independent effects of GERD and obesity on esophageal function, as measured by manometry and impedance. The results may provide insights about the development of postoperative dysphagia after bariatric restrictive procedures, and may help tailor approaches and procedures to avoid this complication.

#### Methods

Between April 2002 and August 2004, we performed manometry, impedance, and 24-h pH monitoring for patients presenting with GERD for fundoplication or bariatric surgery, as well as for asymptomatic volunteers (with no GERD symptoms). Acid suppression therapy was stopped 5 to 7 days before testing. No patients were taking narcotics, anticholinergics, or promotility agents. Tests were performed in the morning after at least 8 h of fasting. Each participant also completed a standardized GERD symptom questionnaire.

#### Symptom questionnaire

Symptoms were rated by patients and subjects on both frequency and severity scales. The frequency scale had the following range of choices: 0 (never), 1 (once per month), 2 (once per week), 3 (once per day), and 4 (several times per day). Any frequency that fell between two numbers was upgraded to the higher number. The severity score was a visual analog scale from 0 to 10 as follows: 0 (never), 2 (mild), 5 (moderate), 8 (severe), and 10 (worst imaginable). We posed questions on 22 symptoms: 11 gastrointestinal symptoms (heartburn, regurgitation, abdominal pain, belching, dysphagia to liquids and solids, bloating, nausea, chest pain, odynophagia, globus) and 11 extraesophageal symptoms (coughing, hoarseness, wheezing, laryngitis, aspiration, choking, dyspnea, sore throat, asthma, bronchitis, pneumonia).

#### Study groups

#### Asymptomatic subjects

Volunteers were recruited using the Web advertising site for University of Washington research recruitment. The control group consisted of 10 volunteers who, when questioned, answered "0" for each of the 22 symptoms on the questionnaire. This portion of the study was approved separately hy the University of Washington's Human Subjects review board (HSD-02-4684-D02).

#### GERD (nonobese) group (GERD)

The study enrolled 22 consecutive nonobesc patients (body mass index [BMI] less than 35) referred to our esophageal function laboratory with symptomatic GERD under evaluation for an antireflux operation. We excluded patients with potential anatomic esophageal obstructions; peptic strictures, previous antireflux operations, or paraesophageal hernias. All these patients had abnormal distal esophageal (5 cm above the lower esophageal sphincter [LES]) acid exposure on 24-h pH monitoring (> 4%).

#### Morbidly obese patients with GERD (MO-GERD)

The study enrolled 22 consecutive morbidly obese patients with symptomatic GERD under evaluation for bariatric surgery at the University of Washington and referred to our esophageal function laboratory for esophageal motility and acid exposure evaluation. All these patients had a BMI (kg/cm<sup>2</sup>) greater than 35. We excluded patients with potential anatomic esophageal obstructions: peptic strictures, previous antireflux operations, or paraesophageal hernias. All the patients had abnormal distal esophageal (5 cm above the LES) acid exposure on 24-h pH monitoring (> 4%).

#### Esophageal function testing

A specially designed solid state catheter with five manometric sensors and four pairs of impedance sensors separated by 5-cm intervals was used to assess esophageal pressures and impedance with the patient in the supine position (Sandhill Scientific Inc, Highlands Ranch, CO, USA). The LES was examined with the distal circumferential manometric sensor. A station pull-through measurement of the LES pressure (LESP) determined the length and proximal position of the sphincter. The catheter then was placed with impedance and manometry sensors 5, 10, 15, and 20 cm above the LES. The esophageal body was assessed over 10 episodes of deglutition with 5-ml aliquots of water, followed by 10 additional swallows of viscous material (Sandhill Scientific Inc).

#### Impedance definitions

A pair of impedance sensors detected the presence and exit of the bolus in a small segment of the esophagus at each of four levels. A bolus was considered to have entered a pair of impedance sensors when the esophageal impedance (in ohms) dropped 50% from its baseline. Exit of the bolus occurred if and when the impedance rose above this 50% drop. Data were analyzed using BioVicw analysis software (Sandhill Scientific Inc), and each tracing was personally reviewed by the investigators. For each swallow, complete (effective) bolus transit had occurred when the holus entered the first pair of sensors and exited the second, third, and fourth pairs of sensors. The study was considered abnormal if complete bolus transit (CBT) occurred less than 80% of the time. Bolus transit time is the time in seconds from the entrance of the bolus in the proximal most channel (channel 1) to its exit in the distal most channel (channel 4). These definitions are accepted on the basis of multi-institutional trials [17].

#### Manometry definitions

Ineffective esophageal motility was defined to mean that more than 30% of swallows were either nonperistaltic or had distal esophageal amplitudes less than 30 mmHg. Nutcracker csophagus was defined as peristaltic contractions with average distal esophageal amplitudes exceeding 180 mmHg.

This study was approved by the University of Washington's Human Subject's Division review board (HSD- #02-2447-D 03).

#### Results

#### Symptoms

The GERD and MO-GERD groups reported similar symptoms. Heartburn was present daily for all patients. There was no difference in the median severity scores between the GERD (8; range, 5–10) and MO-GERD (8; range, 6–10) groups. At presentation, 8 GERD patients and 7 MO-GERD patients demonstrated dysphagia. The mean frequency and severity scores are shown in Table 1.

Table 1. Presenting symptom frequency and severity

	Heartburn frequency <sup>a</sup>	Heartburn severity <sup>b</sup>	Regurgitation frequency <sup>a</sup>	Regurgitation severily <sup>b</sup>	Dysphagia frequency <sup>a</sup>	Dysphagia severity <sup>6</sup>
Asymptomatic	0	0	0	0	0	0
GERD	3.4	7.6	2.2	5.6	2.2	4.8
MO-GERD	3.8	8	3.6	8.6	2.0	2.0

GERD, gastroesophageal reflux disease; MO-GERD, morbidly obese patients with GERD

<sup>a</sup> Frequency: 0 (never), 1 (once a month), 2 (once a week), 3 (once a day), 4 (several times per day)

<sup>b</sup> Severity: 0 (never), 10 (worse)

Table	2.	24-h	pH	monit	oring

	DeMcester score <sup>a</sup>	Distal acid exposure <sup>b</sup>	Distal acid exposure supine % <sup>b</sup>	Distal acid Exposure upright % <sup>b</sup>	Proximal acid exposure % <sup>c</sup>
Asymptomatic	$4.7 \pm 3.9$	$2.2 \pm 1.2$	$1.5 \pm 1.8$	$0.9 \pm 1$	0
GERD	$64.5 \pm 49$	$16 \pm 14$	$13 \pm 13$	$16 \pm 16$	$4.8 \pm 4.7$
MO-GERD	$46.8 \pm 33^{d}$	$13 \pm 10^{e}$	$13 \pm 8$	$11 \pm 11$	6 ± 7

Values expressed as mean  $\pm$  standard deviation

<sup>a</sup> Normal < 14.7

<sup>b</sup> 5 cm above the LES

<sup>c</sup> 15 cm above the LES

<sup>d</sup> vs. Asymptomatic p < 0.01 vs GERD p = 0.09

<sup>e</sup> vs. Asymptomatic p = 0.011 vs GERD p = 0.25

#### 24-h pH monitoring

Aeid exposure was normal for all asymptomatic subjects and abnormal for all GERD and MO-GERD patients. There was no statistical difference in esophageal aeid exposure between the GERD and MO-GERD groups (Table 2).

#### Manometry

The median LESP was 20.2 mmHg (range, 12–33.9 mmHg) in the asymptomatic group; 13.2 mmHg (range, 4.2–53.4 mmHg) for the GERD patients, and 16.7 mmHg (range, 0–43 mmHg) for the MO-GERD patients. All the asymptomatic subjects had normal peristalsis. Five patients in the GERD group had abnormal manometry: two had ineffective esophageal motility, two had nuteraeker esophagus, and one had a hypertensive LES (>50 mmHg). Five patients in the MO-GERD group had abnormal manometry: two had ineffective esophageal motility, two had nuteraeker esophagus, and one had a hypertensive LES (>50 mmHg). Five patients in the MO-GERD group had abnormal manometry: two had ineffective esophageal motility, two had nuteraeker esophagus, and one had diffuse esophageal spasm (Table 3).

#### Impedance data

The asymptomatic subjects had CBT for 98% of swallows. The GERD patients had CBT for 88% of swallows. Effective CBT was experienced by 66% of the MO-GERD patients, a rate significantly lower than that observed for the asymptomatic subjects (p < 0.01) or the GERD patients (p = 0.01) (Table 4).

We then analyzed each patient according to whether he or she could be considered to have normal bolus transit, defined as CBT for at least 80% of swallows. By this criteria, 9 GERD (41%) and 13 MO-GERD (59%)

Table 3. Manometry results

	LES pressure (mmHg)	Distal esophageal amplitude (mniHg)	Normal <sup>a</sup>
Asymptomatic	$20.2 \pm 7$	70.7 ± 18.9	100%
GERD	$13.2 \pm 10$	$74.9 \pm 49.4$	82%
MO-GERD	$16.7 \pm 11^{b}$	$91.9 \pm 46^{\circ}$	82%

Values expressed as mean ± standard deviation

<sup>a</sup> % of patients with normal peristalsis

<sup>b</sup> vs. asymptomatic p = 0.45 vs GERD p = 0.88

<sup>c</sup> vs. asymptomatic p = 0.08 vs GERD p = 0.26

Table 4. Impedance results

	CBT (%)	Bolus transit time (s)		
Asymptomatic	98	7.0 ± 1.4		
GERD	88	$6.75 \pm 1$		
MO-GERD	66 <sup>a</sup>	$6.87 \pm 1.8^{b}$		

CBT, complete bolus transit; GERD, gastroesophageal reflux disease; MO-GERD, morbidly obese patients with GERD

<sup>a</sup> vs. Asymptomatic p < 0.01 vs GERD p = 0.01

<sup>b</sup> vs. Asymptomatic p = 0.85 vs GERD p = 0.47

patients had abnormal impedance (<80% CBT). All the patients that showed ineffective esophageal motility on manometry had abnormal impedance results. There were 7 GERD and 8 MO-GERD patients with abnormal impedance despite normal manometry (Table 5).

For the swallows with CBT, there was no significant difference in bolus transit time (from channel 1 to 4) for the three groups: asymptomatic subjects, 7.0 s; GERD, 6.75 s; MO- GERD, 6.87 s.

Table 5. Patient motility results; impedance (MII) vs manometry

	GERD			MO-GERD			
		Manometry					Manometry
		Normal	Abnormal			Normal	Abnormal
MII	Normal	10	3 2 nutcracker I hypertensive LES	MII	Normal	9	0
	Abnormal	7	2 2 IEM		Abnormal	8	5 2 IEM 2 nutcracker 1 esophageal spasm

GERD, gastroesophageal reflux disease; MO-GERD, morbidly obese patients with GERD; LES, lower esophageal sphincter; IEM, ineffective esophageal motility

#### Correlation between dysplugia and motility

To determine the relationship between dysphagia and esophageal motility in patients without an anatomic restriction, we analyzed the relationship between dysphagia and manometry/impedance. At presentation, 8 GERD patients and 7 MO-GERD patients demonstrated dysphagia. There was no correlation between dysphagia and the presence of abnormal esophageal motility in either group. Among the 8 GERD patients with dysphagia, 3 (38%) had abnormal manometry and 3 (38%) had abnormal impedance, whereas among the 14 patients without dysphagia, 2 (14%) had abnormal manometry and 8 (57%) had abnormal impedance.

Among the 7 MO-GERD patients with dysphagia, 1 (17%) had abnormal manometry and 4 (57%) had abnormal impedance, whereas among the 15 patients without dysphagia, 4 (27%) had abnormal manometry and 9 (60%) had abnormal impedance.

#### Discussion

This study confirmed previous reports [11] that patients with GERD have a relatively high prevalence of abnormal motor function, as defined by manometry. Using a new technique (impedance), we showed that for patients with GERD, these abnormalities (and others not detected with traditional manometry) lead to substantial impairments in esophageal clearance. In fact, impedance demonstrated that 41% of patients with GERD had defective bolus transit through the esophagus, and in that sense, impedance proved to be a more sensitive test than manometry. Finally, we were able to show, for the first time, that the defective motor function seen in patients with GERD is significantly worse when GERD is associated with morbid obesity.

#### Dysmotilty and GERD

The association between GERD and motility disorders is well described, because both ineffective esophageal motility [11] and hypermotility disorders [1] have been detailed. Indeed, in this investigation, we found both types of disorders in our patients with GERD. The role that this abnormal motor function plays in the pathogenesis of reflux disease still is unclear. On the other hand, considering that most antireflux operations increase the resistance to flow through the gastroesophageal junction, the test has been recommended because it identifies, albeit indirectly, patients in whom dysphagia may develop postoperatively. Several studies have shown, however, that abnormal motility, as documented by manometry, does not predict the development of dysphagia after Nissen fundoplication [2, 5]. One reason may be that manometry, by measuring only the character and strength of peristalsis, may not adequately define the sufficiency of esophageal motility.

In the search for a more reliable tool to assess esophageal function, esophageal impedance was developed and recently been validated [17]. Impedance is the measurement of electrical resistance. By attaching pairs of electrical sensors to a manometry catheter, impedance can detect the presence of gas, liquid, or mixed substances in the esophageal lumen. When these sensors are placed sequentially on a catheter that spans the esophagus, impedance can measure the clearance of a swallowed bolus through the esophagus and estimate the effectiveness of the esophagus to clear in a more direct fashion than manometry. Moreover, it can be combined with manometric sensors on the same catheter, providing a more comprehensive assessment of esophageal function.

We were pleasantly surprised to find that whereas manometry identified motility abnormalities in approximately one-fourth of GERD patients, impedance found that all these, as well as some additional patients in whom manometry results appeared normal, had defective bolus clearance. The fact that none of our asymptomatic subjects had any abnormalities detected in the transit or clearance of the bolus strongly suggests that the abnormalities we found are real. The ultimate significance of this relatively high prevalence of defective clearance in the pathogenesis of dysphagia or GERD as well as its potential impact on patients who undergo operations at the cardia remains to be determined.

#### Dysmotility and Obesity

A major finding of this study was that morbid obesity is an independent risk factor of impaired esophageal function for patients with GERD. The association of obesity and esophageal dysmotility has been described previously using manometry [9, 10, 12]. Our study confirmed the existence of manometrie abnormalities in about 25% of patients with MO-GERD. However, unique to our study is the finding that more than half of these patients present with substantial abnormalities in the transit of a bolus through the esophagus, and that this abnormality is significantly worse in morbidly obese than in nonobese GERD patients. Whether GERD is aggravated by the morbid disease state, resulting in a greater ineidence of esophageal motor dysfunction, or whether obesity has an independent impact on esophageal function is unclear. However, given the increase in obesity and the increase in gastric restrictive procedures, which increase resistance to flow through the gastroesophageal junction, it appears that this should be the focus of greater attention.

Currently, esophageal motility testing is not routinely performed before bariatrie operations. This is surprising because motility testing is routinely recommended before antireflux surgery, and most bariatric operations impose an increased resistance to flow at least to the same extent as antireflux surgery. Although this restriction is important to the desired outcome of bariatrie procedures (i.e., restriction of food intake), in some patients there can be a pathologic degree of impaired clearance. This is especially true for purely restrictive procedures such as the LapBand (BioEnteries/Inamed Corp., Santa Barbara, CA, USA) technique. Esophageal dilation and even pseudoaehalasia after these procedures have been described [3]. The results of our study suggest that impaired csophageal clearance may, at least to some degree, play a role in the development of these complications. In fact, our data suggest that because the incidence of dysmotility is so high in these patients (nearly 60%), there may be a place for measuring esophageal function before bariatrie surgery; at least for patients with GERD.

#### Conclusion

Complete esophageal motility testing with manometry and impedance shows a high incidence of esophageal motility abnormalities in patients with GERD. Many of the abnormalities are identified by impedance only. These motility impairments are even more severe in morbidly obese patients with GERD. Although more investigation is needed, these data suggest that patients with GERD, especially the obese, may be at risk for clinically significant impairments in esophageal clearance after restrictive operations of the cardia. *Acknowledgment*. This research was support by the Dennis and Mary Wise Fund for esophageal research.

#### References

- Barreca M, Oelseblager BK, Pellegrini CA (2002) Outcomes of laparoscopic Nissen fundoplication in patients with the "bypereontractile esophagus." Arch Surg 137: 724–728
- Bootb M, Stratford J, Debn TC (2002) Preoperative esophageal body motility does not influence the outcome of laparoscopic Nissen fundoplication for gastroesophageal reflux disease. Dis Esophagus 15: 57–60
- DeMaria EJ, Sugerman HJ, Meador JG, Doty JM, Kellum JM, Wolfe L, Szues RA, Turner MA (2001) High failure rate after laparoseopie adjustable silieone gastrie banding for treatment of morbid obesity. Ann Surg 233: 809–818
- Drossman DA, Li Z, Andruzzi E, Temple RD, Talley NJ, Tbompson WG, Wbitebead WE, Janssens J, Funcb-Jensen P, Corazziari E, et al. (1993) U.S. bouseholder survey of functional gastrointestinal disorders: prevalence, sociodemography, and healtb impact. Dig Dis Sei 38: 1569–1580
- Fibbe C, Layer P, Keller J, Strate U, Emmermann A, Zornig (2001) Esophageal motility in rellux disease before and alter fundoplication: a prospective, randomized, elinical, and manometric study. Gastroenterology 121: 5–14
- Fisher BL, Pennatbur A, Mutnick JL, Little AG (1999) Obesity correlates with gastroesophageal reflux. Dig Dis Sei 44: 2290– 2294
- Greenstein RJ, Rabner JG, Taler T (1994) Bariatrie surgery vs eonventional dieting in the morbidly obese. Obes Surg 4: 16– 23
- Hinder RA, Libbey JS, Gorecki P, Bammer T (1999) Antireflux surgery. Indications, preoperative evaluation, and outcome. Gastroenterol Clin North Am 28: 987–1005 viii
- Hong D, Khajanebee YS, Pereira N, Lockhart B, Patterson EJ, Swanstrom LL (2004) Manometric abnormalities and gastroesopbageal reflux disease in the morbidly obese. Obes Surg 14: 744–749
- Jaffin BW, Knoepflmacher P, Greenstein R (1999) High prevalence of asymptomatic esophageal motility disorders among morbidly obese patients. Obes Surg 9: 390–395
- Leite LP, Jobnston BT, Barrett J, Castell JA, Castell DO (1997) Ineffective esophageal motility (IEM): the primary finding in patients with nonspecific esophageal motility disorder. Dig Dis Sci 42: 1859–1865
- Mereer CD, Rue C, Hanelin L, Hill LD (1985) Effect of obesity on esopbageal transit. Am J Surg 149: 177–181
- 13. National Institutes of Healtb (2000) The practical guide: identification, evaluation, and treatment of overweight and obesity in adults. National Institutes of Healtb, National Heart, Lung, and Blood Institute, and Nortb American Association for the Study of Obesity, Betbesda, MD, NIH Publieation Number 00-4084
- Sebauer PR, Ikramuddin S, Gourash W, Ramanatban R, Luketich J (2000) Outcomes after laparoscopic Roux-en-Y gastric bypass for morbid obesity. Ann Surg 232: 515–529
- Srinivasan R, Tutuian R, Schoenfeld P, Vela MF, Castell JA, Isaae T, Galaria I, Katz PO, Castell DO (2004) Profile of GERD in the adult population of a Northeast urban community. J Clin Gastroenterol 38: 651–657
- Suter M, Dorta G, Giusti V, Calmes JM (2004) Gastroesophageal reflux and esophageal motility disorders in morbidly obese patients. Obes Surg 14: 959–966
- Tutuian R, Vela MF, Balaji NS, Wise JL, Murray JA, Peters JH, Sbay SS, Castell DO (2003) Esophageal function testing with combined multicbannel intraluminal impedance and manometry: multicenter study in bealtby volunteers. Clin Gastroenterol Hepatol 1: 174–182

Surg Endose (2006) 20: 744–747 DOI: 10.1007/s00464-005-3008-y

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

# The MISTELS program to measure technical skill in laparoscopic surgery

#### **Evidence for reliability**

M. C. Vassiliou,<sup>1</sup> G. A. Ghitulescu,<sup>1</sup> L. S. Feldman,<sup>1</sup> D. Stanbridge,<sup>1</sup> K. Leffondré,<sup>2</sup> H. H. Sigman,<sup>1</sup> G. M. Fried<sup>1</sup>

<sup>1</sup> Steinberg-Bernstein Centre for Minimally Invasive Surgery, Department of Surgery, McGill University Health Centre, 1650 Cedar Avenue, #L9.309, Montreal, Quebec, Canada H3G 1A4

<sup>2</sup> Department of Social and Preventive Medicine, University of Montreal, Pavillon 1420 boul. Mont-Royal, Montreal, Quebec, Canada H2V 4P3

Received: 17 June 2005/Accepted: 17 July 2005/Online publication: 27 February 2006

#### Abstract

*Background:* The McGill Inanimate System for Training and Evaluation of Laparoscopic Skills (MISTELS) is a series of five tasks with an objective scoring system. The purpose of this study was to estimate the interrater and test-retest reliability of the MISTELS metrics and to assess their internal consistency.

*Methods:* To determine interrater reliability, two trained observers scored 10 subjects, either live or on tape. Test-retest reliability was assessed by having 12 subjects perform two tests, the second immediately following the first. Interrater and test-retest reliability were assessed using intraclass correlation coefficients. Internal consistency between tasks was estimated using Cronbach's alpha.

*Results:* The internater and test-retest reliabilities for the total scores were both excellent at 0.998 [95% confidence interval (CI), 0.985–1.00] and 0.892 (95% Cl, 0.665–0.968), respectively. Cronbach's alpha for the first assessment of the test-retest was 0.86.

*Conclusions:* The MISTELS metrics have excellent reliability, which exceeds the threshold level of 0.8 required for high-stakes evaluations. These findings support the use of MISTELS for evaluation in many different settings, including residency training programs.

**Key words:** Laparoscopic training — Simulation — Education — Evaluation — Reliability

Correspondence to: G. M. Fried

The current method of evaluating technical skill in surgical trainees is a subjective assessment performed by practicing surgeons. This is relied on during the licensing process and in the selection of medical students for entry into surgical residency programs. Acquisition of skill has traditionally taken place almost exclusively in the operating room, where resources are limited. Recently, the widespread acceptance of minimally invasive techniques has required the development of new and complex technical skills. Important skills required in the practice of laparoscopic surgery, such as hand-eye coordination and three-dimensional visual spatial perception while viewing through a monocular viewing system, are easily reproduced in an inanimate system. In addition, inanimate systems allow practice with the same instruments used in the operating room but in a more relaxed setting without the time constraints present in the operating room. Various inanimate and virtual reality systems have been designed to facilitate improvement of laparoscopic techniques as well as evaluation of skill in trainees and surgeons [10-12]. We have previously described the McGill Inanimate System for Training and Evaluation of Laparoscopic Skills (MISTELS) program [4, 8]. It consists of a series of standardized tasks performed in a trainer box using a laparoscopic optical system and scored for speed and precision.

For an instrument of measure to become a useful tool in the selection and monitoring of trainees, it must show reproducibility of results when assessed by different testers (interrater reliability) and consistency of performance when the same trainec is evaluated on different occasions in similar conditions (test-retest reliability). It is also useful to have information about the internal consistency of a test in order to ensure that all of the components are measuring the same thing. The aim of the current study is to provide evidence for interrater and test-retest reliability and internal consis-

Poster presented in part as at the annual meeting of the Society of American Gastrointestinal Endoscopic Surgeons (SAGES), Denver, Colorado, USA March 2004

tency of the MISTELS system. This would allow use of MISTELS in multiple institutions and for assessment of a trainee's progress over time.

#### Materials and methods

#### Subjects

Twelve volunteers from McGill University, including medical students, surgical residents, and attending surgeons, were tested. Each participant was required to view an instructional video demonstrating proper performance of each of the five tasks prior to being examined. Each participant performed the entire series of tasks twice on the same day. For interrater reliability, 10 subjects were scored by one of two ohservers during the live performance and videotaped. The second blinded observer then scored the performances on videotape using the same materials as the first observer. The materials were kept intact during the first measurement so as not to hias the second rater. For test-retest reliability, subjects were chosen on the hasis that their performance had reached a plateau after prior practice, and each had achieved a total score of more than 230 when tested. Each of these volunteers was asked to perform the five MISTELS tasks twice on the same day. Twelve pairs of scores were used to assess test-retest reliability.

#### MISTELS

The MISTELS system has been previously described in detail [7]. Briefly, the simulator consists of a laparoscopic trainer box with two 12-mm trocars placed at convenient and standard working angles on either side of a zero-degree laparoscope. A large plastic clip and two alligator clips were used to suspend materials in standardized positions for the exercises. The laparoscope and camera (Storz endoskope; telecam) were mounted on a stand at a fixed focal length, thus allowing the examinee to work independently. The optical system consists of the laparoscope, camera, light source, and video monitor (19 in. Sony Trinitron). The video monitor was placed in line with the operator. Some subjects were tested with a gooseneck camera attached directly to a 21-in. television monitor or to a video-digitizing card inserted into the PCMCIA slot of a laptop computer allowing the image to be displayed on the screen.

Performance of each task is graded objectively, taking into account hoth precision and speed. The score for an individual task is calculated hy subtracting the time required (seconds) from a preset cutoff time. A penalty score for inaccuracy is also deducted according to a predetermined system.

#### Tasks

#### Task 1: peg transfer

Using two graspers, the operator is required to lift each of six pegs from one hoard, transfer it to the other grasper, and then place it on the second board. The procedure is then reversed. Cutoff time is 300 sec.

#### Task 2: pattern cutting

The participant is required to cut a 4 cm predrawn circular pattern out of a  $10 \times 10$  cm piece of double-ply suspended gauze. Cutoff time is 300 sec.

#### Task 3: placement of ligating loop (Endoloop)

A pretied slip knot (Surgitie, US Surgical Corporation, Norwalk, CT, USA) is placed on a circumferential line marked on a tuhular foam appendage. Cutoff time is 180 sec.

**Table 1.** Internal consistency and test-retest reliability of the MISTELS program (n = 12)

	Test		Retest			
Task	Mean	SD	Mean	SD	ICC	
1. Peg transfer	82.6	16.5	84.1	15.6	0.591	
2. Pattern cutting	66.1	18.9	66.7	20.5	0.657	
3. Endoloop	70.0	24.1	85.4	17.4	0.374	
4. EC knot	78.4	34.9	80.2	34.7	0.806	
5. 1C knot	69.6	31.7	70.6	31.1	0.577	
Total score	366.9	105.6	387.7	89.7	0.892	
Cronbach's alpha	0.86		0.77			

EC, extracorporeal; IC, intracorporeal; ICC, intraclass correlation coefficient

#### Task 4: extracorporeal knot

A simple suture 120 cm in length is placed through two premarked points in a longitudinally slit Penrose drain. The suture is then tied with an extracorporeal technique using a knot pusher. The cutoff time is 420 sec.

#### Task 5: intracorporeal knot

This is similar to the previous task, except that the suture is 12 cm and an intracorporeal knot is used. The cutoff time is 600 sec.

#### Normalization

The score for each task was normalized by dividing the score obtained by a predetermined standard value that was derived from the maximum score achieved by a chief resident for that task (task 1 = 237, task 2 = 280, task 3 = 142, task 4 = 297, and task 5 = 520) and then multiplying by 100.

#### Statistical analysis

Internal consistency of the MISTELS system was estimated using Cronhach's alpha [3]. Cronhach's alpha was calculated for each of the four data sets corresponding to interrater (n = 10, rater 1 and rater 2) and test-retest (n = 12, test 1 and test 2). In addition, the impact of deleting each of the five MISTELS tasks on Cronhach's alpha was tested on the first test of the test-retest data set using SAS PROC CORR (SAS for Windows, SAS Institute, Cary, NC, USA).

Intraclass correlation coefficients (ICCs) were used to assess interrater and test-retest reliability of the MISTELS metrics. ICCs were calculated for each of the five tasks separately, as well as for the total score. Each ICC was performed using variance components from a two-way random effect model derived from SAS PROC GLM (SAS for Windows, SAS Institute). For interrater reliability, this model assumes that the raters are randomly selected from a larger population, and that each observer scores each participant. For test-retest reliability, it assumes that the two occasions of measurement represent a random sample of all measurement occasions, and that each participant is assessed at each occasion. We calculated 95% confidence intervals (Cls) according to Shrout and Fleiss [13] for each of the ICCs for the total scores.

#### Results

Cronbach's alpha for the first assessment of the testretest was of 0.86 (Table 1), which suggests that the internal consistency of MISTELS is satisfactory [2].

Correlation with total	Cronbach's alpha
0.81	0.83
0.73	0.84
0.74	0.82
0.77	0.83
0.62	0.86
	Correlation with total 0.81 0.73 0.74 0.77 0.62

EC, extracorporeal; IC, intracorporeal

Each task correlated highly with the total score (correlations, 0.62-0.81; Table 2), and internal consistency could not be improved with the elimination of any task (Table 2). Cronbach's alphas corresponding to interrater and retest data sets were all lower than the one obtained with the first assessment of the test-retest, but all remained acceptable (alpha = 0.77-0.78; Tables 1 and 3).

The ICCs for interrater reliability were all above 0.95 (Table 3), which indicates that more than 95% of the variance in scores can be attributed to a "true" difference between participants. The ICCs for test-retest reliability (Table 1) were all lower than those for interrater reliability. However, the ICCs for the total scores for interrater and test-retest were both excellent at 0.998 (95% C1, 0.985–1.00) and 0.892 (95% CI, 0.665–0.968), respectively. It should be noted that three of the 12 subjects who performed the test-retest obtained a score of zero in one of the tests for task 4 (extracorporeal knot) or task 5 (intracorporeal knot). Accordingly, the analysis was also performed excluding these thrce subjects, with even higher reliability (task 4: ICC = 0.88 vs 0.81; Task 5: ICC = 0.96 vs 0.58).

#### Discussion

There is an increasingly recognized need for tools to objectively measure technical skills. Various systems have been devised [10–12], which consist of structured tasks and objective grading systems. The MISTELS system has been devised in a similar fashion, with the intent to assist trainees in the acquisition of laparoscopic skills and to provide an objective way to measure those skills. For an instrument of measure to be used effectively, it must be shown to be valid and reliable [9].

Validity, as defined by the 1985 Standards for Educational and Psychological Testing, is "the appropriateness, meaningfulness, and usefulness of the specific inferences made from test scores" [1]. For a scale to be valid, it must measure what it intends to measure. There are various types of validity, including face, content, construct, and criterion validity [9]. Demonstrating validity is a complex undertaking, usually requiring a series of experiments. Prior to embarking on such an endeavor, it is necessary to show that the test is measuring something in a reproducible fashion. There is no point to painstakingly proving a system to be valid only to recognize later that it is not reliable. Reliability refers

**Table 3.** Internal consistency and interrater reliability of the MISTELS program (n = 10)

	Rater 1		Rater 2			
Task	Mean	SD	Mean	SD	ICC	
1. Peg transfer	81.3	14.9	80.8	15.3	0.991	
2. Pattern cutling	59.7	22.2	58.8	21.5	0.984	
3. Endoloop	67.3	26.0	64.8	27.7	0.985	
4. EC knol	68.1	37.7	67.8	37.4	0.999	
5. IC knot	66.9	30.3	67.1	30.3	1.000	
Total score	343.3	97.8	339.3	100.4	0.998	
Cronbach's alpha	0.77		0.78			

EC, extracorporeal; IC, intracorporeal; ICC, intraclass correlation coefficient

to the consistency, stability, and precision of test scores [14]. It is an indication of the extent to which measurements of individuals in different circumstances or by different observers yield similar results. The necessity to establish evidence for reliability arises from measurement error, which is a characteristic of all instruments of measure.

Types of measurement error can be grouped into three general categories: those associated with the observer (different observers using different methods or the same observer using different methods), the test (variability in the materials used for the test), and the subject (different performances by the subject on different occasions caused by fatigue, distraction, or other reasons). In the MISTELS system, differences in scoring technique between observers could occur in the measurement of time and/or precision. Both of these procedures have been standardized, and little variation is expected between observers. The ICC for interrater reliability for the total score was 0.998 (95% CI, 0.985– 1.00), which confirms our expectation of low variability among observers.

Measurement of test-retest reliability is less straightforward because the MISTELS system is also conceived as an instructional tool. We have previously shown that practice in the simulator improves performance in the simulator [5, 6, 8]. Specifically, we have shown that scores increase significantly after a single repetition, and that the improvement is significantly greater in the less experienced group. To evaluate testretest reliability, the subject must repeat the test, and the results can be biased toward the null when a less experienced subject improves significantly. To counterbalance this bias, we chose subjects who had attained a minimum total score of 230. MISTELS was designed as a coherent system for the training and evaluation of a set of laparoscopic skills. There may be some individuals who are better at certain tasks than others, but the overall performance carries the most practical significance. The cutoff criterion allowed us to include junior-level trainees, such as medical students and junior residents. We believe that if any bias would result from the inclusion of inexperienced subjects, it would be in the direction of low reliability due to the effect of practice on performance. We found that despite this

inclusion, the test-retest reliability for the total is excellent, with an ICC of 0.892 (95% CI, 0.665–0.968).

#### Conclusion

The MISTELS metrics meet the standards of reliability for high-stakes examinations. The data for interrater and test-retest reliability support the use of the MISTELS system in any institution seeking to improve and measure technical skills in trainees and practicing surgeons.

Acknowledgments. This work was funded by an unrestricted educational grant from Tyco Healthcare Canada.

#### References

- 1. American Psychological Association (1985) Standards for educational and psychological testing American Psychological Association, Washington, DC
- 2. Bland JM, Altman DG (1997) Cronbach's alpha. Br Med J 314: 572
- Cronbach L (1951) Coefficient alpha and the internal structure of tests. Psychometrika 16: 297–302

- Derossis AM, Antoniuk M, Fried GM (1999) Evaluation of laparoscopic skills: a 2-year follow-up during residency training. Can J Surg 42: 293–296
- Derossis AM, Bothwell J, Sigman HH, Fried GM (1998) The effect of practice on performance in a laparoscopic simulator. Surg Endose 12: 1117–1120
- Derossis AM, Fried GM, Abrahamowicz M, Sigman HH, Barkun JS, Meakins JL (1998) Development of a model for training and evaluation of laparoscopic skills. Am J Surg 175: 482–487
- Fraser SA, Klassen DR, Feldman LS, Ghitulescu GA, Stanbridge D, Fried GM (2003) Evaluating laparoscopic skills: setting the pass/ fail score for the MISTELS system. Surg Endose 17: 964–967
- Fried GM, Derossis AM, Bothwell J, Sigman HH (1999) Comparison of laparoscopic performance in vivo with performance measured in a laparoscopic simulator. Surg Endosc 13: 1077– 1082
- 9. Gall M, Borg W, Gall J (1996) Educational research: an introduction. Longman, White Plains, NY
- Hanna GB, Drew T, Clinch P, Hunter B, Cuschieri A (1998) Computer-controlled endoscopic performance assessment system. Surg Endosc 12: 997–1000
- Reznick R, Regehr G, MacRae H, Martin J, McCulloch W (1997) Testing technical skill via an innovative "bench station" examination. Am J Surg 173: 226–230
- Rosser JC, Rosser LE, Savalgi RS (1997) Skill acquisition and assessment for laparoscopic surgery. Arch Surg 132: 200–204
- Shrout P, Fleiss J (1979) Intraclass correlations: uses in assessing rater reliability. Psych Bull 86: 420–428
- Streiner D, Norman G (1996) Health measurement scales: a practical guide to their development and use. Oxford University Press, New York

Surg Endosc (2006) 20: 748–752 DOI: 10.1007/s00464-005-0429-6

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

### Laparoscopic subtotal splenectomy in hereditary spherocytosis

To preserve the upper or the lower pole of the spleen?

C. Vasilescu,<sup>1</sup> O. Stanciulea,<sup>1</sup> S. Tudor,<sup>1</sup> D. Stanescu,<sup>2</sup> A. Colita,<sup>3</sup> R. Stoia,<sup>3</sup> D. Coriu,<sup>3</sup> A. Colita,<sup>4</sup> C. Arion<sup>4</sup>

<sup>1</sup> Department of General Surgery and Liver Transplantation, Fundeni Clinical Institute, 258 Fundeni Street, Bucharest, Romania RO-72434

<sup>2</sup> Department of Nuclear Medicine and Ultrasounds, Fundeni Clinical Institute, 258 Fundeni Street, Bucharest, Romania RO-72434

<sup>3</sup> Department of Hematology, Fundeni Clinical Institute, 258 Fundeni Street, Bucharest, Romania RO-72434

<sup>4</sup> Department of Pediatries, Fundeni Clinical Institute, 258 Fundeni Street, Bucharest, Romania RO-72434

Received: 16 June 2005/Accepted: 26 November 2005/Online publication: 16 March 2006

#### Abstract

*Background:* Clinical manifestations of hereditary spherocytosis can be controlled by splenectomy. The use of this procedure has been restricted due to concerns regarding exposure of patients to a lifelong risk of overwhelming infections. Subtotal splenectomy, which removes 85–90% of the enlarged spleen, is a logical alternative. In the first cases performed by laparoscopy we have chosen to preserve the upper pole. However, this technique showed some disadvantages, especially concerning the correct intraoperative evaluation of the splenic remnant volume. Therefore, we developed a new variant of the procedure by preserving the lower pole of the spleen.

*Methods:* Based on the authors' experience in laparoscopy (176 laparoscopic splenectomies), 10 laparoscopic subtotal splenectomies were performed in patients with hereditary microspherocytosis, preserving either the upper or the lower splenic pole.

*Results:* Patient age ranged between 5 and 35 years. The mean volume of the remnant spleen was  $41.4 \text{ cm}^3$ . There were no complications, and no transfusions were needed. Follow-up for 1–30 months was available.

*Conclusion:* Subtotal splenectomy appears to control hemolysis while maintaining splenic function. The laparoscopic approach is safe and effective and should be considered the procedure of choice in hereditary microspherocytosis. Laparoscopic subtotal splenectomy presents an advantage over open subtotal splenectomy, resulting in decreased blood loss, shorter hospital stay, no conversions, fewer operative and postoperative

Correspondence to: C. Vasilescu

complications, and excellent remission rates. On the basis of our experience, the preservation of the lower pole of the spleen seems to be a first-linc option for the optimal evaluation of the residual splenic mass.

Key words: Subtotal splenectomy — Partial splenectomy — Subtotal laparoscopic splenectomy — Hereditary spherocytosis — Laparoscopic splenectomy

Hereditary spherocytosis (HS) is the most common inherited hemolytic anemia in which a primary deficiency in an erythrocyte membrane skeleton protein leads to surface area loss. Erythrocytes with decreased surface area and impaired deformability are trapped in the splenic pulp and phagocytized [19]. The result is chronic hemolysis and an increased propensity for gallstone formation.

The clinical severity of HS varies from symptom-free carrier to severe hemolysis. Patients with HS should be graded by their severity of disease (bascline hemoglobin, reticulocyte count, and jaundice level of activity). This predicts the clinical course and the need for splenectomy [4].

Splencctomy is very effective in reducing hemolysis, leading to a significant prolongation of the red cell life span. The clinical manifestations and complications (anemia and gallstones) are much reduced in severe HS and abolished in milder cases, but at the price of an increased risk of life-threatening sepsis, usually caused by pneumococcal species. This risk is age related, being highest in the youngest children and within the first years after surgery.

Current guidelines recommend pre-splenectomy vaccination against pneumococcus, haemophilus, and meningococcus together with long-term (lifelong)

The paper is based on material presented at the 13th International Congress of the European Association for Endoscopic Surgery (EAES), Venice, 2005

postoperative penicillin prophylaxis. These measures do not completely eliminate the risk because of the serotypes that are not represented in vaceines, penicillinresistant pneumococeal strains, and issues relating to lifelong compliance. Recent studies have raised additional concerns regarding long-term complications after splenectomy, such as atheroselerotic events and pulmonary hypertension [2, 19].

Overwhelming postsplenectomy infections (OPS1) are primarily caused by *Streptococcus pneumoniae*, *Neisseriae meningitides*, or *Haemophilus influenzae*. Despite appropriate antibioties and intensive therapeutie intervention, the overall mortality in published studies for established cases of OPS1 varies from 50 to 70%. Of those patients who die, > 50% die within the first 48 h of hospital admission. In those patients who survive, other sequelae include gangrene leading to amputations, deafness associated with meningitis, or mastoid osteomyelitis and aortic insufficiency secondary to endocarditis [5].

Open subtotal splenectomy has been proven to be a therapeutic alternative in HS. In order to add the advantages of minimally invasive surgery, we began to perform this procedure by laparoscopy. Based on the authors' experience in laparoseopie surgery (176 laparoscopie splenectomies) and after performing five open subtotal spleneetomies with good hematologieal outcomes, we performed laparoscopic subtotal splenectomy in 10 patients. In open surgery, subtotal spleneetomy was performed preserving either the upper [11] or the lower pole of the spleen [2, 16, 18]. In the first eases [22] performed by laparoscopy, we chose to preserve the upper pole. However, this technique showed some disadvantages, especially eoneerning the correct intraopcrative evaluation of the splenic remnant volume. Therefore, we developed a new variant of the procedure by preserving the lower pole of the spleen.

#### Patients and methods

#### Patients

Between June 2002 and January 2005, 10 consecutive patients underwent laparoscopic subtotal splenectomy. The patients were six children (four girls and two boys; age range, 5-12 years; mean age, 7.5 years) and four young adults (age range, 19-35 years). The patients' characteristics are detailed in Table I. The body mass index ranged between 17 and 31. The diagnosis was based on clinical features (variable degree of anemia, jaundice, and splenomegaly) and laboratory findings (spherocytes, raised red cell hemoglobin concentration, and an increase in reticulocytes). Spleen size and volume were determined using ultrasounds. Spleen volume ranged between 243 and 760 cm<sup>3</sup> (mean, 429 cm<sup>3</sup>). When small accessory spleens were detected they were preserved. The indication for surgery was severe anemia with recurrent need for transfusions, and surgery was performed after clinical monitoring of a number of variables (the degree of anemia, the impact of chronic fatigue on life quality, and the presence of gallstones). After discussions with the parents or the patients and after explaining the potential benefits and risks, subtotal laparoscopic splenectomy was performed. The procedure was feasible in all patients, and at the time of the intervention none required final total splenectomy. The mean duration of follow-up for the 10 patients was 13.1 months (range, 1-30).

#### Surgical technique

The objective of the surgical procedure was to remove approximately 85% of the splenic tissue while preserving either the upper or the lower pole of the spleen. Based on the authors' experience and standard approach in laparoscopic splenectomy, the upper pole of the spleen was preserved in the first seven procedures according to the recommendations of Petroianu et al. [13]. In the last three cases, the lower pole of the spleen was preserved based on our experience in open surgery (five open subtotal splenectomies). Under general anesthesia, the patient was placed in the right lateral decubitus position. Four troears were used, two 12 mm ports (one for the laparoscope and the other for the hemostatic instruments) and two 5 mm ports for retractors. Trocar position was similarly utilized with any laparoscopic splenectomy. A 30° laparoscope was inserted into the abdominal eavity and a limited diagnostic laparoscopy was performed in order to detect accessory spleens (an accessory spleen was observed in one case and was preserved).

# Subtotal laparoscopic splenectomy preserving the upper pole

Initially, the splenocolic and phrenosplenic ligaments were partially incised using a harmonic scalpel, and the branches of the gastroepiploic vessels to the lower pole were divided (Fig. 1A). The splenic hilum was carefully dissected, preserving the pancreatic tail intact. The branches of the splenic artery and vein were divided using Ligasure Atlas Sealer/Divider (Tyco Healthcare, Boulder, CO, USA). The last one or two short gastric vessels were preserved by observing the size of the future splenic remnant, which provided a clear line of demarcation on the spleen surface. Standard monopolar electrocautery was used to transect the spleen, ensuring that a 5 mm rim of devascularized splenic tissue remained in situ. Monopolar electrocautery was used in favor of the Ligasure or the harmonic shears in order to visualize splenic remnant vascularization. After tbat, the Ligasure Atlas Sealer/Divider provided excellent hemostatic control. A TachoComb (Nycomed GmbH, Linz, Austria) was used to complete haemostasis. The splenic fragment was placed in an endobag for removal. The most lateral 12 mm port site was used to extract the morcellated tissue, or the specimen was extracted via a Pfannenstiel incision. Two drains were placed at the end of the procedure. The postoperative courses were uneventful, and all patients were discharged within 1 week after surgery.

# Subtotal laparoscopic splenectomy preserving the lower pole

This technique is similar to that used in the open approach (Fig. 1B). After dividing the branches of the splenic hilum and the short gastric vessels using Ligasure Atlas Sealer/Divider, a small fragment of the lower pole was preserved. The vascularization was based on the ascending branch of the left gastrocpiploic artery. The advantage of this technique was better visualization of the remnant size.

#### Results

The mean surgery time was 95 min. The mean operative bleeding was < 70 ml. No intraoperative or postoperative blood transfusions were required. In six eases, the specimen was extracted using an endobag after moreellation. In these cases, weight of the specimen was calculated using the following formula: intact weight (g) = moreellated weight (g) × 1.34 + 45 [23]. In four eases, the specimen was extracted via a Pfannenstiel ineision. The mean weight of the resected specimen was 600 g.

Table 1. Demographic data of patients and values of hemoglobin and reticuloeytes before and after surgery

Patient No.			Spleen volumetry before/ after surgery (em <sup>3</sup> )	Hemoglobi	n value (g/dl)	Reticulocyte value, post-splencctomy (%)	
	Age at surgery (yr)	I ransfusions before surgery		Pre-spleneetomy	Post-spleneetomy		Postsurgical follow-up (mo)
1	16	Yes	664/71	8.4	11	16.2	32
2	19	Yes	760/79	7.7	11	12.7	31
3	5	Yes	336/35	8.6	10.7	18.5	20
4	35	Ycs	603/80	8	10.2	11.3	19
5	7	Yes	307/21	7.5	9.4	8.7	13
6	5	Yes	373/32	8	10.6	6.3	13
7	8	Yes	317/29	8.9	11.8	19.6	12
8	12	Ycs	278/21	7.4	10.4	8.7	4
9	22	Yes	408/26	8.9	12.1	9.8	4
10	10	Yes	243/20	7.6	9.5	6.3	3



Fig. 1. (A) Surgical technique used for laparoscopic subtotal spleneetomy: the remnant upper pole of the spleen with its vascular supply (a short gastric vessel). (B) Surgical technique used for laparoscopic subtotal spleneetomy: the remnant lower pole of the spleen with its vascular supply (a pedicle arising from the left gastroepiploie vessels).

In all patients, the hemoglobin values were increased and the reticulocyte number was decreased following subtotal splenectomy (Table 1), as a direct consequence of an increase in the life span of red blood cells. During a follow-up period that ranged from 1 to 30 months, none of the patients required blood transfusions to maintain adequate hemoglobin levels. In all patients, a Doppler ultrasound performed after surgery showed the vascularization of the splenic remnant. Comparison of the spleen volumetry before and after surgery showed that the splenic remnant represented 15% of the initial volume of the spleen. Following Stoehr et al. [18], we assessed the phagocytic activity of the remnant spleen by scintigraphy with 99-technetium, which showed radionuclide uptake in the splenic area. Also, the percentage of pitted erythrocytes was < 2%. Howell–Jolly bodies appeared transiently after surgery in six of 10 patients. One patient required splenic remnant removal 11 months after the initial surgery due to persistent mild hemolytic anemia. In this case, due to the adhesions of the splenic remnant, conversion to open surgery was necessary.

#### Discussion

#### Indications for surgical treatment

Splenectomy should be performed in children with severe transfusion-dependent HS. The clinical manifestations of HS include anemia, chronic jaundice, and cholelithiasis. Total splenectomy that abrogates hemolysis is the most attractive treatment option, but this exposes the patient to a life-long risk of potentially lethal infections. Splenectomy leads to a significant prolongation of the red cell life span (although this is not necessarily extended to a normal life span). By eliminating the need for regular blood transfusions in severe cases, this surgical procedure also eliminates the associated risks of blood-borne viral infections and iron overload [20]. In addition, the impact on the patient's quality of life is significant, with a major improvement in the patient's physical and professional abilities after surgery.

Unfortunately, spleen removal leads to many inconveniences because the spleen represents approximately 25% of the lymphatic tissue and is the major repository of mononuclear phagocytic cells. As a result, it performs an efficient filtration function for the bloodstream [14]. OPS1 is a well-defined clinical condition resulting from the inability of the splenectomized host to combat infections with encapsulated bacteria. Because of this lethal complication, a number of alternatives to splenectomy have been studied, including autotransplantation of the removed spleen and partial splenectomy [1, 6, 14]. Partial splenectomy is preferable to splenic autotransplantation because it is associated with higher antibody titers after immunization, better pneumococcal splenic uptake, and improved survival rates.

Subtotal splenectomy provides a potentially effective alternative to total splenectomy in patients with HS. It is likely that contact between antigens and lymphoid cells as well as maturation of the pre-B cells can proceed in the remnant spleen.

In the study by Bader-Meunier et al. [2], approximately 15% of the enlarged spleen was preserved in order to decrease the sequestration of red blood cells as much as possible while at the same time maintaining an adequate amount of remnant spleen to preserve its phagoeytic function.

#### Key points in the surgical technique

Open subtotal splenectomy has been suggested as a feasible alternative for severe forms of HS [2, 16, 18]. The potential benefits of this method were first proposed by a group of French surgeons, who published between 1993 and 2001 a study based on 40 cases. Since then, the technique has been applied by others, and additional studies have shown that this method provides good results [16]. It has also recently been hypothesized that a more radical approach to open resection could permanently decrease recurrent hemolysis and at the same time conserve a remnant spleen of 10 cm<sup>3</sup> [18]. Seshadri et al. [17] performed a partial laparoscopic splenectomy for a 2 cm splenie cyst. In the current study, after five open subtotal splenectomies were performed with good hematological outcomes, the potential benefits of the laparoscopic approach were analyzed because laparoscopic splenectomy is currently considered the best approach in patients without splenomegaly [22].

The development of partial splenectomy is based on the knowledge of the splenic vascular pattern [13, 21]. This surgical approach is feasible because of the unique anatomical features of this organ, which is divided into a variable number of almost independent vascular territories. Splenic vessels, including branches of the main splenic pediele, short gastric vessels, and left gastroepiploie vessels, are divided outside the spleen before entering the splenic hilum as terminal vessels. There are three types of anastomoses between the individual branches of the splenic artery: hilar, intraparenchymatous, and subcapsular [3, 7, 8]. The intraparenchymatous anastomoses do not maintain the viability of the externally devascularized regions of the spleen, a fact that makes it possible to obtain a safe hemostasis of the splenic remnant. Subtotal splenectomy can be performed after primary ligation of the main pedicle of the splenic artery and vein, preserving either a short gastric artery [10] or a branch of the left gastroepiploic artery [20].

In order to establish the best therapeutic option, it is essential to determine the amount of splenic parenchyma that needs to be removed to effectively reduce red blood eell destruction. In addition, it is important to evaluate the minimal amount of residual parenchyma necessary for maintaining adequate phagocytic function [2, 19, 20]. Previous studies have shown that this goal can be achieved by preserving approximately 15% of the enlarged spleen. This is the key point of the surgical procedure.

In the current study, the technique described by Tchernia et al. [2, 19, 20] was used during open surgery to preserve the ascending branch of the left gastroepiploic artery. This solution was initially difficult to apply in the laparoscopic approach because in this case the procedure is conducted after manipulation of the lower splenic pole. As a result, Petroianu's technique was initially used as a part of the laparoscopic approach. This technique has been used successfully in a large number of operations to preserve the splenogastric vessels, which arc considered to be enough for the upper splenic pole [9, 11, 12, 15].

In the current study, the lower splenic pole was preserved in the last three patients with good results. In the first procedures, according to the recommendation of Bader-Meunier et al. [2], approximately 15% of the enlarged spleen was preserved, a target that was more difficult to achieve using the laparoscopic approach than with open surgery. In the first procedures the main difficulty was to approximate 15% of the enlarged spleen. The evaluation of the splenic remnant volume was quite difficult in the cases with preservation of the upper pole of the spleen. Therefore, in the last cases we chose to preserve the lower pole of the spleen. This technique allowed us a better mobilization of the splenic remnant and a more accurate evaluation of the volume. The length of the remnant ranged between 3.5 and 6 cm, depending on the initial size of the spleen, and was measured using a 10-cm flexible band. Based on the 10 cases in this study and recent publications [18], it appears that a smaller remnant spleen (a remnant volume between 5 and 10%) is sufficient to preserve the splenic immune and phagocytic functions. Laparoscopic subtotal splenectomy combines the advantages of minimally invasive surgery with the maintenance of phagocytic function of the splenic remnant.

#### Hematologic results

The hemoglobin values increased and the reticulocyte count decreased in the first several days after surgery. This was the result of increased bone marrow production, evidenced by 20% reticulocytes.

Studies published between 1993 and 2001 [2, 19, 20] based on 40 cases that were followed for 14 years after surgery showed that the mean increase in hemoglobin value after partial splenectomy was 3 g/dl, whereas the decrease in the number of reticulocytes averaged approximately  $300 \times 10^9$ /L [2]. This occurred as a direct result of an increase in the red blood cell life span, with an average increase of 6.5 ± 1 days—a result far from normal.

Long-term postsurgical follow-up results show that beneficial clinical effects are sustained over a long period. However, in one study, one patient experienced complications that led to a secondary total splenectomy [2]. A male patient who underwent surgery at the age of 5 years and maintained a hemolytic state after surgery required a secondary splenectomy (clinical symptoms included recurrence of jaundice, chronic fatigue, and mild anemia). Laparoscopic surgery was attempted, but it was necessary to convert to open surgery due to phrenosplenie adhesions. Other complications that also require secondary splenectomy include bleeding and torsion of vascular pedicles.

Several studies found that secondary splenectomy was required in 10% of patients at 5 years and in 33% at 10 years [2, 20]. The size of the remnant spleen increased in all cases. The growth was assessed by ultrasound and CT sean. Significant regrowth of the splenic remnant was noted during the first year after surgery, sometimes four and a half times its postoperative size [18], which did not correlate with the measured hemoglobin values or retieulocyte counts [2]. Although the current study is based on a small scries of patients and the follow-up is short, our aim was to present a new approach in patients with HS. Of course, long-term results are needed in order to prove that the effects are sustained over a long period.

The outcome of patients with HS with total splenectomy was compared to that of patients with subtotal splenectomy [20] performed by open surgery. The results showed that subtotal splenectomy appears to be the first-line treatment option for patients with HS [2]. Regarding the phagocytic activity of the remnant, the percentage of pitted cells after partial splenectomy was similar to that seen in normal individuals, whereas after total splenectomy their numbers were markedly inereased [20].

#### Conclusions

Subtotal splenectomy has been shown to provide a persistent decrease in hemolytic rate while preserving the integrity of splenie phagoeytic function. Subtotal splenectomy should be considered preferable in two situations. First, in children younger than 5 years of age who need regular transfusions, subtotal splencetomy deereases the risk of both severe sepsis and transfusionrelated viral infections. Second, subtotal splenectomy should be eonsidered for eases in which mild or moderate anemia is present and the patient has ehronie discomfort, significant splenic enlargement, or biliary lithiasis but does not require transfusions or requires them only sporadically. Laparoseopic subtotal spleneetomy presents an advantage over open subtotal spleneetomy, resulting in decreased blood loss, shorter hospital stay, no conversions, fewer operative and postoperative complications, and excellent remission rates. On the basis of our experience, the preservation of the lower pole of the spleen seems to be a first-line option for the optimal evaluation of the residual splenic mass. The laparoscopic approach is safe and effective, with encouraging short-term results, but a much longer follow-up (up to 20 years) is needed to be certain that the beneficial effects will be sustained.

Acknowledgment. We thank Leslie Cizmas, Ph.D., Tcxas A&M University.

#### References

- Alvarez FE, Greeo RS (1980) Regeneration of the spleen after eetopic implantation and partial splenectomy. Arch Surg 115: 772–775
- Bader-Meunier B, Gauthier F, Arehambaud F, Cynober T, Mielot F, Dommergues JP, Warszawski J. Mohandas N, Tchernia G (2001) Long-term evaluation of the beneficial effect of subtotal spleneetomy for management of hereditary spherocytosis. Blood 97: 399–403
- 3. Bergman RA, Afifi AK, Miyauchi R (2003) Splenic artery branches: posterior gastrie, superior polar of spleen and gastrosplenie artery. In Illustrated encyclopedia of human anatomic variation. II: eardiovascular system. http://www.anatomyatlases.org
- Bolton-Maggs PH, Stevens RF, Dodd NJ, Lamont G, Tittensor P, King MJ (2004) Guidelines for the diagnosis and management of hereditary spheroeytosis. Br J Haematol 126: 455–474
- Brigden ML, Pattullo AL (1999) Prevention and management of overwhelming postspleneetomy infection—an update. Crit Carc Med 27: 836–842
- Drew PA, Clayer MT, Jamieson GG (1990) The value of splenie autotransplantation. Arch Surg 125: 1224
- Holibkova A, Maehalek L, Houserkova D, Ruzieka V (1998) A contribution to the types of branching and anastomoses of the splenie artery in human spleen. Aeta Univ Palaeki Olomuc Fac Med 141: 49–52
- Maehalek L, Holibkova A, Tuma J, Houserkova D (1998) The size of the splenie hilus, diameter of the splenie artery and its branches in the human spleen. Acta Univ Palaeki Olomuc Fac Med 141: 45–48
- Petroianu A (1988) Treatment of portal hypertension by subtotal spleneetomy and central splenorenal shunt. Postgrad Med J 64: 38-41
- Petroianu A (1992) Splenie viability after segmental devaseularization. Rev Paul Med 110: 39–41
- Petroianu A (1996) Subtotal spleneetomy for treatment of patients with myelofibrosis and myeloid metaplasia. Int Surg 81: 177–179
- Petroianu A, da Silva RG, Simal CJ, de Carvalho DG, da Silva RA (1997) Late postoperative follow-up of patients submitted to subtotal spleneetomy. Am Surg 63: 735–740
- Petroianu A, Ferreira VL, Barbosa AJ (1989) Morphology and viability of the spleen after subtotal splenectomy. Braz J Med Biol Res 22: 491–495
- Petroianu A, Simal CJ (1993) Shifts in the reticuloendothelial system uptake pattern induced by earbon colloid in the rat. Med Sei Res 21: 311–312
- Resende V, Petroianu A (2003) Functions of the splenic remnant after subtotal spleneetomy for treatment of severe splenie injuries. Am J Surg 185: 311–315
- Rice HE, Oldham KT, Hillery CA, Skinner MA, O'Hara SM, Ware RE (2003) Clinical and hematologic benefits of partial splenectomy for congenital hemolytic anemias in children. Ann Surg 237: 281–288
- Seshadri PA, Poulin EC, Mamazza J, Schlachta CM (2000) Technique for laparoscopic partial splenectomy. Surg Laparose Endosc Percutan Tech 10: 106–109
- Stochr GA, Stauffer UG, Eber SW (2005) Near-total splenectomy: a new technique for the management of hereditary spheroeytosis. Ann Surg 241: 40–47
- Tehernia G, Bader-Meunier B, Bertcrottierc P, Eber S, Dommergues JP, Gauthier F (1997) Effectiveness of partial splencetomy in hereditary spherocytosis. Curr Opin Hematol 4: 136–141
- Tehernia G, Gauthier F, Mielot F, Dommergues JP, Yvart J, Chasis JA, Mohandas N (1993) Initial assessment of the hencficial effect of partial splenectomy in hereditary spheroeytosis. Blood 81: 2014–2020
- Treutner K, Klosterhalfen B, Winkeltau G, Moench S, Schumpelick V (1993) Vascular anatomy of the spleen: the basis for organ-preserving surgery. Clin Anat 6: 1–9
- Vasilescu C, Staneiulea O, Colita A, Stoia R, Moieean A, Arion C (2003) Laparoscopic subtotal spleneetomy in the treatment of hereditary spherocytosis. Chirurgia (Bucur) 98: 571–576
- Walsh RM, Chand B, Brodsky J, Heniford BT (2003) Determination of intact splenic weight based on morcellated weight. Surg Endose 17: 1266–1268

Surg Endose (2006) 20: 753–756 DOI: 10.1007/s00464-005-0177-7

© Springer Science+Business Media, Inc. 2005



and Other Interventianal Techniques

# Laparoscopic cholecystectomy using a newly developed laparoscope manipulator for 10 patients with cholelithiasis

K. Tanoue,<sup>1</sup> T. Yasunaga,<sup>2,3</sup> E. Kobayashi,<sup>4</sup> S. Miyamoto,<sup>3</sup> I. Sakuma,<sup>4</sup> T. Dohi,<sup>5</sup> K. Konishi,<sup>1</sup> S. Yamaguchi,<sup>2</sup> N. Kinjo,<sup>6</sup> K. Takenaka,<sup>6</sup> Y. Maehara,<sup>6</sup> M. Hashizume<sup>2</sup>

<sup>1</sup> Department of Innovative Medical Technology, Graduate School of Medical Sciences, Kyushu University, 3-1-1, Maidashi, Higashi-ku, Fukuoka, 812-8582, Japan

<sup>2</sup> Department of Disaster and Emergency Medicine, Graduate School of Medical Sciences, Kyushu University, 3-1-1, Maidashi, Higashi-ku, Fukuoka, 812-8582, Japan

<sup>3</sup> Healthcare System Business Development Department, Hitachi, Ltd., Marunouchi 1-6-6, Chiyoda-ku, Tokyo, Japan

<sup>4</sup> Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan

<sup>5</sup> Graduate School of Information Science and Technology, The University of Tokyo, Tokyo, Japan

<sup>6</sup> Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu University, 3-1-1, Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

<sup>7</sup> Department of Surgery, Fukuoka City Hospital, Yoshizuka, Hakata-ku, Fukuoka, Japan

Received: 30 March 2005/Accepted: 27 September 2005/Online publication: 6 December 2005

#### Abstract

*Background:* Laparoscopic surgery has continued to gain popularity in almost all fields of abdominal surgery, and robotic systems have been introduced in general surgery. Naviot is a new remote-controlled laparoscope manipulator system controlled by the operator's hand. This study assessed its introduction into clinical practice.

*Methods:* A group of 10 consecutive patients with cholelithiasis underwent laparoseopic cholecystectomy assisted by the Naviot system (Naviot group). Another group of 41 patients who underwent laparoscopic cholecystectomy with a conventional human camera holder (human camera group) were selected for a comparison of their operative results with those of the Naviot group.

*Results:* The operative time of  $89.3 \pm 27.1$  min for the Naviot group was significantly longer than that of 74.8  $\pm$  28.1 min for the human camera group (p < 0.05). However, when the setup time for the Naviot system was excluded, the operative time was not significantly different from that for the human camera group. Other operative results showed no significant difference between the two groups.

*Conclusion:* The authors believe that the new Naviot system is feasible for clinical use, and that it enables surgeons to perform solo gastrointestinal surgery.

Key words: Laparoscopic cholecystectomy — Laparoscopic manipulator — Naviot

Correspondence to: K. Tanoue

Laparoseopie surgery has continued to gain popularity in almost all fields of abdominal surgery. Laparoscopic cholecystectomy eurrently is the main standard treatment for patients with cholelithiasis and polyp of the gallbladder.

In conventional laparoscopic surgery, a human assistant controls the laparoscopic image by directing the laparoscope on the operative field as instructed by the surgeon. The task requires ongoing active communication between the surgeon and the assistant, and confusion or physical space conflicts may arise. Because the surgeon must focus on directing the assistant, he or she is distracted from performing the operation. Human camera images may be suboptimal due to lack of control because of tremor, off-center drift, or loss of horizontal orientation. Therefore, frequent correction is required. Also, inadvertent contact with tissue may result in the need to clean the lens frequently. These problems tend to diminish the concentration of the surgeon and impede the flow of the operation [5].

To alleviate such problems attributable to a human camera assistant, robotic camera systems such as AE-SOP (Computer Motion, Inc., Goleta, CA, USA) or EndoAssist (Armstrong Healthcare Ltd., Loudwater, United Kingdom) [1, 9] have been developed. Studies eonducted to compare the differences between human and robotic control of the laparoscope have shown the robotic system to be superior [1–3, 7].

We recently developed a new robotic laparoscope manipulator that can follow a surgeon's wishes without obstructing his manipulation of the instruments [4, 8]. Anticipating that the manipulator would enable us to overcome such difficulties, we used it during a laparoscopic cholecystectomy of patients with cholelithiasis.

Table 1. The operative data of the Naviot and human camera groups

	Naviot $(n = 10)$	Human-camera $(n = 41)$	p value
Gender (M/F)	3/7	17/24	NS
Age (y.o)	$61.1 \pm 14.8$	$56.9 \pm 12.6$	NS
Operative time (min)	$89.3 \pm 27.1$	$74.8 \pm 28.1$	< 0.05
Blood loss (g)	$7.1 \pm 13.5$	$15.1 \pm 52.4$	NS
Conversion to laparotomy	0	1	NS
Postopertive Hospital stay (day)	$5.7 \pm 13.5$	$8.1 \pm 2.2$	NS
Frequency of analgegia for pain	$0.5 \pm 0.7$	$0.8 \pm 0.7$	NS

#### Patients and methods

#### Patients

From October 2002 to March 2003 at Fukuoka City Hospital, 10 consecutive patients (3 men and 7 women with a mean age of  $61.1 \pm 14.8$  years) with cholelithiasis underwent laparoscopic chole-cystectomy assisted by the new Naviot system (Hitachi Ltd., Tokyo, Japan) (Naviot group) (Table 1). The details of the new system were carefully explained to all the patients, and their informed consent was obtained.

Another group of 41 patients who underwent laparoscopic cholecystectomy with a conventional human camera holder at Fukuoka City Hospital from May 2002 to October 2002 (human camera group) were selected for a comparison of their operative results with those of the Naviot group.

#### The new system: NAVIOT

The Naviot system is a robotic laparoscope manipulator that consists of a special laparoscope with an optical zoom, a flexible arm, an actuator, a five-bar linkage mechanism, and a hand controller with two buttons (Fig. 1). The flexible arm has three joints, and the actuator contains a motor that enables the arm to move. A five-bar linkage mechanism links the actuator and the arm. A specific laparoscope (only about 15 cm long) and half of the conventional laparoscope (with an optical zoom lens) are moved with the arm. The movement range is 45° in horizontal directions and 25° in vertical directions. The zoom can provide 6.0 magnifications. A hand controller consisting of two buttons (one for moving and the other for zooming) is attached to the holding part of a laparoscopic forceps (Fig. 2A). By pressing these buttons, the operator can control the zoom lens and move the laparoscope in eight directions. This new system enables the operator to move the operation field more quickly to the site he or she wants to see. The mark of the moving direction and the zooming are superimposed on the video image. Thus, the operator can always identify the movement of the laparoscope.

#### **Operative** procedure

All operations, including those using a conventional human camera holder, were performed by one individual operator familiar with endoscopic surgery and one or two assistants. The operations with Naviot were performed as follows. First, the actuator and the five-bar linkage mechanism were set up on the right side of the patient opposite the operator so as not to prevent forceps movement by the operator. The flexible arm was covered with a sterilized plastic bag, then connected to the five-bar linkage mechanism.

A skin incision about 12 mm long was made on the upper side of the patient's umbilicus. The laparoscope was inserted through a trocar applied to the incision. Then the laparoscope was connected to the flexible arm, and the arm was fixed. With the use of a 5 mm trocar in the upper midline for the right hand of the operator and two 2 mm trocars in the right upper quadrants for the left hands of the operator and the assistant, the cholecystectomy was performed. The controller of the Naviot was put on the 2 mm forceps close to the grip of the operator's left hand, and the operator controlled the direction or zoom of the laparoscope by pushing the buttons with the left forefinger



Fig. 1. Outlook of the Naviot system.

(Fig. 2B). The trocars were inserted, and the surgical procedures, such as clipping the cystic artery and duct and separating the gallbladder from the liver bed, were the same as in an operation with a conventional human camera holder.

#### Clinical analysis

Each patient's medical record was reviewed for the following information: age; sex; dates of admission, surgery, and discharge; surgical procedure; operative time; intraoperative blood loss; intraoperative and postoperative complications; and postoperative clinical course. The operative time was recorded in terms of minutes from incision to closure. The patient's postoperative physiologic state was evaluated as the frequency of analgesia (pentazocine) used during the 48 h after surgery, the duration of fever ( $>38^{\circ}$ C), the day liquid food was initiated, and the duration of the postoperative hospital stay.

During the operation, we measured the movement range of the laparoscope (in both vertical and horizontal directions), the movement velocity, and the zoom magnification. We also measured the operating time, which was divided into three phases: (A) the time from skin incision to insertion of the laparoscope, (B) the time from insertion of the laparoscope to fixation of the arm, and (C) the time from fixation of the arm to the end of the operation.



Fig. 2. A The controller of Naviot. B The operator controlled the direction or zoom of the laparoscope by pushing the buttons with the left forefinger.



Fig. 3. The mean of the setup time and operative time for each patient including (A) the time from skin incision to insertion of the laparoscope, (B) the time from insertion of the laparoscope to fixation of the arm, and (C) the time from fixation of the arm to the end of the operation.

#### Statistical analysis

The Naviot group and the human camera holder group were compared for significant differences in age, sex, intraoperative and postoperative complications, length of hospitalization, operative time, intraoperative blood loss, and postoperative physiologic state. Data are represented as the mean  $\pm$  standard deviation. The Mann–Whitney U test was used to calculate statistical significance ( $p \le 0.05$ ).

#### Results

#### Operative procedures

Between October 2002 and Mareh 2003, 10 patients underwent laparoseopie cholecysteetomy with the Naviot system at Fukuoka City Hospital. Each patient's mean setup and operative times for phases A, B, and C are shown as Fig. 3.

At this writing, no major complications related to the Naviot system have been encountered in any procedures. The patient in ease 5 with a severe history of cholecystitis had severe adhesion around the gallbladder. Because sufficient vision, even with the highest zooming, could not be obtained due to the long distance between the eamera and the gallbladder, we changed the camera of the Naviot system to the human-held eamera for safe performance of the operation. After this ease, we used the new eamera of the Naviot system with improved zooming. A conversion lens was added to the eamera between the zooming lens and the charged-coupled devices (CCD), which enabled us to obtain a clear vision of the object, which existed in the deeper place. In the first four cases, the controller of the Naviot system, which was fastened to the 2 mm forceps with the clip close to the grip of the operator's left hand, easily slipped down during the operation. To remedy this problem, the controller was fixed at a comfortable position on the 2 mm forceps by driving serces after case 4 (Fig. 3).

The operative time of 89.3  $\pm$  27.1 min in the Naviot group was significantly longer for laparoscopic surgery than the 74.8  $\pm$  28.1 min for the human eamera group (p < 0.05). However, when the setup time (time of B) for the Naviot system was excluded, there was no significant difference in operative time between the Naviot group and the human eamera group. There also was no significant difference in estimated blood loss between the Naviot group (15.1  $\pm$  52.4 g).

#### Postoperative physiologic state

There were no significant differences in the frequency of analgesia (pentazoeine 15 mg) administered for postoperative abdominal pain during the 24 h after surgery, the duration of postoperative fever, the start of liquid food, or the duration of postoperative hospital stay between the two groups (Table 1).

#### Discussion

A new remote-controlled laparoscope manipulator system (Naviot) was introduced for 10 cases of laparoscopic cholecystectomy, and all the procedures were performed safely and efficiently.

The search to find an alternative to a human camera holder for both economy and efficiency first led to the development of mechanical manipulators such as Omnitract, Bookwalter, Boonpong holder, and the like [6]. With these manipulators, the operator must release instruments and interrupt the procedure to make operating field adjustments. Robotic systems designed to avoid this weakness have been introduced in general surgery. A series of 20 patients undergoing laparoscopic cholecystectomy using a self-guiding robot has been reported by Omote et al. [7], who found no significant difference in operating time between their series and historical controls. Aiono et al. [1] reported that the EndoAssist robotic camera holder performed effectively and reduced operating time, as compared with a human camera holder, during laparoscopic cholecystectomy.

With the Naviot system, the operator can move the laparoscope in eight directions smoothly, and can obtain a fine visual field to his or her satisfaction. The hand control adopted as the man-machine interface with this system is superior to voice control in terms of intuitive and delicate control of the laparoscope [4]. What is more, the operator also can move the laparoscopic forceps simultaneously because the hand controller is attached to the holding part of the forceps. Using this system, we were able to perform the laparoscopic cholecystectomy more efficiently.

A zoom lens control also can provide a fine view at some distance from the target. Furthermore, a better view of the surgical field was obtained after improvement of the zooming function. The optical zoom with the laparoscope also is very beneficial. The operator does not need to insert the laparoscope deeply to see the operative field. This means that there is less risk of injuring the organs in the abdominal cavity. A wide workspace also can be made in the abdominal cavity because the laparoscope is 15 cm in the length, which is half that of the conventional laparoscope. Because the tip of this short laparoscope does not protrude from the trocar in the peritoneal cavity, accidental collision with tissue can be avoided in the operation. Additionally, the zoom lens used with the laparoscope is specially made not to collect moisture. Because the operator does not need to pull the laparoscope out to elean the lens so often during the operation, time is saved.

We divided the total operation time into three phases: A, B, and C. Preparation of the system occurred in phase B. Although the overall operative time for the Naviot group was longer than for the human camera group, the net operative time (overall operative time – sctup time [B]) was not significantly longer. Furthermore, the setup time for the Naviot system could be shorter with acquisition of more experience. There was no significant difference in the postoperative physiologic state (frequency of analgesia, duration of postoperative fever, start of liquid food, and duration of postoperative hospital stay) between the two groups. Accordingly, use of the Naviot system for laparoscopic cholecystectomy can be comparable with the human-held camera.

In the first four cases, we had some trouble with insufficient zooming of the camera and frequent slipping of the hand controller from the forceps. We resolved these problems by adding a conversion lens between the zooming lens and the CCD in the camera, and by fixing the controller on the forceps with driving screws. After these improvements, we could obtain a better surgical view, and could be released from the irritation of pushing the wrong buttons and of refixing the controller on the forceps with interruption of surgery.

#### Conclusion

We managed 10 cases of laparoscopic cholecystectomy using the Naviot, a remote-controlled laparoscopic manipulator system. Using the Naviot, we accomplished all the procedures safely and efficiently. We believe this new system is feasible for clinical use. It enables surgeons to perform "solo surgery" in the field of gastrointestinal surgery.

Acknowledgments. This study was partly supported by the Research for the Future Program (JSPS-RFTF 99100902) and the Japan Association for the Advancement of Medical Equipment.

#### References

- Aiono S, Gilbert JM, Soin B, Finlay PA, Gordan A (2002) Controlled trial of the introduction of a robotic camera assistant (EndoAssist) for laparoscopic cholccystectomy. Surg Endosc 16: 1267– 1270
- Hashizume M, Shimada M, Tomikawa M, Ikeda Y, Takahashi I, Abe R, Koga F, Gotoh N, Konishi K, Maehara S, Sugimachi K (2002) Early experiences of endoscopic procedures in general surgery assisted by a computer-enhanced surgical system. Surg Endose 16: 1187–1191
- Kavoussi LR, Moore RG, Adams JB, Partin AW (1995) Comparison of robotic versus human laparoscopic camera control. J Urol 154: 2134–2136
- Kobayashi E, Masamune K, Sakuma I, Dohi T, Hashimoto D (1999) A new safe laparoscopic manipulator system with a five-bar linkage mechanism and an optical zoom. Comp Aid Surg 4: 182–192
- Kondraske GV, Hamilton EC, Scott DJ, Fischer CA, Tesfay ST, Taneja R, Brown RJ, Jones DB (2002) Surgeon workload and motion efficiency with robot and human laparoscopic camera control. Surg Endosc 16: 1523–1527
- Kornsuthisopon S (1999) New release of the Boonpong laparoscopic camera holder. Aust N Z J Surg 69: 141–143
- Omote K, Feussner H, Ungcheuer A, Arbter K, Guo-Qing W, Siewert JR (1999) Self-guided robotic camera control for laparoscopic surgery compared with human camera control. Am J Surg 177: 321–324
- Tomikawa M, Hashizume M, Kobayashi E, Yamaguchi S, Sakuma I, Fujie M, Nakamura F, Dohi T, Shimada M, Sugimachi K (2002) Merits of a newly developed laparoscope manipulator: experiences with 4 cases. In: Lemke U, Vannier MW, Inamura K, Farman AG, Doi K, Reiber JHC (eds) CARS2002-H. The book is the proceeding of CARS 2002. CARS/Springer pp 320–323
- 9. Unger SW, Unger HM, Bass RT (1994) AESOP robotic arm. Surg Endose 8: 1131

Surg Endose (2006) 20: 757–762 DOI: 10.1007/s00464-005-0275-6

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

# Modified needlescopic video-assisted thoracic surgery for primary spontaneous pneumothorax

The long-term effects of apical pleurectomy versus pleural abrasion

#### Y.-C. Chang,<sup>1</sup> C.-W. Chen,<sup>1</sup> S.-H. Huang,<sup>1</sup> J.-S. Chen<sup>2</sup>

<sup>1</sup> Department of Surgery, Far Eastern Memorial Hospital, No. 21, Nan-ya South Road, Sec 2, Banciao City, Taipei, Taiwan, 220
<sup>2</sup> Division of Thoracic Surgery, Department of Surgery, National Taiwan University Hospital and National Taiwan University College of Medicine, No. 7, Chung-Shan South Road, Taipei, Taiwan, 100

Received: 18 April 2005/Accepted: 9 August 2005/Online publication: 25 January 2006

#### Abstract

*Background:* The objective of this study was to evaluate the feasibility and safety of modified needleseopie videoassisted thoracic surgery (VATS) for treating primary spontaneous pneumothorax. The efficacy between apical pleureetomy and pleural abrasion through this technique was also compared.

*Methods:* Between 2001 and 2003, 65 patients with primary spontaneous pneumothorax underwent modified needlescopic VATS procedures. The blebs were resected with endoscopic linear staplers. Pleurodesis was achieved by apical pleurectomy before September 2002 (n = 30) and by pleural abrasion for the remainder of the study period (n = 35).

*Results:* Mean operation time was 103 min in the pleurectomy group and 78 min in the abrasion group (p = 0.001). Complications developed in four patients (6.2%): prolonged air leaks in three patients and wound infection in one patient. The mean postoperative hospital stay was  $3.8 \pm 1.8$  days. The two groups had comparable doses of requested analgesics, complication rates, postoperative chest tube and hospital stays, and postoperative pulmonary function test. Ipsilateral recurrence did not occur in any of the pleurectomy group patients after a mean follow-up of 31 months, but it occurred in three patients (8.6%) in the abrasion group after a mean followup of 19 months.

*Conclusions:* Modified needlescopie VATS provides a feasible and safe procedure for treating primary spontaneous pneumothorax. In terms of efficacy, apical pleurectomy is more effective in preventing ipsilateral recurrence than pleural abrasion.

**Key words:** Needlescope — Video-assisted thoracic surgery — Spontaneous pneumothorax — Pleurectomy — Pleural abrasion

Optimal surgical management of primary spontaneous pneumothorax has been a matter of debate, especially regarding the type of surgical approach and the method of pleurodesis [2, 8]. Recent advances in video-assisted thoracic surgery (VATS) that combine bullectomy with pleural abrasion or apical pleurectomy provide a feasible alternative for treatment of recurrent primary spontaneous pneumothorax and have been chosen as the preferred management by many physicians [11, 17]. However, after conventional VATS, postoperative chest pain has not been completely alleviated [3, 13], and healing of the surgical wounds of conventional VATS has not been cosmetically excellent.

Needlescopic equipment and instruments have been applied to thoracic procedures. Initial reports suggested that needlescopic VATS was feasible in treating primary spontaneous pneumothorax and resulted in better cosmesis and less postoperative pain [4, 9, 18]. However, the number of patients in these studies is small, and the long-term effectiveness of this new technique remains unknown. Because of the inferior vision of the needlescope and inadequate strength of the instruments, pleurodesis is usually not performed or it is accomplished by pleural abrasion [4, 9, 18]. Therefore, the recurrence rate of needlescopie VATS could be higher compared to that of conventional VATS.

In previous studies, apical pleureetomy performed by open thoraeotomy or conventional VATS was shown to be feasible and resulted in a very low ineidence of recurrence [1, 15]. However, apical pleurectomy has not been performed using the needlescopic VATS technique.

Correspondence to: J.-S. Chen



In this study, the technique and experience of modified needlescopic VATS for the treatment of primary spontaneous pneumothorax are reported. Also, the long-term effects and pulmonary functional results of pleural abrasion and apical pleurectomy through this minimally invasive approach are compared.

#### Materials and methods

From June 2001 to October 2003, consecutive patients undergoing a modified needlescopic VATS procedure for the treatment of spontaneous pneumothorax were retrospectively identified through a review of records at the Far Eastern Memorial Hospital, Taipei County, Taiwan. The indications for operation included ipsilateral recurrence, continuous air leaks for more than 72 h, and contralateral recurrence. Patients older than 55 years of age or with preexisting pulmonary diseases were excluded from this study to avoid patients with secondary pneumothorax. In addition, patients who underwent a conventional VATS technique or had previous VATS procedures on the involved side were also excluded from the analyses.

#### Operative techniques of modified needlescopic VATS

The operation techniques for modified needlescopic VATS have been described previously [4]. In brief, under general anesthesia using intubation with a double-lumen endotracheal tube, patients were placed in a lateral decubitus position and the ipsilateral lung was deflated. Two sets of independent videothoracoscopic equipment and monitors (Karl Storz, Tuttlingen, Germany), one for needlescopic videothoracoscopy and the other for 10-mm videothoracoscopy, were used simultaneously and placed near the patient's head. The needle scopic procedures were modified hy using the 10 mm videothoracoscopy of for most of the surgical steps. A needlescope was indicated only when we needed to use the chest tube wound to insert the endoscopic stapler, to extract the specimen, or to perform apical pleurectomy or pleural abrasion.

A 10-mm 30° telescope (Karl Storz) was first inserted through the previous chest tube wound to examine the pleural cavity. If the chest tuhe wound was not available, a 12 mm port was made at the sixth or seventh intercostal space. Three small skin punctures were made, and mini-ports were inserted for needlescopic instruments (3 mm instruments, Olympus, Tokyo). The inferior mini-port for the needlescope was located at the seventh or eighth intercostal space of the middle axillary line, one intercostal space lower than the level of the chest tube wound, to avoid a mirror image during surgical procedures. The two superior mini-ports were located at the fourth and fifth intercostal

Fig. 1. Operative technique of apical pleurectomy via modified needlescopic video-assisted thoracic surgery. (A) Allocation of the skin incisions for chest tube wound and mini-ports: a and b, 3-mm superior mini-ports for mini-endograspers; c, 12-mm chest tube wound for conventional or 10-mm thoracoscopic instruments; and d, 3-mm inferior mini-port for needlescope. (B) Under the vision of the needlescope, a mini-endograsper was used to lift the parietal pleura while it was hluntly stripped off the inner chest wall by inserting the curved dissector with a pledget through the chest tuhe wound (R5, the fifth rib).

space of the anterior and posterior axillary lines, respectively. Initially, the 10 mm telescope and two mini-endograspers were used to identify the blebs. When a bleb was identified, it was fixed by mini-endograsper at one of the superior mini-ports. A 3 mm 30° needlescope was introduced at the inferior mini-port to visualize the bleb. The 10-mm telescope was then withdrawn, and a 45 mm endoscopic linear stapler was introduced to resect the bleb. If no air leakage was identified, apical stapling was routinely performed at the most suspicious area. The surgical specimen was retrieved from the chest tube wound. The 10 mm telescope was inserted again to check the stapling line. Electrocautery was used when bleeding from the stapling line was noted.

Apical pleurectomy or pleural abrasion was performed under the vision of needlescope. Between June 2001 and September 2002, pleurodesis was performed by apical pleurectomy. Between October 2002 and October 2003, pleural abrasion was performed because of its shorter operation time and easier operative technique. Informed consent was obtained from each patient after thorough explanation of the procedures.

For apical pleurectomy, a 3-mm 30° needlescope was introduced through the inferior mini-port. The inferior border of the pleural strip was marked by mini-endohook electrocautery at the upper margin of the fifth or sixth rib, depending on the location of the blebs. The miniendograspers from the superior mini-ports were used to lift and steady the pleura while it was bluntly stripped off the inner chest wall by inserting the curved dissector with a pledget through the chest tube wound (Fig. 1). The dissection was made in an apical direction. The longitudinal limit of the resection ran along the sympathetic trunk posteriorly and the internal mammary artery anteriorly to the height of the left subclavian artery on the left side or the brachiocephalic trunk on the right side.

Pleural abrasion was also performed under the vision of needlescope. The upper half of the parietal surface was abraded hy inserting the curved dissector with a diathermy scratch pad through the chest tube wound. We used the 10 mm videothoracoscope intermittently to ascertain that the upper half of the pleural surface was thoroughly and evenly rubbed.

After the operation, normal saline solution was instilled, and the 10-mm videothoracoscope was introduced to check for air leaks and bleeding. A chest tube was placed in the apex through the chest tube wound. The punctures sites of the 3 mm port were closed with a 4-0 nylon suture. The wounds were then secured with adhesive tape (Steri-Strips). The surgical specimens were routinely sent for pathological examination.

#### Postoperative care

The patients were extubated in the operating theater and observed for 1 or 2 h in the recovery room. The tube was connected to a low-pressure suction system of approximately -10 to -20 cm H<sub>2</sub>O. Post-operative analgesics included routine oral nonsteroid analgesics and

Table 1. Clinical characteristics of patients treated by needlescopic apical pleurectomy or pleural abrasion

	Apical pleurectomy $(n = 30)$	Pleural abrasion $(n = 35)$	<i>p</i> value
Age (vr) <sup>a</sup>	27.5 ± 11.0	$24.2 \pm 7.0$	0.165
Gender (male)	27 (90%)	33 (94%)	0.655
Smoker	11 (37%)	13 (37%)	1.000
Side involved (left)	13 (43%)	22 (63%)	0.140
Surgical indications			0.391 <sup>b</sup>
Ipsilateral recurrence	17 (57%)	17 (49%)	
Persistent air leaks	10 (33%)	14 (40%)	
Contralateral recurrence	3 (10%)	4 (11%)	
Bleb number			0.391 <sup>b</sup>
No bleb	4 (13%)	4 (11%)	
One or two blebs	17 (57%)	23 (66%)	
Multiple blebs (≥3 blebs)	9 (30%)	8 (23%)	
Blood loss <sup>a</sup>	$94.1 \pm 23.9$	$85.0 \pm 35.1$	0.236
Surgery duration (min) <sup>a</sup>	$102.5 \pm 31.5$	77.5 ± 28.3	0.001

<sup>a</sup> Mean  $\pm$  standard deviation

<sup>b</sup> Analyzed by Pearson chi-square test

acctaminophen. Intramuscular meperidinc hydrocbloride (Demerol, 50 mg/ampule) was administered every 4–6 h according to the patient's request if the pain became intolerable or could not be relieved by oral analgesics. The tube was removed when the lung was fully expanded and no air leaks could be noted in a 24-b period. Patients were discbarged I day after the removal of the drainage tube if no complications occurred.

#### Postoperative pulmonary function

Postoperative pulmonary function tests were performed for patients able to attend a hospital outpatient appointment at least 6 months after surgery. Forced vital capacity (FVC) and forced expiratory volume in 1 s ( $FEV_{1,0}$ ) were measured using a Microspiro HI-298 spirometer (Chest Corporation, Tokyo) with the patients seated.

#### Data collection and analyses

The clinical data, operative findings, operation time, duration of postoperative cbest drainage, length of bospital stay, and complications were collected from the medical records. The requested doses of meperidine were collected from the nursing records. Patients were followed up by clinical visits for at least 12 months and by telephone conversation thereafter. Continuous variables such as age or weight were expressed as mean  $\pm$  standard deviation and analyzed by the two-sample *t*-test. Categorical variables such as gender or smoking status were presented by frequency (%) and were analyzed by Fisher's exact test or by Pearson's cbi-square test. The rates of freedom from recurrence were estimated using the Kaplan–Meier metbod. The logrank test was used to analyze differences in recurrence. Statistical analyses were performed using SPSS 10.0 software. A *p* value of less than 0.05 was considered significant.

#### Results

Sixty-five patients who had undergone a needleseopie VATS procedure for primary spontaneous pneumothorax were identified at our institution. The mean age was 25.5 years (range, 16–52; 60 males and five females); 24 patients (37%) were smokers. The indications for operation were ipsilateral recurrence in 34 (52.3%), persistent air lcaks in 24 (36.9%), and contralateral recurrence in seven (10.8%). Necdlescopie VATS was unilateral in all eases. Blebs or bullae were identified in 57 patients (87.7%), and multiple blebs ( $\geq$ 3 blebs) were visualized in 17 of these patients (26.2%). The most common site for blebs was the upper lobe (51 patients, 78.5%).

The operative findings and management are summarized in Table 1. The two groups of patients had eomparable elinical characteristics in terms of age, sex, smoking status, surgieal indication, bleb number, and estimated blood loss. The mean operation time was 103 min in the pleureetomy group and 78 min in the abrasion group (p = 0.001). The results of treatment are summarized in Table 2. Intramuscular meperidine was requested by 58% of the patients, mainly on postoperative day 1; the mean accumulated dosage of injection was eomparable for both groups. No operative deaths and no major complications were reported. Three patients (4.7%) had air leaks lasting longer than 5 days; one patient had infection of the ehest tube wound. These patients were managed eonservatively. The eomplication rate for both groups was eomparable. The mean postoperative hospital stay was 3.9 days (range, 2–10) in the pleurectomy group and 3.8 days (range, 2-12) in the abrasion group.

The patients underwent a postoperative follow-up ranging from 12 to 39 months, with a median follow-up of 25.4 months. Patients in the pleurectomy group had a significantly longer mean follow-up than the abrasion group (31.2 vs 19.4 months). Ipsilateral recurrence of pneumothorax occurred in three patients (8.6%) treated by pleural abrasion but in none of the patients treated by apieal pleureetomy. The recurrences occurred 4, 6, and 24 months after the operation, respectively. The rates of freedom from recurrence after the operation for both groups were determined using the Kaplan-Meyer method and are plotted in Fig. 2 and compared by the log-rank test. Patients who underwent apical pleureetomy had a significantly lower rate of ipsilateral recurrence (p = 0.024). When the operative findings were reviewed, no leaking site could be identified in one patient, and multiple small blebs were noted in two other patients. Two of three recurrent patients underwent a reoperation by eonventional VATS. Missed blebs were identified in the vieinity of the previous staple line. No

Table 2. Results of patients treated by needlescopic apical pleurectomy or pleural abrasion

	Apical pleurectomy $(n = 30)$	Pleural abrasion $(n = 35)$	p value
Meperidine hydrochloride (mg) <sup>a</sup>	$56 \pm 56$	51 ± 69	0.746
Postoperative chest tube duration (d) <sup>a</sup>	$3.2 \pm 1.6$	$3.1 \pm 1.1$	0.812
Postoperative hospital stay (d) <sup>a</sup>	$3.9 \pm 1.7$	$3.8 \pm 1.5$	0.860
Complications			
Air leaks $> 5 d$	2 (6.7%)	1 (2.9%)	0.591
Wound infection	0 (0%)	1 (2.9%)	1.000
Follow-up (mo) <sup>a</sup>	$31.2 \pm 5.3$	$19.4 \pm 3.2$	< 0.001
Recurrence	0 (0%)	3 (8.6%)	0.243
Pulmonary function test <sup>b</sup>			
FVC $(\%)^{c}$	$86.3 \pm 9.8$	$88.5 \pm 11.8$	0.574
$FEV_{1,0}$ (%) <sup>e</sup>	94.9 ± 11.2	$99.0 \pm 13.9$	0.378

FVC, forced vital capacity; FEV<sub>1.0</sub>, forced expiratory volume in 1 s

<sup>a</sup> Mean  $\pm$  standard deviation

<sup>b</sup> Fifteen patients in the pleurectomy group and 25 patients in the abrasion group were available for pulmonary function test.

<sup>c</sup> Represented as percentage of predicted value



obvious adhesions were noted in the whole pleural eavity, even in the areas of previous pleural abrasion. The blebs were excised, and pleurodesis was accomplished by apical pleurectomy. The remaining patient was treated by chest tube insertion.

#### Discussion

This study demonstrates that modified needlescopic VATS is a feasible and safe procedure for treating primary spontaneous pneumothorax. The results also show that apieal pleurectomy is more effective than pleural abrasion in obtaining pleural symphysis and freedom from recurrence without increased pain, complications, or comorbidities. Fig. 2. Freedom from recurrent pneumothorax in each group after modified needlescopic videoassisted thoracic surgery (p = 0.024 by log-rank test). The number of patients at risk for each 6month period is indicated under the corresponding time.

Needlescopic surgery generally refers to procedures that use instruments with an external diameter  $\leq 3 \text{ mm}$ [6, 10]. Although needlescopic surgery represents the next logical step in the evolution of VATS, there are technical limitations associated with the needlescopic instruments [12, 18]. Scopes with a small needle size provide poor visibility. Furthermore, the mini-instruments are inconvenient for grasping the bulky lung parenchyma and performing pleurodesis because of their small jaws and inadequate strength.

For patients with spontaneous pneumothorax, an ineision of at least 10–12 mm for a ehest tube was usually present before the operation or indicated after the operation. This incision made it possible for us to use a 10-mm thoracoscope for most of the surgical steps. In addition, the endoscopic linear stapler for pulmonary

resection and the curved ring forceps or dissector for pleural abrasion and pleureetomy can be introduced through this wound. Taking advantage of this wound, modified needlescopic VATS can be safely and easily performed with no compromise of the surgical procedures [4].

Although modified needlescopic VATS is feasible and safe, our experience showed that needlescopic bullectomy and pleural abrasion carry a higher rate of recurrence (8.6%) compared to the same procedures performed by conventional VATS [11, 17]. The early postoperative recurrences of pneumothorax are generally thought to be due to overlooked existing pulmonary blebs. It is possible that needlescopic VATS leads to a higher risk of missing blebs than conventional VATS. In this situation, a strong and adequate pleural adhesion is indicated to prevent recurrence of the pneumothorax. Our long-term recurrence rate is 0 in the pleurectomy group, indicating that apical pleurectomy can provide reliable and durable pleural symphysis to prevent further recurrence, even using needlescopic procedures.

Since the introduction of surgical intervention for spontaneous pneumothorax, pleurectomy has been a reliable method for pleurodesis. The procedure has a very low ineidence of recurrence, ranging from 0 to 1%, by open thoracotomy, transaxillary minithoraeotomy, or VATS [1, 15]. British Thoracic Society guidelines suggest that pleural abrasion is associated with a higher rate of recurrence than pleureetomy [8]. Compared to pleurectomy, pleural abrasion is a procedure that has problems associated with standardization of the material used as well as the force and time needed for abrasion. Postoperatively, we usually found that the parietal pleura were unevenly abraded. In the two patients who underwent reoperation after recurrence, only scanty adhesions were noted in the pleural eavity. In contrast, parietal pleurectomy created a dense and even inflammatory surface between the pleura and the chest wall [6]. Several authors have since described pleurectomy as the most secure procedure to obtain permanent pleurodesis [6, 14]. Since the disease is almost always located at the apex of the lung, we believe that in young people a limited apical pleurectomy combined with local excision of the diseased lung is sufficient for definitive control of recurrences. In addition, if lung blebs are found and resected in the middle or lower lobes, pleurectomy can also be extended to a lower level.

Although dense plcurodesis may limit the expansion of the lungs, there is no evidence that apical pleurectomy significantly affects pulmonary function. Previous studics showed that the mean vital capacity of patients with previously documented pneumothorax who did not have surgery was 87 or 88% of the normal predicted values [5, 16]. By comparison, our data showed that the vital capacity following either pleural abrasion or apical pleurectomy did not alter significantly. In addition, FVC and FEV<sub>1.0</sub> were comparable for both groups of patients.

The intensity of postoperative pain caused by apical pleureetomy or pleural abrasion was also evaluated by comparing the requested doses of meperidine. Our data showed that the incidence and mean doses of meperidine injection were comparable for both groups. This suggests that apical pleurectomy may not induce stronger pain than pleural abrasion.

Some researchers do not favor pleurectomy as a routine pleurodesis procedure due to the following pitfalls. The surgery is more technically demanding, and the operation time is significantly longer. Thus, the cost associated with the operation is higher. In addition, apical pleurectomy makes further operations more difficult. In our experience, the operation time can be significantly shortened after the learning curve, and the expense of difficult further operations can be compensated by the decrease in short-term recurrence.

We acknowledge that this is a retrospective study and the comparison was based on a historical control. However, our results suggest that needlescopic VATS is feasible and safe, and it may provide satisfactory results if apical pleurectomy is used. Further prospective research that includes randomized design is needed to rigorously test the effects of this treatment.

#### References

- Ayed AK, Al-Din HJ (2000) The results of thoracoscopic surgery for primary spontaneous pneumothorax. Chest 118: 235–238
- Baumann MH, Strange C, Heffner JE, Light R, Kirby TJ, Klein J, Luketich JD, Panacek EA, Sahn SA: AACP Pneumothorax Consensus Group (2001) Management of spontaneous pneumothorax: an American College of Chest Physicians Delphi consensus statement. Chest 119: 590–602
- Bolotin G, Lazarovici H, Uretzky G, Zlotnick AY, Tamir A, Saute M (2000) The efficacy of intraoperative internal intercostals nerve block during video-assisted thoracic surgery on postoperative pain. Ann Thorac Surg 70: 1872–1875
- Chen JS, Hsu HH, Kuo SW, Tsai PR, Chen RJ, Lee JM, Lee YC (2003) Needlescopic versus conventional video-assisted thoracoscopic surgery for primary spontaneous pneumothorax: a comparative study. Ann Thorac Surg 75: 1080–1085
- De Troyer A, Yernault JC, Rodenstein D, Englert M, De Coster A (1978) Pulmonary function in patients with primary spontaneous pneumothorax. Bull Eur Physiopathol Respir 14: 31–39
- Gaensler EA (1956) Parietal pleurectomy for recurrent spontaneous pneumothorax. Surg Gynecol Obstet 102: 293–308
- Gagner M, Carcia-Ruiz A (1998) Technical aspects of minimally invasive abdominal surgery performed with needlescopic instruments. Surg Laparosc Endosc 8: 171–179
- Henry M, Arnold T, Harvey J; Pleural Diseases Group, Standards of Care Committee, British Thoracic Society (2003) BTS guidelines for the management of spontaneous pneumothorax. Thorax 58(Suppl 2): 39–52
- Ikeda Y, Miyoshi S, Seki N, Kobayashi S, Umezu H, Tamura M (2003) Needlescopic operation for partial lung resection. Ann Thorac Surg 75: 599–601
- Mamazza J, Schlachta CM, Seshadri PA, Cadeddu MO, Poulin EC (2001) Needlescopic surgery: a logical evolution from conventional laparoscopic surgery. Surg Endosc 15: 1208–1212
- Nakamoto K, Maeda M, Okamoto T, Kameyama K, Sugita A, Hayashi E (1998) Preoperative diagnosis with video-assisted thoracoscopy with miniaturized endoscopes in general thoracic surgery: a preliminary study. Chest 114: 1749–1755
- Naunheim KS, Mack MJ, Hazelrigg SR, Ferguson MK, Ferson PF, Boley TM, Landreneau RJ (1995) Safety and efficacy of video-assisted thoracic surgical techniques for the treatment of spontaneous pneumothorax. J Thorac Cardiovasc Surg 109: 1198–1204
- Passlick B, Born C, Sienel W, Thetter O (2001) Incidence of chronic pain after minimal-invasive surgery for spontaneous pneumothorax. Eur J Cardiothorac Surg 19: 355–358

- Saha SP, Arrants JE, Kosa , Lee WH Jr (1975) Management of spontaneous pneumothorax. Ann Thorac Surg 19: 561–564
- Weeden D, Smith GH (1983) Surgical experience in the management of spontaneous pneumothorax, 1972–82. Thorax 38: 737–743
- Williams MH, Kane C (1965) Pulmonary function in patients who have recovered from spontaneous pneumothorax. Dis Chest 47: 153–156
- Yim AP, Ho JK (1995) One hundred consecutive cases of videoassisted thoracoscopic surgery for primary spontaneous pneumothorax. Surg Endosc 9: 332–336
- Yoon YH, Kim KH, Han JY, Baek WK, Lee CS, Kim JT (2000) Management of persistent or recurrent pneumothorax with a two millimeter mini-videothoracoscope. J Korean Med Sci 15: 507– 509

Surg Endosc (2006) 20: 763–769 DOI: 10.1007/s00464-004-2231-2

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

## Systematic evaluation of different approaches for minimizing hemodynamic changes during pneumoperitoneum

T. Junghans,<sup>1</sup> D. Modersohn,<sup>2</sup> F. Dörner,<sup>1</sup> J. Neudeeker,<sup>1</sup> O. Haase,<sup>1</sup> W. Sehwenk<sup>1</sup>

<sup>1</sup> Department of General, Visceral, Vascular, and Thoracic Surgery, Charité Universitary Medicine, Campus Mitte, Schumannstrasse 20/21, 10117 Berlin, Germany

<sup>2</sup> Department of Cardiovascular Surgery and Experimental Cardiac Surgery, Charité Universitary Medicine, Campus Mitte, Schumannstrasse 20/21, 10117 Berlin, Germany

Received: 23 September 2004/Accepted: 22 September 2005/Online publication: 25 January 2006

#### Abstraet

*Background:* Capnoperitoneum (CP) compromises hemodynamic function during laparoseopy. Three therapeutic concepts were evaluated with an aim to minimize the hemodynamic reaction to CP: First, a controlled increase of intrathoracic blood volume (ITBV) by intravenous fluids; second, partially reduced sympathetic activity by the  $\beta_1$ -blocker esmolol; and third, a decrease in mean arterial pressure (MAP) by the vasodilator sodium nitroprusside.

*Methods:* For this study, 43 pigs were assigned to treatment with fluid and sodium nitroprusside (group A) or with esmolol (group B). In both groups, the pigs were assigned to head-up, head-down, or supine position, resulting in three different subgroups. Invasive hemo-dynamic monitoring was established including left heart eatheter and cardiac oxygen lung water determination (COLD) measurements. Measurements were documented before CP with the animals in supine position, after induction of a 14-mmHg CP with the animals in each body position, after a 10% reduction in MAP by vasodilation, and after an increase in 1TBV of about 30% by infusion of 6% hydroxyethylstareh solution.

*Results:* Increasing ITBV improved hemodynamic function in all body positions during CP. Esmolol reduced eardiae output and myoeardial contractility. Sodium nitroprusside did not improve hemodynamic function in any body position.

Correspondence to: T. Junghans

Conclusions: Optimizing volume load is effective for minimizing hemodynamic changes during CP in the head-up and in head-down positions. In general,  $\beta_1$ blockers cannot be recommended because they might additionally compromise myocardial contractility and suppress compensatory reaction of the sympathetic nerve system. Vasodilation has not improved hemodynamic parameters during CP.

Key words: Fluid Management — Hemodynamic — Laparoseopy — Pathophysiology — Penumoperitoneum — Therapy

Capnoperitoneum (CP) during laparoseopie surgery eauses specifie hemodynamic ehanges. Furthermore, eardiae function during laparoseopy might be additionally impaired in patients with preexisting cardiovaseular diseases. With knowledge of the physiologic background and the therapeutic approaches available to minimize hemodynamic ehanges, the number of patients excluded from laparoseopie surgery can be reduced to a minimum.

During CP, a decrease in eardiae stroke volume and eardiae output often has been described as well as an increase in mean arterial pressure or heart rate [2, 6, 9, 11, 17, 24]. These changes are eaused by an increase in the intraabdominal pressure (IAP), which is supposed to reduce venous blood return to the heart [4, 7, 18–20] followed by a decrease in intrathoracie blood volume (ITBV) or left ventrieular end diastolie volume (LVEDV) [5, 14, 16, 25]. Either ITBV or LVEDV is useful for assessing eardiac preload. Cardiae preload determines cardiae function according to the Frank– Starling mechanism. A lower eardiae preload results in less filling of the ventrieles, with a consecutively deereased cardiac stroke volume. The reduction in stroke

Parts of this article were presented at the 119th and 120th meetings of the German Surgical Society, 2002 in Berlin, Germany, and 2003 in Munich, Germany, at the Congress of Anaesthesiology and Surgical Intensive Care Medicine-HAI 2003 in Berlin, Germany, and at the meeting of the European Society of Surgery (ESS) 2003 in Prague, Czech Republik. Parts of this article also were published as an abstract in *Langenbecks Arch Surg* (Supp) 2002;31:499 and *Langenbecks Arch Surg* (Supp) 2003;32:503.

volume is recognized by receptors in the carotid sinus, increasing sympathetic nerve activity. Thus, mean arterial pressure or heart rate is increased during CP.

Experimental and clinical studies have suggested that hemodynamic function during laparoscopy might be improved by increasing intravascular volume or medication with esmolol or nitroglycerine [3, 10, 13], but these different therapeutic strategies have never been evaluated systematic. Thus, according to the physiologic causality, different therapeutic approaches were compared. In this animal trial, an increase in intrathoracic blood volume by infusion therapy, a reduction in mean arterial pressure or heart rate by  $\beta_1$ -blockers, or vasodilation by sodium nitroprusside was induced during CP. Because body position is well known to influence hemodynamic function during CP, the animals were divided into different groups to represent head-up, headdown, and supine positions.

#### Materials and methods

The hemodynamic reactions of 43 pigs with a body weight of  $30 \pm 2.6$  kg were observed during CP with an 1AP of 14 mmHg. The pigs were assigned randomly to treatment with fluid and sodium nitroprusside (group A) or with esmolol (group B). In botb groups, the animals were divided according to three different body positions:  $30^{\circ}$  head-up,  $30^{\circ}$  head-down, and supine. Cardiovascular monitoring included measurement of parameters to assess eardiae preload, eardiac afterload, and myocardial contractility. The study was approved by the animal welfare committee of the city of Berlin, Germany (Reg.Nr. 0026/00).

#### Anesthetic regimen

The day before the experiment, the pigs were allowed to drink water ad libitum. The animals were premedicated with a 400-mg intramuscular metomidate (bypnodil). With the animals in supine position general anesthesia was induced with etomidate 6 mg, fentanyl dihydrogencitrat 0.1 mg, and eisatracurium 10 mg. Anesthesia was maintained by ventilation with 1% to 2% isofluran and by intermittent injections with cisatracurium and fentanyl. Blood gases were analyzed during every measurement as well as end-tidal carbon dioxide (CO<sub>2</sub>). The frequency of mechanical ventilation was adjusted to end-tidal CO<sub>2</sub> to maintain normocapnia. No hyperventilation was necessary during the measurements. The animals received 0.2 ml/min/kg of electrolyte solution. To avoid blood clotting around the vascular catheters, a bolus with 5,000 1.U. heparin followed by a continuous rate of 1,000 1.U. per hour was given intravenously. After all the measurements, the animals were killed by intravenous injection of T61 (0.3 ml/kg).

#### Monitoring

After induction of general anestbesia, a central venous catbeter was placed in the superior cava vein to measure central venous pressure (CVP). Heart rate was counted by electrocardiogram (ECG). A 4-F oximetry-thermodye-dilution catbeter (Pulsion Medical Systems Inc, Minieb, Germany) was placed in the descending aorta to measure mean arterial pressure MAP. The eatheters were connected to the cardiae oxygenation lung water determination (COLD) monitoring system (PULSION COLD Z-021; Pulsion Medical Systems Inc). The ITBV which mainly influences cardiae preload was calculated by the monitoring system as well as cardiac output (CO) and cardiac stroke volume. Systemic vascular resistance (SVR) was calculated as SVR = (MAP - CVP)  $\times$  79.9 / CO. Myocardial contractility, independent of load conditions and heart rate, is best described by myocardial contractility (Emax) [21, 22], and was measured via left beart conductance catheter (SPC 572 Millar Mikro-Tip catbeter;

Krauth Medical Cardiovascular, Hamburg, Germany). The catheter consists of two pressure sensors placed in the left ventricle and the ascending aorta via carotid arteriotomy. Volumes were calculated by 12 electrodes using the technique of conductance. Emax is characterized as the slope of the straight line that represents the relation between intraventricular pressure and volume at the end of the systole after different loaded cardiac cycles. The pressure–volume relation was modified using transient vena eaval and descending aortic occlusion to imitate acute changes of eardiac pre- and afterload.

#### Measurements

Before all measurements, the animals received 1,000 ml of 0.9% NaCl solution intravenously to avoid bypovolemia. After placement of all the catheters and an adaptation of 20 min, baseline values were recorded. The animals were assigned randomly to be placed in supine, 30° head-up, or 30° head-down position. In group A, a CP of 14 mmHg was established and kept by an insufflator (Thermoflator, Storz, Germany) via a port in the right abdomen. Values during CP were recorded after another adaptation of 15 min. The following measurement of lemodynamic parameters was conducted after a 10% decrease in MAP by continuous infusion with the vasodilatator so-dium nitroprusside. The vasodilation was stopped after the measurement. The CP was released, and the animals were placed back in supine position.

After 30 min of adaptation, ITBV was increased by 30% up to a maximum infusion of 1.5 1 6% bydroxyetbyl stareb solution. The animals were again placed in their previous position combined with a CP of 14 mmHg. Measurements were repeated after another adaptation of 15 min. In group B, a bolus with 1mg/kg of 1be ultrasbort effective  $\beta_1$ -blocker esmolol was given intravenously after the baseline measurements just before CP was established. Esmolol medication was adjusted to a rate of 200 µg/kg/min related to the balf-life period. Values during CP were recorded after another adaptation of 15 min.

#### Statistical analysis

The hypothesis was that increasing intravascular volume during CP improves the bemodynamic function. The main end point regarding bemodynamic function was stroke volume. The number of animals was restricted to a maximum of nine animals per group by the animal welfare committee. Eight animals were excluded from the study because of bemodynamic decompensation (n = 5), bleeding (n = 2), or embolic complication (n = 2) in the initial phase of the experiment. Because completion of eight animals per group was not possible with the remaining animals, the trial was closed with seven animals per group except for cight animals in group A (bead-down).

Periodically calculated parameters are given as the mean of three measurements. Values are presented as median and 5 to 95 percentile. The Wilcoxon test was used to assess differences between the measurements. The relative differences between the baseline values and the measurements were compared between the three body positions using the Kruskal-Wallis-test. Differences between group A and B are presented as box and whisker plots, and were compared using the Mann-Whitney-U test (Figs. 1–5). All p values less than 0.05 were considered significant. For statistical analysis, the SPSS 11.0 (Chicago, Illinois, USA) software was used.

#### Results

Baseline measurements were not different between the three body positions. In the supine position, CP was well tolerated (Table 1), whereas CP in the head-down position caused some negative effects. Intrathoracic blood volume and stroke volume decreased, whereas systemic vascular resistance increased (Table 2). The negative


Fig. 1. Differences between groups A and B with regard to changes in heart rate during capnoperitoneum, as compared with baseline values.



	baseline M1	CP M2	CP + nipruss M3	CP + fluids M4
HR per min	101 (73-127)	97 (73–139)	87 (72–135)	91 (73-104)
CO (l/min)	4 (2.8-5.7)	4 (2.8-4.9)	$3.2(2.4-3.9)^{a}$	$5(4.3-6.4)^{a}$
SV (ml)	49 (36-65)	48 (27-58)	$38(25-51)^{a}$	$63(58-78)^{a}$
ITBV (ml)	557 (485-670)	575 (477-597)	$491 (411 - 534)^{a}$	$718(612-811)^{a}$
SVR $(dyn/s/cm^{-5})$	1370 (926-1876)	1537 (1.078-1.602)	1388 (1,047-1,710)	999 (742-1162) <sup>a</sup>
MAP (mmHg)	78 (67-109)	78 (65-99)	$62(54-72)^{a}$	$84(78-107)^{a}$
Emax mm Hg per ml	3.9 (2.9-4.4)	3.6 (2.7-4.3)	3.8 (2.6-4)	3.2 (2.7-4)

nipruss, nitroprusside; HR, heart rate; CO, cardiac output; SV, stroke volume; ITBV, intrathoracic blood volume; SVR, systemic vascular resistance; MAP, mean arterial pressure; Emax, myocardial contractility; Hg, mercury; M, measurement  $^{a} p \leq 0.05$  compared with M2, Wilcoxontest

Table 2 Hemodynamic of	changes during ea	ipnoperitoneum (CP)	in group A head-down	nosition mediau	1 (5-95 percentile)

	baseline M1	CP M2	CP + uipruss M3	CP + fluids M4
HR per min	90 (83-115)	95 (78-119)	111 (73–126)	105 (85-168)
CO (l/min)	4.2 (3.9-5.6)	3.9 (3.3-5.8)	$3.5(2.9-5.4)^{a}$	$5.2(4.4-9.3)^{a}$
SV (ml)	56 (48-73)	49 (37-69) <sup>b</sup>	$43(31-51)^{a}$	$61(58-67)^{a}$
ITBV (ml)	624 (597-906)	584 (519-833) <sup>b</sup>	$520 (470 - 747)^{a}$	756 (721–945) <sup>a</sup>
SVR $(dyn/s/cm^{-5})$	1034 (788-1,259)	1199 (936–1,572) <sup>b</sup>	1045 (791-1455)	774 (608-1,010) <sup>a</sup>
MAP (mmHg)	69 (51-78)	77 (61-88)	$62 (45-70)^{a}$	81 (73-96)
Emax mm Hg per ml	4 (3.1-4.3)	4.1 (3.2-4.6)	4.5 (3.1-4.8)	4.2 (3-5.4)

nipruss, nitroprusside; HR, heart rate; CO, cardiae output; SV, stroke volume; 1TBV, intrathoracic blood volume; SVR, systemie vascular resistance, MAP, mean arterial pressure; Emax, myocardial contractility; Hg, mercury; M, measurement

 $p^{a} < 0.05$  compared with M2, Wilcoxontest

 $p = p^{\prime} < 0.05$  compared with M1, Wileoxontest

influences were more pronounced during CP in the head-up position (Table 3).

During CP in the head-down position, intrathoracie blood volume was reduced about 7%, with a consecutive decrease in stroke volume of about 13%, whercas systemic vascular resistance increased about 16%. In spite of the reduced stroke volume in combination with an unchanged heart rate, eardiac output remained unchanged as compared with base line measurements (Table 2).

During CP in the head-up position, intrathoraeie blood volume was decreased by 16% and stroke volume by 29%. Systemic vascular resistance increased by 12%, and MAP slightly decreased by 6%. Cardiac output also was reduced by 23% (Table 3). The following parameters were different between the body positions during CP: intrathoracic blood volume (p < 0.01), MAP (p < 0.05), stroke volume (p < 0.05), and eardiac output (p < 0.05). All these parameters were significantly improved by an inerease in intrathoracie blood volume during CP in all body positions because MAP increased to normal ranges and stroke volume as well as eardiae output were elevated. With the better hemodynamie function, systemie vaseular resistance decreased (CP + fluids in Tables 1–3).

The reduction in MAP during CP by infusion of sodium nitroprusside did not improve hemodynamic parameters independently of the body position. Mean arterial pressure was decreased without differences be-

	baseline M1	CP M2	CP + nipruss M3	CP + fluids M4
HR per min	95 (56-110)	99 (76-131)	95 (86-151)	107 (82-144)
CO (l/min)	4.7 (3.4-5.5)	$3.6(2.7-4.4)^{a}$	2.9 (2.5-3.9)	$6.1(5.2-7.3)^{b}$
SV ml	55 (45-69)	$39(30-59)^{a}$	33 (23-44)	55 (34-86) <sup>b</sup>
ITBV ml	630 (542-843)	527 (439–769) <sup>a</sup>	497 (397-685)	755 (669–937) <sup>b</sup>
$SVR(dvn/s/cm^{-5})$	1135 (944-1454)	$1274(1.042-1.762)^{a}$	1324 (983-1,690)	$1053 (912 - 1, 173)^{b}$
MAP (mmHg)	71 (60-85)	67 (55–76) <sup>a</sup>	58 (47-69) <sup>b</sup>	96 (80–101) <sup>b</sup>
Emax mm Hg per ml	3.9 (2.8-4.8)	3.6 (2.9-4.6)	$3.2(2.6-4.7)^{\rm b}$	3.5 (3-4.3)

Table 3. Hemodynamic changes during capnoperitoneum (CP) in group A head-up position, median (5-95 percentile)

nipruss, nitroprusside; HR, heart rate; CO, cardiac output; SV, stroke volume; 1TBV, intrathoracic blood volume; SVR, systemic vascular resistance; MAP, mean arterial pressure; Emax, myocardial contractility; Hg, mercury; M, measurement

 $p^{a} < 0.05$  compared with M1, Wilcoxontest

<sup>b</sup> p < 0.05 eompared with M2, Wilcoxontest



Fig. 2. Differences between groups A and B with regard to changes in cardiac output during eapnoperitoneum, as compared with baseline values.



tween the body positions (p = 0.17). Vasodilation induced a reduction in intrathoracic blood volume during CP in the supine and in head-down positions, as compared with CP alone. The increased reduction in intrathoracic blood volume caused a greater fall in stroke volume and cardiac output in these positions. Changes in intrathoracic blood volume, systemic vascular resistance, or cardiac output did not occur in the head-up position (CP + nitroprusside in Table 1–3). Fig. 3. Differences between groups A and B with regard to changes in stroke volume during capnoperitoneum, as compared with baseline values.

The medication with the  $\beta_1$ -blocker esmolol had negative effects on hemodynamic parameters during CP in all body positions because heart rate (Fig. 1) and consecutively cardiac output (Fig. 2) decreased, whereas stroke volume (Fig. 3) and mean arterial pressure (Fig. 4) remained unchanged. Emax was compromised during CP in the head-up and in head-down positions (Fig. 5). With the decrease in cardiac output, systemic vascular resistance increased in the head-up position (Fig. 6).

mean arterial pressure [MAP] M2 in % of M1



Fig. 4. Differences between groups A and B with regard to changes in mean arterial pressure during capnoperitoneum, as compared with baseline values.



Fig. 5. Differences between groups A and B with regard to changes in myocardial contractility during capnoperitoneum, as compared with baseline values.





Fig. 6. Differences between groups A and B with regard to changes in systemic vascular resistance during capnoperitoneum as compared with baseline values.

# Discussion

# Increasing intrathoracic blood volume

Intrathoracic blood volume decreased during CP in the head-up and in head-down positions. An increase in intrathoracic blood volume by infusion of colloidal fluids improved hemodynamic function in all the body positions. This therapeutic approach was suitable for counteracting the decrease in stroke volume during CP caused by a compromised venous blood return and consecutively reduced intrathoracic blood volume. These mechanisms seem to be the most predominant determining factors of hemodynamic function during laparoscopic procedures. Thus, sympathetic reactions such as an increase in MAP or heart rate above normal values were avoided and stroke volume as well as cardiac output were improved. An earlier experimental study from our group showed that all the following conditions during laparoscopy independently influence cardiac output and stroke volume: the increase in intravascular volume, the body position, and the IAP [10].

In the current trial, the hemodynamic system was negatively influenced only in the head-up and in headdown positions, but not in the supine position. However, during almost all laparoscopic procedures, headup or head-down positioning is necessary to optimize the intraperitoneal view, so the exact combination of IAP and body position is of major importance regarding hemodynamic changes. Therefore, the interpretation of different hemodynamic data during laparoscopy must always consider body position, amount of IAP, and presumed vascular volume load.

Although 1 I saline solutions were infused preoperatively, animal hypovolemia during the measurements without an additionally increased preload cannot be excluded because normal ranges of intravascular volumes in pigs are not defined. Nevertheless, such a slight hypovolemia may very well reflect the clinical preoperative volume status in patients after preoperative fasting in combination with bowel-cleaning procedures.

The positive effect of increased venous blood return during laparoscopic surgery in the head-up position also was described in a clinical observational study. These authors of the study used intermittent pneumatic compression devices of the lower extremities to improve venous blood return by calf muscle compression. Usually, these devices, are used for prophylaxis of deep vein thrombosis. With the use of these devices venous blood return and cardiac index increased significantly during CP in the head-up position [1]. Increasing intravascular volume by infusing of 500 ml of hydroxyethyl starch increased cardiac output in another nonrandomized clinical trial during gynecologic laparoscopy in the case of presumed decreased volume preload [23].

# Reducing sympathetic nerve activity

Medication with esmolol during CP decreased cardiac output in all body positions, as compared with CP alone. Because stroke volume remained unchanged, two effects of esmolol contributed to that reduction: the decrease in heart rate and myocardial contractility. Only in the head-up position, did systemic vascular resistance increase. If the sympathetic reaction during CP is considered to compensate for some negative hemodynamic effects such as the fall in stroke volume, esmolol might additionally compromise the hemodynamic system by locking this compensatory effect. However, the effects of esmolol during CP were interpreted differently in a blinded randomized clinical trial. Perioperative medication with csmolol reduced both heart rate and MAP during laparoscopic cholecystectomy in the head-up position. With the decrease in MAP, both the adrenaline and the antidiuretic hormone release were increased in the esmolol group [13]. Although the authors interpreted the fall in MAP and heart rate as improvement in the hemodynamic function they could not exclude, the possibility that esmolol compromises an increased sympathetic activity during CP. Medication with cloni-

dinc during CP may have effects comparable with those of esmolol. During laparoscopic cholecystectomy in head-up position, the catecholamine release decreased after medication with clonidine whereas both vasopressin and cortisol levels remained unchanged. Hemodynamic parameters were presumed to be improved because MAP, heart rate, and systemic vascular resistance were decreased. Cardiac output was not changed in the clonidine group during CP, but decreased after release of CP [8]. Clonidine also was effective in reducing MAP and heart rate during laparoscopic cholecystectomy in the head-up position in another clinical trial, but no data were given regarding stroke volume or cardiac output [15]. Again, the possibility cannot be excluded that both esmolol as well as clonidine compromise physiologic reaction to compensate for hemodynamic changes of CP.

# Effect of vasodilation during CP

An increased MAP also was reduced during laparoscopic surgery by the administration of nitroglycerine to patients at cardiac risk of American Society of Anesthesiology (ASA) 3 and 4. Simultaneously with the fall in MAP, the systemic vascular resistance and the cardiac index were significantly improved in the aforementioned clinical study. These patients were monitored by invasive measurement using the pulmonary artery catheter [3]. In the current animal study, the hemodynamic parameters were not improved by vasodilation, perhaps because MAP was relatively low and did not increase during CP. Thus, the trial did not reflect the clinical situation of patients with a high MAP during CP. In healthy patients vasodilation during CP might additionally compromise venous blood return because of peripheral pooling. This effect was shown in the current study because intrathoracic blood volume was more decreased during CP in the head-down and supine positions after medication with sodium nitroprusside, as compared with CP alone.

# Conclusions

Intraoperative normovolemia and fluid management seem to be most important for minimizing the hemodynamic side effects of CP. However, with standard hemodynamic monitoring, it is difficult to assess intravascular volume status [12]. On the basis of experimental data, no recommendation can be given about the amount of volume that might be necessary to avoid the negative hemodynamic effects of CP. However, invasive monitoring is not required for healthy patients undergoing laparoscopic surgery because they usually toleratc CP without major problems. Medication with  $\beta_1$ blockers or vasodilators may compromise the compensatory effects of CP, but also may be helpful for patients with severe cardiac comorbidity. For these patients, invasive monitoring should be considered. Further rescarch investigating the influence of CP on cardiovascular changes and the effects of  $\beta_1$  blockade, vasodilation, or volume loading must be performed using animal models of chronic heart failure to simulate the clinical situation of elderly multimorbid patients undergoing laparoscopic surgery.

Acknowledgment. The study was supported by the German Research Foundation (DFG JU 364/2-1).

#### References

- Alishahi S, Francis N, Crofts S, Duncan L, Bickel A, Cuschieri A (2001) Central and peripheral adverse hemodynamic changes during laparoscopic surgery and their reversal with a novel intermittent sequential pneumatic compression device. Ann Surg 233: 176–186
- Dexter SP, Vucevie M, Gibson J, McMahon MJ (1999) Hemodynamic consequences of high- and low-pressure capnoperitoneum during laparoscopic cholecystectomy. Surg Endose 13: 376– 381
- Feig BW, Berger DH, Dougherty TB, Dupuis JF, Hsi B, Hickey RC, Ota DM (1994) Pharmaeologic intervention can reestablish baseline hemodynamic parameters during laparoscopy. Surgery 116: 733–741
- Goodale RL, Beehe DS, McNevin MP, Boyle M, Letourneau JG, Ahrams JH, Cerra F (1993) Hemodynamie, respiratory, and metaholic effects of laparoscopic cholecystectomy. Am J Surgery 166: 533–537
- Hachenherg T, Ehel C, Czorny M, Thomas H, Wendt M (1998) Intrathoracic and pulmonary blood volume during CO<sub>2</sub>-pneumoperitoneum in humans. Aeta Anaesthesiol Scand 42: 794– 798
- Haxby EJ, Gray MR, Rodriguez C, Nott D, Springall M, Mythen M (1997) Assessment of cardiovascular changes during laparoscopic hernia repair using ocsophageal Doppler. Br J Anaesth 78: 515–519
- Jorgensen JO, Lalak NJ, North L, Hanel K, Hunt DR, Morris DL (1994) Venous stasis during laparoscopie cholecystectomy. Surg Laparosc Endosc 4: 128–133
- Joris J, Chiche JD, Canivet JL, Jaquet NJ, Legros JJ, Lamy ML (1998) Hemodynamic changes induced by laparoscopy and their endoerine correlates: effects of clonidinc. JACC 32: 1389–1396
- Joris JL, Noirot DP, Legrand MJ, Jacquet NJ, Lamy ML (1993) Hemodynamic changes during laparoscopic cholecystectomy. Anesth Analg 76: 1067–1071
- Junghans T, Böhm B, Boucke T, Huscher D (2000) Die intravenöse Volumensuhstitution vermindert die hämodynamischen Auswirkungen des Pneumoperitoneums. Minim Invasive Chirurg 9: 127–130

- Junghans T, Böhm B, Gründel K, Schwenk W (1997) Effects of pneumoperitoneum with carhon dioxide, argon, or helium on hemodynamic and respiratory function. Areh Surg 132: 272–278
- Junghans T, Böhm B, Haase O, Fritzmann J, Zuckermann-Beeker H (2002) Conventional monitoring and intravascular volume measurement can lead to different therapy after upper gastrointestinal tract surgery. Intensive Care Med 28: 1273–1275
- Koivusalo AM, Scheinin M, Tikkanen I, Yli Suomu T, Ristkari S, Laakso J, Lindgren L (1998) Effects of esmolol on haemodynamic response to CO<sub>2</sub> pneumoperitoneum for laparoscopic surgery. Acta Anaesthesiol Scand 42: 510–517
- Kraut EJ, Anderson JT, Safwat A, Barhosa R, Wolle BM (1999) Impairment of eardiac performance hy laparoscopy in patients receiving poitive end-expiratory pressure. Arch Surg 134: 76–80
- Laisalmi M, Koivusalo AM, Valta P, Tikkanen I, Lindgren L (2001) Clonidine provides opioid-sparing effect, stable hemodynamics, and renal integrity during laparoscopic cholecystectomy. Surg Endose 15: 1331–1335
- Marathe US, Lilly RE, Silvestry SC, Schauer PR, Davis JW, Pappas TN, Glower DD (1996) Alterations in hemodynamics and left ventricular contractility during earbon dioxide pneumoperitoneum. Surg Endosc 10: 974–978
- McLaughlin JG, Scheeres DE, Dean RJ, Bonnell BW (1995) The adverse hemodynamic effects of laparoscopic cholecystectomy [see comments]. Surg Endosc 9: 121–124
- Millard JA, Hill BB, Cook PS, Fenoglio ME, Stahlgren LH (1993) Intermittent sequential pneumatic compression in prevention of venous stasis associated with pneumoperitoneum during laparoscopic cholecysteetomy. Arch Surg 128: 914–919
- Schwenk W, Bohm B, Junghans T, Hofmann H, Müller JM (1998) Intermittent sequential compression of the lower limbs prevents venous stasis in laparoseopie and eonventional colorectal surgery. Dis Colon Rectum 40: 1056–1062
- Schwenk W, Böhm B, Fügener A, Müller JM (1998) Intermittent pneumatic sequential compression (ISC) of the lower extremities prevents venous stasis during laparoscopic choleeysteetomy: a prospective randomized study. Surg Endosc 12: 7–11
- Takeuehi M, Odake M, Takaoka H, Hayashi Y, Yokohama M (1992) Comparison hetween preload reeruitable stroke work and the end systolic pressure-volume relationship in man. Eur Heart J 13(Suppl E): 80–84
- Takeuehi M, Takaoka H, Odake M, Hayashi Y, Hata K, Yokohama M (1992) Assessment of left ventricular function using a conductance eatheter in the human heart. Jpn Circ J 56: 730–734
- 23. Tuppurainen T, Mäkinen J, Salonen M (2002) Reducing the risk of systemic embolization during gynecologic laparoscopy. Acta Anaesthesiol Scand 46: 37–42
- Wallace DH, Serpell MG, Baxter JN, O'Dwyer PJ (1997) Randomized trial of different insufflation pressures for laparoscopic cholecystectomy. Br J Surg 84: 455–458
- Zuckerman R, Gold M, Jenkins P, Rauscher LA, Jones M, Heneghan S (2001) The effects of pneumoperitoneum and patient position on hemodynamics during laparoscopic cholecystectomie. Surg-Endose 15: 562–565

Surg Endosc (2006) 20: 770–772 DOI: 10.1007/s00464-004-9115-3

© Springer Science+Business Media, Inc. 2006

SUNDIGA .

and Other Interventianal Techniques

# Linear stapling of the short gastric vessels reduces blood loss and splenectomy at oesophageal and gastric surgery

# J. F. Dowdall, O. J. McAnena

Department of Surgery, University College Hospital, New Castle, Galway, Ireland

Rcceived: 19 May 2004/Accepted: 8 October 2004/Online publication: 19 January 2006

#### Abstract

*Background:* Increased operative blood loss, blood transfusion and nontherapeutic splenectomy negatively influence postoperative morbidity and mortality following esophageal or gastric resection. A critical point at which blood loss and iatrogenic splenic injury occurs is at the time of division of the short gastric vessels. We examined the efficacy of using a laparoscopic linear cutting stapler (developed for minimal access surgery) to divide with the short gastric vessels at open surgery.

*Methods:* Fifty-six patients were included. In 28 consecutive patients the linear stapler was used when dividing the short gastric vessels. These were compared to 28 matched controls (short gastric vessels were divided between hemostats and ligated). In the two patient groups, patient age, body mass index, and preoperative hemoglobin levels were similar.

*Results:* Operation time, splenectomy rates, blood transfusion, and mean transfusion volume were all significantly reduced in the group where the stapler was used.

*Conclusions:* Use of a linear cutting stapler reduced operation time, blood product use, and incidental splenectomy in patients undergoing radical open esophageal and gastric surgery.

**Key words:** Spleen conservation — Short gastric vessels — Transfusion requirements

gastric junction, regional nodal dissection (celiac axis, splenic hilum and upper pancreas) is required [5]. The significant factors associated with perioperative complications are intra-operative blood loss, blood transfusion, splenectomy, and prolonged operation time rather than the extent of nodal dissection [1, 3, 5, 8, 9]. Extended nodal dissection in itself contributes to operation time and blood loss, but so too does inadvertent splenic injury. As the latter is theoretically avoidable, there is potential for reducing perioperative morbidity by avoiding incidental splenectomy. In our own experience, a critical point at which splenic injury occurs is when the short gastric vessels are divided. The principal difficulty when dividing the vasa brevia is the limited access created by the overhanging costal margin and diaphragm, especially in obese patients and in those with an acute subcostal angle.

(cardia) and III (subcardia) cancers of the esophago-

With the advent of minimally invasive surgery, technology has been developed to overcome the difficulties due to limited access. We have previously described a technique to divide the short gastric vessels at gastric and esophageal surgery using a linear vascular cutting stapler developed for laparoscopic surgery [7]. We assessed whether use of the linear vascular cutting stapler could facilitate division of the short gastric vessels at conventional gastrectomy/esophagectomy and reduce operation time, blood loss, and the need for splenectomy.

Resection of the stomach or esophagus for malignant disease is associated with significant postoperative morbidity and mortality. It has been suggested that performing extended lymphadenectomy may contribute to this [6], but at gastrectomy D2 dissection remains the standard of care (>20% of patients with "early" gastric cancer have involved nodes) [3]. Similarly for types II

# Methods

#### Patients

Fifty-six patients are included in this report. In 28 consecutive patients the linear stapler was used when dividing the short gastric vessels, and these are matched to the previous 28 consecutive patients who underwent identical operations (the short gastric vessels were divided by the conventional technique between hemostats). In the two patient groups, patient age, body mass index, and preoperative haemoglobin levels were not significantly different (data not shown). The distribu-

Correspondence to: O. J. McAnena

Table 1. Transfusion and splencetomy rates broken do	lown by	operation type
--	---------	----------------

		Linear stapler used				Linear stapler not used			
Procedure	п	Splenectomy	Number of patients Iransfused	Units transfused (mean ± SEM)	n	Splenectomy	Number of patients transfused	Units transfused (mean ± SEM)	
Transhiatal	13	0	3	$0.5 \pm 0.3$	13	1	10	$2.2 \pm 0.5$	
Three-slage	8	0	4	$1.1 \pm 0.4$	8	1	7	$3.0 \pm 0.6$	
Gastrectomy	7	0	2	$0.7 \pm 0.5$	7	2	5	$3.0 \pm 0.9$	
Total	28	$0^{a}$	9 <sup>a</sup>	$0.8 \pm 0.6^{a}$	28	4	22	$2.4 \pm 0.7$	

<sup>a</sup> Significant versus group where stapler was not used (p < 0.05)

tion of procedures performed was as shown in Table 1. Data recorded included anesthesia time, transfusion requirements, incidental splenectomy rate (no therapeutic splnectomics were performed), and the rate of postoperative respiratory sepsis (delined as positive sputum culture or chest x-ray changes consistent with pneumonia).

# **Operative** Technique

All patients who underwent gastrectomy had standard D2 dissection, while those with type II (cardia) and III (subcardia) cancers of the esophago-gastric junction underwent nodal dissection of the celiac axis, splenic hilum, and upper pancreas. In those patients where the linear stapler was used, the operative technique was altered slightly. Following detachment of the greater omentum from colonic and parietal attachments, the operator's left hand was placed on the greater curve of the stomach, with the gastrosplenic omentum sandwiched between the index and middle fingers. This allowed placement of the linear stapler (Endo-GIA; Autosuture, U.S. Surgical, Norwalk, CT) across the gastrosplenic ligament as previously described [7]. Two to three applications were usually sufficient.

#### Results

The male:female ratio was 1.15:1. The mean age was 67 years (range 43–86 years). There was one perioperative (30-day) mortality in each group (no difference between groups).

In the group where the conventional technique was used to divide the short gastries, there were four ineidental splenectomies (14.3%). This was reduced to zero in the group in which the stapler was used (p < 0.05, paired-sample *t*-test). The number of patients transfused in the nonstapled group was significantly higher (22 of 28) than in the stapled group (nine of 28) (p < 0.05, paired-sample *t*-test). The total number of units transfused was also significantly higher (2.4  $\pm$  0.7 units versus 0.8  $\pm$  0.6 units, p < 0.05, ANOVA, Tukey post hoe) (Table 1).

Anesthesia time was also shorter in the stapled versus the nonstapled group ( $174 \pm 42$  versus  $228 \pm 48$  min, p < 0.05). The ineidence of postoperative respiratory sepsis was also significantly lower in the stapled group (5 versus 12, p < 0.05, paired-sample *t*-test).

# Discussion

Prolonged operating time, excessive blood loss, and spleneetomy contribute to the morbidity associated with gastroesophageal surgery [3, 5, 9]. A recent analysis of

more than 12,000 gastreetomies for gastrie eareinoma demonstrated significantly poorer outcomes in patients who had nontherapeutie splenectomy [8]. In our experienee, injury to the spleen oceurred most frequently when ligating and dividing the short gastric vessels. Anatomieally, the intimate relationship between the gastrie fundus and the eephalad portion of the spleen eontributes, while the friable and vaseular nature of the spleen itself is also a factor. In the 28 eases where the short gastrie vessels were dealt with by standard ligation and division, four nontherapeutie spleneetomies were performed. Patient factors are unlikely to have played a role, as individual body mass indiees were similar in both groups. When the vasa brevia were divided by stapling, the need for spleneetomy was obviated in all eases.

With the advent of minimally invasive surgery, stapling devices were developed specifically to overcome limited access. Our premise was that such a device would have significant application at open surgery. The longer instrument shaft length allowed for ease of placement, while the overlapping rows of staples effectively sealed vessels. Operation time was significantly shorter as eompared to the conventional approach. Another factor that is undoubtedly involved in the reduction of operation time is the lesser number of splenectomies performed. The lower incidence of respiratory morbidity in the stapled group was an interesting and unexpected finding. Presumably shorter operating time and lower incidence of transfusion contributed to this.

We found the technique particularly helpful in the obese and those with a narrow substernal angle. In these, patients gastroesophageal surgery is technically more demanding, as the vasa brevia to the upper pole of the spleen are obscured by overhanging of the eostal margin, and also because of the difficulty in identifying the vessels through the fat of the gastrosplenie omentum and where intraoperative blood loss and transfusion requirements are traditionally greater [8].

Hemorrhage either from the vasa brevia themselves or from iatrogenie splenie injury (oeeurring at the time of division) is difficult to control, adds significantly to the operating time, and may increase perioperative blood product requirements as demonstrated by the results. Although perioperative allogeneie blood transfusion does not affect long-term survival after esophagogastreetomy for earcinoma, it does have a significant association with short-term survival in a group whose overall survival is often limited after resection [2].

The current cost of the stapler in Ireland is  $\in 270$ . This rises to  $\in 560$  when two refills are used. This is considerably more expensive than the use of hemostats and ties. However, we feel that this additional cost could be offset by operating-room time saved, blood products saved, and reduced bed days used as a result of a lower rate of respiratory complications.

One might suggest that the patients in the control group used for comparison were historical and therefore open to criticism. However, they were procedure matched and taken consecutively from those patients who had surgery performed immediately before the introduction of the stapling device. There were no differences in demographics, body mass index, or perioperative mortality between the groups. Therefore, although the comparison is not a randomized one, it is legitimate as a representative control group was used. As incidental splenectomy confers no oncological benefit [9] but does result in long-term immune compromise [2, 4, 8], it is reasonable to suggest that the use of the linear stapler may influence patient outcome.

# References

- Brady MS, Rogatko A, Dent LL, Shiu MH (1991) Effects of spleneetomy on morbidity and survival following curative gastrectomy for carcinoma. Arch Surg 126: 359–364
- Craig SR, Adam DJ, Yap PL, Leaver HA, Elton RA, Cameron EW, Sang CT, Walker WS (1998) Effect of blood transfusion on survival after esophagogastreetomy for carcinoma. Ann Thorac Surg 66(2): 356–361
- Cuscheri A, Fayers P, Fielding J, Craven J, Baneewicz J, Joypaul V, Cook P (1996) Postoperative morbidity and mortality after D1 and D2 resections for gastric cancer: preliminary results of the MRC randomised controlled surgical trial. Lancet 347: 995–999
- Fughs CS, Mayer RJ (1995) Gastrie eareinoma. N Engl J Med 333: 32–41
- Griffith JP, Sue-Ling HM, Martin I, Dixon MF, McMahon MJ, Axon AT, Johnston D (1995) Preservation of the spleen improves survival after radical surgery for gastrie cancer. Gut 36: 684–690
- Kranenbarg EK, van de Velde CJ (1998) Gastrie eaneer in the elderly. Eur J Surg Oncol 24(5): 384–390
- Power C, Maguire D, McAnena OJ (2000) A technique for dealing with the short gastric vessels during gastric surgery. Eur J Surg Oneol 26(1): 80–81
- Wanebo HJ, Kennedy BJ, Winehester DP, Stewart AK, Fremjen AM (1997) Role of spleneetomy in gastrie eaneer surgery: adverse effect of elective spleneetomy on long-term survival. J Am Coll Surg 185: 177–184
- Whiting JL, Fielding JWL (1998) Radical surgery for early gastrie cancer. Eur J Surg Oneol 24: 263–268

Surg Endose (2006) 20: 773–782 DOI: 10.1007/s00464-005-0331-2

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

# Zoom endoscopic monitoring of small bowel allograft rejection

T. Kato,<sup>1</sup> J. J. Gaynor,<sup>1</sup> S. Nishida,<sup>1</sup> N. Mittal,<sup>2</sup> G. Selvaggi,<sup>1</sup> D. Levi,<sup>1</sup> J. Moon,<sup>1</sup> J. Thompson,<sup>2</sup> P. Ruiz,<sup>3</sup> J. Madariaga,<sup>1</sup> A. G. Tzakis<sup>1</sup>

<sup>1</sup> Department of Surgery, University of Miami, School of Medicine, 1801 NW 9th Avenue, 5th Floor, Miami, FL 33136, USA

<sup>2</sup> Department of Pediatrics, University of Miami, School of Medicine, Miami, FL 33136, USA

<sup>3</sup> Department of Pathology, University of Miami, School of Medicine, Miami, FL 33136, USA

Received: 29 April 2005/Accepted: 8 November 2005/Online publication: 16 March 2006

#### Abstract

*Background:* The small bowel has been successfully transplanted in patients with irreversible intestinal failure. This report aims to describe endoscopic monitoring of small bowel rejection.

*Methods:* A magnification endoscope (zoom endoscope) was used in this study. In the first part of the study (Oetober 1998 to Mareh 2000, 271 endoscopy sessions), the specific endoscopic findings that correlated with rejection were determined. An analysis then was performed on data from the second period (Mareh 2001 to November 2002, 499 sessions) to evaluate the zoom endoscope's aceuracy in monitoring rejection.

Results: Specifie endoscopie findings of rejection found in the first period included background erythema, villous congestion, blunted villous tip, and shortened villous height. When the rejection was successfully treated, endoscopie appearance returned to normal. On the basis of these findings, five endoscopie criteria (villous shortening, villous blunting, background erythema, villous congestion, and mueosal friability) were used to seore endoscopic sessions in the second period. Endoscopie diagnosis of rejection was compared with histology. Adult patients showed a sensitivity of 45%, a specificity of 98%, a positive predictive value of 82%, and a negative predictive value of 88%. In pediatrie patients, these values were, respectively, 61%, 84%, 57%, and 86%. On 59 distinct oceasions (30 in period 1 and 29 in period 2) in which the results were endoscopy negative yet biopsy positive (mild) for rejection, we elected not to treat these rejections on the basis of elinical evaluation, and 58 (98%) resolved without further therapy.

*Conclusions:* With the use of magnification, endoscopy is a useful tool for monitoring acute rejection in the small bowel allograft.

bowel transplantation — Zoom endoscopic monitoring

Key words: Small bowel allograft rejection — Small

With recent improvements in surgical techniques and immunosuppressive regimens, small bowel transplantation has become a desirable and potentially life-saving alternative for patients with short gut syndrome or other problems eausing intestinal failure. Worldwide experienee with this procedure has grown relatively fast beeause elinieal outeomes after intestinal transplantation have been improving over time [1, 5, 7, 17, 21]. However, the teehnology and clinical protoeols for monitoring graft function and treating acute rejection still are in the early stages of evolution. In contrast to known serum markers routinely used for monitoring liver or kidney transplant recipients, serum makers for bowel graft rejection still are under investigation [1, 5, 7, 17, 21, 22]. Therefore, frequent biopsies of the small bowel graft are eritieal for detecting the presence of acute cellular rejection. Early detection of rejection also is extremely important because rapid progression to severe rejection can occur without proper treatment, and advaneed stages of rejection are associated with poorer outeomes [11, 14]. Whereas random mucosal biopsy provides detailed information about a very small area, the endoscopie view can provide information about large areas. Endoseopy provides the quickest method for assessing the overall health of graft mucosa and is essential in obtaining specimens for histologic evaluation [6, 9, 13, 26]. Unfortunately, surveillance with standard endoscopy has not been sufficiently eapable of monitoring rejection [8, 29-31]. Standard endoseopy is just not powerful enough to provide an aceurate view of the mild changes that oeeur in the early phase of small bowel rejection.

The use of magnification endoscopy has been reported for evaluation of Barrett's esophagus and celiac disease [25, 28]. Subsequently, in 1999, we published the

Part of this work was presented during the annual meeting of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES), 29 March to 1 April 2000, Atlanta, GA *Correspondence to:* T. Kato

first case report featuring magnification with a zoom video endoscope (ZVE) for evaluation of intestinal graft mucosa [16]. In this initial experience, the ZVE enabled us to visualize the villi and crypt areas directly in almost microscopic detail, thereby providing a more accurate overall assessment of graft mucosa health than with a standard endoscope. In addition, the ZVE enabled us to take a biopsy from the most representative area of the graft. We describe our experience using zoom endoscopy to monitor patients for small bowel rejection.

# Patients and methods

We used zoom endoscopes (Pentax EC-3830LZ, EG 3430Z; Pentax Precision Instrument, Inc, Orangeburg, NY; and Olympus GIFQ160Z; Olympus America, Inc, Melville, NY, USA) for this study. Pentax equipment was used during the first period of the study, and Olympus equipment was used during the second period (see below). The reason for the change in equipment was solely administrative. With either zoom endoscope, the operator has the capability to change the image on a standard 14-in. video monitor from the standard endoscopic view (magnification  $\times 12.5$ ) to a highly magnified view (magnification  $\times 105$ ). This versatility is clearly seen in Fig. 1A through D, which shows normal through near maximum ZVE views of normal small intestinal mucosa. The degree of magnification available with this scope allows the endoscopist to visualize both the entire length of villi and the surrounding crypt area. Normal villous architecture seen on ZVE was similarly found in the histologic specimen taken at the same time (Fig. 1E and F).

During the first period of the study, a total of 271 endoscopic sessions involving 16 intestinal transplant recipients were evaluated over a 17-month period (October 1998 to March 2000). As a pilot study, we retrospectively analyzed our experience using the zoom endoscope to perform surveillance for rejection among these 16 patients. For each patient, the endoscopic procedure was performed through a surgically created ileostomy. The endoscope was advanced at least 5 cm from the ostomy outlet to avoid changes related to the ostomy site. In most cases, ZVE surveillance was performed 10 to 20 cm from the ostomy.

All procedures were performed at the patient's bedside without sedation. To correlate the visual impression of the villi and crypts under high magnification with the histology of the small bowel mucosa, the endoscopist obtained biopsies through the working channel of the endoscope during each endoscopic session. All endoscopies were performed hy a single endoscopist (T.K.) during this period.

The demographics of the patients during the two study periods are summarized in Table 1. The transplantations were performed as previously described [17]. All the patients during the first period received daclizumab (Zenapax; Roche Laboratories, Nutley, NJ, USA) for induction immunosuppression along with standard doses of tacrolimus and corticosteroids for maintenance. Steroid bolus injection (1 g of methylprednisolonc) and/or cycle (200 mg of methylprednisolonc per day followed by a tapered schedule of 160 mg, 120 mg, 80 mg, 40 mg, and 20 mg per day for a total of 5 days) or OKT3 (Muromonab-CD3; Ortho Biotech Inc., Raritan, NJ, USA) was used to treat rejection. Sirolimus (Rapamycin; Wyeth Ayerst Research, Philadelphia, PA, USA) was added in most cases to the baseline immunosuppressive regimen after treatment of refractory rejection (usually recurrent rejection after the use of OKT3).

During the second period of the study (March 2001 to Novemher 2002), we evaluated the capability of the zoom endoscope to provide surveillance for small bowel rejection. Specifically, a scoring system to identify rejection was defined from the results of the pilot study in the first period. A total of five criteria were chosen for endoscopic scoring of bowel rejection. Four of these criteria were based on the mucosal imaging results of the first period: shortening of the villous length, blunting of the villous tip, erythema at the background of the mucosa, and villous congestion. A fifth criterion, mucosal friability, representing the intestinal tissue's sensitivity to the actual use of the ZVE equipment (not its visualized imagery), was selected because our experience showed that it also provided a strong diagnostic sign for acute bowel rejection.

In keeping with the international criteria established for scoring individual histologic components of acute rejection in both kidney and liver transplantation [4, 27], a scoring system of 0 to 3 (0, normal; 1, mildly abnormal; 2, moderately abnormal; 3, severely abnormal) was chosen for each of the five endoscopic criteria, with a higher score denoting worse appearance. An overall ZVE small bowel rejection score was defined as the sum of the five individual component scores. Thus, a total ZVE score of 0 to 15 was possible. Scoring guidelines for each criterion are summarized in Table 2.

Next, the total score was categorized to define a ZVE grading scheme as follows: a total score of 0 was normal (grade 0), 1 to 5 was indeterminate for rejection (grade 1), 6 to 10 was mild (grade 2), 11 to 14 implied moderate rejection (grade 3), and 15 implied severe rejection (grade 4). Finally, for each sample, the ZVE grade of rejection was compared with the histologic grade of rejection: 0 (no rejection), 1 (indeterminate for rejection), 2 (mild rejection), 3 (moderate rejection), and 4 (severe rejection). Because we do not treat histologic grade 1 rejection, histologic grades 0 and 1 were considered negative for rejection. It must be noted that according to the recently revised histologic criteria of the Eighth International Small Bowel Transplant Symposium held in September, 2003 in Miami, Florida, code grade 1 denotes mild rejection (previously grade 2), and the code 1ND denotes an indeterminate grade (previously grade 1) [23]. However, because we were using the previous grading scale throughout the study, we are reporting the results as such in this article.

The planned schedule for performing endoscopies (along with histologic surveillance) during hoth periods of the study was as follows: once every 2 to 4 days during the first 2 weeks after transplantation, once weekly during the next 3 to 8 weeks after transplantation, and once a month during months 3 to 6 (or until stoma closure). During the course of any rejection episode, endoscopies were performed for the patient every 2 to 4 days.

The ZVE evaluations were performed by a total of four endoscopists (T.K., S.N., and G.S. for adults; T.K., G.S., and N.M. for children). Each endoscopist had a large experience in transplant endoscopy, and each had experience using the ZVE before the beginning of the second study period. Each biopsy was taken from the most representative area of the graft.

The ZVE evaluation and score were recorded at the time of endoscopy before histologic evaluation. The hiopsy tissue was embedded in formalin, then processed and stained in a standard hematoxylineosin stain. Each biopsy was read by a single, expert pathologist (P.R.), who was blinded to the endoscopic score while reading the biopsy. Data were obtained from the following 47 patients who received intestinal transplants at the University of Miami and agreed to participate during the second period of the study: 26 adults (median age, 33 years; range, 20–59 years) and 21 children (median age, 6 years; range, 0.8–15 years). Because of technical limitations with the use of zoom endoscopy on very small intestines, the recipients of transplants from donors younger than 2 years were not included in the study.

A total of 499 ZVE sessions involved the 47 patients (308 in adults and 191 in children). For 20 patients (15 adults and 5 children), an induction protocol with Alemtuzumab (Campath-1H, 1LEX Oncology, Inc., San Antonio, TX, USA) was used [34, 35]. The remaining 27 patients received the same immunosuppression protocol as those in the first period.

Institutional review board approval was obtained for this study along with written consent from each subject who chose to participate. Tests of association were performed using the (uncorrected) Pearson chi-squared test. A p value less than 0.05 was considered to indicate a statistically significant finding.

#### Results

#### Period 1

Figure 2 illustrates crypt erythema seen during the course of mild acute rejection. The space between villi has expanded and appears erythematous (Fig. 2B and C). In the standard, unmagnified endoscopic view, it is not clear whether this same area appears normal or displays minimal to mild mucosal erythema (Fig. 2A).



Fig. 1. Zoom videoendoscopic view of normal small bowel graft mucosa (standard [A], moderate [B], and maximum [C, D] magnification), along with corresponding histology taken at the same time (E, F), showing a well-preserved villous architecture with sparse cellularity present in the lamina propria. (hematoxylin-cosin; E magnification ×100, F magnification ×400)

Villi were relatively preserved at this mild rejection stage. Histologic examination of the mucosal biopsies taken at the same time confirmed the morphologie changes visualized by ZVE. Typically, the pathology demonstrated the presence of a mild lymphocytic infiltrate in the lamina propria, the presence of activated lymphocytes, and scattered single cell apoptosis (Fig. 2D and E). Villi may have been mildly edematous, but were otherwise morphologically normal.

As rejection progressed to a moderate grade, the blunting and shortening of villi became more obvious and were accompanied by a mild to moderate degree of erythema (Fig. 3A and B). Some of the villi in Fig. 3A and B appear considerably shorter and irregular in length, as compared with the villi of the mild rejection case seen in Fig. 2A to C. Such distinctions were not possible with standard magnification. Pathologic examination of mucosal biopsies taken simultaneously during ZVE demonstrated that the cellular infiltrate in the lamina propria had markedly inereased, and the villi had become even more blunted (Fig. 3C and D). Crypt injury was increasingly prominent, with multiple clusters of apoptotic cells and crypt abseesses.

# Table 1. Patient demographics

	Period 1 $(n = 16)$	Period 2 $(n = 47)$
Age (years)	Adults ( $n = 15$ ; range, 21–48; median, 37)	Adults ( $n = 26$ ; range, 20–59; median 33)
Condon	Children $(n = 1; 15)$	Children ( $n = 21$ ; range, 0.77–15; median, 6)
Gender	Mate  (n = 7) $ Female  (n = 9)$	Female (n = 26)
Diagnosis	Mesenteric thrombosis $(n = 6)$	Mesenterie thromhosis $(n = 8)$
Dagaoon	Gardner's syndrome $(n = 2)$	Pseudoobstruction $(n = 7)$
	Pseudoobstruction $(n = 2)$	Gastrosehisis $(n = 6)$
	Trauma $(n = 2)$	Trauma $(n = 6)$
	Others $(n = 4)$	Volvulus $(n = 4)$
		Intestinal atresia $(n = 4)$
		Crohn's disease $(n = 3)$
		Others $(n = 9)$
Graft types	lsolated intestine graft $(n = 7)$	Isolated intestine graft $(n = 22)$
	Liver and intestine graft $(n = 0)$	Liver and intestine graft $(n = 4)$
	Multiviseeral graft $(n = 8)$	Multiviseeral graft ( $n = 18$ )
	Modified without the liver $(n = 1)$	Modified without the liver $(n = 3)$
Immunosuppression	Daelizumab $N = 16$	Daelizumab ( $n = 27$ )
protoeols		Alemtuzumah ( $n = 20$ )

Table 2. Zoom endoscopie seores

		Seore
Height of villi	Normal	0
0	Mildly shortened	1
	Moderately shortened	2
	Flat	3
Villous blunting	Normal	0
0	Mildly blunted	1
	Moderately blunted	2
	Flat	3
Background		
mueosal erythema	No erythema	0
	Mild erythema	1
	Moderate erythema	2
	Severe erythema	3
Villus eongestion	Normal vaseular pattern	0
C	Mildly eongested	1
	Moderately eongested	2
	Severely eongested	3
Friahility	Normal	0
5	Easily bleeds with biopsy forceps	1
	Easily bleeds with endoseopie trauma	2
	Continuous oozing or ulceration	3

The severe form of acute rejection was manifested by complete denudation (i.e., loss of surface epithelium). Although easily recognized in the standard endoscopic view (Fig. 4A), the change was even more clearly identified in the magnified ZVE image (Fig. 4B). The biopsy taken at the same time showed histologic changes of severe cellular infiltrate, destruction of crypts, and disappearance of surface epithelium (Fig. 4C and D).

Among the 271 endoscopic and histologic examinations performed on these patients were 58 histologic diagnoses of rejection (1 to 6 diagnoses per patient; average, 3.6) including mild (n = 46), moderate (n = 8), severe (n = 3), and vascular rejection (n = 1). Mucosal changes observed by ZVE at the same time included background erythema and blunted and/or shortened villi (n = 8), crypt erythema only (n = 7), blunted and/or shortened villi only (n = 3), flattened villi/denuded mucosa (n = 3), villus congestion (n = 1), and no noteworthy changes (n = 36). Each of the three histologic diagnoses of severe rejection corresponded with a ZVE reading of flattened villi/denuded mucosa, and the single case of histologic vascular rejection coincided with a ZVE reading of villus congestion. In fact, the single episode of vascular rejection occurred in a patient with a positive cytotoxic crossmatch. Histologic examination showed diffuse submucosal hemorrhage in hematoxylin-cosin stain. The immunofluorescent stain for complement and immune complex yielded positive results [12, 24]. It should be noted that 18 of the 58 rejection diagnoses were based on repeat endoscopies and biopsics performed on the same patient during the same rejection episode and during the same course of therapy. Figure 5 compares the ZVE picture of moderate rejection seen in one patient 13 days after transplantation (A) and the normal ZVE picture 1 month later in this same patient after treatment for the rejection (B). In all cases, when the histologic appearance became normal after treatment for rejection, the mucosal abnormalities visualized by the ZVE also disappeared.

Normal mucosal appearance on ZVE was found for 78% (36/46) of the biopsies histologically diagnosed as mild rejection. In 30 of these instances, endoscopy and biopsy had been performed as routine surveillance without clinical symptoms. Despite the mild rejection present on the histologic examination of the mucosal biopsies, we elected to withhold additional immunosuppressive treatment, because i) the mucosal villi and crypt appeared normal on ZVE, and ii) there were no clinical symptoms. All these 30 instances of rejection reversed without treatment. Conversely, on the basis of clinical judgment, six histologically indicated instances of mild rejection were treated despite their relatively normal appearance on ZVE. All these rejections reversed with steroid treatment.



Fig. 2. Zoom videoendoseopic (ZVE) view along with corresponding histology of the forceps biopsy taken at the same time showing mild acute allograft rejection in a multivisceral graft recipient 6 days after transplantation. Standard (A) and moderate (B, C) ZVE magnification shows relatively well-preserved villus stalk with increased erythema visible in the background between villi. The histologic specimens show minimal blunting of villi with mildly increased inflammatory infiltrate in the lamina propria correlating with ZVE findings (D) and mild cryptitis with scattered apoptotic bodies at high power (E). (hematoxylineosin; D magnification  $\times 100$ , E magnification  $\times 400$ )

On all but one occasion, positive findings on ZVE were supported by histologic readings of rejection. In the one exception, ZVE showed erypt erythema, whereas histologic readings were ineonelusive for rejection. Given our confidence in the ZVE findings together with a clinical symptom of rejection (fever), we administered treatment for rejection. After the treatment, the ZVE-visualized appearance of erypt erythema reversed.

With the exception of acute cellular rejection, no pathologic conditions, such as cytomegalovirus enteritis or posttransplant lymphoproliferative disease, were encountered in these patients. There were no complications associated with ZVE and biopsies.

#### Period 2

A comparison between the ZVE and histologic grades is shown in Tables 3 to 5. During 499 overall ZVE sessions, 87 were positive for rejection (grade 2 or greater), and 412 were negative (grades 0 to 1). By histology, 111 were positive for rejection, and 388 were negative (Table 3). When ZVE was correlated with histology scores in adults, 28 were both endoscopy and histology positive for rejection, 6 were endoscopy positive but histology negative, 34 were endoscopy negative but histology positive, and 240 were both endoscopy and histology negative (Table 4). For children, 30 were both endoscopy and histology positive for rejection, 23 were



Fig. 3. Zoom vidcoendoscopic (ZVE) view along with corresponding histology of the forceps biopsy taken at the same time showing moderate allograft rejection in an isolated bowel graft recipient 9 days after transplantation. Low (A) and near maximum (B) ZVE magnification of increased erythema with villi of irregular length. Some villi are markedly shortened. Low-power histologic examination (C) demonstrates a notable inflammatory infiltrate in the lamina propria with obviously blunted villi, and high-power histologic examination (D) shows advanced stage of crypt injury with numerous apoptotic bodies involving the crypts. (hematoxylin-cosin; C magnification ×100, D magnification ×400)

endoscopy positive but histology negative, 19 were endoscopy negative but histology positive, and 119 were both endoscopy and histology negative (Table 4). The condition of endoscopy positive but histology negative for rejection was significantly higher for children than for adults (12% vs 2%; p < 0.00001). Therefore, positive predictive value and specificity were lower for children. For adults, there was an estimated sensitivity of 45%, a specificity of 98%, a positive predictive value of 82%, and a negative predictive value of 88%. The corresponding values for children were, respectively, 61%, 84%, 57%, and 86%. The overall estimates were 52% for sensitivity, 93% for specificity, 67% for positive predictive value, and 87% for negative predictive value.

In 29 cases that were endoscopically negative but histologically positive for rejection (17 among adults and 12 among children), treatment was withheld because the patient lacked clinical symptoms of rejection such as diarrhea or fever. This approach was based on our experience during period 1, in which all these episodes resolved spontaneously. Actually, in 28 (97%) of 29 cases during period 2, the subsequent biopsy also turned negative for rejection without therapy. Only one case was persistently read as rejection, with subsequent treatment given. Thus, if we exclude these 28 histologically false-positive cases (16 among adults and 12 among children) from the calculations, then the estimated sensitivity and negative predictive value increase, respectively, to 61% and 93% in adults, and to 81% and 94% in children (Table 5).

# Discussion

Control of acute allograft rejection has been one of the most important and challenging dilemmas in clinical intestinal transplantation [1, 5, 7, 11, 14, 17, 21]. Endoscopic surveillance with multiple biopsies has been performed as the primary tool for detecting rejection and instituting timely and appropriate treatment [6, 9, 13, 26]. An ileostomy normally is created at the time of transplantation to facilitate endoscopic access to the graft mucosa [17]. Theoretically, the endoscopic examination allows quick assessment of a large area of graft, as compared with a histologic specimen that includes an extremely limited area of mucosa. Despite this advantage, standard endoscopy alone is not reliable in determining the degree of rejection and has been used primarily for sampling.

Mucosal changes in intestinal allograft rejection observed with standard endoscopy consist of erythema, granularity, and friable hemorrhagic mucosa during episodes of mild to moderate rejection, and deepithelialized, so-called "denuded," mucosa with scvcre rejection. The loss of the mucosal surface epithelium is easily







recognized during standard endoscopy. However, once the rejection has progressed to this stage, it has rarely been reversible [11, 14]. Therefore, initiation of appropriate treatment is essential before the rejection progresses to this very likely irreversible stage. Given standard endoscopy's lack of accuracy in successfully diagnosing mild to moderate rejection [8, 29–31], elinicians have not relied on it as a surveillance tool.

The histologic changes that occur in the intestinal mucosa during the course of rejection have been well described in both experimental and clinical intestinal transplantation [3, 10, 18, 19, 36]. At an early stage of rejection, the villi become edematous, slightly shortened, or both, and crypt areas expand with a mildly increased cellular infiltrate in the lamina propria. As the rejection progresses, the villi become more obviously blunted as eellular infiltrate in the lamina propria increases, leading to a total loss of surface epithelium and severe eellular infiltrates at the end stage of severe rejection. These changes in the microscopic architecture of the mucosa are more easily recognized in whole-thickness biopsy. However, they can be difficult to assess in a standard forceps biopsy because this type of biopsy includes only a limited number of villi and crypts.

The use of videomicroscopic monitoring was reeently demonstrated to be effective in detecting rejection in an animal model [32]. We also published the first case study describing the use of ZVE for evaluation of intestinal graft mueosa [16]. Magnification endoscopy (ZVE) enabled us to see the microscopic architecture of

ZVE grade	Adulis	Children	Histologic grade	Adults	Children
Grade 0–1 (score 0–5)	274	138	Grade 0–1	246	142
Grade II (score 6-10)	28	51	Grade 2	51	37
Grade III-IV (score 11-15)	6	2	Grade 3-4	11	12
Total	308	191	Total	308	191

 Table 4. Cross-tabulation of the zoom video endoscope and histology scores in period 2

	Adults	Children		Adults	Children
Endoscopy +, pathology +	28	30	Endoscopy +, pathology +	28	30
Endoscopy +, pathology -	6	23	Endoscopy +, pathology -	6	23
Endoscopy –, pathology +	34	19	Endoscopy –, pathology + (excluding clinically insignificant rejection)	18	7
Endoscopy -, pathology -	240	119	Endoscopy –, pathology –	240	119
Total	308	191	Total	292	179

Table 5. Sensitivity, specificity, positive and negative predictive values for ZVE as a predictor of histologic grade, by patient age

	Adults	Children	Overall	Adults (adjusted)*	Children (adjusted)*
Sensitivity	45%	61%	52%	61%	81%
Specificity	98%	84%	93%	98%	84%
Positive predictive value	82%	57%	67%	82%	57%
Negative predictive value	88%	86%	87%	93%	94%

\* Adjusted by removing the 28 histologically false positive cases from the calculations

the intestinal mucosa in this patient, who experienced a mild acute rejection. Our current report describes a series of 16 consecutive patients in whom ZVE was used to evaluate the small bowel graft mucosa followed by a prospective study that demonstrated reasonable accuracy of ZVE for 47 additional patients.

Given our experience with the scries of 16 patients, we were able to correlate the visual morphologic changes seen on the ZVE with the various histologically determined grades of rejection. The use of ZVE rather than standard endoscopy permits greater recognition of changes in the villi and crypt areas. It more clearly identifies the mucosal changes that occur as the rejection progresses. We now believe that the endoscopic assessment of mucosa afforded by ZVE provides a fast and reliable confirmation of the histologic diagnosis.

The patchy, segmental nature of intestinal rejection warrants multiple random sampling [1, 5, 7, 21, 37]. Such random sampling may no longer be required because the visualization provided by the ZVE allows biopsy of the most representative area of the intestinal mucosa. Additionally, graft intestinal mucosa often shows inflammatory changes attributable to surgical trauma. Biopsy of these inflamed areas of the stoma can distort the histologic diagnosis [36]. We believe that by using the ZVE as a tool to identify appropriate areas for biopsy, inaccurate diagnoses of rejection attributable to sampling error will be minimized.

In a total of 59 instances (30 in period 1 and 29 in period 2), histologically diagnosed mild rejection was not treated because there were no clinical symptoms of rejection and the mucosa appeared normal on ZVE. All but one of these instances of rejection reversed without treatment. These may have been subclinical rejection episodes that occur in the graft mucosa. Another possible explanation is that an overdiagnosis of changes in limited sampling areas tends to occur. The changes seen in a small sampled area from a biopsy may not always represent the status of the entire graft. Rather, these focal changes may be self-limited. The ZVE, along with the concurrent clinical status of the patient, appears to provide reliable support for deciding whether to treat such histologic diagnoses of rejection or not.

The specificity and negative predictive value of the ZVE score were very high (respectively, 93% and 87% overall). However, the sensitivity was low for both children (61%) and adults (45%), and the positive predictive value was low for children (57%). The low sensitivity appeared to be attributable, in part, to histologic rejection that does not require treatment. When we excluded these (histologically false-positive) rejection episodes, the sensitivity increased for both children (81%) and adults (61%). The reason for a low positive predictive value for children is not clear. However, children are more likely to have infectious enteritis. It is possible that enteritis mimics rejection in endoscopic appearance, and that some cases were undocumented.

We did not encounter apparent infectious enteritis during these ZVE sessions. Cytomegalovirus (CMV) enteritis has been previously reported to cause significant morbidity in clinical intestinal transplantation [20, 33]. The initial preference for CMV-negative grafts along with the subsequent improvement in CMV prophylaxis with the use of gangeiclovir has reduced the incidence of CMV enteritis. In our series of more than 150 clinical intestinal transplantations performed since 1994, CMV enteritis was encountered in only two cases. Even if we include other types of infectious enteritis such as those attributable to adenovirus and nontuberculous mycobacterium-associated enterocolitis, the incidence still is limited to less than 10% of all cases, as compared with rejection, which has a distinct possibility of occurring in nearly every case after intestinal transplantation [2, 15, 17]. Thus, none of these types of infections can explain completely via differential diagnosis the large number of histologically false-positive ZVE readings observed in this study.

Posttransplant lymphoproliferative disease (PTLD) is known to occur rather frequently in the intestinal transplant recipient [1, 5, 7, 21]. Specifically, PTLD can occur in the submucosal lymphoid tissue of the bowel allograft. Although we did not encounter any gastro-intestinal-related PTLD during the follow-up period of the patients enrolled in this study, we have seen cases of PTLD occurring in the intestinal allografts of other patients. The appearance of PTLD in intestinal allograft can be clearly distinguished from that of rejection, because it shows a distinctive area of abnormality (ulcer, lesion) with the surrounding mucosa appearing normal.

As we collect prospective data using the ZVE in combination with histologic study, we plan to use standard stepwise and logistic regression techniques for each of the five criteria defined in this report (villous length, villous blunting, background erythema, villous congestion, and friability) to investigate whether they contain equal or dissimilar predictive value for the histologic diagnosis of rejection. We also will investigate whether the individual component scoring of 0 to 3 used in this study is optimal for predicting the histologic diagnosis, or whether another set of numeric scores used for the normal, mild, moderate, and severe categories would provide greater predictive value. In addition, measurement of inter- and intraobserver variability also will require future investigation.

In conclusion, histologic examination continues to remain the gold standard for a diagnosis of rejection. However, the ZVE provides information on a larger mucosal surface area than can be evaluated with a forceps biopsy, and we currently believe that ZVE as a diagnostic tool should be used concomitantly with the biopsy. The enhanced ability afforded by ZVE to assess the health of graft mucosa has been shown in this study to provide useful information toward a rapid diagnosis and, along with the presenting clinical symptoms, to help in deciding whether to treat histologically diagnosed mild rejection. Clearly, the important question about whether to withhold treatment in all eases involving histologic diagnosis of mild rejection, but without corresponding clinical symptoms, must await confirmation by further studies. Finally, as a sampling tool, ZVE will help in identifying the most appropriate areas for biopsy.

Acknowledgments. This work was supported in part by grant no. DK061445 from the National Institutes of Health, and by an educational grant from Fujisawa Healthcare. In addition, the zoom endoscopic equipment used during the first period of the study was supplied free of eharge by Pentax Precision Instruments, Inc.

### References

- Abu-Elmagd K. Reyes J, Bond G, Mazariegos G, Wu T, Murase N, Sindhi R, Martin D. Colangelo J, Zak M, Janson D, Ezzelarab M, Dvorchik I, Parizhskaya M. Deutsch M, Demetris A, Fung J, Starzl TE (2001) Clinical intestinal transplantation: a decade of experience at a single center. Ann Surg 234: 404–417
- Berho M, Torroella M, Viciana A, Weppler D, Thompson J, Nery J, Tzakis A, Ruiz P (1998) Adenovirus enterocolitis in human small bowel transplants. Pediatr Transplant 2: 277–282
- Cohen Z, Nordgren S, Lossing A, Cullen J, Craddock G, Langer B (1984) Morphologie studies of intestinal allografi rejection: immunosuppression with cyclosporine. Dis Colon Rectum 27: 228–234
- 4. Demetris A, Adams D, Bellamy C, and other members of an International Panel. (2000) Update of the International Banff Schema for Liver Allograft Rejection: working recommendations for the histopathologic staging and reporting of chronic rejection. An International Panel. Hepatology 31: 792–799
- Fishbein TM, Gondolesi GE, Kaufman SS (2003) Intestinal transplantation for gut failure. Gastroenterology 124: 1615– 1628
- Garau P, Orenstein SR, Niegut DA, Putnam PE, Reyes J, Tzakis AG, Kocoshis SA (1994) Role of endoscopy following small intestinal transplantation in children. Transplant Proc 26: 136–137
- Goulet O, Jan D, Brousse N, Revillon Y, Ricour C (1997) Intestinal transplantation. J Pediatr Gastroenterol Nutr 25: 1–11
- Hassanein T, Schade R, Soldevilla-Pico C, Tabasco-Minguillan J, Abu-Elmagd K, Furukawa H, Kadry Z, Demetris A, Tzakis A, Todo S (1994) Clinical and endoscopic features of rejection in small bowel transplant recipients. Transplant Proc 26: 1413
- Hassanein T. Seahde RR, Soldevilla-Pico C, Tabasco-Minguillan J, Abu-Elmagd K, Furukawa H, Kadry Z, Demetris A, Tzakis A, Todo S (1994) Endoscopy is essential for early detection of rejection in small bowel transplant recipients. Transplant Proc 26: 1414–1415
- Holmes JT, Klein MS, Winawer SJ. Fortner JG (1971) Morphological studies of rejection in canine jejunal allographs. Gastroenterology 61: 693–706
- Ishii T, Mazariegos GV, Bueno J, Ohwada S, Reyes J (2003) Exfoliative rejection after intestinal transplantation in children. Pediatr Transplant 7: 185–191
- 12. Jordan SC, Quartel AW, Czer LSC, Admon D, Chen G, Fishbein MC, Schwieger J, Steiner RW, Davis C, Tyan DB (1998) Post-transplant therapy using high-dose human immunoglobulin (intravenous gammaglobulin) to control acute humoral rejection in renal and cardiac allograf1 recipients and potential mechanism of action. Transplantation 66: 800–805
- Kato T (2000) New technique for prevention and treatment of rejection in intestinal transplantation. Curr Opin Organ Transplant 5: 284–289
- Kato T, Berho M, Weppler D, Ruiz P, Pinna AD, Khan FA, Nery JR, Tzakis AG (2000) Is severe rejection an indication for retransplantation? Transplant Proc 32: 1201
- Kato T, Dowdy L, Weppler D, Ruiz P, Thompson J, Raskin J, Tzakis A (1998) Nontuberculous mycobacterial-associated enterocolitis in intestinal transplantation. Transplant Proc 30: 2537–2538
- 16. Kato T, O'Brien CB, Nishida S, Hoppe H, Gasser M, Berho M, Rodriguez MJ, Ruiz P, Tzakis A (1999) The first case report of the use of a zoom videoendoscope for the evaluation of small bowel graft mucosa in a human after intestinal transplantation. Gastrointest Endosc 50: 257–261
- Kato T, Ruiz P, Thompson JF, Eskind LB, Weppler D, Khan FA, Pinna AD, Nery JR, Tzakis AG (2002) Intestinal and multiviseeral transplantation. World J Surg 26: 226–237
- Lee RG, Nakamura K, Tsamandas AC, Abu-Elmagd K, Furukawa H, Huston WR, Reyes J, Tabaseo-Minguillan JS, Todo S,

Demetris AJ (1996) Pathology of human intestinal transplantation. Gastroenterology 110: 1820–1834

- Madara JL, Kirkman RL (1985) Structural and functional evolution of jejunal allograft rejection in rats and the ameliorating effects of cyclosporine therapy. J Clin Invest 75: 502–512
- Manez R, Kusne S, Green M, Abu-Elmagd K, Irísh W, Reyes J, Furukawa H, Tzakis A, Fung JJ, Todo S, Starzl TE (1995) Incidence and risk factors associated with the development of cytomegalovirus disease after intestinal transplantation. Transplantation 59: 1010–1014
- Mittal NK, Tzakis AG, Kato T, Thompson JF (2003) Current status of small bowel transplantation in children: update 2003. Pediatr Clin North Am 50: 1419–1433
- 22. Pappas PA, Tzakis AG, Gaynor JJ, Carreno MR, Ruiz P, Huijing F, Kleiner G, Rabier D, Kato T, Levi DM, Nishida S, Gelman B, Thompson JF, Mittal N, Saudubray JM (2004) An analysis of the association between scrum citrulline and acute rejection among 26 recipients of intestinal transplant. Am J Transplant 4: 1124–1132
- 23. Ruiz P, Bagni A, Brown R, Cortina G, Harpaz N, Magid MS, Reyes J (2004) Histological criteria for the identification of acute cellular rejection in human small howel allografts: results of the pathology workshop at the VIII International Small Bowel Transplant Symposium. Transplant Proc 36: 335–337
- Ruiz P, Garcia M, Pappas P, Berney T, Esquenazi V, Kato T, Mittal N, Weppler B, Levi D, Nishida S, Nery J, Miller J, Tzakis A (2003) Mucosal vascular alterations in isolated small bowel allografts: relationship to humoral sensitization. Am J Transplant 3: 43–49
- 25. Siegel LM, Stevens PD, Lightdale CJ, Green PHR, Garcia-Carrasquillo RJ, Rotterdam H (1997) Combined magnification endoscopy with chromoendoscopy in the evaluation of patients with suspected malahsorption. Gastrointesinal Endose 46: 226–230
- Sigurdsson L, Reyes J, Putman PE, Del Rosario JF, DiLorenzo C, Orenstein SR, Todo S, Kocoshis SA (1998) Endoscopies in pediatric small intestinal transplant recipients: five-year experience. Am J Gastroenterol 93: 207–211
- 27. Solez K, Axclsen RA, Benediktsson H, Burdick JF, Cohen AH, Colvin RB, Croker BP, Droz D, Dunnill MS, Halloran PF, Hayry P, Jennette JC, Keown PA, Marcussen N, Mihatsch MJ, Morozumi K, Myers BD, Nast CC, Olsen S, Racusen LC, Ramos EL, Rosen S, Sachs DH, Salomon DR, SanFilippo F, Verani R, VonWillebrand E, Yamaguchi Y (1993) International standardi-

zation of criteria for the histologic diagnosis of renal allograft rejection: the Banff working classification of kidney transplant pathology. Kidney Int 44: 411–422

- Stevens PD, Lightdale CJ, Green PHR, Siegel LM, Garcia-Carrasquillo RJ, Rotterdam H (1994) Combined magnification endoscopy with chromoendoscopy for the evaluation of Barret esophagus. Gastrointestinal Endose 40: 747–749
- Tabasco-Minguillan J, Hutson W, Weber K, Lec RG, Furukawa H, Abu-Elmagd K, Todo S, Rakela J (1996) Endoscopic features of acute cellular rejection. Transplant Proc 28: 2765–2766
- Tabasco-Minguillan J, Hutson W, Weber K, Lee RG, Dcmetris AJ, Furukawa H, Abu-Elmagd K, Todo S, Rakela J (1996) Prospective evaluation of endoscopy in acute cellular rejection and cytomegalovirus infection. Transplant Proc 28: 2778–2779
- Tabasco-Minguillan J, Weber K, Nelson F, Hutson W, Furukawa H, Abu-Elmagd K, Todo S, Rakela J (1996) Variability in the interpretations of endoscopic findings in patients with intestinal transplantation. Transplant Proc 28: 2775–2776
- 32. Timmermann W, Hoppe H, Otto C, Gasser M, Vowinkel T, Gassel AM, Meyer D, Gassel HJ, Ulrichs K, Thiede A (1999) Videomicroscopic imaging of graft mucosa for monitoring immunosuppressive therapy after small intestinal transplantation in rats. Transplantation 67: 1555–1561
- Todo S, Reyes J, Furukawa H, Abu-Elmagd K, Lee RG, Tzakis A, Rao AS, Starzl TE (1995) Outcome analysis of 71 clinical intestinal transplantations. Ann Surg 222: 270–282
- 34. Tzakis AG, Kato T, Nishida S, Levi DM, Tryphonopoulos P, Madariaga JR, DeFaria W, Nery JR, Regev A, Vianna R, Miller J, Esquenazi V, Weppler D, Ruiz P (2003) Alemtuzumab (Campath 1-H) combined with tacrolimus in intestinal and multivisceral transplantation. Transplantations 75: 1512–1517
- 35. Tzakis AG, Kato T, Nishida S, Levi DM, Madariaga JR, Nery JR, Mittal N, Regev A, Cantwell P, Gyamfi A, Weppler D, Miller J, Tryphonopoulos P, Ruiz P (2003) Preliminary experience with Campath 1H (C1H) in intestinal and liver transplantation. Transplantation 75: 1227–1231
- White FV, Reyes J, Jaffe R, Yunis EJ (1995) Pathology of intestinal transplantation in children. Am J Surg Pathol 19: 687– 698
- Yamataka A, Miyano T, Fukunaga K, Kobayashi H, Nozawa M, Sasaki K (1992) Patchy distribution of rejection changes in small intestinal transplantation. J Pediatr Surg 27: 602–603

Surg Endosc (2006) 20: 783–786 DOI: 10.1007/s00464-005-0257-8

© Springer Science+Business Media, Inc. 2006



and Other Interventionol Techniques

# Bravo capsule induction of esophageal hypercontractility and chest pain

C. Tharavej, J. A. Hagen, G. Portale, C.-C. Hsieh, T. A. K. Gandamihardja, J. C. Lipham, J. H. Peters, S. R. DeMeester, P. F. Crookes, C. G. Bremner, T. R. DeMeester

Division of Thoracic and Foregut Surgery, Department of Surgery, University of Southern California, Los Angeles, CA, USA

Received: 15 April 2005/Accepted: 8 November 2005/Online publication: 16 March 2006

#### Abstract

Background: The Bravo catheter-free pH monitoring system uses a capsule attached to the esophageal mucosa to detect acid exposure. Placement of the Bravo capsule is associated with intermittent chest pain in 50% of normal volunteers. The authors hypothesized that ehest pain in this setting may be attributable to hypertensive esophageal contractions induced by the Bravo eapsule. Methods: The study population consisted of 40 consecutive patients with reflux symptoms who had stationary esophageal manometry within 1 h after Bravo eapsule placement. The control group consisted of 40 patients with symptomatic gastroesophageal reflux disease (GERD) from a population of patients with foregut symptoms who were computer matched to the study group for age, sex, lower esophageal sphineter (LES) pressure, LES length, and 24-h pH composite score. The patients in the control group had manometry before Bravo eapsule placement. The occurrence of ehest pain was assessed before and during the monitoring period by interview and review of the patient's diary. Mean contraction amplitudes in the distal third of the csophagus after 10 wet swallows were averaged. The prevalence of patients with esophageal contraction amplitudes in the distal third that exceeded the 95th percentile of normal (180 mmHg) and the mean amplitude of distal third esophageal contractions in the study and control populations were compared. In the study group, the incidence of chest pain among the patients with hypercontractility of the esophagus was compared with the ineidence among those without hypercontractility.

*Results:* The mean contraction amplitude was higher in the study group (144.7 vs 105.5 mmHg; p = 0.002). The

number of patients with a mean distal esophageal contraction amplitude exceeding the 95th percentile of normal also was significantly higher in the study group (13/40 vs 5/40; p = 0.03). A total of 10 patients experienced new onset of ehest pain with the Bravo eapsule in place, and 6 patients experienced hypertensive esophageal contractions.

*Conclusions:* The intraesophageal Bravo capsule can eause hypertensive esophageal contractions, which may lead to chest pain.

Key words: Bravo capsule — Catheter-free pH monitoring system — Chest pain — Hypertensive esophageal contraction

Until recently, the standard method for measuring esophagcal acid exposure has been the eatheter-based ambulatory 24-h pH test. Although this test is the "gold standard" for assessing intraesophageal pH, it has limitations such as discomfort, inconvenience, and interference with normal activity [1]. These limitations lead patients to modify their diet and daily activities, which may indirectly affect the results of the test [2].

Recently, the Bravo eatheter-free pH monitoring system was introduced to overcome these shortcomings. A randomized trial has shown that the Bravo pH monitoring system is more convenient for the patient and interferes less with normal activity than the conventional catheter based system [3]. It also allows comfortable recording as long as 48 h for 89% of patients [2]. As a result, the new Bravo system is gaining popularity as an alternative to the eatheter technique.

We and others noted that ehest pain is frequently reported during the Bravo pH monitoring period, especially during swallowing. This led us to hypothesize that the Bravo capsule may cause chest pain by inducing esophageal dysmotility.

Presented at the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) meeting, Hollywood, FL, USA, 13–16 April 2005 *Correspondence to:* J. A. Hagen

# Materials and methods

# Patient population

From January 2003 to Fehruary 2004, 40 consecutive patients (median age, 45.5 ycars; range, 25–74 years) with gastroesophageal reflux symptoms had stationary csophageal manometry within 1 h *after* endoscopic Bravo capsule placement. This study group was compared with a control group consisting of 40 patients (median age, 49 years; range, 25–86 years) who had csophageal manometry 1 h *before* Bravo capsule placement. The control patients were matched to the study group for age, gender, lower esophageal sphincter (LES) pressure, LES length, and pH composite score. Patients with a history of achalasia, benign or malignant stricture, or a previous antireflux esophageal or gastric surgery were excluded from the study.

## Esophageal manometry

Standard esophageal motility studies were performed after an overnight fast. A 12-Fr, 8-lumen water-perfused motility catheter was passed through the anesthetized nostril into the stomach. The catheter had side holes located at 5-cm intervals, with 4 holes arranged radially (90° apart) at one level (Arndorfer Medical Specialties, Greendale, WI, USA). Each channel was connected to a pressure transducer, which was constantly perfused with hubble-free distilled water at 0.6 mL/min by a low-compliance pneumohydraulic pump (Arndorfer Medical Specialties).

The LES parameters including the resting pressure at the respiratory inversion point (RIP), the total LES length, and the length below the RIP (abdominal LES length) were measured using the stationed pull-through technique. Esophageal hody motility was assessed by positioning the catheter with the most proximal side hole 1 cm below the lower border of the upper esophageal sphincter and the other four side holes trailing at 5 cm intervals.

To assess esophageal hody contractility, 10 swallows of water (5 ml) were given at 25-s intervals. Data regarding wave progression, morphology, and contraction amplitude were recorded and analyzed using Polygram software version 4.2 (Medtronics, Inc., Shoreview, MN, USA). The amplitude of contraction in the distal third of the esophagus was defined as the mean amplitude in the lower two channels. Hypertensive esophageal contractions were defined as a mean amplitude above the 95th percentile determined in normal subjects (>180 mmHg).

### Bravo capsule and pH monitoring system

The Bravo capsule  $(6 \times 5.5 \times 25 \text{ mm})$  was attached to the csophageal mucosa using a prepackaged assembly (Fig. 1). The capsule has a well (diameter, 4 mm; depth, 3.5 mm) connected to an external vacuum unit during placement for attaching a small pin to the esophageal mucosa. The capsule contains antimony pH and reference electrodes located at the distal end. By means of radiotelemetry, pH data are transmitted to a receiver worn on the belt. After 48 h, the patient returned the receiver, and the data were uploaded to a computer for analysis.

# Placement of the Bravo capsule

# Study group

The Bravo capsule was placed during endoscopy with the patient under conscious sedation using intravenous Demerol and Versed. The distance between the upper limit of the gastric rugal folds (the gastroesophageal junction) and the incisor teeth was measured. The endoscope was removed, and the delivery system was passed transorally into the esophagus. There it was advanced until the pH sensor of the capsule was located at a point 6 cm above the upper limit of the rugal folds [4]. Capsule attachment was confirmed by reintroduction of the endoscope (Fig. 2). A motility catheter then was passed into the stomach, and esophageal motility was performed as described earlier.



Fig. 1. Bravo capsule (left) and delivery system with receiver (right).



Fig. 2. Endoscopic view of an attached Bravo capsulc.

### Control group

After standard esophageal manometry, the position of the upper border of the LES was identified. The Bravo delivery system was passed transnasally, and the capsule was attached 5 cm above the upper border of the LES. A chest x-ray was obtained to confirm attachment of the capsule.

#### Chest pain symptom assessment

Chest pain was assessed in the study group by interview hefore placement of the Bravo capsule. After placement, each patient was given a standardized symptom questionnaire. The presence or absence of chest pain before and during the monitoring period was recorded. Patients with a history of chest pain before the Bravo study and those with incomplete self-reported questionnaires were excluded from the chest pain assessment analysis.

# Ethical considerations

This study protocol was approved hy Health Sciences Campus, the Institution Review Board of the University of Southern California.

# Statistical analysis

Chi-square and Fisher's exact tests were used to compare categorical variables, and the Student's *t*-test was used for continuous variables. Significant differences were assumed at a p value less than 0.05.

Table 1. Characteristics of the study and control groups

	Study group $(n = 40)$	Control group $(n = 40)$	p Value	
Age (years): <i>n</i> (range)	45.5 (40-53) <sup>a</sup>	49 (44-63) <sup>a</sup>	0.18	
Gender (M/F)	15/25	15/25	1.0	
Composite pH score	$19.27 \pm 2.89$	$19.21 \pm 3.41$	0.99	
Total length (cm)	$3.04 \pm 0.16$	$2.94 \pm 0.12$	0.62	
Ahdominal length (cm)	$1.77 \pm 0.13$	$1.62 \pm 0.10$	0.38	
LES pressure (mmHg)	$16.95 \pm 1.44$	$17.96 \pm 1.64$	0.64	

LES, lower esophageal sphincter

<sup>a</sup> Interquartile range (IQR)



Fig. 3. Comparison of mean distal esophageal contraction amplitude (DEA) between groups. The mean DEA was significantly higher in the study group (147.7  $\pm$  9.9 vs 105.5  $\pm$  8.8 mmHg; p = 0.002).

#### Results

Each group consisted of 15 male and 25 female patients. There were no significant differences in demographie data, LES parameters, and pH composite score between the two groups (Table 1).

The distal esophageal contraction amplitude was significantly higher in patients with a Bravo capsule in place (147.7  $\pm$  9.98 vs 105.5  $\pm$  8.84 mmHg; p = 0.002) (Fig. 3). A total of 13 patients (33%) in the study group had distal esophageal contraction amplitudes exceeding the 95th percentile of normal, as compared with only 4 patients (10%) in the control group (Fig. 4). The total number of hypertensive esophageal contractions also was significantly higher among the patients with the Bravo capsule in place (134/400 vs 55/400; p < 0.0001) (Fig. 5).

The self-report questionnaire was completed by 20 patients in the study group who had no history of chest pain. New onset chest pain developed in 10 of these patients (50%) after Bravo eapsule placement. In 6 of these patients, the pain was associated with hypertensive esophageal contractions. All the patients with new onset chest pain had hypertensive esophageal contractions. In contrast, hypertensive contractions were present in only 4 of the 14 patients (29%) without chest pain (p = 0.003) (Fig. 6).

#### Discussion

A recent randomized control trial has shown that the Bravo catheter-free pH monitoring system is better tol-



Fig. 4. Comparison of the number of patients with distal esophageat contraction amplitude (DEA) exceeding 180 mmHg hetween groups. The number of patients with DEA exceeding 180 mmHg was higher in the study group (13/40 vs. 5/40; p = 0.03).



Fig. 5. Comparison of the number of swallows with distal esophageal contraction amplitude (DEA) exceeding 180 mmHg between the groups. The total number of wet swallows in each group was 400. The number of swallows with a DEA exceeding 180 mmHg was higher in the study group (134/400 vs 55/400;  $p \le 0.0001$ ).



Fig. 6. Comparison of the prevalence of new chest pain hetween patients with (n = 6) and those without (n = 14) hypertensive esophageal contractions. The prevalence of new chest pain was higher among the patients with hypertensive contractions [6/6 (100%) vs 4/14 (28.5%); p = 0.003].

erated by patients and allows more normal activity than the conventional catheter-based system. As a result, it is rapidly gaining popularity among patients with GERD. As experience with this technique has grown, it has become evident that patients commonly report chest pain with the Bravo capsule in place. In a recent randomized trial comparing Bravo and conventional pH monitoring systems [3], chest pain was experienced by 60% of the patients monitored with the Bravo capsule. Pandolfino et al. [2] also have reported a significant frequency of chest pain after Bravo placement. In this smaller series, 34.5% of the patients experienced chest pain, and in 2 of 10 patients, the chest pain was severe enough for the patient to request immediate capsule removal. The mechanism of chest pain in these patients is not clear.

We hypothesized that esophageal motility abnormalities may be the cause of this chest pain. This hypothesis was based on previous studies demonstrating the strong relationship between esophageal motor abnormalities and the symptom of chest pain [5]. In particular, esophageal hypercontractility has been reported in 27% to 59% of patients with (noncardiac) chest pain [6–9].

We recently completed a study to assess the clinical utility of the Bravo system. This study included patients evaluated at two clinical sites. At one site, the Bravo capsule was placed transnasally at a location determined by manometry performed before Bravo placement. At the other site, the majority of patients had the Bravo capsule placed transorally at a location determined by endoscopy. In these patients, esophageal manometry often was performed after Bravo placement. This allowed us to identify the study and control groups included in this report. These patients, matched for age, sex, LES characteristics, and composite pH score, provided comparable groups for studying the possible effects of a Bravo capsule on motility.

Our results confirm previous observations regarding the common occurrence of new onset chest pain after Bravo insertion. In our experience, this symptom develops after Bravo placement in 50% of patients previously free of chest pain. We also have shown that 60% of the patients in whom chest pain develops will have esophageal hypercontractility when motility is performed with a Bravo capsule in place. Furthermore, even in patients without a new onset chest pain, hypertensive contractions frequently occur in the distal esophagus. One-third of all patients studied with the Bravo capsule in place had hypertensive contractions, which was more than three times the frequency in the control group. In addition, more than one-third of all contractions measured in the Bravo group were in the hypertensive range. Together, these observations support our hypothesis, suggesting that the Bravo capsule may indeed induce contraction abnormalities in the esophagus known to be associated with chest pain.

For the patients in the study group, the motility studies were performed after Bravo capsule insertion for the purposes of patient comfort and convenience. This approach not only allowed the endoscopy, motility study, and Bravo monitoring all to be accomplished in one session, but it also allowed placement of the Bravo capsule with the patient under sedation. Clearly, this strategy is attractive to patients. The results of this study suggest the need for caution in using this approach because assessment of esophageal motility can be complicated. The duration of the contraction abnormalities associated with Bravo placement is unknown, but it seems prudent to allow sufficient time to pass between Bravo monitoring and esophageal manometry for detachment of the capsule. In the largest study on Bravo monitoring reported to date, capsule detachment occurred within 14 days for 99% of those studied [10].

### Conclusion

In conclusion, the Bravo capsule frequently causes hypertensive contractions in the distal esophagus, and these contractions are commonly associated with chest pain. Consequently, manometry should not be performed with a Bravo capsule in place, and patients scheduled for Bravo pH monitoring should be informed that chest pain may develop.

#### References

- Fass R, Hell R, Sampliner RE, Pulliam G, Graver E, Hartz V, Johnson C, Jaffe P (1999) Effect of ambulatory 24-hour esophageal pH monitoring on provoking activities. Dig Dis Sci 44: 2263– 2269
- Pandolfino JE, Richter JE, Ours T, Guardino JM, Chapman J, Kahrilas PJ (2003) Ambulatory esophageal pH monitoring using a wireless system. Am J Gastroenterol 98: 740–749
- Wong WM, Bautista J, Dekel R, Malagon IB, Tuchinsky I, Green C, Dickman R, Esquivel R, Fass R (2005) Feasibility and tolerability of transnasal/per oral placement of the wireless pH capsule vs traditional 24-h oesophageal pH monitoring: a randomised trial. Aliment Pharmacol Ther 21: 155-163
- Kahrilas PJ, Lin S, Chen J, Manka M (1999) The effect of hiatus hernia on gastroesophageal junction pressure. Gut 44: 476–482
- Pilhall M, Borjesson M, Rolny P, Mannheimer C (2002) Diagnosis of nutcracker esophagus, segmental or diffuse hypertensive patterns, and clinical characteristics. Dig Dis Sci 47: 1381–1388
- Brand DL, Martin D, Pope CE II (1977) Esophageal manometrics in patients with angina-like chest pain. Am J Dig Dis 22: 300–304
- Chobanian SJ, Benjamin SB, Curtis DJ, Catlau EL Jr (1986) Systematic esophageal evaluation of patients with noncardiac chest pain. Arch Intern Med 146: 1505–1508
- Katz PO, Dalton CB, Richter JE, Wu WC, Castell DO (1987) Esophageal testing of patients with noncardiac chest pain or dysphagia. Ann Intern Med 106: 593–597
- Traube M, Albibi R, McCallum RW (1983) High-amplitude peristallic esophagesl contractions associated with chest pain. JAMA 250: 2655–2659
- Lin E, Waring JP, Ramaswamy A (2003) Analysis of 245 consecutive studies using 48-hour wireless pH(Bravo) probes for GERD evaluation. Gastroenterology 124: A689

Surg Endosc (2006) 20: 787–790 DOI: 10.1007/s00464-004-2186-3

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

# Laparoscopic hepatic resection

# G. Borzellino, A. Ruzzenente, A.-M. Minicozzi, F. Giovinazzo, C. Pedrazzani, A. Guglielmi

First Department of General Surgery, University of Verona, Piazzale Stefani 1, 37126 Verona, Italy

Received: 25 July 2004/Accepted: 10 June 2005/Online publication: 16 March 2006

#### Abstract

Background: Although laparoscopy in general surgery is increasingly being performed, only recently has liver surgery been performed with laparoscopy. We critically review our experience with laparoseopie liver resections. Methods: From January 2000 to April 2004, we performed laparoseopie hepatic resection in 16 patients with 18 hepatic lesions. Nine lesions were benign in seven patients (five hydatid eysts, three hemangiomas, and onc simple eyst), five were malignant in five patients (five hepatoearcinoma), and four patients had an uncertain preoperative diagnosis (one suspected hemangioma and three suspected adenomas). The mean lesion size was 5.2 cm (range, 1-12). Twelve lesions were located in the left lobe, three were in segment VI, one was in segment V, one was in segment IV, and one was in the subcapsular part of segment VIII.

*Results:* The conversion rate was 6.2%; intraoperative bleeding requiring blood transfusions occurred in two patients. Mean operative time was 120 min. Mean hospital stay was 4 days (range, 2–7). There were no major postoperative complications and no mortality.

*Conclusions:* Hepatie resection with laparoscopy is feasible in malignant and benign hepatic lesions located in the left lobe and anterior inferior right lobe segments (IV, V, and VI). Results are similar to those of the open surgical technique in carefully selected cases, although studies with large numbers of patients are necessary to drawn definite conclusions.

Key words: Laparoscopy — Liver resection

During the past few decades, the acquisition of new teehnology and the improved surgieal experience have increased indications for laparoscopic abdominal surgery. In hepatie surgery, laparoscopy has been confined to the staging of tumors, biopsy, and the treatment of simple scrous cysts. Only recently has laparoscopic hepatic resection been successfully performed. A review of

Correspondence to: G. Borzellino

the literature reports only 200 cases of hepatie resection published up to 2001 [3].

Surgieal skills in both liver open surgery and laparoseopie surgery are required. Moreover, preoperative and intraoperative assessment is necessary to define the location of the tumor and its relationships with vascular structures. Proper technological support is required for the dissection of the parenehyma for adequate control of hemostasis [6].

This article reports our experience with laparoseopie hepatic resection and critically reviews indications and results.

### Materials and methods

From January 2000 to April 2004, 16 patients underwent laparoscopic hepatic resection for 18 lesions. Patients were positioned in the supine position, with the legs abducted, and the surgeon stood hetween the legs with the first assistant to the patient's right. The first 10 mm trocar was inserted in the umbilical site after creating pneumoperitoneum. After exploration of the abdominal cavity, three trocars were positioned: one in the epigastrium; one in the right hypochondrium for retraction of the liver and aspiration; and one 10 mm trocar in the left hypochondrium for clips.

Intraoperative ultrasonographic evaluation of the hepatic lesion and its relations with vessels was performed in all cases. For lesions located in segments II and III, the left lobe was completely mobilized. In lesions located in segments V and VI, we performed a mobilization of the right lobe when necessary.

In patients with hydatid cysts, removal of the cysts was achieved by cystopericystectomy in all cases. In one case with an 8 cm cyst, we used intraoperative PAIR (puncture, aspiration, injection, reaspiration) under laparoscopic control before cystopericystectomy.

For patients with malignant lesions, we performed atypical resection in one patient with a peripheral exophytic lesion and typical segmentectomy or bisegmentectomy in four patients with lesions located deeper in the parenchyma. In all cases of malignant lesions, surgical margins were larger than 2 cm.

Laparoscopic liver resection was also indicated in three patients with uncertain diagnosis (hepatic adenoma vs focal nodular hyperplasia) and in another case for atypical hemangioma in a patient with previous breast cancer.

In the first three cases [hepatocarcinoma (HCC) of segment 11, hydatid cyst of segment V1, and hemangioma of segment 11], we used the Kellyclasia technique for hepatic parenchymal dissection, monopolar cautery, and/or application of vascular clips for hemostasis. In other patients, ultrasound scalpel was utilized for parenchymal section.

For extraction of the surgical specimen, we performed a suprapubic transverse 8 cm incision for solid lesions or enlargement of the periumbilical incision in cystic lesions. In all cases, a plastic bag was used.

## Results

During the study period, we performed laparoscopic liver resection in 16 patients (five male and 11 female) with an average age of 51 years (range, 28–74). Indications for laparoscopic resection, the characteristics of the lesions, and their locations are summarized in Table 1.

Among 18 resected lesions, nine were benign (five hydatid cysts, two symptomatic hemangiomas, one hemangioma associated with cholclithiasis, and one simple cyst), five were malignant (HCC in Child–Pugh class A eirrhosis), and lour had uncertain preoperative diagnosis (one atypical hemangioma and three suspected adenomas).

As shown in Table 1, lesions were mainly located in lcft segments II and III and right inferior segments V and VI. Only in one case was the lesion located at the surface of the anterior part of segment VIII.

Surgery was performed using the laparoscopic approach in 15 of 16 patients, with a 6.2% of conversion rate (one case of hemangioma of segment II). Intraoperative hemorrhage requiring blood transfusion (2 units) occurred in two cases. In the first case, hemorrhage was successfully managed by laparoscopy, and in the second ease it resulted in conversion to open surgery. The two eases of intraoperative hemorrhage occurred during the first period when the Kelly-clasia technique was utilized for dissection of parenchyma. Use of the ultrasonic sealpel improved results in the following patients, with a mean blood loss of 150 ml (range, 50–200). Resection was performed without vascular clamping in all cases.

The mean operative time was 120 min (range, 60– 180). Mean hospital stay after surgery was 4 days (range, 2–7). There were no major surgical complications and no operative mortality. In two patients, minor surgical complications occurred (surgical wound infection in one and persistent abdominal pain in another).

# Discussion

Among patients for whom liver surgery is indicated, those for which a laparoscopic approach can be proposed are yet limited [3, 10, 13]. Only a few can be submitted to laparoscopic liver resection. In our experience of 187 liver resections performed during the study period, we chose the laparoscopic approach for only 16 patients. Despite the small number of patients, our experience confirms the feasibility of laparoscopic hepatic resection, with results similar to those reported in the literature.

Berthio et al. [3] reviewed 200 cases of resection for both benign and malignant lesions and reported a conversion rate of 7%, mean hospital stay of 7.7 days, postsurgieal morbidity of 16%, and mortality of 0.5%. Two different studies on laparoscopic hepatic resection, one for benign lesions and one for malignant lesions, reported similar results in terms of conversion rates (10 and 13.5%, respectively), mean hospital stay (5 and 7 days, respectively), and mortality (0% in both studics). On the contrary, morbidity was 5% for patients with resection for benign lesions and 22% for malignant lesions.

In comparative studies of open liver surgery, mortality and morbidity were not increased in laparoscopy, operative time was similar or slightly longer, blood loss was similar or lower, and mean hospital stay was shorter for the laparoscopy group [12, 15, 20, 23, 25].

Control of hemostasis is one of the major topics of liver surgery and use of ultrasonic or radiofrequency scalpels, used more commonly during laparoscopic resection, helps to reduce blood loss, as reported in comparative studies between laparoscopie and open hepatic surgery. However, failure to control bleeding during laparoscopic surgery is one of the most important eauses of conversion to the open teehnique [2, 7, 9, 10, 13, 15, 22]. In our experience, after introduction of ultrasound scissors blood loss was reduced to a mean of 150 ml and the conversion rate was reduced to 0%.

Argon beam coagulation is effective for hemostasis, but gas embolism is reported in the literature and it has been related to the increase in intraabdominal pressure [8, 14]. For this reason, the use of argon bean coagulation is not recommended for laparoscopic liver resections.

The most important factor for laparoscopic resection is selection of patients. Size and site of lesion and its relationship with a major vessel must be accurately evaluated during preoperative staging.

Benign lesions are generally referred for surgical resection for the presence of symptoms or for uncertain diagnosis. Hemangiomas and nodular focal hyperplasia are rarely submitted to surgical resection, and few cases have been reported in the literature [10, 16, 17, 19, 22].

In our experience, hemangiomas were referred for surgical resection for preoperative uncertain diagnosis in one ease, whereas in the other ease, the patient sulfered from cholelithiasis and the hemangioma, located on the surface of segment V, was resected during laparoscopic cholecystectomy.

The role of hepatic resection for hydatid cysts is controversial, and the introduction of laparoscopy has only added a new element to the debate. The treatment of choice is still total cystopericystectomy, which reduces the frequency of intraoperative anaphylactic reactions and relapses [5]. Laparoscopic partial cystectomy or PAIR lead to results similar to those of open surgical and percutaneous techniques: anaphylactic reaetions in 3% of cases and relapse in 3–10% of patients after a follow-up of 17 or 18 months [4, 11, 24]. Manterola et al. [21] published a series of eight eases of laparoscopic total cystopericystectomy, with anaphylactic reactions and relapse in none of the patients after a mean follow-up of 30 months.

Laparoscopic treatment of malignant lesions is still a matter of debate, even though laparoscopic liver resection was successfully applied in cirrhotic patients with primary tumors [7]. Comparative studies with open surgery have shown a shorter hospital stay with the Table 1. Site, size, and histological features of lesions

Preoperative diagnosis	No. of patients	Site (segment)	Size (cm)	Histological features
Benign lesions	7			
		11-111	8	Hydatid cyst
		VI	4	Hydatid cyst
		VI-IV-VIII	3, 2.5, 1	Hydatid cyst
		VI	9	Hemangioma
		11	8	Hemangioma
		V	4.5	Hemangioma <sup>a</sup>
		11-111	12	Simple cyst
Malignant lesions	5			
		111	6	HCC
		11	5	HCC
		11-111	3	HCC
		11-111	3	HCC
		11–111	2	HCC
Uncertain diagnosis	4			
		11-111	7	Hemangioma <sup>b</sup>
		II-III	5	FNH vs adenoma
		11-111	7.5	FNH vs adenoma
		11-111	6	FNH vs adenoma

FHN, focal nodular hyperplasia; HCC, hepatocarcinoma

<sup>a</sup> Associated with cholelitiasis

<sup>b</sup> Patients with previous breast cancer

laparoscopic approach, with similar morbidity, mortality, and long-term results [20, 25]. However, a multicenter study reported discouraging results, with surgical margins < 1 cm in 30% of resected HCC and in 20% of resected metastasis; moreover, positive surgical margins were found in 6.7% of patients [13].

In cirrhotic patients, laparoscopic liver resection has been performed with good results also with coexisting poor hepatic function [1, 18, 26]. The authors concluded that saving the abdominal wall vascular collaterals in the laparoscopic approach reduced postoperative decompensation in patients with portal hypertension [7]. These preliminary data suggest an extension of indications for surgery of primary malignancy in cirrhotic patients, but comparative studies with less invasive treatments are necessary.

Currently, the laparoscopy approach is employed mainly for minor hepatic resections, wedge resections, segmentectomy, and bisegmentectomy. Experience in laparoscopic major liver resections is limited to sporadic cases in the literature, and usually the approach is not fully laparoscopic [3, 10, 15].

In conclusion, laparoscopic hepatic resection is technically feasible in properly selected cases. Indications for benign lesions are limited, and the role of laparoscopic liver resection in malignant lesions is still controversial. Studies with large numbers of patients and long-term follow-up must be performed before more precise conclusions on the role of laparoscopy in liver surgery can be drawn.

#### References

 Abdel-Atty MY, Farges O, Jagot P, Belghiti J (1999) Laparoscopy extends the indications for liver resection in patients with cirrhosis. Br J Surg 86: 1397–1400

- Berends FJ, Meijer S, Prevoo W, Bonjer HJ, Cuesta MA (2001) Technical considerations in laparoseopic liver surgery. A solid organ easily forgotten. Surg Endose 15: 794–798
- Berthio L, Waage A, Gagner M (2002) Hépatectomies sous laparoscopie. Ann Chir 127: 164–170
- Bickel A, Loberant N, Singer-Jordan J, Goldfeld M, Daud G, Eitan A (2001) The laparoscopic approach to abdominal hydatid eysts. Arch Surg 136: 789–795
- Casado OA, Gonzales EM, Segurola CL, Calvo AG, Pinto IG, Saborido BP. Crespo GP, Johansson CO (2000) Results of 22 years of experience in radical surgical treatment of hepatic hydatid cysts. Hepato-Gastroenterology 48: 235–243
- Cherqui D (2002) Résections hépatiques par voie coelioscopique. Utiles ou futiles? Ann Chir 127: 171–174
- Cherqui D, Husson E, Hammoud R, Malassagne B, Stéphan F, Bensaid S, Rotman N, Fagniez PL (2000) Laparoscopic liver resections: a feasibility study in 30 patients. Ann Surg 6: 753–762
- Croce E, Azzola M, Russo R, Golia M, Angelini S, Olmi S (1994) Laparoscopic liver resection with the argon beam. Endosc Surg Allied Technol 2: 186–188
- Descottes B, Glineur D, Lachaehi F, et al. (2003) Laparoscopic liver resection of benign liver tumors. Surg Endosc 17: 23–30
- Descottes B, Lachachi F, Durand-Fontanier S, Sodji M, de Pech Laclause B, Valleix D (2000) Traitement laparoscopique des tumeurs solides ete kystiques du loie. Etude de 33 cas. Ann Chir 125: 941–947
- Ertem M, Uras C, Karahasanoglu T, Erguney S, Alemdaroglu K (1998) Laparoscopic approach to hepatic hydatid disease. Dig Surg 15: 333–336
- Farges O, Jagot P, Kirstetter P, Marty J, Belghiti J (2002) Prospective assessment of the safety and benefit of laparoscopic liver resections. J Hepatobiliary Pancreat Surg 9: 242–248
- 13. Gigot JF, Glineur D, Santiago Azagra J, et al. (2002) Hepatobiliary and Pancreatic Section of the Royal Belgian Society of Surgery and the Belgian Group for Endoscopic Surgery Laparoscopic liver resection for malignant liver tumors: preliminary results of a multicenter European study. Ann Surg 236: 90–97
- Hashizume M, Takenaka K, Yanaga K, Ohta M, Kajiyama K, Shirabe K, Itasaka H, Nishizaki T, Sugimachi K (1995) Laparoscopic hepatic resection for hepatocellular carcinoma. Surg Endose 9: 1289– 1297
- Huscher CGS, Lirici MM, Chiodini S (1998) Laparoscopic liver resection. Semin Laparoscopic Surg 5: 204–210
- Kalil AN, Mastalir ET (2002) Laparoscopic hepatectomy for benign liver tumors. Hepato-Gastroenterology 49: 803–805

- Katkhouda N, Mavor E (2000) Laparoscopic management of benign liver disease. Surg Clin North Am 80: 1203–1211
- Ker CG, Chen HY, Juan CC, Chang WS, Tsai CY, Lo HW, Yau MT (2000) Laparoscopic subsegmentectomy for hepatocellular carcinoma with cirrhosis. Hepato-Gastroenterology 47: 1260–1263
- La Porte L, Hamy A, Paineau J, Visset J (1996) Chirurgie coelioscopique pour tumeur bénigne du foie. J Chir 9/10: 432– 436
- Mala T, Edwin B, Gladhaug I, Fosse E, Soreide O, Bergan A, Mathisen O (2002) A comparative study of the short-term outcome following open and laparoscopic liver resection of colorectal metastases. Surg Endosc 16: 1059–1063
- Manterola C, Fernandez O, Munoz S, Vial M, Losada H, Carrasco R, Bello N, Barroso M (2002) Laparoscopic pericystectomy for liver hydatid cysts. Description of results observed during long follow-up period. Surg Endosc 16: 521–524

- Mouiel J, Katkhouda N, Gugenheim J, Fabiani P (2000) Possibilities of laparoscopic liver resection. J Hepatobiliary Pancreat Surg 7: 1–8
- Rau HG, Buttler E, Meyer G, Schardey HM, Schildberg FW (1998) Laparoscopic liver resection compared with conventional partial hepatectomy—a prospective analysis. Hepato-Gastroenterology 45: 2333–2338
- 24. Seven R, Berber E, Mercan S, Eminoglu L, Budak D (2000) Laparoscopic treatment of hepatic hydatid cysts. Surgery 128: 36-40
- Shimada M, Hashizume M, Maehara S, Tsujita E, Rikimaru T, Yamashita Y, Tanaka S, Adachi E, Sugimachi K (2001) Laparoscopic hepatectomy for hepatocellular carcinoma. Surg Endosc 15: 541–544
- Yamanaka N, Tanaka T, Tanaka W, Yamanaka J, Yasui C, Ando T, Takada M, Maeda S, Okamoto E (1998) Laparoscopic partial hepatectomy. Hepato-Gastroenterology 45: 29–33

Surg Endose (2006) 20: 791–793 DOI: 10.1007/s00464-004-2157-8

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

# The "stamp method": a new treatment for perforated peptic ulcer?

M. J. O. E. Bertleff,<sup>1</sup> R. S. B. Liem,<sup>2</sup> H. L. Bartels,<sup>3</sup> P. H. Robinson,<sup>1</sup> J. F. A. Van der Werff,<sup>4</sup> H. J. Bonjer,<sup>5</sup> J. F. Lange<sup>6</sup>

<sup>1</sup> Department of Plastic Surgery. Academic hospital Groningen, Hanzeplein 1, P.O. Box 30001, Groningen, 9700 RB . The Netherlands

<sup>2</sup> Department of Cell Biology, Section Electron Microscopy, University Groningen, The Netherlands

<sup>3</sup> Lahoratory for Animal Research, University of Groningen, The Netherlands

<sup>4</sup> Department of Plastic Surgery, Slotervaart Hospital, Amsterdam, The Netherlands

<sup>5</sup> Erasmus Medical Center Rotterdam, The Netherlands

<sup>6</sup> Department of Surgery, Medical Center Rijnmond Zuid, Rotterdam, The Netherlands

Received: 1 July 2004/Accepted: 11 Fehruary 2005/Online publication: 16 March 2006

# Abstract

*Background:* The aim of this study was to develop a simple method for closure of a perforated peptic uleer, making it more accessible for laparoscopic surgery.

*Methods:* An experimental pilot study was performed using five male Wistar rats. The perforation was closed by a bioabsorbable patch made of lactide–glyeolid– caprolactone fixed with glue onto the outside of the stomach.

*Results:* Postoperatively, there were no signs of leakage or other complications. Histologically, there were no signs of inflammation on the inside of the stomach, and there was a 50% reduction of the perforation each successive postoperative week. No adverse reactions beeause of the degradable material or gluc were observed. *Conclusions:* Treatment of a perforated peptie ulcer by placing a patch of biodegradable material like a "stamp" on the outside of the stomach is a feasible option.

Key words: Perforated peptic ulcer — Laparoscopy — Biodegradable patch — Glue

# Introduction

Laparoscopic correction still is not the gold standard for management of a perforated peptic uleer [4] although many advantages of laparoscopic procedures have been demonstrated during the recent years with regard to postoperative morbidity and pain [1, 8, 9]. During a multicenter Dutch trial (LAMA trial), we compared laparoseopie elosure of perforated peptie uleer with the eonventional method using laparotomy. It seemed that one of the problems with the laparoscopic procedure involves the suturing teehnique [2]. It is especially difficult for surgeons to take big bites, to prevent cutting out of the sutures [9], and to exert sufficient tension on the knot during intra- or extraeorporal knotting.

The operation time for laparoscopic correction often is prolonged [2, 6], which is mainly due to the learning curve with regard to suturing technique. This may be the reason why many surgeons (especially during the night) are not even starting up laparoscopically or soon convert to laparotomy. A new method has been developed in which perforation of the stomach in rats was elosed with a biodegradable patch fixed with glue, similar to putting a "stamp" on the outside of the stomach. In the current eurrent experiment, this "stamp" method was evaluated.

# Methods

Five adult male Wistar rats with an average weight of 255 were used in this trial. With the rats general anesthesia (isoflurangas 2% with oxygen), a median laparotomy was performed. At a fixed point on the ventral side of the stomach, a perforation was created with a diameter of 0.5 cm. A patch, made of lactide–glycolid–caprolactone (LGC) (Polyganics, B.V., Groningen, The Netherlands) was cut into a circle with a diameter of 1 cm and an overlap of 0.5 cm all around the perforation. The patch was glued on the outside of the stomach with Gluhran 2 (n-hutyl [2] cyanoacrylate, methaeryloxysulfolane). The addomen was elosed in two layers with Polysorh 4-0.

After lweek, the first rat underwent relaparotomy under general anesthesia. After inspection of the abdomen, the patch, including the full thickness of the stomach wall, was resected and sent for histology. Each successive postoperative week, one rat underwent the same procedure, resulting in a total elinical and histologie follow-up period of 5 weeks.

For histology, tissue specimens were rinsed in saline and placed into a fixative containing 2% glutardialdehyde huffered with 0.1 mol/ phosphate huffer, pH 7.4. Then the specimens were dehydrated through a graded concentration of ethanol and emhedded in glycol methaerylate. From all samples, 2-µm-thick sections were prepared using a disposable histoknife and a Reichert–Jung "2050 supercut"

Correspondence to: M. J. O. E. Bertleff



Fig. 1. Histology at week 1.

microtome. The sections were mounted on glass slides and stained with toluidine blue and alkaline fuchsin (Merck, Darmstadt, Germany). All the sections were evaluated and photomicrographed using a Olympus BX-50 microscope (Olympus Optical Co., Japan)

# Results

All the rats survived the first operation without complications. The mean operating time was 10 min. At relaparotomy, there were neither signs of leakage nor evidence of peritonitis. There were small adhesions, mainly to the liver, and in the first two rats there also were small adhesions between the spleen and the stomach. From week 1, the patch was covered by omentum. Biodegradation of the patch was visible at week 5. The diameter of the perforation, as observed from the inside of the stomach, decreased by 50% cvcry week. After 5 weeks, only a pinpoint perforation could be found. All the rats gained weight during their weeks of follow-up evaluation, with an average weight of 348 g (range, 313-392 g), on the day of re-operation meaning they were in good condition.

# Histology

At 1 weck postoperatively, no inflammatory cells were detected on the inside of the stomach. Bacteria were found in the superficial mucous layer of the epithelium, and among the microvilli of cpithelial cells. They were distributed irregularly, patchy and with heavy colonization, in some areas but did not invade the epithelium. The mucosa consisted of dense connective tissue and numerous blood vessels. The basement membrane underlying the epithelial basal cells was clearly visible. The muscular layer and submucosal glands were present, but did not continue, as was to be expected.

At 2 weeks postoperatively, cellular ingrowth of inflammatory cells, especially granulocytes, was seen lying against the LGC patches. Multinucleated giant cells also were scen at the interface of the patches.

At 3 weeks postoperatively, the LGC patches were covered by a capsule of fibrotic tissue. This capsule consisted of 14 to 15 layers of fibroblasts, collagen fibcrs, extracellular matrix, and numerous blood vessels. Still, some multinucleated giant cells were observed in the patches, and degradation of the LGC patches had begun.

At 4 weeks postoperatively, the amount of inflammatory infiltrate had increased, while other cells such as macrophages and multinucleated giant cells were seen infiltrating the LGC-patches. The capsule of fibrotic layer became ticker. Fibroblasts as well as collagen fibers and blood vessels were found more frequently and denser, as compared with findings 3 weeks postoperatively.

At 5 weeks postoperatively, the fibrotie layer on the patch had increased Macrophages and giant cells still were found infiltrating and phagocytosing the LGC patches. There were epithelial cells close to the perforation. The muscular layer still showed perforation. There were no signs of rejection (Figs. 1 and 2).

# Discussion

A new method for closure of peptie uleer perforation was tested in rats, and the first results seem promising. Sealing of the perforation with a glued biodegradable patch seemed sufficient because no leakage occurred. Furthermore, no inflammation or other side effects to the abdominal wall were observed. It was decided that at this phase, the use of an iatrogenically made perforation would be sufficient, because we wanted only to evaluate whether this new technique would work.

Of eourse, this test did not eompletely mimie the elinical situation. There was no edema, no fibrin deposition, and no inflammation of surrounding tissue. However, no elinical evidence so far had proved that a perforation of any longer duration with fibrin deposition has a worse outcome with regard to healing of the perforation itself and a higher risk for reperforation. It eould even be suggested that fibrin deposition helps in sealing the perforation, but more research on this topic is needed.

Closure of a perforated peptie uleer by a Graham omental patch or mere sutures has been performed for many years [7, 8]. Several alternative techniques have been tried [7, 9]. The incentive for introducing these new operations was to simplify the procedure and make it suitable for minimal invasive therapy [9]. A few procedures ean be accomplished by endoscopy, but often it still is necessary to combine it with laparotomy or laparoscopy [3, 6]. Lau et al. [5] described a method for closing the perforation using spongostan fixed with fibrin glue. This seemed to be suitable only for smaller perforations. The patch used in this study can be introduced through a trocar and unfolded with ease because it has no memory. Glubran 2, the glue used in this trial, is a synthetic surgical glue european conformity (CE) certified for internal use. It is liquid, does not need any preparation and ean be applied to the patch inside the abdomen using a laparoscopic needle. The size of the perforation does not matter because the patches ean be cut easily into any desired size.

In eonelusion we propose a simple technique for elosure of a perforated peptie uleer, making laparoseopie correction of a perforated peptic uleer more accessible. A randomized elinical trial will be initiated in due course.

*Acknowledgement*. The authors thank Polyganies, BV (Gronigen, The Netherlands) for providing the patches and (Global Entrepreneurship Monitor) GEM (Italy) for providing the glue and Mr. G. Kors for the technical assistance.

#### References

- Agresta F, Michelet I, Coluci G, Bedin N (2000) Emergency laparoseopy: a community hospital experience. Surg Endose 14: 484– 487
- Bertlelf MJOE, Bemelman W, Oei I, Smulders J, vanderHam A, Bonjer H, Lange J (2002) Laparoscopic correction versus open surgery of perforated peptic ulcer: first results of the LAMA trial. Surg Endosc 16: S3
- Hashiba K, Carvaiho AM, Diniz G, Barbosa de Aridrade N, Guedes CAF, Filho LS, Lima CAP, Coetho HE, Oliveira RA (2001) Experimental endoscopic repair of gastric perforations with an omental patch and clips. Gastrointest Endose 54: 500–504
- Lagoo SA (2002) Laparoscopic repair for perforated peptic ulcer. Ann Surg 235: 320–321
- Lau WY, Leung KL, Kwong KH, Davey IC, Robertson C, Dawson JJ, Chung SCS, Li AKC (1996) A randomized study comparing laparoscopic versus open repair of perforated peptic ulcer using suture or sutureless technique. Ann Surg 224: 131–138
- Pescatore P, Halkic N, Calmes JM, Blum A, Gillet (1998) Combined laparoscopic-endoscopic method using an omental plug for therapy of gastroduodenal uleer perforation. Gastrointest Endose 48: 411–414
- Siu WT, Leong HT, Law BKB, Chau CH, Li ACN, Fung KH, Tai YP, Li MKW (2002) Laparoscopic repair for perforated peptic uleer: a randomized controlled trial. Ann Surg 235: 313–319
- Sø JBY, Kum CK, Fernandes ML, Goh P (1996) Comparison between laparoscopic and conventional omental patch repair for perforated duodenal ulcer. Surg Endose 10: 1060–1063
- Wemyss-Holden S, White SA, Robertson G, Lloyd D (2002) Color eoding of sutures in laparoscopic perforated duodenal ulcer: a new concept. Surg Laparose Endosc Percutan Tech 2002 12: 177–179

Surg Endosc (2006) 20: 794–800 DOI: 10.1007/s00464-005-0086-9

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

# Elucidating the relationship between cardiac preload and renal perfusion under pneumoperitoneum

Simon Bergman,<sup>1,2</sup> Arni Nutting,<sup>4</sup> Liane S. Feldman,<sup>1,2</sup> Melina C. Vassiliou,<sup>1,2</sup> Christopher G. Andrew,<sup>1,2</sup> Sebastian Denyttenaere,<sup>1,2</sup> Debbie Woo,<sup>2</sup> Franco Carli,<sup>3</sup> Luc Jutras,<sup>4</sup> Jean Buthieu,<sup>5</sup> Donna D. Stanbridge,<sup>1</sup> Gerald M. Fried<sup>1,2</sup>

<sup>1</sup> Steinberg-Bernstein Centre for Minimally Invasive Surgery, McGill University, Montreal, QC, Canada

<sup>2</sup> Department of Surgery, McGill University, Montreal, QC, Canada

<sup>3</sup> Department of Anesthesia, McGill University, Montreal, QC, Canada

<sup>4</sup> Department of Pediatrics, McGill University, Montreal, QC, Canada

<sup>5</sup> Department of Medicine, McGill University, Montrcal, QC, Canada

Received: 17 February 2005/Accepted: 27 September 2005/Online publication: 16 March 2006

## Abstract

*Introduction:* Pneumoperitoneum is associated with a well-described decrease in renal blood flow, but it remains unclear whether a decrease in cardiac preload is responsible. Our aim was to characterize the relationship between cardiac preload and renal perfusion during pneumoperitoneum.

Methods: Eleven pigs were submitted to three 30 minute study periods: 1) Baseline (n=11): no interventions, 2) Pneumoperitoneum (n=11): 12 mmHg CO2 pneumoperitoneum, 3) Preload Reduction: pneumoperitoneum and nitroglycerin infusion (n=8); or pneumoperitoneum and hemorrhage to a mean arterial pressure (MAP) of 40 mmHg (n=3). Echocardiographic measurements of left ventricular end-diastolic diameter (LVEDD) were used as an index of preload. Renal cortical perfusion (RCP) was measured using laser doppler flowmetry.

*Results:* LVEDD decreased from  $4.2 \pm 0.5$  to  $4.1 \pm 0.6$  cm (p = 0.02) with pneumoperitoneum and then to  $4.0 \pm 0.5$  cm (p = 0.03) with the addition of nitroglycerin. There was no statistically significant change in RCP with pneumoperitoneum ( $33.5 \pm 8.4$  to  $28.5 \pm 8.4$  ml/min/100g tissue, p = 0.2), but it decreased to  $18.5 \pm 11.3$  ml/min/100g tissue (p = 0.001) with the addition of nitroglycerin. The correlation between RCP and LVEDD was weak (0.35, p = 0.003), whereas correlation between RCP and MAP was superior (R = 0.59, p < 0.0001).

*Conclusion:* While decreasing preload under extreme lab conditions also decreases RCP, simply creating a pneumoperitoneum of 12 mmHg does not. The decrease in renal blood flow associated with pneumoperitoneum is likely not solely a function of preload.

Key words: Renal (kidneys) — Pneumoperitoneum — Cardiac preload

The elevated intraabdominal pressure needed to create a laparoscopic workspace causes a varying degree of physiologic impairment unique to the field of minimally invasive surgery. Because this approach has become increasingly complex and currently is performed for patients with diminished cardiorespiratory reserve and renal disease, it is essential that we understand and minimize this impairment.

Multiple animal and human studies have well described the classic systemic hemodynamic effects of increased intraabdominal pressure: tachycardia, increased arterial blood pressure, peripheral vasoconstriction, and decreased stroke volume and cardiac output [19, 26]. It is thought that by virtue of the Frank–Starling mechanism, a decrease in venous return to the heart caused by inferior vena cava compression is responsible for a fall in cardiac preload and subsequent reduction in cardiac output [30]. Oliguria also is a common clinical renal manifestation of pneumoperitoneum. As shown in several animal studies, glomerular filtration rate is decreased in association with a drop in renal blood flow and renal cortical perfusion (RCP) [4], particularly when intraabdominal pressures exceed 15 mmHg.

Many factors are thought to contribute to the modulation of renal blood flow under pneumoperitoneum, including direct compression of the renal vessels and parenchyma, neurohumoral factors, and changes in systemic hemodynamics [9, 29]. According to one theory that has had an impact on clinical practice, the decrease in cardiac preload caused by pneumoperitoneum induces a relative hypovolemic state and subsequent

Correspondence to: Liane S. Feldman

renal hypoperfusion, even if there is no drop in cardiac index. This view is supported by a porcine study showing that aggressive intravenous hydration reverses the impaired renal blood flow caused by a pneumoperitoneum of 15 mmHg [23]. Accordingly, vigorous intravenous hydration during laparoscopic live donor nephrectomy has been recommended by many authors because of its presumed ability to maintain preload and thereby maintain renal perfusion [11]. However, in a pilot study with humans undergoing laparoscopic donor nephrectomy, we previously found that preload, as estimated by esophageal Doppler measurement of flow time, did not in fact decrease with a carbon dioxide ( $CO_2$ ) pneumoperitoneum of 12 to 15 mmHg [10].

Because of the difficulty measuring cardiac preload, the exact relationship between preload and renal blood flow during pncumoperitoneum has never been studied directly and remains unclear. There exists no gold standard for measuring cardiac preload. Pulmonary eapillary wedge pressure is a reliable index only when cardiac compliance is normal and not subject to transmitted abdominal pressures to the chest, as is the case during laparoscopy [28]. In the laboratory setting, preload may be assessed with echocardiography by measurement of the left ventricular end-diastolic diameter. Findings have shown echographic cardiac chamber measurements to be reliable measures of blood loss, suggesting that this technique is a good index of preload [6].

This study aimed to elucidate, in an animal model, the relationship between renal perfusion and cardiac preload, as measured by left ventricular end-diastolic diameter during pneumoperitoneum alone and during an experimentally induced fall in preload in this same context. We hypothesized that although renal blood flow decreases with pneumoperitoneum, and further with the addition of nitroglycerin, there is not a significant relationship between it and cardiac preload.

# Methods

#### Anesthesia and the animal model

For this study, 11 female Landraee pigs (weight, 20–30 kg) were fasted overnight with unrestrieted access to free water. Throughout the experiment, the animals received normal saline at a maintenance rate of 20 ml/h via an auricular vein. The induction protocol consisted of a BAG premix (butorphanol 0.25 mg/kg, acepromazine 0.18 mg/kg, glycopyrrolate 0.009 mg/kg), followed by ketamine 14 mg/kg and variable doses of sodium pentothal 25 mg/kg to maintain adequate sedation and analgesia. During the experimental period, anesthesia was maintained with 2% to 3% isoflurane and paralysis with paneuronium 2 mg, administered every 30 min.

The animals were intuhated and ventilated with a Newffield ventilator (Penlon, Abingdon, UK), set at a fraction of inspired oxygen (FiO<sub>2</sub>) of 40%, which was increased as needed to keep the oxygen saturation above 95%. Fresh gas flow was set at 0.3 to 0.4 l/s, with an inspiration-to-expiration ratio of 1:2. These settings were modified depending on end-tidal carbon dioxide levels using a sensor attached to the endotracheal tube at the mouth (Nellcor, Tyeo Healtheare, Manslield, MA, USA) in an attempt to avoid hyperearbia. To measure the acid–base status, arterial blood gas samples were drawn once hefore the start of the experiment and every 30 min thereafter. A reetal thermometer was inserted, and normothermia was maintained with the use of a heating lamp and warming pad.

This animal model is a modified version of that described hy Chiu et al. [4] for the study of renal hemodynamics during laparoscopy. The animals were placed in the left lateral decubitus position, and a neck eut-down was performed exposing the common earotid artery and external jugular vein, with care taken not to injure the vagus nerve. An 18-gauge vascular eatheter was inserted in the earotid artery and connected to a transducer to measure heart rate and mean arterial pressure (MAP). A Swan-Ganz pulmonary artery catheter (Biosensors International, Singapore) was introduced via an introducer catheter (Arrow International, Reading, PA, USA) placed in the jugular vein and positioned in the pulmonary artery under fluoroscopic guidance (Stenoscop; GE HealthCare Technologies, Waukesha, WI, USA), for eardiae output, central venous pressure (CVP), and pulmonary capillary wedge pressure (PCWP) measurements. A Veress needle was used to ereate a CO<sub>2</sub> pneumoperitoneum of 12 mmHg. Gerota's fascia of the right kidney was ineised laparoscopically, and a superlicial laser Doppler flow prohe (Transonie, Ithaea, NY, USA) was positioned over the renal parenehyma.

#### Experimental protocol

The animals underwent three experimental phases of 30 min each. All 11 animals underwent the baseline phase followed by the pneumoperitoneum phase. Finally, in the preload reduction phase, preload was decreased using either a nitroglycerin infusion (n = 8) or controlled hemorrhage (n = 3).

Phase 1 (baseline): For 30 min, the animal did not have a pneumoperitoneum and did not undergo any interventions.

Phase 2 (pneumoperitoneum): A  $CO_2$  pneumoperitoneum of 12 mmHg was generated and maintained for 30 min.

Phase 3 (preload reduction): The animals in the nitro group remained under pneumoperitoneum and received an intravenous nitroglycerin drip started at 5 mcg/min, with 5-mcg incremental increases every 5 min for 30 min. Several human studies have shown that nitroglycerin reduces preload with hemodynamic effects similar to hemorrhage or vena caval occlusion [1]. To demonstrate that infusion of nitroglycerin in this pneumoperitoneum model has effects similar to those of acute blood loss, a more clinically relevant preload reducer, three animals under the same pneumoperitoneum were hled down to a mean arterial pressure of 40 mmHg and then studied over 30 min [31].

Hemodynamic measurements were taken every 10 min throughout the three experimental phases. Using a Spacelabs (Issaquah, WA, USA) anesthesia monitor, heart rate, MAP, CVP, PCWP, and eardiae output (thermodilution technique that averages three measurements taken at end-expiration) were recorded. Systemic vascular resistance was calculated using the following formula: (MAP-CVP)  $\times$ 80/eardiae output. Left ventricular end-diastolie diameter (LVEDD), recorded three times and averaged every 10 min, was measured hy M-mode in the parasternal short-axis plane using transthoracic echocardiography equipment with a 3-Mhz transducer (Philips Sonos 1500, Andover, MA, USA). Finally, RCP was measured using a laser Doppler flowmeter (Transonie, Ithaca, NY, USA), which measures perfusion in the underlying 1-mm<sup>3</sup> volume of renal parenchyma. Findings have shown that RCP has an excellent correlation with direct flow measurements from the renal artery and is easier to place [4].

#### Statistical analysis

Data are expressed as means  $\pm$  standard deviation. All hemodynamic measurements were recorded every 10 min and averaged over each experimental phase. They were analyzed using the paired Student's *t*test. Univariate linear regression analysis was used to assess individual associations between RCP and the other hemodynamic parameters. A *p* value less than 5% was set as the criterion for statistical significance. Analysis was performed using GB-Stat 6.5.4 (Dynamic Microsystems Inc., Silver Spring, MD, USA).

We estimated the size of the sample assuming that insulflation of the abdomen to a pressure of 12 mmHg would decrease renal perfusion by 30%, from 50  $\pm$  18 to 35 ml/min/100 g tissue, according to the

results of Chiu et al. [4] To demonstrate this difference at a 5% level of significance would require 11 pigs for 80% power.

#### Results

The mean weight of the animals was  $25.5 \pm 2$  kg. Two of the animals were accidentally given one supplemental dose of thiopental at the beginning of the first experimental phase. The hemodynamic changes that occurred arc summarized in Fig. 1. Heart rate, MAP, CVP, and PCWP all increased, respectively, by 5.4% (p = 0.008), 8.6% (p < 0.001), 52% (p <0.001), and 14.2% (p = 0.03) during the pneumoperitoneum phase. During the nitroglycerin infusion, MAP, CVP, and PCWP fell, respectively, by 16.3% (p < 0.001), 30.0% (p < 0.001), and 21.8% (p = 0.03), whereas heart rate increased by 11% (p < 0.001) (Fig. 1a to 1d). There was no statistically significant change in cardiac output during the pneumoperitoneum phase (p = 0.2) or during the nitroglycerin infusion (p = 0.2) (Fig. 1e). Systemic vascular resistance (SVR) did not increase significantly with pneumoperitoneum, but decreased from  $1,132 \pm 312$  to 961  $\pm$  310 dynes  $\cdot$  sec/cm<sup>5</sup> with the addition of the nitroglycerin infusion (p < 0.0001). During pneumoperitoneum, LVEDD decreased by 2.4% (p = 0.02), and during the nitroglycerin infusion, it decreased by 5.6% (p = 0.04) (Fig. 1f). Finally, there was no significant change in RCP from baseline to the pneumoperitoneum phase (p = 0.2), but it decreased by 33.8% (p < 0.001) with the nitroglycerin infusion (Fig. 1g). In the preload reduction phase, only the trends in heart rate, cardiac output, and SVR differed between the nitroglycerin and the hemorrhage groups. The animals in the latter group showed a significant decrease in heart rate (p = 0.01) and cardiac output (p = 0.003), but SVR was unchanged.

There was a poor correlation between RCP and LVEDD (R = 0.35; p = 0.003), as shown on the scatterplot in Fig. 2. The correlation between RCP and cardiac output also was poor (R = 0.37; p = 0.001). Renal cortical perfusion correlated better with PCWP (R = 0.48; p < 0.0001) and best with MAP (R = 0.62; p < 0.0001). Finally, the correlation between RCP and CVP was not significant. When the animals were stratified into those with a baseline LVEDD equal to or lower than the median baseline greater than 4.1 cm (n = 4) and those with a baseline greater than 4.1 cm (n = 4), the former group showed a significant drop in RCP, whereas the latter group did not (Fig. 3).

The partial pressure of CO<sub>2</sub> (PaCO<sub>2</sub>) was 59  $\pm$  12 and the arterial pH was 7.33  $\pm$  0.08 at the start of the baseline period and had remained constant, respectively, at 57  $\pm$  15 (p = 0.4) and 7.35  $\pm$  0.08 (p = 0.1) by the end of the period. After 30 min of pneumoperitoneum, the PaCO<sub>2</sub> had increased to 67  $\pm$  11 (p = 0.01), and the pH had decreased to 7.28  $\pm$  0.09 (p = 0.01). Both remained constant by the end of the preload reduction phase, with the PaCO<sub>2</sub> at 68  $\pm$  10 (p = 0.8) and the arterial pH at 7.29  $\pm$  0.09 (p = 0.6).

### Discussion

To our knowledge, this is the first study to directly assess the physiologic effects of pneumoperitoneum on both cardiac preload and renal perfusion. As expected, MAP, CVP and PCWP all increased with a CO2 pneumoperitoneum of 12 mmHg. Despite a very slight decrease in LVEDD, used as an index of preload, there was no significant change in cardiac output or RCP overall during pneumoperitoneum. When stratified according to baseline preload, however, those animals with underfilling at baseline demonstrated decreased RCP under pneumoperitoneum. When preload was then further reduced, using nitroglycerin administration or hemorrhage, LVEDD dropped further and renal perfusion decreased significantly. After acute hemorrhage, the direction of change in all the parameters except cardiac output followed that of the animals that had received nitroglycerin. Although nitroglycerin is primarily a preload reducer, the expected drop in cardiac output may have been offset by nitroglycerin's afterload-reducing properties, as evidenced by the decrease in SVR during the infusion. This had been noted previously in animals with elevated baseline arterial elastance or abnormal baseline ventricular function [13]. In this study, SVR decreased significantly during the nitroglycerin infusion. Systemic vascular resistance, known to increase under pneumoperitoneum, is thought to contribute to the decrease in cardiac output that may be seen with high-pressure pneumoperitoncum [29].

The impact of pneumoperitoneum on preload and its relationship with cardiac output is complex and not yet fully elucidated. The classical response to pneumoperitoneum is believed to be a decrease in venous return secondary to inferior vena cava compression. This is based on clinical studies demonstrating a drop in cardiac output and a hemodynamic response to pneumoperitoneum similar to that of hypovolemic shock [26]. However, several other studies have demonstrated no change in cardiac output [21], and have even shown increases in cardiac output after pneumoperitoneum [16].

The study examining the impact that pneumoperitoneum has on preload is complicated by the fact that the usual clinical measures of ventricular filling, such as PCWP and CVP, are unreliable during pneumoperitoneum. As the elevated abdominal pressures are transmitted to the chest, pressure measurements taken in the chest risc regardless of true volume status [7, 28]. Transthoracic echocardiography has been used experimentally to evaluate preload by measurement of left ventricular end-diastolic diameter, area, or volume. The only limitation to echocardiographic preload assessment during pneumoperitoneum is the change in cardiac geometry that may occur with upward displacement of the diaphragm [24]. Larsen et al. [21] found LVEDD to be increased with the onset of a 12-mmHg pneumoperitoneum in adults undergoing laparoscopic cholecystectomy. In the same patient population, Gannedahl et al. [12] also found left ventricular end-diastolic area to be increased. In healthy women undergoing exploratory laparoscopic surgery of the lower abdomen, left ven-



Fig. 1. Cardiac and renal hemodynamics at baseline, with pneumoperitoneum and with preload reduction. Cardiac and renal hemodynamics at baseline, with pneumoperitononeum and with preload (a: heart rate, b: mean arterial pressure, c: central venous pressure, d: pulmonary capillary wedge pressure, e: cardiac output, f: left ven-

trieular end-diastolic area was increased at a pneumoperitoneum of 10 mmHg, and further increased when the intraabdominal pressure was raised to 15 mmHg [34].

In contrast, Cunningham et al. [5] and Dorsay et al. [8] reported no change in left ventricular end-diastolic area during pneumoperitoneum in healthy patients [5,

tricular end-diameter, g: renal perfusion. Solid line: all animals, dotted line: nitroglycerin group, dashed line: hemorrhage group. Preload reduction (nitro group / hemorrhage group).  $*p \le 0.05$  as compared with previous value.

8], and one study found no change in atrial natiuretic peptide released in response to increased atrial pressure in healthy patients undergoing laparoscopic surgery [18]. The only study to demonstrate a reduction in left ventricular end-diastolic dimensions also was the only study to include patients with an American Society of Anesthesia (ASA) index above 2 [36].



**Fig. 2.** Correlation between renal cortical perfusion and left ventricular end-diastolic diameter.

left ventricular end-diastolic diameter. Left ventricular end-diastolic diameter (LVEDD)  $\leq 4.1$ cm (dashed line), LVEDD > 4.1 cm (solid line); n = 4 for both groups. \*p < 0.05 within group;  $\dagger p < 0.05$  between groups.

Fig. 3. Renal cortical perfusion at baseline and under pneumoperitoneum, stratified according to

The findings in these studies seem to suggest that in healthy individuals, preload is in fact increased with induction of pneumoperitoneum, perhaps because of splanchnic blood flow redistribution to the chest. In the current animal study, LVEDD decreased only slightly with the onset of pneumoperitoneum, to a degree of questionable clinical significance, and cardiac output did not decline.

Under a  $CO_2$  pneumoperitoneum of 12 mmHg, we found no significant decline in RCP, which is consistent with what has been reported by others for that pneumoperitoneum level [20, 22]. A pneumoperitoneum of at least 15 mmHg was necessary in most studies to demonstrate a decrease in renal blood flow [4, 23]. On the other hand, during the nitroglycerin infusion, RCP diminished by one-third of its previous value, but LVEDD decreased by only about 6%. Hence, the overall correlation between LVEDD and RCP was poor. The correlation of RCP with PCWP and CVP also was poor, which was expected given the limitations of these measures during pneumoperitoneum.

On the other hand, MAP and cardiac output seemed to be moderately well associated with the changes in renal perfusion. This is in keeping with the findings of Chiu et al. [2], who showed that inferior vena cava compression in a dog model led to a significant decrease in cardiac output, but only a mild decrease in renal blood flow. Left ventricle compression caused the same decrement in cardiac output and renal blood flow. Hence, if a drop in preload is postulated to play a significant role in the decrease in renal perfusion under pneumoperitoneum, it would be mainly through its ability to decrease cardiac output. A decrease in cardiac output, however, is clearly not the principal factor involved in renal blood flow modulation. In fact, findings have shown that normalizing cardiac output with plasma expanders at a pneumoperitoneum of 20 mmHg failed to improve renal blood flow and glomerular filtration rate [15]. In addition, several studies have demonstrated decreases in renal perfusion and function without a change in cardiac output [3, 23].

Despite the poor correlation between RCP and LVEDD, the animals with higher baseline preloads did

not drop their renal perfusion, as compared with those that had lower baseline preload values. London et al. [23] showed that intravascular volume expansion in a pig model reversed the drop in renal blood flow that occurred with a 15-mmHg  $CO_2$  pneumoperitoneum, although creatinine elearanee was not improved. The solution may be goal-directed fluid optimization during pneumoperitoneum, taking baseline fluid status into account, as long as clinically relevant, reliable, and easy-to-measure parameters are sought [35].

Another factor for explaining the changes in renal hemodynamics and function seen under pneumoperitoneum is parenchymal compression, as demonstrated by Razvi et al. [32]. These authors inflated a 15-mmHg pressure euff around one kidney and showed a decrease in renal blood flow, glomerular filtration rate, and urine output similar to that seen with pneumoperitoneum. Renal vein compression by increased intraabdominal pressure also may contribute to renal ischemia by elevating back pressure into the renal glomeruli [25]. Finally, neurohormonal effects are being studied as potential factors contributing to decreased renal blood flow during pneumoperitoneum [14, 17, 27].

Some limitations to this study must be addressed. First, respiratory gas exchange was significantly affected during this study. Although oxygen saturation was easily maintained above 95% in all the animals, ventilation was more problematie. The animals were hypercarbic and acidotic at baseline, most likely because of suboptimal anesthetic management. Despite increasing minute ventilation during the pneumoperitoneum and nitro/bleed phases, the animals reached a mean pH of 7.28 and a PaCO<sub>2</sub> of 68 mmHg. Hypercarbia is known to stimulate the sympathetic nervous system, but acidosis also may depress cardiac function. This may be a confounding factor in the interpretation of the physiologic ehanges that occurred from baseline to the pneumoperitoneum phase, although significant changes in eardiae output and RCP were not seen with pneumoperitoneum alone. In addition, when argon and helium are used for insufflation, although hypercarbia and acidosis are minimized, the systemic hemodynamic effects are unchanged. These data suggest that it is the elevated intraabdominal pressure rather than the aeid-base imbalance that contributes most to the physiologic ehanges [20].

Finally, the most important limitation of this study lies in the choice of the model. Although pigs are currently the best and most commonly used species for simulating human physiology, there are nevertheless significant interspecies differences. Pigs seem to have a higher basal adrenergic drive, and their hemodynamics are highly sensitive to fluid status, analgesia, and acid– base status [33]. Furthermore, considering that baseline health status is difficult if not impossible to ascertain and control for, pigs, depending on how they are bred, fed, and raised, will be more susceptible to congenital cardiac abnormalities and to illness. This may explain the degree of baseline physiologic variability noted in this study. As an example, although the pigs all were treated the same in terms of access to fluids in the preoperative period, baseline CVP varied 4 to 15 mmHg, and PCWP varied 8 to 16 mmHg. This has led some authors to state that changes observed in pig models eannot reliably predict changes occurring in humans. Therefore, these data, as in all animal studies, should be interpreted with eaution and simply considered a means for hypothesis generation.

In conclusion, we found that whereas decreasing preload under extreme laboratory conditions also decreased renal perfusion, simply creating a pneumoperitoneum to 12 mmHg did not. Because a pneumoperitoneum of 12 mmHg decreased renal blood flow only in animals with a low baseline preload, vigorous intravenous hydration, as recommended during laparoscopie donor nephrectomy, may benefit only patients who are volume depleted at baseline.

#### References

- Atkins BZ, Silvestry SC, Davis JW, Kisslo JA, Glower DD, Jr (1999) Means for load variation during echoeardiographic assessment of the Frank-Starling relationship. J Am Soc of Echoeardiogr 12: 792–800
- Chiu AW, Azadzoi KM, Hatzichristou DG, Siroky MB, Krane RJ, Babayan RK (1994) Effects of intra-abdominal pressure on renal tissue perfusion during laparoscopy. J Endourol 8: 99– 103
- Chiu AW, Chang LS, Birkett DH, Bahayan RK (1995) The impact of pneumoperitoneum, pneumoretroperitoneum, and gasless laparoscopy on the systemic and renal hemodynamics. J Am Coll Surg 181: 397–406
- Chiu AW, Chang LS, Birkett DH, Babayan RK (1996) A porcine model for renal hemodynamic study during laparoscopy. J Surg Res 60: 61–68
- Cunningham AJ, Turner J, Rosenbaum S, Rafferty T (1993) Transoesophageal echocardiographic assessment of haemodynamic function during laparoscopic cholecystectomy. Br J Anaesth 70: 621–625
- D'Angelo AJ, Kline RG, Chen MH, Halpern VJ, Cohen JR (1997) Utility of transcsophageal echocardiography and pulmonary artery catheterization during laparoscopic assisted ahdominal aortic ancurysm repair. Surg Endose 11: 1099–1101
- Dalibon N, Schlumberger S, Saada M, Fischler M, Riou B (1999) Haemodynamic assessment of hypovolaemia under general anaesthesia in pigs submitted to graded haemorrhage and retransfusion. Br J Anaesth 82: 97–103
- Dorsay DA, Greene FL, Baysinger CL (1995) Hemodynamic changes during laparoscopic cholecystectomy monitored with transesophageal echocardiography. Surg Endose 9: 128–133
- Dunn MD, McDougall EM (2000) Renal physiology. Laparoscopic considerations. Urol Clin North Am 27: 609–614
- Feldman LS, Anidjar M, Metrakos P, Stanhridge D, Fried GM, Carli F (2004) Optimization of cardiac preload during laparoscopic donor nephrectomy: A preliminary study of central venous pressure versus esophageal doppler monitoring. Surg Endose 18: 412–416
- Flowers JL, Jacobs S, Cho E, Morton A, Rosenberger WF, Evans D, Imbembo AL, Bartlett ST (1997) Comparison of open and laparoscopic live donor nephrectomy. Ann Surg 226: 483–489
- Gannedahl P, Odcherg S, Brodin LA, Sollevi A (1996) Effects of posture and pncumoperitoneum during anaesthesia on the indices of left ventricular filling. Acta Anaesthesiol Seand 40: 160–166
- Haber HL, Simek CL, Bergin JD, Sadun A, Gimple LW, Powers ER, Feldman MD (1993) Bolus intravenous nitroglycerin predominantly reduces afterload in patients with excessive arterial elastance. J Am Coll Cardiol 22: 251–257
- Hamilton BD, Chow GK, Inman SR, Stowe NT, Winfield HN (1998) Increased intra-abdominal pressure during pneumoperitoneum stimulates endothelin release in a canine model. J Endourol 12: 193–197

- Harman PK, Kron IL, McLachlan HD, Freedlender AE, Nolan SP (1982) Elevated intra-abdominal pressure and renal function. Ann Surg 196: 594–597
- Hashimoto S, Hashikura Y, Munakata Y, Kawasaki S, Makuuchi M, Hayashi K, Yanagisawa K, Numata M (1993) Changes in the cardiovascular and respiratory systems during laparoscopic cholecystectomy. J Laparoendosc Surg 3: 535–539
- Hazebroek EJ, de Vos tot Nederveen Cappel R, Gommers D, van Gelder T, Weimar W, Steyerherg EW, Bonjer HJ, IJzermans JNM (2002) Antidiuretic hormone release during laparoscopic donor nephrectomy. Arch Surg 137: 600–604
- Hirvonen EA, Nuutinen LS, Vuolteenaho O (1997) Hormonal responses and cardiac filling pressures in head-up or head-down position and pneumoperitoneum in patients undergoing operative laparoscopy. Br J Anaesth 78: 128–133
- Joris JL, Noirot DP, Legrand MJ, Jacquet NJ, Lamy ML (1993) Hemodynamic changes during laparoscopic cholecystectomy. Anesth Analg 76: 1067–1071
- Junghans T, Bohm B, Grundel K, Schwenk W, Muller JM (1997) Does pneumoperitoneum with different gases, body positions, and intraperitoneal pressures influence renal and hepatic hlood flow?. Surgery 121: 206–211
- Larsen JF, Svendsen FM, Pedersen V (2004) Randomized clinical trial of the effect of pneumoperitoneum on cardiac function and haemodynamics during laparoscopic cholecystectomy. Br J Surg 91: 848–854
- Lindberg F, Bergqvist D, Bjorck M, Rasmussen I (2003) Renal hemodynamics during carbon dioxide pneumoperitoneum: An experimental study in pigs. Surg Endosc 17: 480–484
- London ET, Ho HS, Ncuhaus AM, Wolfe BM, Rudich SM, Perez RV (2000) Effect of intravascular volume expansion on renal function during prolonged CO<sub>2</sub> pneumoperitoneum. Ann Surg 231: 195–201
- Marathe US, Lilly RE, Silvestry SC, Schauer PR, Davis JW, Pappas TN, Glower DD (1996) Alterations in hemodynamics and left ventricular contractility during carhon dioxide pncumoperitoneum. Surg Endosc 10: 974–978
- McDougall EM, Monk TG, Wolf JS, Jr., Hicks M, Clayman RV, Gardner S, Humphrey PA, Sharp T, Martin K (1996) The effect of prolonged pneumoperitoncum on renal function in an animal model. J Am Coll Surg 182: 317–328

- McLaughlin JG, Scheeres DE, Dcan RJ, Bonnell BW (1995) The adverse hemodynamic effects of laparoscopic cholecystectomy. Surg Endosc 9: 121–124
- Mikami O, Fujise K, Matsumoto S, Shingu K, Ashida M, Matsuda T (1998) High intra-ahdominal pressure increases plasma catecholamine concentrations during pneumoperitoneum for laparoscopic procedures. Arch Surg 133: 39–43
- Myre K, Buanes T, Smith G, Stokland O (1997) Simultaneous hemodynamic and echocardiographic changes during abdominal gas insufflation. Surg Laparosc Endosc 7: 415–419
- O'Malley C, Cunningham AJ (2001) Physiologic changes during laparoscopy. Anesthesiol Clin North Am 19: 1–19
- Ortega AE, Richman MF, Hernandez M, Pcters JH, Anthone GJ, Azen S, Beart RW, Jr (1996) Inferior vena caval blood flow and cardiac hemodynamics during carhon dioxide pneumoperitoneum. Surg Endosc 10: 920–924
- Radovancevic B, Eichstaedt HC, Tamez D, Patel V, Eya K, Nolden LK, Byler D, Cohen D, Frazier OH (2003) Prolonged controlled hemorrhagic shock in a swine model: Is there a role for mechanical circulatory assistance? ASAIO J 49: 721–726
   Razvi HA, Fields D, Vargas JC, Vaughan ED, Jr., Vukasin A,
- Razvi HA, Fields D, Vargas JC, Vaughan ED, Jr., Vukasin A, Sosa RE (1996) Oliguria during laparoscopic surgery: Evidence for direct renal parenchymal compression as an etiologic factor. J Endourol 10: 1–4
- 33. Richard L, Bures E, Lacoste L, Declerck A, Carretier M, Debaenc B, Fusciardi J (2002) Pigs are not a reliable experimental model in the study of the haemodynamic and respiratory effects of  $CO_2$  pneumoperitoneum. Acta Anaesthesiol Scand 46: 74–79
- 34. Rist M, Hemmerling TM, Rauh R, Siebzchnruhl E, Jacobi KE (2001) Influence of pneumoperitoneum and patient positioning on preload and splanchnic hlood volume in laparoscopic surgery of the lower abdomen. J Clin Anesth 13: 244–249
- Sinclair S, Singer M (1997) Intraoperative intravascular volume optimisation and length of hospital stay after repair of proximal femoral fracture: Randomised control trial. Br Med J 315: 909– 912
- Zuckerman R, Gold M, Jenkins P, Rauscher LA, Jones M, Heneghan S (2001) The effects of pneumoperitoneum and patient position on hemodynamics during laparoscopic cholecystectomy. Surg Endose 15: 562–565
Surg Endosc (2006) 20: 801–805 DOI: 10.1007/s00464-005-0479-9

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

### Do all patients with abnormal intraoperative cholangiogram merit endoscopic retrograde cholangiopancreatography?

S. Varadarajulu,<sup>1</sup> M. A. Eloubeidi,<sup>1</sup> C. M. Wilcox,<sup>1</sup> R. H. Hawes,<sup>2</sup> P. B. Cotton<sup>2</sup>

<sup>1</sup> Division of Gastroenterology-Hepatology, University of Alabama at Birmingham, 410 Lyons Harrison Research Building,

1530 3rd Avenue South, Birmingham, AL 35294-0007, USA

<sup>2</sup> Digestive Disease Center, Medical University of South Carolina, Charleston, SC, USA

Received: 4 July 2005/Accepted: 27 December 2005/Online publication: 16 March 2006

#### Abstract

*Background:* Endoscopic retrograde cholangiopancreatography (ERCP) is commonly used for postoperative evaluation of an abnormal intraoperative cholangiogram (IOC). Although a normal IOC is very suggestive of a disease-free common bile duct (CBD), abnormal studies are associated with high false-positive rates. This study aimed to identify a subset of patients with abnormal IOC who would benefit from a postoperative ERCP.

*Methods:* This prospective study investigated 51 patients with abnormal IOC at laparoscopic eholeeysteetomy who underwent postoperative ERCP at two tertiary referral centers over a 3-year period. Univariate and multivariate logistic regression analyses were performed to determine predictors of CBD stones at postoperative ERCP.

*Results:* For all 51 patients, ERCP was successful. The ERCP showed CBD stones in 33 eases (64.7%), and normal results in 18 eases (35.2%). On univariate analysis, abnormal liver function tests (p < 0.0001) as well as 1OC findings of a large CBD stone (p = 0.03), multiple stones (p = 0.01), and a dilated CBD (p = 0.07) predicted the presence of retained stones at postoperative ERCP. However, on multivariable analysis, only abnormal liver function tests correlated with the presence of CBD stones (p < 0.0001).

*Conclusions:* One-third of patients with an abnormal IOC have a normal postoperative ERCP. Elevated liver function tests can help to identify patients who merit further evaluation by ERCP. The use of less invasive methods such as endoscopic ultrasound or magnetic resonance eholangiopanereatography should be considered for patients with normal liver function tests to minimize unnecessary ERCPs.

Key words: Abnormal intraoperative cholangiogram — CBD — Common bile duct — Endoscopie retrograde eholangiopanereatography — ERCP — IOC — Predietors

Laparoscopic cholecystectomy has replaced open cholecystectomy as the procedure of choice for the vast majority of individuals with symptomatic cholelithiasis. The advantages of the laparoscopic technique include reduced pain, early discharge from the hospital, rapid return to normal activity, and improved cosmetic appearance [26]. The increasing popularity of this minimally invasive procedure has resulted in an increased demand for endoscopic retrograde cholangiopancreatography (ERCP) for CBD stone extraction [21].

Predominantly secondary in origin, most CBD stones are thought to have migrated through the cystic duct from their primary site of origin in the gallbladder. The incidence of CBD stones detected at the time of cholecystectomy ranges from 8% to 15% [1, 4, 8, 14, 15]. Intraoperative chlangiography (IOC) is performed during laparoscopic cholecystectomy to evaluate the CBD for stones using contrast injection through a catheter introduced into the cystic duct. Retained CBD stones then are extracted by postoperative ERCP or laparoscopic CBD exploration.

Laparoscopic CBD exploration is not widely available because the procedure requires special equipment, advanced training, and significant expertise on the part of the surgeon [33]. On the other hand, because ERCP is readily available at most hospitals, it is the method most commonly used to evaluate abnormal IOC findings. The disadvantages of this approach are twofold. First, IOC is associated with a false-positive rate ranging from 2% to 60% [16, 20, 22, 33]. Second, ERCP is an invasive

Correspondence to : S. Varadarajulu

procedure associated with a complication rate of 2.5% to 11% [10, 25].

The purpose of this study was to identify a subset of patients with abnormal IOC who are most likely to benefit from a postoperative ERCP: This includes those patients more likely to have residual stones in the CBD.

#### Materials and methods

This prospective study investigated all patients referred for postoperative ERCP after laparoscopic cholecystectomy for evaluation of abnormal fluorocholangiography findings at two tertiary referral centers over a 3-year period (2001–2004). The patients were referred for ERCP by gastrointestinal surgeons from within the hospital or from outside facilities.

Before ERCP, patient demographic information and indications for laparoscopic cholecystectomy were documented. A white cell count and a liver function panel were drawn on all patients 1 to 2 h before ERCP. Any deviation above the normal reference range was considered abnormal for both liver function tests (LFTs) and white cell count. The interval between laparoscopic cholecystectomy and presentation for ERCP was documented.

The IOC findings were reviewed before ERCP, and the indications were divided into the following seven categories: (a) nonpassage of contrast into the duodenum, (b) isolated CBD filling defect suggestive of small stoncs ( $\leq 4$  mm), (c) isolated CBD filling defect suggestive of large stones ( $\geq$ 5mm), (d) multiple CBD filling defects suggestive of stones, (e) CBD dilation ( $\geq$ 8mm), (f) nonvisualization of distal CBD, and (g) palpable CBD stones at surgery. A cutoff value of 5 mm was chosen to distinguish between small and large CBD stones [1, 11, 39], and a CBD dilation [31], on the basis of published data.

The endoscopist evaluated the radiographs to determine the size of the CBD stones, the diameter of the CBD, and other IOC findings. A radiologist was consulted when necessary to confirm the interpretation and to correct any fluoroscopic magnification. Patients who presented for ERCP without IOC radiographs and those who had undergone prior endoscopic biliary interventions were excluded from the study. Also, patients with an abnormal IOC but with a diagnosis of other diseases such as pancreatic cancer or cholangiocarcinoma by computed tomography (CT) scan or endoscopic ultrasound (EUS) were excluded from the study.

The ERCPs were performed in the hospital endoscopy suites of the two centers with the patient under conscious sedation induced by combinations of midazolam, meperidine, and droperidol administered by the endoscopist. All the procedures were performed by four attending endoscopists or by gastroenterology fellows under the supervision of attending physicians. The ERCP was performed with a standard "therapeutic" duodenoscope (Olympus TJF-100/130/140/ 160, Olympus America Corporation, Inc., Melville, NY) with an outer diameter of 12.5 mm and an accessory channel diameter of 4.2 mm. Successful ERCP was defined as cannulation of the bile duct along with satisfactory performance of a planned diagnostic study or therapeutic intervention. All CBD stones seen at ERCP were extracted using a retrieval balloon or basket after performance of a biliary sphincterotomy. An occlusion cholangiogram then was performed to ensure clearance of the bilc duct. Absence of CBD stones at ERCP was attributed to spontaneous stone passage or a false-positive IOC. The fluoroscopic findings, therapeutic interventions, and complications, if any, were recorded for each patient.

Patient follow-up assessment was through discussions of the patients at fortnightly pancreaticobiliary meetings with nurse coordinators, who called patients after ERCP, or through reference to outpatient clinic visits. Patients were called 6 months after ERCP for clinical follow-up information.

#### Statistical analysis

Continuous variables were reported as medians and ranges because data were not normally distributed. Continuous variables were com 
 Table 1. Indications for endoscopic retrograde cholangiopancreatography (ERCP) based on intraoperative cholangiogram (IOC) findings

IOC finding <sup>a</sup>	n
Nonpassage of contrast into the duodenum	17
Small stones	17
Large stones	12
Multiple stones	9
Dilated CBD	11
Nonvisualization of distal CBD	3
Palpable CBD stones	1

CBD, common bile duct

<sup>a</sup> Individual patients may have more than one IOC finding

pared using the Wilcoxon rank sum test. Categorical variables were compared using the chi-square test. To determine predictors of CBD stones at postoperative ERCP, univariate and multivariable logistic regression analyses were performed. Both stepwise and best model selections were used, providing similar results. Adjusted odds ratios and their corresponding 95% confidence intervals (CI) were reported. All statistical analyses were performed using the SAS (Cary, NC, USA) software version 6.12. A two-sided type 1 error of 0.05 was used for all statistical tests.

All participating patients provided informed consent. The study was approved by the institutional review board of both centers.

#### Results

The indications for laparoscopic cholecystectomy in the 51 patients (33 women) (mean age, 51.9 years; range, 21-85 years) were symptomatic cholelithiasis in 26 (51%), gallstone pancreatitis in 12 (24%), cholecystitis in 10 (20%), and biliary dyskinesia in 3 (5%) patients. Preprocedural LFTs were abnormal in 32 patients (62.7%), and the white blood cell count level was high in 18 patients (35%). The IOC findings that prompted an ERCP are shown in Table 1. Because IOC was performed at multiple centers, procedural technique for IOC was not standardized. For one patient, the laparoscopic approach was converted to open cholecystectomy: This patient had a large palpable CBD stone, but a bile duct exploration was not undertaken, and he was referred for postoperative ERCP. The median time to ERCP after surgery was 2 days (range, 1-7 days).

For all 51 patients (100%), ERCP was successful. The ERCP showed CBD stones in 33 of the 51 patients (64.7%), and no stones in the remaining patients (Table 2). Whereas CBD stones were seen in 29 of 33 patients (87.8%) with abnormal LFTs, only 3 of 18 patients (16.6%) with normal LFTs had CBD stones (p < 0.0001). Sphincterotomy was performed using a pull-type papillotome in 30 patients and a needle-knife in 3 patients. Temporary pancreatic stents to facilitate biliary access were placed in two of three patients who underwent needle-knife sphincterotomy. Stones were extracted using a balloon for 26 patients and a combination of balloon and retrieval basket for 7 patients. All stones were retrieved in a single ERCP session for all patients. No immediate or late complications were encountered in any patient.

On univariate analysis, abnormal liver LFTs (p < 0.0001) as well as IOC findings of large stones

 Table 2. Comparison of demographics, laparoscopic cholecystectomy indications, laboratory investigations, and intraoperative cholangiogram (IOC) findings in patients with and without common bile duct (CBD) stones at endoscopic retrograde cholangiopancreatography (ERCP) (results of univariate analysis)

Variables	Stones at ERCP $(n = 33)$	No stones at ERCP $(n = 18)$	p Value
Demographics			
Age: years (IQR)	54 (44-71)	41 (31-60)	0.21
Female: <i>n</i> (%)	21 (63.6)	12 (66.7)	0.83
Indications for surgery			
Cholelithiasis	20	6	0.42
Cholecystitis	5	5	1.00
Pancreatitis	7	5	0.61
Biliary dyskinesia	1	2	0.59
Median time to ERCP (days) n (range)	2 (1-5)	2 (1-7)	0.29
Laboratory parameters			
Abnormal LFTs: n (%)	29 (87.87)	3 (16.6)	< 0.0001
AST median (IQR)	74 (43-173)	23 (19-31)	0.001
ALT median (IQR)	90 (55-249)	26 (18-43)	0.0006
ALP median (IQR)	162 (121-214)	88 (71-101)	0.0002
TB median (IQR)	1.9(1.1-3)	1 (0.8 - 1.1)	0.001
DB median (IQR)	1.2(0.7-2)	0.5 (0.2–0.8)	0.001
White cell count: n (range)	8.2 (7-10.4)	7 (6.1–10.1)	0.32
IOC findings: $n (\%)^n$			
1. Nonpassage of contrast to duodenum	9 (27.27)	8 (44.44)	0.17
2. Small stones	8 (24.24)	8 (44.44)	0.1
3. Large stones	11 (33.33)	1 (5.55)	0.03
4. Multiple stones	9 (27.27)	0	0.01
5. Dilated CBD	9 (27.27)	1 (5.55)	0.07
6. Poor visualization of distal CBD	4 (12.1)	1 (5.55)	0.2
7. Palpable CBD stones at surgery	I (3.03)	0	0.47

LFT, liver function test; IQR, interquartile range; AST, aspartate transaminase; ALT, alanine transaminase; ALP, alkaline phosphatase; TB, total bilirubin; DB, direct bilirubin

<sup>a</sup> Individual patients may have more than one IOC finding

(p = 0.03), multiple stones (p = 0.01), and dilated CBD (p = 0.07) predicted residual stones at postoperative ERCP (Table 2). However, on multivariable analysis, only abnormal LFTs predicted retained CBD stones (odds ratio [OR], 19.7; CI, 2.8–134.9; p < 0.0001). Elevation of both transaminases and cholestatic enzymes predicted CBD stones. Patients with abnormal LFTs were 20 times more likely to have stones at ERCP than those with normal LFTs. At the 6-month follow-up assessment, all the patients were doing well without any evidence of recurrent biliary obstruction.

#### Discussion

The optimal evaluation of abnormal IOC findings in this era of laparoscopic cholecystectomy still remains undefined. In this study, one-third of the patients with an abnormal IOC had a normal postoperative ERCP. This is consistent with prior reports describing a low rate of stone recovery at ERCP in the postoperative setting [1, 13, 33].

None of the IOC findings in this study accurately predicted the presence or absence of CBD stones. The reason for the poor predictability of IOC is unclear. It is likely that some CBD stones pass spontaneously and that other IOC findings are false-positive, such as air bubbles mimicking stones or failure of contrast to drain into the duodenum, which possibly is related to papillary stenosis rather than an impacted stone. In this study, ERCP was undertaken for a majority of the patients (>80%) within 48 h of cholecystectomy. The rate for spontaneous migration of stones through the duodenal papilla is unknown. In this study, only 1 of 12 patients with large stones at IOC had a normal ERCP, as compared with 8 of 16 patients who had small stones. This is in agreement with a prior study from France that reported a negative predictive value of 95% for spontaneous migration of large stones at a 1 month follow-up assessment [11]. On the other hand, 28% of small stones had spontaneous migration.

Clinicians encounter three temporal situations in the management of CBD stones: (a) patients for whom the stones are suspected preoperatively, (b) patients for whom the stones are found intraoperatively, and (c) patients whose stones present postoperatively. The role of ERCP in managing postoperative stones is well established [23, 24, 35]. On the contrary, optimal management of suspected preoperative and intraoperative stones remains a challenge.

In the preoperative setting, inability to predict accurately which patients have CBD stones is problematic because when stones are suspected on the basis of clinical, biochemical, and imaging tests, normal prcoperative ERCP results still are obtained for 40% to 70% of patients [2, 3, 5, 17, 28, 40]. The optimal technique for biliary ductal clearance should be based on the local availability of expert endoscopists, surgical expertise in laparoscopic CBD surgery, and the general condition of the patient [9, 32, 41]. In the intraoperative setting, if stones are suspected, laparoscopic extraction can be attempted on the basis of equipment and expertise availability. Although findings have shown intraoperative ERCP to be feasible and successful in this setting, it has not been widely used [24]. The other option is the use of postoperative ERCP to evaluate abnormal IOC findings [9, 27, 34, 41]. This concept has gained wide acceptance and has been endorsed by the current National Institutes of Health State-of-the-Science Conference on ERCP [7].

Unfortunately, a significant number of patients undergoing postoperative ERCP in this study still did not have CBD stones. On multivariable analyses, abnormal LFT was the only variable that accurately predicted the presence of CBD stones. The utility of LFTs to predict CBD stones has been demonstrated previously in many studies, particularly for patients undergoing preoperative ERCP [6, 18, 29]. Although we checked LFTs in all patients before ERCP, we did not document details on preoperative LFTs. This precluded any correlation between the presence of CBD stones and fluctuation in enzyme levels.

Given that nearly one-third of patients with abnormal IOC do not have stones and the relative invasiveness and risks associated with ERCP, not all patients with an abnormal IOC should be subjected to routine postoperative ERCP. In a report from Seattle, acute biliary pancreatitis developed during the follow-up period in only 1.7% of patients who underwent laparoscopic cholecystectomy for symptomatic cholelithiasis [37].

Subsequent studies have shown that "on-demand" ERCP would be a better option than routine ERCP for evaluation of IOC abnormalities because most small stones migrate spontaneously [1, 14, 19, 33]. In the absence of clevated LFTs, less invasive tests such as EUS or magnetic resonance cholangiopancreatography (MRCP) should be used for evaluating these patients. The accuracy of both tests, particularly EUS, to evaluate for CBD stones is greater than 95% [12, 30, 36, 38]. Those with definitive CBD stones can then be subjected to ERCP. This selective approach would substantially dccrease ERCP-related morbidity and would be more cost effective [1]. The risk-benefits of a postoperative ERCP and alternative methods of evaluation should be discussed thoroughly with the patient.

A major limitation of our study is the small number of patients enrolled and the short follow-up period. With the advent of EUS, the number of postoperative ERCPs for evaluation of abnormal IOC findings has decreased substantially at both of our institutions. For instance, since the conclusion of this study, 5 of 10 patients referred to our centers for evaluation of an abnormal IOC received a diagnosis of a stone-free duct by EUS. As the application of EUS for evaluation of pancreaticobiliary disorders continues to expand, the role of diagnostic ERCP is expected to diminish further. Also, operating characteristics could not be calculated because we evaluated only the patients with an abnormal IOC. However, performing ERCP for patients with a normal IOC is both impractical and unethical. It also is well known that small stones can be missed at ERCP. However, doing a routine biliary sphincterotomy for all patients undergoing ERCP is potentially dangerous. At the 6-month follow-up assessment, none of our patients had any symptom recurrence, suggesting a low likelihood of a missed stone.

This study was conducted at two tertiary referral centers for ERCP. The number of patients with an abnormal IOC evaluated by methods other than ERCP, such as MRCP or EUS, is unknown.

The ability of abnormal intraoperative cholangiogram findings to predict retained CBD stones is unclear. In this prospective study, one-third of the patients with abnormal IOC had normal postoperative ERCP results. Only abnormal LFTs accurately predicted the presence of CBD stones. The use of less invasive methods such as EUS or MRCP should be considered for patients with normal postoperative LFTs to minimize unnecessary ERCPs.

- Ammori BJ, Birbas K, Davides D, Vczakis A, Larvin M, McMohan MJ (2000) Routine vs "on demand" postoperative ERCP for small bile duct ealculi detected at intraoperative cholangiography: clinical evaluation and cost analysis. Surg Endosc 14: 1123–1126
- Arregui M, Davis CJ, Arkush AM, Nagan RF (1992) Laparoscopic cholecystectomy combined with endoscopic sphineterotomy and stone extraction or laparoscopic choledochoscopy and electrohydraulic lithotripsy for management of cholelithiasis with choledocholithiasis. Surg Endosc 6: 10–15
- Barkun JS, Fried GM, Barkun AN, Sigman HH, Hinchey EJ, Garzon J, Wexler MJ, Mcakins JL (1993) Cholccystectomy without operative cholangiography: implications for common hile duct injury and retained common bile duct stones. Ann Surg 218: 371– 379
- Besten LD, Doty JE (1981) Cholcdocholithiasis. Surg Clin North Am 61: 893–907
- Birkett D (1992) Technique of cholangiography and cystic-duct choledochoscopy at the time of laparoscopic cholecystectomy for laser lithotripsy. Surg Endose 6: 252–254
- Bose SM, Mazumdar A, Prakash S, Koeher R, Katariya S, Pathak CM (2001) Evaluation of the predictors of choledocholithisis: eomprehensive analysis of clinical, biochemical, radiological, radionuclear, and intraoperative parameters. Surg Today 31: 117– 122
- Cohen S, Bacon BR, Berlin JA, Fleischer D, Hecht GA, Loehrer PJ Sr, MeNair AE Jr, Mulholland M, Norton NJ, Rabeneck L, Ransohoff DF, Sonnenberg A, Vannier MW (2002) National Institutes of Health State-of-the-Science Conference Statement: ERCP for diagnosis and therapy. Gastrointest Endose 56: 802–809
- Cranley B, Logan H (1980) Exploration of the common bile duct: the relevance of the elinical picture and the importance of preoperative cholangiography. Br J Surg 67: 869–872
- DePaula AL, Hashiba K, Bafutto M, Zago R, Grecco E (1993) Laparoscopic treatment of choledocholithiasis. Surg Laparosc Endosc 3: 157–160
- Freeman ML (2002) Adverse outcomes of ERCP. Gastrointest Endosc 56(6 Suppl): S273–S282
- Frossard JL, Hadengue A, Amouyal G, Choury A, Marty O, Giostra E, Sivignon F, Sosa L, Amouyal P (2000) Choledocholithiasis: a prospective study of spontaneous common bile duct stone migration. Gastrointest Endose 51: 175–179
- Fulcher AS, Turner MA, Capps GW, Zfass AM, Baker KM (1998) Half-Fourier RARE MR cholangiopancreatography in 300 subjects. Radiology 207: 21–32
- Hainsworth PJ, Rhodes M, Gompertz RH, Armstrong CP, Lennard TW (1994) Imaging of the common bile duet in patients undergoing laparoseopie cholecystectomy. Gut 7: 991–995
- Hawasli A, Lloyd L, Cacucei B (2000) Management of choledocholithiasis in the era of laparoscopic surgery. Am Surg 66: 425– 430

- Hunter JG (1992) Laparoscopic transcystic common bile duct exploration. Am J Surg 163: 53–56
- Ijzermans JN, De Waard P, Merkelbach JW (1989) Cholangiography during cholecystectomy: a plea for selective use. Neth J Surg 41: 79–81
- Johanson JF, Cooper G, Eisen GM, Freeman M, Goldstein JL, Jensen DM (2002) Quality assessment of ERCP: endoscopic retrograde cholangiopancreatography. Gastrointest Endosc 56: 165– 169
- Katz D, Nikfarjam M, Sfakiotaki A, Christophi C (2004) Selective endoscopic cholangiography for the detection of common bile duct stones in patients with cholelithiasis. Endoscopy 36: 1045–1049
- Kondylis PD, Simmons DR, Agarwal SK, Ciardiello KA, Reinhold RB (1997) Abnormal intraoperative cholangiography: treatment options and long-term follow-up. Arch Surg 132: 347–350
- Korman J, Cosgrove J, Furman M, Nathan I, Cohen J (1996) The role of endoscopic retrograde cholangiopancreatography and cholangiography in the laparoscopic era. Ann Surg 223: 212–216
- Legorreta A. Brooks RJ, Staroscik RN, Xuan Y, Costantino GN, Zats S (1995) Increased ERCP rate following the introduction of laparoscopic cholecystectomy. J Laparaendosc Surg 5: 271–278
- Levine SB, Lerner HJ, Leifer ED, Lindheim SR (1983) Intraoperative cholangiography: a review of indications and analysis of age-sex groups. Ann Surg 198: 692–697
- Liberman MA, Phillips EH, Carroll BJ, Fallas MJ, Rosenthal R, Hiatt J (1996) Cost-effective management of complicated choledocholithiasis: laparoscopic transcystic duct exploration or endoscopic sphineterotomy. J Am Coll Surg 182: 488–494
- Lilly MC, Arregui ME (2001) A balanced approach to choledocholithiasis. Surg Endose 15: 467–472
- 25. Loperfido S, Angelini G, Benedetti G, Chilovi F, Costan F, De Berardinis F, De Bernardin M, Ederele A, Fina P, Fratton A (1998) Major early complications from diagnostic and therapeutic ERCP: a prospective multicenter study. Gastrointest Endose 48: 1–10
- McMahon AJ, Russel IT, Baxter JN, Ross R, Anderson JR, Morran CG, Sunderland G, Galloway D, Ramsay G, O'Dwyer PJ (1994) Laparoscopic versus mini laparotomy cholecystectomy: a randomized controlled trial. Lancet 343: 135–138
- Ng T, Amaral JF (1999) Timing of endoscopic retrograde cholangiopancreatography and laparoscopic cholecysteetomy in the treatment of choledocholithisis. J Laparoendosc Adv Surg Tech A 9: 31–37
- O'Doherty D, Neoptolemos J, Carr-Locke D (1986) Endoscopic sphineterotomy for retained common bile duct stones in patients with T-tube *in situ* in the early postoperative period. Br J Surg 73: 454–456

- 29. Onken JE, Brazer SR, Eisen GM, Williams DM, Bouras EP, DeLong ER, Long TT 3rd, Pancotto FS, Rhodes DL, Cotton PB (1996) Predicting the presence of choledocholithiasis in patients with symptomatic cholelithiasis. Am J Gastroenterol 91: 762–767
- Palazzo L, Girollet P-P, Salmeron M, Silvian C, Roseau G, Canard JM, Chaussade S, Couturier D, Paolaggi JA (1995) Value of endoscopic ultrasonography in the diagnosis of common bile duct stones: comparison with surgical exploration and ERCP. Gastrointest Endosc 42: 225–231
- Perret RS, Sloop GD, Borne JA (2000) Common bile duct measurements in an elderly population. J Ultrasound Med 19: 727–730
- 32. Rhodes M, Sussman L, Cohen L, Lewis MP (1998) Randomised trial of laparoscopic exploration of common bile duct versus postoperative endoscopic retrograde cholangiography for common bile duct stones. Lancet 351: 159–161
- Ryberg AA, Fitzgibbons RJ Jr, Tseng A, Maffi TR, Burr LJ, Doris PE (1997) Abnormal cholangiograms during laparoscopic cholecystectomy. Surg Endosc 11: 456–459
- Schmitt CM, Baillie J, Cotton PB (1995) ERCP following laparoscopic cholecystectomy: a safe and effective way to manage CBD stones and complications. HPB Surg 8: 187–192
- Shapiro SJ, Gordon LA, Daykhovsky L, Grundfest W (1991) Laparoscopic exploration of the common bile duct: experience in 16 selected patients. J Laparoendosc Surg 1: 333–341
- 36. Soto JA, Barish MA, Alvarez O, Medina S (2000) Detection of choledocholithiasis with MR cholangiography: comparison of three-dimensional fast spin-echo and single- and multisection half-Fourier rapid acquisition with relaxation enhancement sequences. Radiology 215: 737–745
- Stoker ME (1995) Common bile duct exploration in the era of laparoscopic surgery. Arch Surg 130: 265–268
- Sugiyama M, Atomi Y (1997) Endoscopic ultrasonography for diagnosing choledocholithiasis: a prospective comparative study with ultrasonography and computed tomography. Gastrointest Endose 45: 143–146
- Taylor TV, Armstrong CP (1987) Migration of gallstones. Br Med J (Clin Res Ed) 294: 1320–1322
- Tham TCK, Lichtenstein DR, Vandervoort J, Wong RCK, Brooks D, Van Dam J, Ruymann F, Farraye F, Carr-Locke DL (1998) Role of endoscopic retrograde cholangiopancreatography for suspected choledocholithiasis in patients undergoing laparoscopic cholecystectomy. Gastrointest Endosc 47: 50–56
- 41. Urbach DR, Khajanchee YS, Jobe BA, Standage BA, Hansen PD, Swanstrom LL (2001) Cost-effective management of common bile duct stones: a decision analysis of the use of endoscopic retrograde cholangiopanereatography (ERCP), intraoperative cholangiography, and laparoscopic bile duct exploration. Surg Endosc 15: 4–13

Surg Endosc (2006) 20: 806-808 DOI: 10.1007/s00464-005-0430-0

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

### Urgent laparoscopic cholecystectomy in the management of acute cholecystitis: timing does not influence conversion rate

Y.-C. Wang, H.-R. Yang, P.-K. Chung, L.-B. Jeng, R.-J. Chen

Department of Surgery, China Medical University Hospital, No. 2, Yuh-Der Road, Taichung 404, Taiwan

Received: 16 June 2005/Accepted: 15 November 2005/Online publication: 16 March 2006

#### Abstract

*Background:* The optimal treatment of acute cholecystitis is urgent laparoscopic cholecystectomy. Most reports suggest that a delay of 72 or 96 h from onset of symptoms leads to a higher conversion rate. This study assessed the conversion rate in relation to the timing of urgent laparoscopic cholecystectomy for acute cholecystitis.

*Methods:* During a 12 month period, 112 patients received laparoscopic cholecystectomy for acute cholecystitis at a tertiary care university hospital in central Taiwan. Data were collected prospectively.

*Results:* The overall conversion rate was 3.6% (4/112). Of 62 procedures performed within 72 h from onset of symptoms, 2 were converted, as compared with 2 of 50 procedures after 72 h. Of 76 procedures performed within 96 h from onset of symptoms, 3 were converted, as compared with 1 of 36 procedures after 96 h. There were no mortalities or common bile duct injuries.

*Conclusions:* The conversion rate for urgent laparoscopic cholecystectomy among patients with acute cholecystitis can be as low as 3.6%. The timing of urgent laparoscopic cholecystectomy has no impact on the conversion rate.

Key words: Acute cholecystitis — Laparoscopic cholecystectomy

The introduction of laparoscopic cholecystectomy (LC) in 1988 was followed by a dramatic shift from open cholccystectomy (OC) to the laparoscopic approach. The benefits of laparoscopic surgery include reduced postoperative morbidity and postoperative pain, shortened hospitalization, quicker return to a normal lifestyle, and lower overall health care expenses [1, 7]. With improvement in instruments and technique, acute cholecystitis is no longer considered a contraindication for the laparoscopic approach [10, 17, 19].

It has been suggested that laparoscopic cholecystectomy should be performed within 72 to 96 h, after which there is an increased risk of complications and conversion to an open procedure [6, 9, 13, 18]. Conversion from LC to OC leads to abandonment of all the benefits associated with this minimally invasive operation and markedly increases the hospital length of stay and hospital expenses. Therefore, failure to perform urgent cholecystectomy within 72 to 96 h after onset of symptoms might be an indication for interval cholecystectomy [13]. Unfortunately, the conversion rate tends to be higher with interval surgery because dense fibrotic adhesion is more prevalent among patients who have interval surgery [15].

The aim of treatment for all patients with acute cholecystitis in our hospital is to perform definitive surgery at the initial emergency admission. With the accumulation of surgical experience and progressive improvement in surgical technique, this study reevaluated the feasibility of LC for patients with acute cholecystitis lasting longer than 72 h. The study assessed the conversion rate in relation to the timing of urgent laparoscopic cholecystectomy for acute cholecystitis.

#### Materials and methods

All patients with a diagnosis of acute cholecystitis who underwent cholecystectomy at our hospital in 2004 were enrolled in this study. The exclusion criteria specified septic shock; planned common bile duct exploration attributable to failure of preoperative endoscopic retrograde cholangiopancreaticography (ERCP), with stone extraction for choledocholithiasis; and previous complex upper abdominal surgery. All patients were admitted on an emergency basis with a diagnosis of acute cholecystitis, defined as acute upper abdominal pain with tenderness under the right costal margin; fever exceeding 37.5°C, leukocytosis exceeding  $10.39 \times 10^9/l$  (normal,  $< 10.39 \times 10^9/l$ ), or both; and ultrasonographic findings including thickened gallbladder wall, edematous gallbladder wall, distended gallbladder, presence of gallstones, and ultrasonographic Murphy's sign.

Data were collected prospectively. A standardized data form was used to record information, including demographic details, time

Correspondence to: Y.-C. Wang

**Table 1.** Relationship between duration of preoperative symptoms of acute cholecystitis and the frequency of conversion, operative time, and length of hospital stay in patients received laparoscopic cholecystectomy (Using 72 hours as cut point)

	Within 72 h	After 72 h	р
Case number	62	50	
Number of conversion	2	2	
Conversion rate	3.2%	4%	0.835
Median operative lime (Minute)	100 (50-225)	90 (60-210)	0.933
Median hospital stay (Day)	5 (2-14)	6 (2-18)	0.916

elapsed before surgery, intraoperative findings, complications, and postoperative length of hospital stay. The objective was to perform LC for all patients at their initial admission. All the patients received urgent LC within hours after admission or after preoperative ERCP was completed. Preoperative ERCP was performed in response to the following indications: ultrasonographic evidence of a dilated common duct, presence of common duct stones, or serum elevation in bilirubin. Preoperative sphincterotomy and stone extraction were performed in common duct stone was present. During surgery, gallbladder puncture was routinely used to relieve tension of the gallbladder and facilitate the subsequent manipulation. The gallbladder was removed under the protection of an endobag.

All variables were compared by the unpaired t-test or chi-square test. A probability level less than 0.05 was considered statistically significant.

#### Results

A total of 125 patients with a diagnosis of acute cholecystitis underwent urgent eholecysteetomy. Among the 125 patients, urgent laparoscopic cholecystectomy was performed initially for 112 patients (89.6%). The remaining 13 patients received open cholecystectomy because of earlier upper abdominal surgery (n = 3), planned common bile duct exploration (n = 3), and sepsis with unstable hemodynamic status (n = 7).

Of the 112 patients who received urgent laparoscopie cholecystectomy, 58 (52%) were women. The median age was 55 years (range, 20–87 years). Of the 112 eases, 16 (14.3%) were complicated cholecystitis involving gangrene, empyema, and perforated gallbladder. A total of 19 patients received a preoperative ERCP, and common duet stones were found and extracted after an endoscopie sphincterotomy from 14 patients (12.5%).

The overall conversion rate was 3.6% (4 of 112). The reasons for conversion to open cholecystectomy included unidentified gallbladder because of severe adhesion (n = 1), hemorrhage (n = 2), and Calot's triangle not visualized (n = 2). Tables 1 and 2 show the effect that the timing of urgent LC had on the conversion rate, operative time, and length of hospital stay. There was no statistically significant difference between the conversion rates for surgery performed within and after 72 h from onset of symptoms (2 of 62 vs 2 of 50 procedures; p = 0.835). Similarly, there was no difference in the conversion rates for surgery performed within and after 96 h from onset of symptoms (3 of 76 vs 1 of 36 procedures; p = 0.776). The median time to surgery for all patients was 3 days (range, 1–14 days).

**Table 2.** Relationship between duration of preoperative symptoms of acute cholecystitis and the frequency of conversion, operative time, and length of hospital stay in patients received laparoscopic cholecystectomy (Using 96 hour as cut point)

	Within 96 h	After 96 h	p
Case number	76	36	
Number of conversion	3	1	
Conversion rate	3.9%	2.8%	0.776
Median operative time (Minute)	100 (50-225)	90 (60-180)	0.745
Median hospital stay (Day)	5 (2-15)	6 (2-18)	0.022

The median operative time was 100 min (range, 50-225 min) for all procedures, 100 min (range, 50–225 min) for surgery within 72 h, 95 min (range, 60-210 min) for surgery after 72 h, 100 min (range, 50-225 min) for surgery within 96 h, and 90 min (range, 60–180 min) for surgery after 96 h. There were no differences in operating times between surgery within or after 72 h and that within or after 96 h from onset of symptoms. The median length of hospital stay was 5 days (range, 2-18 days) for all procedures, 5 days (range, 2-14 days) for surgery within 72 h, 6 days (range, 2-18 days) for surgery after 72 h, 5 days (range, 2-15 days) for surgery within 96 h, and 6 days (range, 2-18 days) for surgery after 96 h. There was no difference in the lengths of hospital stay for surgery within or after 72 h, but there was a significantly longer hospital stay for surgery after 96 h from onset of symptoms.

There was no mortality, although complications occurred for six patients (5.4%). Two patients experienced port-site infection. Three patients experienced subhepatic fluid collection, which was subsequently aspirated under ultrasound guidance. Cystic duet leakage was observed in one patient, which was managed with a common bile duct stent. After a follow-up period of 8 weeks, no subsequent complications were observed.

#### Discussion

With the development of technical skill and laparoscopic instruments, urgent laparoscopic cholecystectomy currently is considered the optimum treatment for patients admitted with acute cholecystitis [11]. As compared with successful laparoscopic chlccystectomy, conversion of laparoscopic to open cholecystectomy not only leads to an increase in operative time, patient morbidity, and hospital stay, but also results in a significant increase in overall eost [4]. It has been shown that the conversion rates are high in cases of acute cholecystitis. Pessaux ct al. [18] reported a conversion rate of approximately 27% in a 132 case study in 2000, and Navez et al. [17] obtained a conversion rate of 20% in a 609 patient study in 2001. In a 124 case study, Prakash et al. [19] reported a conversion rate of approximately 18.5% in 2002. These high conversion rates resulted from the technical difficulty of identifying biliary anatomy and managing severe inflammatory adhesions around the acutely inflamed gallbladder.

A number of risk factors for conversion in cases of acute cholecystitis, such as old age, large stones, a history of previous biliary disease, and a nonpalpable gallbladder, are associated with repeated inflammation, which results in a scarred and fibrosed gallbladder [15]. Timing of surgery from onset of symptoms has been considered one of the risk factors. It has been suggested that the optimum timing for urgent cholecystectomy is within 72 h after onset of symptoms. This suggestion was based on the finding that the conversion rate rises sharply after 72 h from onset of symptoms, negating the potential benefits of urgent surgery [3, 13, 15, 18, 21, 22]. Others have found the optimum timing for surgery to be within 96 h after onset of symptoms, with longer delays leading to a rising conversion rate [2, 5, 6, 9, 16, 17].

Furthermore, Koo and Thirlby [13] suggested that failure to perform urgent cholecystectomy within 72 or 96 h after onset of symptoms might be an indication for interval cholccystectomy. Unfortunately, interval surgery with initial conservative treatment for acute cholecystitis may result in failure. Fowkes and Gunn [8] reported that 20% of their patients failed conservative treatment during the initial hospitalization for acute cholecystitis and required surgery. Another 27% of patients who have their cholecystectomies delayed are readmitted for emergency cholecystectomy before their scheduled elective procedure. In addition, an increased proportion of shrunken gallbladder and chronic cholecystitis with dense fibrotic adhesions is encountered in interval laparoscopic cholecystectomy patients, which renders laparoscopic dissection impossible and sometimes unsafe. Several reports have demonstrated that the conversion and complication rates tend to be higher with interval surgery [14, 15, 20, 23].

The current results do not support the finding of earlier studies with regard to the timing of urgent laparoscopic cholecystectomy. The conversion rates were similar for surgery performed within 72 and 96 h after onset of symptoms, and did not vary significantly for cholecystectomy performed after these times. On the basis of this result, interval surgery for acute cholecystitis with symptoms after 72 or 96 h would not be considered. This is consistent with findings of recent studies showing that a delay in urgent surgery did not increase the conversion rate [2, 12].

The absence of major complications, such as bile duct injury, suggests that LC can be performed safely in patients with acute cholecystitis. The duration of surgery was similar in both groups. The major advantage of carly LC (within 96 h) is the reduction in the total hospital stay. However, the total hospital stay was similar for surgery performed within and after 72 h from onset of symptoms.

Surgeons often must decide whether to try a laparoscopic approach for patients with acute cholecystitis or not, especially for patients who have experienced symptoms longer than 72 or 96 h. The current study demonstrated that urgent laparoscopic cholecystectomy for acute cholecystectomy is safe and associated with a low conversion rate. The timing of surgery did not influence the outcome.

- Berggren U, Gordh T, Grama D, Haglund U, Rastad J, Arvidsson D (1994) Laparoscopic versus open cholecystectomy: hospitalization, sick leave, analgesia, and trauma responses. Br J Surg 81: 1362–1365
- Bhattacharya D, Senapati PS, Hurle R, Ammori BJ (2002) Urgent versus interval laparoscopic cholecystectomy for acute cholecystitis: a comparative study. J Hepatobiliary Pancreat Surg 9: 538–542
- Chandler CF, Lane JS, Ferguson P, Thompson JE, Ashley SW (2000) Prospective evaluation of early versus delayed laparoscopic eholecystectomy for treatment of acute cholecystitis. Am Surg 66: 896–900
- De Pouvourville G, Rihet-Reinhart N, Fendriek M, Houry S, Testas P, Huguier M (1997) A prospective comparison of the costs and morbidity of laparoscopic versus open cholecystectomy. Hepatogastroenterology 44: 35–39
- Eldar S, Eitan A, Biekel A, Sabo E, Cohen A, Abrahamson J, Matter I (1999) The impact of patient delay and physician delay on the outcome of laparoscopic cholecystectomy for acute cholecystitis. Am J Surg 178: 303–307
- Eldar S, Sabo E, Nash E, Abrahamson J, Matter I (1997) Laparoscopic cholecystectomy for acute cholecystitis: prospective trial. World J Surg 21: 540–545
- Escaree JJ, Chen W, Sehwartz JS (1995) Falling eholecystectomy thresholds since the introduction of laparoscopic cholecystectomy. JAMA 273: 24–31
- Fowkes FG, Gunn AA (1980) The management of acute cholecystitis and its hospital cost. Br J Surg 67: 613–617
- Garher SM, Korman J, Cosgrove JM, Cohen JR (1997) Early laparoseopie cholecysteetomy for aeute cholecystitis. Surg Endosc 11: 347–350
- Glavic Z, Begic L, Simlesa D, Rukavina A (2001) Treatment of acute cholecystitis: a comparison of open vs laparoscopic cholecystectomy. Surg Endose 15: 398–401
- Hunter JG (1998) Acute cholceystitis revisited: get it while it's hot. Ann Surg 227: 468–469
- Knight JS, Mercer SJ, Somers SS, Walters AM, Sadek SA, Toh SK (2004) Timing of urgent laparoscopic cholecystectomy does not influence conversion rate. Br J Surg 91: 601–604
- Koo KP, Thirlby RC (1996) Laparoscopic cholecystectomy in acute cholecystitis: what is the optimal timing for operation? Arch Surg 131: 540–544, discussion 544–545
- Lai PB, Kwong KH, Leung KL, Kwok SP, Chan AC, Chung SC, Lau WY (1998) Randomized trial of early versus delayed laparoscopie cholecysteetomy for acute cholecystitis. Br J Surg 85: 764–767
- Lo CM, Liu CL, Fan ST, Lai EC, Wong J (1998) Prospective randomized study of early versus delayed laparoscopic cholecystectomy for acute cholecystitis. Ann Surg 227: 461–467
- Lo CM, Liu CL, Lai EC, Fan ST, Wong J (1996) Early versus delayed laparoscopic cholecystectomy for treatment of acute eholecystitis. Ann Surg 223: 37–42
- Navez B, Mutter D, Russier Y, Vix M, Jamali F, Lipski D, Cambier E, Guiot P, Leroy J, Mareseaux J (2001) Safety of laparoscopic approach for acute cholecystitis: retrospective study of 609 eases. World J Surg 25: 1352–1356
- Pessaux P, Tueeh JJ, Rouge C, Duplessis R, Cervi C, Arnaud JP (2000) Laparoscopic cholecystectomy in acute cholecystitis: a prospective comparative study in patients with acute vs ehronie cholecystitis. Surg Endosc 14: 358–361
- Prakash K, Jacob G, Lekha V, Venugopal A, Venugopal B, Ramesh H (2002) Laparoscopic cholecystectomy in acute choleeystitis. Surg Endosc 16: 180–183
- Schrenk P, Woisctschlager R, Rieger R, Wayand WU (1998) A diagnostic score to predict the difficulty of a laparoseopie cholecystectomy from preoperative variables. Surg Endosc 12: 148–150
- Serralta AS, Bucno JL, Planells MR, Rodero DR (2003) Prospective evaluation of emergency versus delayed laparoscopie cholecystectomy for early cholecystitis. Surg Laparose Endosc Pereutan Tech 13: 71–75
- Uchiyama K, Onishi H, Tani M, Kinoshita H, Ueno M, Yamaue H (2004) Timing of laparoscopic cholecystectomy for acute cholecystitis with cholecystolithiasis. Hepatogastroenterology 51: 346–348
- Willsher PC, Sanabria JR, Gallinger S, Rossi L, Strasherg S, Litwin DE (1999) Early laparoscopic cholecystectomy for acute eholeeystitis: a safe procedure. J Gastrointest Surg 3: 50–53

Surg Endosc (2006) 20: 809–811 DOI: 10.1007/s00464-005-0426-9

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

## It is not worthwhile to perform ileoscopy on all patients

#### K. K. Y. Yoong,<sup>1</sup> T. Heymann<sup>2</sup>

<sup>1</sup> Oldchurch Hospital, Waterloo Road, Romford RM7 OBE, UK

<sup>2</sup> Tanaka Business School, Imperial College London, South Kensington Campus, London SW7 2AZ, UK

Received: 15 June 2005/Accepted: 21 October 2005/Online publication: 21 January 2006

#### Abstract

*Background:* It remains controversial whether ileoseopy should be attempted in all patients. Although the ease of ileoscopy and the diagnostic yield have been well described, there have been no studies describing the value of the diagnostic yield in altering elinical management. We carried out a study to ascertain whether it is worthwhile to perform ileoseopy in all patients having a colonoscopy.

*Methods:* We carried out a retrospective study of all patients who had a colonoscopy between January 1, 2002, and December 31, 2003. The patient details, indications, findings, and complications of the procedure were recorded, together with the histopathology reports of colonic and terminal ileal biopsies. Clinical ease note of patients with a positive diagnosis was reviewed to ascertain whether there was a change to the patient's management following an abnormal biopsy result.

*Results:* A total of 2,149 colonoscopies were performed. In 346 patients (16.1%), the terminal ileum was intubated. There were 16 abnormal findings on histology, which gave a diagnostic yield of 4.6% of all ileoscopies. A change to management occurred in only half of these patients.

*Conclusion:* Ileoscopy should only be attempted in situations in which the indication is warranted and that would alter management. It is not cost-effective to carry out ileoscopy on all patients.

Key words: Costs — General — Endoseopy

A useful investigation can be defined as one in which the result will alter the management or add confidence to the clinician's diagnosis. Unnecessary investigations increase waiting times, waste resources, and expose patients to unnecessary risk.

Correspondence to: K. K. Y. Yoong

Ileoscopy as an adjunct to colonoscopy was first described three decades ago [5]. It helps to establish completeness of the examination, and in some cases small bowel inflammatory disease can only be established by ileoscopy [2, 8]. Some prospective studies have described success rates of >90% for ileal intubation [1, 9]. One of these studies was carried out in a tertiary referral center with state-of-the-art equipment and demonstrated a median time of 55 sec to pass from ceeum to terminal ileum [1].

It remains controversial whether ileoscopy should be attempted in all patients. Although the ease of ileoscopy and the diagnostic yield have been well described, there have been no studies describing the value of the diagnostic yield in altering elinical management. The aim of this study is to ascertain whether ileoscopy is a useful, cost-effective investigation for all patients having a colonoscopy.

#### Patients and methods

We performed a retrospective analysis of all patients who had a colonoscopy in the medical gastroenterology department at Kingston Hospital between January 1, 2002, and December 31, 2003, which is a 500-bed district general hospital serving a population of approximately 320,000. The hospital's medical gastroenterology department performs approximately 1,000 colonoscopies per year. Three endoscopists and three trainees currently staff the department. Data were obtained from medical records and our computerized endoscopy datahase. The patient details, indications, and findings of the procedure were recorded. Histopathology reports of colonic and terminal ileal biopsies were also recorded. Clinical case notes of patients with abnormal appearance on histology were reviewed to establish how such findings had altered management. We defined abnormal findings as clinically significant if this resulted in a new diagnosis, initiation or change in drug treatment, or further investigation.

Colonoscopy was performed using Olympus CF-240L or CF-20HL video colonoscopies.

#### Results

A total of 2,149 colonoscopies were performed by eight medical gastroenterologists or their advanced trainees

Table 1. Indications for colonoscopy

Table 2. Indications, diagnostic yield, and clinical significance of patients who had ileal intubation

Indication	No. of patients	
Diarrhea	450	
Rectal bleeding (overt/occult)	407	
Abdominal pain	327	
Anemia	320	
Follow-up for polyp	252	
Follow-up for inflammatory bowel disease	254	
Constipation	178	
Intermittent constipation and diarrhea	104	
Family history of colorectal eancer	104	
Weight loss	76	
Abdominal mass	29	

Indication	No. of patients	Diagnostic yield	Clinically significant
Diarrhea	119	11 (9.2%)	4 (3.3%)
Abdominal pain	74	2 (2.7%)	0 (0%)
Rectal blecding (overt/oecull)	62	5 (8.1%)	3 (4.8%)
Follow-up for inflammatory bowel disease	33	5 (15.2%)	2 (6.1%)
Anemia	36	4 (11.1%)	2 (5.6%)
Abdominal mass	3	1 (33.3%)	1 (33.3%)

Table 3. Indications, diagnosis, and management change of patients with abnormal ileal histology

Patient No.	Indications	Diagnosis	Management change
1	Anemia, diarrhea	Crohn's disease	Started on new drugs
2	Diarrhea, rectal bleeding	Crohn's disease	Started on new drugs
3	Follow-up for IBD	Crohn's disease	Started on new drugs
4	Diarrhea	Crohn's disease	Starled on new drugs
5	Follow-up for IBD	Crohn's disease	Started on new drugs
6	Abdominal mass	High-grade lymphoma	Chemotherapy and surgery
7	Anemia, diarrhea	Tubereulosis	Antitubereulous drugs
8	Anemia	High-grade dysplastic villous adenoma	Right hemicolectomy

IBD, inflammatory bowel discase

from January 2002 to December 2003, and the indications are summarized in Table 1. Most patients had more than one indication. The most common indications were diarrhea, rectal bleeding (over/occult), abdominal pain, and anemia. Approximately one-fourth of the patients (252 + 254 = 516) were asymptomatic and had colonoscopy carried out for surveillance of previous polyps or chronic inflammatory bowel disease (IBD).

In 346 cases (16.1%), the terminal ileum was successfully intubated. Of these, 115 biopsies of the terminal ileum were taken for histology. Table 2 summarizes the indications of patients with successful ileal intubation, with diagnostic yield and its clinical significance.

Of the 346 ileal intubations, 16 were found to have an abnormal finding on histology, which gave a diagnostic yield of 4.6%. Of the 16 patients who had abnormal ileal histology, the findings prompted a change to the clinical management in eight patients (Table 3).

#### Discussion

In 2001, the British Society of Gastroenterology working party suggested that the average district general hospital should plan for an annual workload of 800– 1,000 colonoscopies per 100,000 population [7]. With a population of approximately 60 million, this would equate to 600,000 colonoscopies per year [6]. Assuming ilcoscopy takes an additional 55 sec per procedure and that it was performed on all colonoscopies, this would mean an additional 10,000 h of procedure time [1]. Therefore, although the additional time taken to perform an ileoscopy during colonoscopy may seem short, taken in a national context this would mean an additional 2,000 colonoscopies annually. There will also be an impact on the histopathology services.

The diagnostic yield with clinical significance was highest in patients who had colonoscopy performed for investigation of an abdominal mass, follow-up of IBD, anemia, and diarrhea. Our study confirms the findings of previous studies, which demonstrated no benefit of ileoscopy for unselected patients undergoing colonoscopy [4, 9], suggesting that it may be better to perform ileoscopy only on patients in whom there is an abnormal finding and also in whom treatment is likely to be altered. With the time saved, an additional 2000 patients could be colonoscoped.

The current analysis has some limitations. There is a selection bias in patients who had ileoscopy and ileal biopsies. Although the ileoscopy rate of 16.1% is much lower than that of other studies, the diagnostic yield of 4.6% is similar. This suggests selective ileoscopy has not compromised diagnostic yield. A prospective study would have been preferable, but this study reflects the practice of a district general hospital and not a tertiary referral center, in which ileoscopy would be selected in patients most likely to provide a diagnostic yield. However, with a positive diagnostic yield, in only half of the patients were the ileoscopy findings clinically significant to warrant a change in management. Another argument against performing ileal biopsies on all patients is the risk of transmitting variant Creutzfeldt-Jakob discase from one patient to another [3].

Ileoscopy is an important part of colonoscopy, but our study highlights the cost implication related to performing ileoscopy on all patients when most patients will not have their clinical management altered following the procedure. However, some have argued that an ileoscopy revealing a normal mucosa of the terminal ileum may be helpful [9], and it must be emphasized that a colonoscopist needs to become proficient in this technique to perform it when necessary.

In conclusion, ileal intubation should only be attempted when the indication is most likely to have an abnormal finding and alter management.

#### References

- Ansari A, Soon SY, Saunders BP, Sanderson JD (2003) A prospective study of the technical feasibility of ileoscopy at colonoscopy. Scand J Gastroenterol 38: 1184–1186
- 2. Batres LA, Maller ES, Ruchelli E, et al. (2002) Terminal ileum intubation in pediatric colonoscopy and diagnostic value of

conventional small bowel contrast radiography in pediatric inflammatory bowel disease. J Pediatr Gastroenterol Nutr 35: 320-323

- Bramble MG, Ironside JW (2002) Creutzfeldt–Jakob disease: implications for gastroenterology. Gut 50: 888–890
- Kundrotas LW, Clement DJ, Kubik CM, et al. (1994) A prospective evaluation of successful terminal ileum intubation during routine colonoscopy. Gastrointest Endose 40: 544–546
- Nagasako K, Yazawa C, Takemoto T (1972) Biopsy of the terminal ileum. Gastrointest Endosc 19: 7–10
- 6. Office for National Statistics (2001) Census 2001. Stationary Office, London
- 7. Working Party of the British Society of Gastroenterology Endoscopy Committee (2001) Provision of endoscopy related services in district general hospitals, working party report
- Zwas FR, Bonheim NA, Berken CA, Gray S (1994) Ileoscopy as an important tool for the diagnosis of Crohn's disease: a report of seven cases. Gastrointest Endosc 40: 89–91
- Zwas FR, Bonheim NA, Berken CA, Gray S (1995) Diagnostic yield of routine ileoscopy. Am J Gastroenterol 90: 1441–1443

Surg Endosc (2006) 20: 812–814 DOI: 10.1007/s00464-005-0580-0

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

### **Rigid sigmoidoscopy**

#### A potential hazard for cross-contamination

#### D. Z. Lubowski, G. L. Newstead

Department of Colorectal Surgery, St. George and Prince of Wales Hospitals, Sydney, Australia

Received: 23 August 2005/Accepted: 26 October 2005/Online publication: 17 January 2006

#### Abstract

*Background:* Rigid sigmoidoscopy using a disposable or nondisposable sigmoidoscope is a common outpatient procedure. It has been assumed that the nondisposablc bellows and light head of the sigmoidoscope remain frec from enteric organisms so that the procedure is sterile if a disposable or nondisposable (metal) sigmoidoscope shaft is used. The aim of this study was to identify the presence of organisms within the bellows or light head of the sigmoidoscope.

*Methods:* Of 21 patients undergoing rigid sigmoidoseopy with a disposable instrument, bacterial eultures were taken from the inside of sterile Jackson-Pratt bulbs in 12 patients, with the bulbs being used to simulate the nondisposable insufflation bellows. In an additional nine patients, swabs were taken for eulture from the inside of the nondisposable light head.

*Results:* Enterie gram-negative *Escherichia coli* and mixed anaerobic organisms were eultured from the Jackson-Pratt bulbs in two eases, and gram-positive organisms were cultured in another case. Gram-negative organisms, including *Bacillus, Proteus mirabilis, Klebsiella*, and *Enterococcus faecalis*, were cultured from the inside of the light head in two eases.

*Conclusion:* Sigmoidoscopy using a disposable instrument is not a sterile procedure and may pose a risk of patient-to-patient cross-contamination by potentially harboring organisms in the bellows or light head.

Key words: Sigmoidoscopy — Cross-infection — Bacteriology

Sigmoidoscopy is a frequently performed procedure, usually in the outpatient setting. The term "sigmoidoscopy," used in Europe and Australasia, is perhaps a misnomer since although the instrument may sometimes be passed beyond the rectosigmoid junction, generally the procedure examines only the rectum; indeed, "proctoscopy" is the terminology used for the procedure of sigmoidoscopy in the United States. The instrument can be inserted up to its limit in approximately 40% of examinations [6] and a good view of the rectal mueosa can usually be obtained. If a large amount of stool is encountered in the rectum on initial digital examination, then an enema may be administered prior to sigmoidoscopy. In some eolorectal clinies, patients routinely receive an enema prior to the initial examination, which facilitates the process of examination of the rectum. A small amount of solid stool may not preclude a satisfactory view of the reetal mueosa.

Clinical indications for rigid sigmoidoseopy are varied. At the patient's first visit, the authors always carry out sigmoidoscopy in patients with rectal bleeding if there is no painful or tender anal lesion. This may be the definitive investigative procedure in the young patient with minor reetal bleeding and identifiable anal pathology. Identification of rectal polyps or cancer at sigmoidoscopy is important if there is otherwise to be a delay before colonoscopy. The distribution of adenomatous polyps in the colon is now well-known from colonoscopy studies. Approximately two-thirds occur distal to the splenic flexure, with 40–55% in the rectum and sigmoid [5, 12, 13]. In cases in which there is a solitary adenoma, 56% are found in the rectum and sigmoid [4].

Cancer most commonly occurs in the rectum, although during the past four decades there has been a shift in distribution toward the right side of the colon. In 1980, 41.9% of patients who died from eolorectal cancer in England and Wales had rectal cancer. The proportions in Australia were similar [8]. In the United States, a number of studies, including that from the National Cancer Institute (1966) [1], have documented a shift to the right side of the colon and hence an increase in the ratio of colon eancer to rectal cancer [1, 2, 10]. Interestingly this ehange has apparently been less marked in the United Kingdom [7] and Australia [3].

Correspondence to: D. Z. Lubowski

Sigmoidoscopy will diagnose proctitis, benign rectal ulceration as part of the solitary rectal ulcer syndrome, and is useful to identify rectal prolapse in the patient who has not been aware of external prolapse by observing the reetum while asking the patient to strain down. The procedure is very helpful in monitoring the progress of diseases such as colitis, radiation proctitis, or the mucosal changes in the solitary rectal ulcer syndrome, and a stool specimen may be collected during sigmoidoscopy for microscopy and culture.

Initially, rigid sigmoidoseopes were metal and reusable. There has been a gradual endange to disposable plastic instruments, thus avoiding the need to elean and sterilize instruments after each use. Apart from the labor costs, in a busy elinie it is therefore necessary to have a number of expensive instruments. Most elinicians thus choose to use disposable sigmoidoseopes. Only the sigmoidoscope shaft is disposable, whereas the bellows and light source are reusable. Throughout the years, we have noted that watery stool may oceasionally reflux into the bellows during the examination, particularly if the patient raises abdominal pressure during the procedure, and this has necessitated eleaning the bellows. The reusable light source may be affected in a similar way. Thus, concern was raised about sterility of the subsequent procedure and, in particular, the possibility of transmitting baeteria or other organisms harbored in the bellows or tubing, or in the light source, to the next patient during air insufflation.

The aim of this study was to examine the hypothesis that bacteria may be found on the internal surfaces of the bellows and light source after sigmoidoscopy, thus creating a potential risk for cross-contamination.

#### Materials and methods

The study was approved by the South East Area Health Service (Southern Section). Twenty-one patients undergoing rigid sigmoidoscopy as part of their routine clinical examination were studied. Sigmoidoscopy was carried out after a digital anorectal examination with the patient lying comfortably in the left lateral position on the examination couch. A disposable sigmoidoscope (Weleh Allyn, New York, USA) was used in each case; no hiopsies were taken.

In the first 12 patients, the nondisposable bellows and tuhing of the Welch Allyn sigmoidoscope set were replaced with a sterile Jackson-Pratt bulb (Baxter Healthcare, Deerfield, IL, USA). This bulh is usually used as a closed-wound suction drainage system hut functions effectively for air insufflation. The egg-shaped bulh has an inlet port with an antireflux valve and a drainage port with removable plug, and it is supplied in a sterile plastic pack. The Jackson-Pratt hulh was removed from its container using ascptic technique. The drainage port was connected to a length of sterile soft connection tuhing, which in turn was connected to the nondisposable Welch Allyn sigmoidoscope light head. In the first six of these 12 patients, the light head was cleaned and then sterilized in the autoclave at 120°C under pressure for 15 min before each use. In the last six of the 12 patients, the light head was not sterilized and was reused as in normal practice.

The instrument was withdrawn on completion of the sigmoidoscopy and, using aseptic technique, the hulh was disconnected from the tubing and 20 ml sterile saline was instilled via the drainage port. The plug was then inserted into the drainage port and the saline was gently swirled in the hulh. A sterile Steri-Strip tape was placed to seal the inlet port. The hulh was then sent to the pathology laboratory for microbiological examination of the contents of the saline. Cultures were set up on blood agar, MacConkey agar, and chocolate agar plates for gram-positive and gram-negative aerobic and anaerobic organisms and on LJ slope for mycohacteria. Incuhation was continued for 48 h, and once a growth was identified, incubation was continued to allow organism identification.

Tests were carried out on the light head of the sigmoidoseope in an additional nine patients. For this part of the study, the normal reusable Weleh Allyn hellows was used. The light head was cleaned and sterilized as described previously at the heginning of each consulting session and then used in several consecutive sigmoidoscopic examinations. In each case, the sigmoidoseope was inserted, the ohturator withdrawn, and the light head then connected to the sigmoidoscope. At the end of the examination, new sterile gloves were used and swahs were taken from the inside surface of the light head and cycpiece. The swahs were placed in Amies' transport medium and subsequently plated on blood agar, chocolate agar, and Colistin/naladixic acid hlood agar plates.

#### Results

Cultures from the Jackson-Pratt bulbs identified no organisms in nine of the 12 patients. In three patients, the following cultures were found, respectively: 100-1,000 colonies/ml of *Escherichia coli* and mixed anaerobie organisms; 100-10,000 colonies of *E. coli* and mixed anaerobie organisms; and a small count (<10 colonies/ml) of *Staphylococcus epiderunidis*. The first of these three patients was in the group in which the light head had been sterilized immediately before each use.

Cultures from the light heads in the second group of nine patients showed no growth in seven patients. In two patients, the following organisms were found: (1) one colony of *Bacillus* species; and (2) *Proteus mirabilis*, *Klebsiella pneumoniae*, and *Enterococcus faecalis*.

#### Discussion

Sterilization of instruments is an important eonsideration in modern surgieal practice. In recent years, coneern has been heightened by the advent of transmissible life-threatening diseases, for both elinical and medicolegal reasons. The processes applied to the cleaning of instruments used for flexible sigmoidoseopy or eolonoseopy are now carefully regulated, usually with disinfection in 2% glutaraldehyde for 20 min or 0.2% paracetic acid using an automated system (Steris, Mentor, OH, USA). Detailed guidelines for infection control in surgery from the Royal Australasian College of Surgeons [11] and state health authorities within Australia [9] are in place, including rules governing sterilization of instruments and use of single-use surgical items.

This study has shown that feeal bacteria may be found in the bellows of the rigid sigmoidoscope. Baeteria were cultured from a previously sterile bulb after a single use. The inside surface of the bellows and, in this experimental study, the bulb are in free communication with the rectum via gas under pressure. Relatively small bacterial counts were isolated in this study, but it is presumed that higher counts are likely to be found in the nondisposable bellows after use in multiple patients. Bacteria were identified in the bellows in 25% of patients after only a single use and may therefore be expected more frequently and in greater numbers with multipleuse bellows. The study was designed to ensure that bacteria had not reached the bellows by contamination from the air or by direct patient contact, thus demonstrating that bacteria can easily be introduced into the bellows during sigmoidoscopy. The culture of *Staph. epidermidis* was most likely to have been a skin contaminant from the operator despite the aseptic technique, a phenomenon known to occur in other circumstances such as when inserting an intravenous cannula under aseptic conditions, although it is possible that the organisms originated from the anal skin of the patient and entered the bellows during the sigmoidoscopy.

Bacteria were also isolated from the inside surface of the light head. The sigmoidoscope design is such that there is free communication between the light head, bellows, sigmoidoscope shaft, and the rectal gas, which has been insufflated under pressure. The light head is therefore exposed to organisms in the same way as the inside of the bellows. Although the design of the sigmoidoscope attempts to prevent contamination of the light head by providing a plate at the proximal end of the shaft, hence separating the light head from the shaft, the shaft may become contaminated during withdrawal of the obturator. Thus, organisms may also reach the light head after withdrawal of the obturator followed by coupling the light head onto a contaminated sigmoidoscope shaft. Placement of the light head in direct contact with the potentially contaminated proximal end of the shaft is inconsistent with the rigid principles of aseptic technique and instrument sterilization, which are mandatory in all surgical and other endoscopic procedures.

It has been proposed that a disposable filter be used, interposed between the light head and the bellows to prevent organisms from reaching the bellows. Filters are now available for use with some sigmoidoscopes (Welch Allyn; Seward, Norfolk, UK); although an appropriate filter will prevent passage of bacterial and viral particles, penetration of the filter by fluid will render it ineffective. Filters should therefore only be considered for single use, but even then they may become ineffective during a procedure if exposed to liquid stool. In addition, a filter does not prevent contamination of the internal surfaces of the light head.

We are not aware of any documented cases in which clinical infection occurred as a result of cross-contamination during sigmoidoscopy. Organisms may potentially gain access if mucosal biopsies are taken or polyps are removed during sigmoidoscopy, but minor trauma to the mucosa resulting from insertion of the instrument may also be a potential site of entry. Guidelines from the Australian Therapeutics Goods Administration recommend sterilization of all parts of the rigid sigmoidoscope to prevent cross-infection [14]. Current disposable sigmoidoscope designs do not provide the same sterile conditions as with other surgical instruments or flexible endoscopes, and prevention of cross-contamination cannot be ensured unless the bellows, tubing, and light head are autoclaved before each use.

- Axtell LM, Chiazzi L (1966) Changing relative frequency of cancer of the colon and rectum in the United States. Cancer 19: 750–754
- Cady B, Pearson AV, Manson DO, Maunz DL (1974) Changing patterns of colorectal carcinoma. Cancer 33: 422–426
- Chapuis PH, Newland RC, Macpherson JG, Dent O, Payne J, Pheils MT (1981) The distribution of colorectal carcinoma and the relationship of tumour site to survival of patients following resection. Aust N Z J Surg 51: 127–131
- 4. Ekelund G (1963) Cancer and polyps of colon and rectum. Acta Pathol Microbiol Scand 59: 165–170
- Gillespie PE, Chambers TJ, Chan KW, Doronzo F, Morson BC, Williams CB (1979) Colonic adenomas: a colonoscopic survey. Gut 20: 240–245
- Keighley MRB, Williams NS (1993) Polypoid disease and polyposis syndromes (anatomical distribution). In Surgery of the anus, rectum and colon. Saunders, London, pp 760–771
- Keighley MRB, Williams NS (1993) Colorectal cancer: cpidemiology, aetiology, pathology, clinical features and diagnosis. In Surgery of the anus, rectum and colon. Saunders, London, pp 830–885
- McDermott FT, Hughes ESR, Pihl H, Milne BJ, Price AB (1981) Comparative results of surgical management of single carcinomas of the colon and rectum: a series of 1939 patients managed by one surgeon. Br J Surg 68: 850–855
- NSW Health Department. Infection control policy document circular 2002/45, pp 1–42
- Rosato FE, Marks G (1981) Changing site distribution patterns of colorectal cancer at Thomas Jefferson University Hospital. Dis Colon Rectum 24: 93–96
- Royal Australasian College of Surgeons (1998) A guide to infection control. Infection control in surgery policy document
- Shinya H, Wolff W1 (1979) Morphology, anatomic distribution and cancer potential of colonic polyps. An analysis of 7000 polyps. Ann Surg 190: 679–683
- Tedesco FJ, Waye SD, Avclla JR, Villabos MM (1980) Diagnostic implications of the spatial distribution of colonic mass lesions (polyps and cancer). Gastrointest Endosc 26: 95–97
- Therapeutics Goods Administration (1995) Australian Therapeutics Device Bulletin No. 28

Surg Endosc (2006) 20: 815–819 DOI: 10.1007/s00464-005-0428-7

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

### Preconditioning-like amelioration of erythropoietin against laparoscopy-induced oxidative injury

E. Ates,<sup>1</sup> S. Yilmaz,<sup>2</sup> E. Ihtiyar,<sup>1</sup> B. Yasar,<sup>1</sup> E. Karahuseyinoglu<sup>1</sup>

<sup>1</sup> General Surgery Department, Osmangazi University, The Faculty of Medicine, Eskischir, Turkey
<sup>2</sup> General Surgery Department, Afyon Kocatepe University, The Faculty of Medicine, Afyon, Turkey

Received: 16 June 2005/Accepted: 24 October 2005/Online publication: 21 February 2006

#### Abstract

*Background:* Laparoscopic surgery has gained wide acceptance for almost every kind of surgical procedure, although it has produced significant oxidative injury to intraabdominal organs depending on the pressure level and the kind of the gas used. The literature describes several preventive measures for decreasing the postla-paroscopic oxidative injury such as low intraabdominal pressure, gasless laparoscopy, and laparoscopic preconditioning. Erythropoietin was shown previously to decrease ischemia–reperfusion injury to the liver. The current study evaluated the effect of erythropoietin against laparoscopy-induced oxidative injury, as compared with laparoscopic preconditioning.

Methods: For this study, 64 male Spraque-Dawley rats were randomly assigned to one of the following groups. The control group was subjected to a sham operation. The laparoscopy group was subjected to 60 min of pncumoperitoneum. The laparoscopic preconditioning plus laparoscopy group was subjected to 5 min of insufflation and 5 min of desufflation followed by 60 min of pneumoperitoncum. The crythropoietin plus laparoscopy group was subjected to a subcutaneous injection of erythropoictin as a single 1,000-U/kg dose followed by 60 min of pncumoperitoneum. After 45 min of desufflation subsequent to cessation of pneumoperitoneum, blood, liver, and kidney samples were obtained from half of the rats. The other half of the rats were observed for a reperfusion period of 24 h. Tissue and blood samples also were obtained after this period.

*Results:* Laparoscopy produced significant oxidative injury, as compared with the sham treatment. Laparoscopic preconditioning produced significant amelioration of the ischemic injury. Although crythropoietin administration during the prelaparoscopic period decreased the pneumoperitoneum-induced oxidative injury, the beneficial effect of laparoscopic preconditioning was more pronounced.

Correspondence to: S. Yilmaz

*Conclusion:* Laparoscopic preconditioning is more effective than the preischemic administration of erythropoietin in reducing laparoscopy-induced oxidative injury.

Key words: Erythropoietin — Ischemia/reperfusion injury — Laparoscopic preconditioning — Laparoscopy — Malondialdehyde — Tumor necrosis factor-a

Laparoscopic surgery has gained wide acceptance for almost every kind of surgical procedure because it results in less pain and a shorter hospital stay. However, the pneumoperitoneum itself is not a completely harmless procedure, as evidenced by several structural and functional alterations observed after its use. Pneumoperitoneum carries the risk of splanchnic ischemia as well as metabolic and infectious consequences depending on the pressure level used for pneumoperitoneum and the kind of gas used [11, 21, 22]. The adverse alterations also have been attributed to the increased carbon dioxide (CO<sub>2</sub>) load [10, 16].

As a consequence of these responses, pneumoperitoneum produces ischemia during insufflation and reperfusion during desufflation. Therefore, laparoscopic procedures actually produce ischemia-reperfusion injury.

The induction of free oxygen radicals after the restoration of blood flow during the desufflation phase is one of the most important mechanisms of organ dysfunction after laparoscopy. We previously demonstrated that laparoscopy produces severe oxidative injury to the liver, kidney [22], small intestine [24], and peritoneal [3] tissues. The mechanisms activated after ischemia-reperfusion actually prolong the hypoxia that has already begun during the ischemia. The aggravated hypoxia is followed by increased inflammatory cytokines and reactive oxygen species, thereby increasing the tissue injury [4].

We also have reported that tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interleukin-6 (1L-6) contribute at least

partially to oxidative injury observed after laparoscopy [1, 3]. The oxidative injury represents the loss of normal balance between the oxidative agents and natural antioxidant scavengers. It is not only an indicator, but also a contributor to organ damage, especially damage to the liver and kidney [7, 17, 20]. We have previously shown that ischemic preconditioning (Pre), originally reported as beneficial for myocardial tissues [8], can be applicable in laparoscopic procedures, with brief periods of pneumoperitoneum and desufflation administered before the sustained pneumoperitoneum period [22, 24].

Laparoscopic preconditioning (L-Pre), one of the most important preventive methods against laparoscopy-induced oxidative injury, produces its beneficial effect by decreasing the cytokine responses [1]. It also is more effective than low-pressure pneumoperitoneum in reducing ischemic insult associated with laparoscopy [3]. During laparoscopic procedures, the intraabdominal organs are exposed to ischemic injury not only during the insufflation phase, but also during the deflation phase. The liver and the kidneys are among the splanchnic organs most severely affected by laparoscopic procedures, as implied by increased blood levels of liver enzymes noted in animals and humans, and by structural alterations including liver necrosis as well as renal failure and necrosis [6, 15].

Erythropoictin (EPO) is a hypoxia-inducible hemopoictic growth factor expressed mainly in the kidney. It has an antioxidant property and a neuroprotective effect against ischemia in cell culture and animals [5, 18]. We previously demonstrated that the administration of EPO before ischemia protects the liver from ischemiareperfusion injury caused by pedicular clamping, as depicted by decreased blood values of liver enzymes, IL-2, and TNF- $\alpha$ , as well as attenuated liver tissue oxidation [23]. Considering that the ischemia-reperfusion injury actually is a process triggered by hypoxia, in the current study, we evaluated the effect of EPO against laparoscopy-induced oxidative injury, as compared with the L-Pre method.

#### Materials and methods

The current study was conducted with the approval of the University Ethical Committee. The experiment protocol was designed in accordance with the 1996 revised edition of *The Guide for the Care and Use of Laboratory Animals* published by the U.S. National Institutes of Health.

After overnight fasting with free access to tap water, 64 male Spraque–Dawley rats weighing  $280 \pm 15$  g were enrolled into the study and randomly assigned to one of the following groups, with 16 animals in each group. The control group was subjected to a sham operation without pneumoperitoneum. The laparoscopy group was subjected to 5 min of insufflation and 5 min of desufflation followed by 60 min of pneumoperitoneum. The EPO plus laparoscopy group was subjected to a subcutaneous injection of EPO as a single dose of 1,000 U/kg followed by 60 min of pneumoperitoneum.

#### Surgical technique

The rats were anesthetized with an intramuseular injection of ketamine, and a 1-em midline laparotomy was performed below the umbilicus. After that, a eatheter was placed into the peritoneum as the vehicle of pneumoperitoneum created in all the animals except the sham group using a technique described by Polat et al. [12]. Pneumoperitoneum was obtained for 60 min by connecting the other side of the tube to a  $CO_2$  insufflator (Nortech Model 3-315-00, Fribourg, Switzerland) fixed at 15 mmHg.

Before insufflation, the aforementioned incision was closed with a tight purse-string suture to prevent leakage. After 45 min of desufflation subsequent to the cessation of pneumoperitoneum, laparotomy was performed and blood samples were obtained from half of the rats via intraaortic cannulation, along with liver and kidney samples. The other half of the rats were observed for the 24-h reperfusion period. The aforementioned tissue and blood samples also were obtained after this period.

#### Biochemical analysis

The blood samples were centrifuged immediately at 3,000 rpm for 5 min. Alanine aminotransferase (ALT) and laetie dehydrogenase (LDH) values were measured with a Hitaehi 911 automatie analyzer (Boehringer, Mannheim, Germany).

Plasma concentrations of TNF- $\alpha$  were determined by enzymelinked immunoassay (ELISA) using the commercially available kits for rat ELISA TNF- $\alpha$  (R&D system, Quantkin M Murine, Minneapolis, MN, USA). Plasma malondialdehyde (MDA) levels were measured as an indicator of lipid oxidation using the thiobarbituric acid method of Ohkawa et al. [9].

The liver and kidney tissues were rinsed with ice cold saline and frozen at  $-70^{\circ}$ C until the assay. The tissues were homogenized with an Ultra Turrax homogenizer (IKA T18 basic, Wilmington, NC, USA). After the centrifugation of the homogenates, the supernates were removed and used for MDA analysis by the method described previously [9].

#### Statistical analysis

Data are expressed as mean  $\pm$  standard error of the mean. Values lower than 0.05 were accepted as statistically significant. The betweengroup parameters obtained were evaluated using the Kruskal–Wallis nonparametric test. The Mann–Whitney *U*-test was used to compare the data between the paired groups.

#### Results

The results in terms of plasma LDH, ALT, TNF- $\alpha$ , and MDA values as well as the liver and kidney tissue MDA levels are graphically displayed in Figs. 1 to 4.

#### Effect of laparoscopy

The comparison of the results between the sham and laparoscopy groups showed that laparoscopy produced significantly increased plasma levels of LDH, ALT, TNF- $\alpha$ , and MDA, as well as increased liver and kidney tissue MDA values, after 45 min of reperfusion. Whereas the ALT levels declined to the normal level 24 h after desufflation, the LDH levels remained still higher that of the sham group. Therefore, laparoscopy produced significant ischemia–reperfusion injury in terms of plasma and tissue oxidative markers, especially after 45 min of desufflation.

#### Effect of L-Pre on laparoscopy-induced oxidative injury

The addition of the preceding L-Pre procedure to laparoscopy produced significant amelioration in terms of



The ALT values of the groups

a

120

100 80

45 mm

75 60

30

15

0

TNF-alpha

Sham

micmol/l. 45



Fig. 2. The alanine aminotransferase (ALT) values of the groups after 45 min and 24 h of reperfusion. **a**  $p \le 0.05$  for the sham and laparoscopy groups. **b** p < 0.05for the L-Pre + laparoscopy and laparoscopy groups.



LPre+LaparoscopyEPO+Laparoscopy

Fig. 3. The plasma tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and maloudialdehyde (MDA) values of the groups after 45 min of reperfusion. a  $p \le 0.05$  for the laparoscopy versus L-Pre + laparoscopy and the erythropoietin (EPO) laparoscopy groups. **b** p < 0.05 for the L-Pre + laparoscopy versus the EPO + laparoscopy group.

decreased plasma levels of LDH, ALT, TNF-a, and MDA, as well as decreased liver and kidney tissue MDA values, after 45 min of desufflation. The results for ALT and LDH after 24 h of desufflation showed lower levels than in the laparoseopy group. However, although L-Pre produced a significant beneficial effect on oxidative injury, as compared with the laparoscopy group, the LDH and TNF- $\alpha$  values were higher than those of the sham group.

Laparoscopy

Groups

#### Effect of EPO on laparoscopy-induced oxidative injury

MDA

The administration of EPO before the pneumoperitoneum significantly decreased the plasma levels of LDH, TNF- $\alpha$ , and MDA, as compared with those of the laparoseopy group. However, the comparison of the results between the sham and EPO groups showed that the levels of LDH, ALT, TNF- $\alpha$ , and MDA still were higher than those of the sham group. Although EPO adminis-



The MDA levels of kidney and liver tissues of the groups

Fig. 4. The liver and renal tissue levels of malondialdehyde (MDA) in the groups. a p < 0.05 for the laparoscopy versus the sham group. b p < 0.05 for the laparoscopy versus the L-Pre + laparoscopy group.

tration during the prelaparoscopic period decreased the pneumoperitoneum-induced oxidative injury, the beneficial effect of L-Pre was more pronounced than that of EPO in terms of LDH, TNF- $\alpha$ , and MDA levels.

#### Discussion

Laparoscopy represents an ischemia-reperfusion phenomenon that produces splanchnic ischemia. Pneumoperitoneum is used to facilitate the visualization of intraabdominal organs intraoperatively. However, not only pneumoperitoneum itself, but also the deflation phase carries the risk of ischemic potential for splanchnic organs [22, 24]. Augmented productions of free oxygen radicals and other hazardous compounds occur during insufflation and also during the deflation phase. Therefore, laparoscopy itself is an ischemiareperfusion phenomenon that produces splanchnic ischemia and related adverse alterations in intraabdominal organs, which may have a particular importance, especially for elderly patients, who have limited functional capacity [14]. The liver and kidney are among the splanchnic organs affected by laparoscopic procedures.

Increasing recognition of the adverse effects from laparoscopy has led to different alternative approaches such as laparoscopy with low intraabdominal pressure or the gasless abdominal wall–lifting method. We have previously shown that L-Pre is a beneficial method for decreasing the laparoscopy-related oxidative injury [22, 24]. It also has been reported that L-Pre may be more effective than low-pressure pneumoperitoneum in reducing the ischemic insult associated with laparoscopy [3].

The L-Pre technique is performed by administering short periods of insufflation and deflation sequence(s) at the beginning of laparoscopy. It is an endogenous protective mechanism that prepares the tissue against subsequent laparoscopic injury. Some cellular proteins released during L-Pre are known to make the cell resistant to the harmful effect of free oxygen radicals delivered after the ischemia–reperfusion phase.

O'Malley et al. [10] previously reported that  $CO_2$ , the gas most frequently used for pneumoperitoneum, led to local and systemic acidosis, aggravating ischemic organ injury. However, an ideal gas for establishing pneumoperitoneum still is under investigation. It previously was found that helium seems to limit the postoperative oxidative response after laparoscopy [25]. The end result of laparoscopic injury is increased levels of inflammatory mediators, vasoactive kinins, fibrins, cytokines, and free oxygen radicals [19]. The lipid peroxidation induced by these mediators is associated with altered ischemic and hepatic enzymes. The plasma MDA values also reflect the oxidative modification of the lipid layer of the cell membrane.

Erythropoietin is a hypoxia-inducible growth factor expressed mainly in the kidney. It has multiple protective effects against oxidants and apoptosis [13]. We also reported previously that preischemic administration of EPO had protective effects on hepatic ischemiareperfusion injury [23]. It significantly decreased the postischemia-reperfusion elevations of ALT, TNF- $\alpha$ , IL-2, and tissue MDA in an experimental hepatic ischemia model in rats.

It was reported previously that preischemic administration of EPO was beneficial in reducing the ischemic injury to neuronal and myocardial tissues. We also reported recently that the preischemic administration of EPO in a dose of 1,000 U/kg prevented postischemic injury in rat kidney tissues. This beneficial effect was suggested to result from tyrosine kinase pathway activation because that effect was blocked by the administration of Genistein, which is a tyrosine kinase inhibitor [2]. Therefore, it may be speculated that EPO plays its role in laparoscopy-induced ischemia-reperfusion injury via tyrosine kinase enzyme activation, but this assumption still requires further experimental studies.

Because the eventual result of laparoscopy is related to organ ischemia, we tried in the current study to evaluate the effect of preischemic administration of EPO on laparoscopy-induced tissue injury, as compared with L-Pre. In this model, we evaluated ALT and LDH as contributors to organ parenchymal deterioration, plasma TNF- $\alpha$  as a cytokine mediator, and plasma, renal, and hepatic tissue MDA as lipid peroxidation markers. The comparison of the sham and laparoscopy groups showed that laparoscopy produced significantly increased levels of these parameters. Moreover, the harmful effect of laparoscopy continued partially 24 h after the deflation. However, the addition of L-Pre to laparoscopy significantly decreased the elevated parameters and almost declined them to sham group values. Although EPO administration significantly deereased the LDH, TNF-a, and MDA values, as compared with those of the laparoscopy group, they still were higher than those of the L-Pre plus laparoscopy group. Therefore, the beneficial effect L-Pre in terms of these parameters was more pronounced than that resulting from EPO administration. The results of this study show that EPO, an anti-ischemie agent, produces significant protection against laparoseopy-induced tissue injury. However, the beneficial effect of L-Prc was more significant than that of preischemically administered EPO. These results therefore suggest that L-Pre produced a protective effect against laparoscopy-related injury with the help of several mechanisms. It may be that L-Pre aets via meehanisms other than its antiischemic property, which is the most important protective mechanism of EPO. In conclusion, L-Pre and EPO decrease laparoscopic tissue injury probably with the help of different unique mechanisms.

- 1. Altindis M, Yilmaz S, Polat C, Serteser M (2004) Sequential periods of preconditioning decrease laparoscopy-related elevations in hepatic TNF- $\alpha$  and 1L-6 levels in rats. J Laparoendose Adv Surg Tech 6: 380–383
- Ates E, Yalein AU, Yilmaz S, Koken T, Tokyol C. Protective effect of erythropoietin on renal ischemia and reperfusion injury. ANZ J Surg (in press)
- Cevrioglu AS, Yilmaz S, Koken T, Tokyol C, Yilmazer M, Fenkei V (2004) Comparison of the effects of low intraahdominal pressure and ischemic preconditioning on the generation of oxidative stress markers and inflammatory cytokines during laparoscopy in rats. Hum Reprod 9: 2144–2151
- Colletti LM, Kunkel SL, Walz A, Burdick MD, Kunkel RG, Wilke CA, Strieter RM (1996) The role of cytokine networks in the local liver injury following hepatic ischemia–reperfusion in the rat. Hepatology 23: 506–514
- Grasso G (2001) Neuroprotective effect of recombinant human erythropoietin in experimental subarachnoid hemorrhage. J Neurosurg Sci 45: 7–14
- Kirsch AJ, Hensle TW, Chang DT, Kayton ML, Olsson CA, Sawczuk IS (1994) Renal effects of CO<sub>2</sub> insufflation: oliguria and acute renal dysfunction in a rat model. Urology 43: 453–459

- Licher CS (1997) Role of oxidative stress and antioxidant therapy in alcoholic and nonalcoholic liver disease. Adv Pharmacol 38: 601–628
- Murry CE, Jenning RB, Reimer KA (1986) Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. Circulation 74: 1124–1136
- Ohkawa H, Ohishi N, Yagi K (1979) Assay for lipid peroxidase in animal tissues hy thiobarbiturie acid reaction. Anal Biochem 95: 351
- O'Malley C, Cunningham AJ (2001) Physiologic changes during laparoscopy. Anesthesiol Clin North Am 19: 1–19
- Polat C, Aktepe OC, Akbulut G, Yilmaz S, Arikan Y, Dilek ON, Gokce O (2003) The effects of increased intraabdominal pressure on hacterial translocation. Yonsei Med J 44: 259–264
- Polat C, Arikan Y, Vatansev C, Akbulut G, Yilmaz S, Dilek FH, Gokee O (2002) The effects of increased intraabdominal pressure on colonic anastomoses. Surg Endosc 16: 1314–1319
- Sakanaka M, Wen TC, Matsuda S, Masuda S, Morishita E, Nagao M, Sasaki R (1998) In vivo evidence that erythropoietin protects neurons from ischemic damage. Proc Natl Acad Sci U S A 95: 635–640
- Sato K, Kawamura T, Wakusawa R (2000) Hepatic blood flow and function in elderly patients undergoing laparoscopic cholecystectomy. Anesth Analg 90: 1198–1202
- Schachtrupp A, Toens Ch, Hoer J, Klosterhalfen B, Lawong AG, Schumpelick V (2002) A 24-h pneumoperitoneum leads to multiple organ impairment in a poreine model. J Surg Res 106: 37–45
- Schafer M, Sagesser H, Reichen J, Krahenbuhl L (2001) Alterations in hemodynamics and hepatic and splanchnic circulation during laparoscopy in rats. Surg Endosc 15: 1197–1201
- Serracino-Inglott F, Habih NA, Mathie RT (2001) Hepatic ischemia-reperfusion injury. Am J Surg 181: 160–166
- Siren AL, Ehrenreich H (2001) Erythropoietin: a novel concept for neuroprotection. Eur Arch Psychiatry Clin Neurosci 251: 179–184
- Vittimherga FJ, Foley DP, Meyers WC, Callery MP (1998) Laparoscopic surgery and the systemic immune response. Ann Surg 227: 326–334
- Walker LM, York JL, Imam SZ, Ali SF, Muldrew KL, Mayeux PR (2001) Oxidative stress and reactive oxygen species generation during renal ischemia. Toxicol Sci 63: 154–148
- Windberger U, Siegl H, Ferguson JG, et al. (1995) Hemodynamic effects of prolonged addominal insufflation for laparoscopic procedures. Gastrointest Endosc 41: 121–129
- Yilmaz S, Ates E, Polat C, Koken T, Tokyol C, Akbulut G, Gokce O (2003) Ischemic preconditioning decreases laparoscopy-induced oxidative stress in small intestine. Hepatogastroenterology 50: 979–982
- Yilmaz S, Ates E, Tokyol C, Pehlivan T, Erkasap S, Koken T (2004) The protective effect of erythropoietin on ischemia-reperfusion injury of liver. HPB 3: 169–173
- Yilmaz S, Koken T, Tokyol C, Kahraman A, Akbulut G, Serteser M, Polat C, Gokce C, Gokce O (2003) Can preconditioning reduce laparoscopy induced tissue injury? Surg Endosc 17: 819–822
- 25. Yilmaz S, Polat C, Kahraman A, Koken T, Arikan Y, Dilek ON, Gokee O (2004) The comparison of the oxidative stress effects of different gases and intraabdominal pressures in an experimental rat model. J Laparoendose Adv Surg Tech. 3: 165–168

Surg Endose (2006) 20: 820–823 DOI: 10.1007/s00464-005-0369-1

© Springer Seienee+Business Media, Inc. 2006

SUNUIGA

and Other Interventianal Techniques

# Laparoscopic ventral hernia repair using expanded polytetrafluoroethylene-polyester mesh compound

M. Farrakha

General Surgery Department, Mafraq Hospital, P.O. Box 2951, Ahu Dhabi, United Arah Emirates

Received: 14 May 2005/Accepted: 25 August 2005/Online publication: 27 February 2006

#### Abstract

*Background:* Many biomaterials and composites have been used in clinical and experimental laparoscopic ventral hernia repair. The ideal prosthesis should allow firm binding to the abdominal wall without adhesion to the bowel.

*Methods:* A compound prosthesis was made by circumferentially suturing a Gore-Tex mesh as visceral interface to a smaller polyester mesh as parietal interface, and it was used in 46 laparoscopic ventral hernia repairs between January 2000 and December 2004.

*Results:* Average operative time was 65 min, with no intraoperative complications. Mean hospital stay was 2.2 days. Postoperative complications were five seromas, two hematomas, and one recurrence after a mean follow-up of 32.2 months.

*Conclusion:* The prosthesis used was made of two biomaterials that have been tested and tried over the years. The polyester layer is known to induce sufficient tissue ingrowth, whereas Gore-Tex minimizes adhesion formation. The memory of the compound was high enough to allow easy laparoscopic unrolling and handling.

Key words: Laparoscopy — Ventral — Incisional — Hernia — Repair

The published results of laparoscopic ventral hernia repair are satisfactory, with minimal perioperative morbidity and low early recurrence rates [3, 17, 19, 27, 28, 31, 32]. The type of mesh prosthesis used has been the matter of much debate. Firm binding to the abdominal wall with sufficient fibrocollagenous tissue ingrowth, on the one hand, and no or minimal liability to adhesion formation with underlying bowel, on the other hand, are the main characteristics of the ideal prosthesis required. Polypropylene and polyester biomaterials are known to form severe adhesions if put in direct contact with bowel, and Gore-Tex (ePTFE), which is known to have minimal liability to form adhesions, has raised concern about the adequacy of tissue ingrowth. The formation of a compound made of a Gore-Tex mesh and a polyester mesh will accomplish the two required benefits. In a previous article [14, 15], we presented our early experience with the use of polyester mesh and cPTFE soft tissue patch for bilayer laparoscopic repair of ventral hernia. Further refinement and modification of the technique in a larger number of patients is presented herein.

#### Materials and methods

A total of 46 patients (35 females and 11 males) with ventral hernias were treated hetween January 2000 and December 2004. Their age ranged from 23 to 75 years (median, 37.6). Thirty patients had ineisional hernia, 14 had spontaneous paraumhilieal hernias, and two had umbilieal and lower abdominal incisional hernias. In 14 patients, hernias were recurrent. The long axis of defect ranged from 4 to 12 em.

#### Technique

Following abdominal insufflation hy the use of a Veress needle introdueed away from any previous ineision, access to the abdominal cavity was obtained by using a 10-mm visiport (Tyeo Healthcare). A 30° telescope was used and two 5-mm ports were introduced under vision. Their sites were chosen to give proper orientation and suitable working length and angle. Hernia contents were reduced and all adhesions divided using either electrosurgical power or ultracision (Ethicon Endo-Surgery). The defect edges were clearly exposed and the extent of overpass was then marked on the skin (at least 4 em). Compound prosthesis was made by suturing a Gore-Tex mesh (W. L. Gore and Associates, Flagstaff, AZ, USA) to a thin polyester mesh using multiple 2-0 sutures passed through the two meshes 2 cm from the edge of the Gore-Tex mesh. Excess polyester mesh was then eut 1-1.5 em short of the edge of the Gore-Tex mesh to eliminate the possibility of bowel adhesions to the edge of the prosthesis. Four prolene loops were left in each of the borders of the prosthesis. The prosthesis was rolled and introduced into the abdominal eavity. A 10 × 15-em prosthesis was passed through a 10-mm port. The port was removed to introduce a larger prosthesis through the port site. Unrolling and spreading eould readily he obtained due to the higher memory acquired hy the addition of the polyester layer to the ePTFE layer. The prosthesis spread on the abdominal viscera with the polyester surface facing the defect was raised to the abdominal wall by anchoring the prolene loops consecutively to a suture passer introduced through the abdominal wall at the proposed site of overpass to ensure proper orientation of the prosthesis across the defect. The loops were held hy artery forceps followed hy lixation of prosthesis using helical coils from a 5-mm Protack (Tyco Healthcare) aided hy counterpressure on the abdominal wall to face the tip of the Protack and compensate for its lack of angulation. Prolene loops were then cut and removed. This technique aided in proper orientation of the opaque compound across the defect. Carc should be taken during fixation of the prosthesis to the defect border on the same side of the ports. Reverse movement of instruments along the eve-hand-target axis may lead to lixation of the prosthesis short of the required overpass. One 5-mm port may oceasionally be required to allow proper overpass and fixation. A 10 × 15-cm prosthesis was used in 38 cases, and a  $15 \times 20$ -cm prosthesis was used in seven eases. In one other patient, two 10 × 15-cm prostheses were required to cover multiple midline defects from umhilical and lower abdominal incisional hernia. Pressure dressing was then applied on the hernia sae.

#### Results

The technique was successfully completed in all patients. Two patients had laparoseopie eholeeysteetomy performed during the same session, and one patient had interval appendicectomy with repair of a hernia seeondary to drainage of appendicular abseess. Operative time ranged from 30 to 85 min (average, 65), with no intraoperative complications apart from troear injury of the left inferior epigastric artery in one ease. All but one patient tolerated oral fluids the next morning following surgery. Abdominal CT sean was occasionally required during follow-up to rule out recurrence. Postoperative complications were five seromas that were relatively small and resolved spontaneously. Two patients had persistent hematoma at the hernia sae. One small, firm, and lobulated hematoma was suspected for recurrence and excluded by ultrasonography. In another patient, hematoma was large and surrounded by bluish skin discoloration. The patient was conservatively treated and discharged 9 days following surgery. In this case, we easily inverted the hernia sac into the abdomen after lyses and reduction of hernia contents. The sac was excised by electroeautery with a pair of seissors. Division of the hernia sae should be avoided. One patient had early postoperative recurrence. The patient had an epigastrie incisional hernia following coronary bypass. Recurrence was detected 2 months postoperatively at the upper end of the delect. The patient had ehronie persistent cough and low ejection fraction. Reoperation was denied by the anesthetists. Abdominal CT sean was selectively used to differentiate recurrence from postoperative seroma or hematoma at the hernia site. Hospital stay ranged from 1 to 9 days (mean, 2.2). No other late complications occurred in this series after a mean follow-up of 32.2 month (range, 6-53).

#### Discussion

Selection of the mesh prosthesis has been the subject of much concern among surgeons performing laparoscopie repair of ventral hernias. Firm binding to the abdominal wall with sufficient fibrocollagenous tissue ingrowth, on the one hand, and no or minimal liability to adhesion formation with underlying bowel, on the other hand, are the characteristics of the ideal prosthesis for the laparoseopie approach to ventral hernias.

ePTFE is the biomaterial selected by most researchers as an intraperitoneal prosthesis due to its known minimal liability to form adhesions with the bowel [24, 25]. Dual mesh is a form of ePTFE with different pore sizes on either surface that was recently introduced to enhance the tissue integration of an otherwise inert biomaterial. It has been used by many surgeons [3, 10, 13]. Despite the known effect on adhesion formation, polypropylene mesh has been used by some authors in the intraperitoneal position [16, 20]. The very low recurrence rate of 1.1% reported by Franklin et al. [16] was attributed to the superior tissue ingrowth with the polypropylene mesh. Two-thirds of their patients who had relaparoseopy had mild to severe adhesions to the mesh.

The characteristics of polypropylene and Gore-Tex biomaterials have been compared in many experimental studies. Whereas ePTFE led to the formation of thin, flimsy adhesions and a thin connective tissue capsule covering, polypropylene resulted in more dense adhesions and disorganized dense connective tissue covering. Resistance to traction was shown to be higher with polypropylene [6, 12, 35]. Abdominal wall defects repaired with ePTFE resulted in significantly more frequent herniations, observed at the faseia and patch interface, than with polypropylene (60 vs 0%). Insulfieient ingrowth of fibroeollagenous tissue into the ePTFE patch, with insufficient anchorage of the patch to the fascia, was histologically confirmed, whereas the polypropylene mesh was completely incorporated into libroeollagenous tissue, which was continuous with the adjacent fascia [23, 29]. The effect of the structure of three ePTFE prostheses (Soft Tissue Patch, Dual Mesh, and Myero-Mesh) on the searring process in an abdominal wall defect experimental model was studied. The three types of PTFE prosthesis induced a low ineidence of adhesion formation between biomaterial and viscera. The integration mechanisms of the three prostheses were similar and culminated in the eneapsulation of the PTFE by the neoformed tissue with a macrophage response similar to that of any reparative process in the absence of biomaterial [5, 8].

Conflicting results led to continued studies to find the proper prosthetic biomaterial. Eliminating adhesions to mesh by mechanical or other means during a eritical time (first 7 days) is assumed to control adhesions to the mesh and subsequent mesh-related complications. Laparoseopic adhesiolysis at 1 week was shown to minimize subsequent adhesion formation to polypropylene and ePTFE mesh during a 4-month follow-up in an experimental setup [26].

Composite prosthesis made by integrating a biodegradable extracellular matrix analog or an absorbable mesh barrier with a permanent structural biomaterial to reduce adhesion formation was tried in both experimental and elinical studies Barie et al. [4] used a polyester/polyglactin 910 (Vicryl) composite. This composite led to intestinal adhesions and fistulation in both clinical and experimental studies [30, 34]. Polypropylene mesh/ polypropylene sheeting and polypropylene mesh/silastic composites were shown to reduce adhesion formation to bowel compared to other biomaterials, including polypropylene, polyester, expanded polytetrafluoroethylene, polypropylene/polyglactin, polypropylene/polyglycolic acid, and polypropylene/fibrin [2]. Although silastic implantation (Dacron-reinforced silicone rubber) caused no adhesions, graft extrusion and evisceration were common [22]. polypropylene/Gelfilm composite lead to a similar adhesion response as that of PP alone after the gelatin dissolved in 1 week. Sepramesh (a polypropylene coated with sodium hyaluronate), carboxymethyl cellulose film, Parietex mesh (a polyester/collagen composite), and polypropylene collagen composite have similar effects in minimizing adhesion formation [1, 7, 9, 18, 21, 33]. However, in one study infection was more prevalent with Parietex composite mesh, with concurrent increased mesh surface covered by adhesions [33].

The addition of a layer of polyester to the inert ePTFE mesh would make the repair stronger and better longterm results would then be expected. The facial interface layer of polyester mesh stimulates fibrocollagenous tissue ingrowth, leading to firm binding of the prosthesis to the abdominal wall, whereas the ePTFE layer minimizes the possibility of adhesion formation to the underlying bowel. The fixation of the two biomaterials was found to modify its physical characteristics, making its unfurling and spreading in the operative field easier than with the Gore-Tex biomaterial alone [14]. With proper orientation and firm fixation to healthy surrounding tissue, an ideal repair is expected. Repair of hernias near bony prominences, such as the xiphoid process or symphesis pubis, requires special attention. Effort should be made to reduce all adherent hernia contents to avoid persistent postoperative swelling that may be mistaken for recurrence, leading to unsatisfactory results for the patient. Very large prosthesis should have reasonable tension to avoid postoperative bulge and suspected recurrence.

The biomaterials used in this technique are currently available, and their combined use in our technique did not increase difficulties or cost compared to the use of other composite biomaterials that still require extensive experimental and clinical studies. Fixation of the prosthesis with helical coils from a 5-mm Protack instrument (Tyco Healthcare) was found to be reliable and technically feasible. Tissue ingrowth will lead to further, lasting integration of the prosthesis with the abdominal wall. Transabdominal sutures used for mesh fixation in laparoscopic ventral hernia repair were reported to be an occasional source of prolonged postoperative pain requiring postoperative local anesthetic injection [11]. Neither early nor late postoperative pain specifically related to tacks occurred in this series.

#### Conclusion

Laparoscopic ventral hernia repair has many advantages over the open technique. It enables dissection of all intraabdominal adhesions commonly present in incisional hernia cases, clearly exposes healthy edges of the hernia defect and the presence of multiple defects, and provides the ability to fix a large prosthesis to accomplish adequate overpass, thus reducing the incidence of recurrence. The technique allows for other intraabdominal pathologies to be dealt with without the need for large or multiple incisions. Morbidity is thus reduced and hospital stay and recovery period are shortened. Adoption of the laparoscopic approach should not compromise the principles of successful mesh repair of ventral hernia. The prosthetic biomaterial should stimulate sufficient tissue ingrowth to restore abdominal wall integrity with a minimal tendency for adhesion formation. A compound prosthesis, ePTFE-polyester, was formed and successfully used in this series. It combines the benefits and the known, tested characteristics of its two components. In addition, polyester biomaterial modifies the memory of Gore-Tex, making its manipulation in the operative field easier. Clear exposure of the defect edges, proper orientation of the prosthesis, sufficient overpass, firm fixation to the abdominal wall, and division of all adhesions are essential requirements for a successful repair.

- Alponat A, Lakshminarasappa SR, Yavuz N, Goh PM (1997) Prevention of adhesions by Seprafilm, an absorbable adhesion barrier: an incisional hernia model in rats. Am Surg 63: 818– 819
- Amid PK, Shulman AG, Lichtenstein IL, Sostrin S, Young J, Hakakha M (1994) Experimental evaluation of a new composite mesh with the selective property of incorporation to the abdominal wall without adhering to the intestines. J Biomed Mater Res 28: 373–375
- Aura T, Habib E, Mekkaoui M, Brassier D, Elhadad A (2000) Laparoscopic tension-free repair of anterior abdominal wall incisional and ventral hernias with an intraperitoneal Gore-Tex mesh: prospective study and review of the literature. J Laparoendosc Adv Surg Tech A 12: 263–267
- Barie PS, Mack CA, Thompson WA (1995) A technique for laparoscopic repair of herniation of the anterior abdominal wall using a composite mesh prosthesis. Am J Surg 170: 62–63
- Bellon JM, Bujan J, Contreras LA, Carrera-San Martin A, Jurado F (1996) Comparison of a new type of polytetrafluoroethylenc patch (Mycro Mesh) and polypropylene prosthesis (Marlex) for repair of abdominal wall defects. J Am Coll Surg 183: 11–18
- Bellon JM, Contreras LA, Bujanm J, Carrera San Martin A (1997) The use of biomaterials in the repair of abdominal wall defects: a comparative study between polypropylene meshes (Marlex) and a new polytetra-fluoroethylene prosthesis (Dual Mesh). J Biomater Appl 12: 121–135
- Besim H, Yalcin Y, Hamamci O, Arslan K, Sonisik M, Korkmaz A, Erdogan S (2002) Prevention of intraabdominal adhesions produced by polypropylene mesh. Eur Surg Res 34: 239–243
- Bujan J, Contreras LA, Carrera-San Martin A, Bellon JM (1997) The behavior of different types of polytetrafluoroethylene (PTFE) prostheses in the reparative scarring process of abdominal wall defects. Histol Histopathol 12: 683–690
- Butler CE, Navaro FA, Orgill DP (2001) Reduction of abdominal adhesion using composite collagen–GAG implants for ventral hernia repair. J Biomed Mater Res 58: 75–80
- Carbajo MA, del Martin Olmo JC, Blanco J (2000) What is the appropriate mesh for laparoscopic intraperitoneal repair of abdominal wall hernia? Surg Endosc 14: 408
- Carbonell AM, Harold KL, Mahmutovic AJ, Hassan R, Matthews BD, Kercher KW, Sing RF, Heniford BT (2003) Local injection for the treatment of suture site pain after laparoscopic ventral hernia repair. Am Surg 69: 688–692

- Christoforoni PM, Kim YB, Preys Z, Lay RY, Montz FJ (1996) Adhesion formation after incisional hernia repair: a randomized porcine trial. Am Surg 62: 935–938
- DeMaria EJ, Moss JM, Sugerman HJ (2000) Laparoscopic intraperitoneal polytetrafluorocthylene (PTFE) prosthetic patch repair of ventral hernia. Prospective comparison to open prefascial polypropylene mcsh repair. Surg Endose 14: 326–329
- Farrakha M (2000) Laparoscopic treatment of ventral hernia. A bilayer repair. Surg Endose 14: 1156–1158
- Farrakha M (2001) Laparoscopic treatment of ventral hernia. Bilayer repair (Reply to Comment). Surg Endosc 15: 1243
- Franklin ME, Dorman JP, Glass JL, et al. (1998) Laparoscopic ventral and incisional hernia repair. Surg Laparosc Endose 8: 294– 299
- Goodney PP, Birkmeyer CM, Birkmeyer JD (2002) Short-term outcomes of laparoscopic and open ventral hernia repair: a metaanalysis. Arch Surg 137: 1161–1165
- Greenawalt KE, Butler TJ, Rowe EA, Finneral AC, Garlick DS, Burns JW (2000) Evaluation of sepramesh biosurgical composite in a rahhit hernia repair model. J Surg Res 94: 92–98
- Heniford BT, Ramshaw BJ (2000) Laparoscopic ventral hernia repair: a report of 100 consecutive cases. Surg Endosc 14: 419–423
- Holzman MD, Purut CM, Reintgen K, Eubanks S, Pappas TN (1997) Laparoscopic ventral and incisional hernioplasty. Surg Endose 11: 32–35
- Hooker GD, Taylor BM, Driman DK (1999) Prevention of adhesion formation with use of sodium hyaluronate-based bioresorbable membrane in a rat model of ventral hernia repair with polypropylene mesh—a randomized, controlled study. Surgery 125: 211–216
- Jenkins SD, Klamer TW, Parteka JJ, Condon RE (1983) A eomparison of prosthetic materials used to repair abdominal wall defects. Surgery 94: 392–398
- Law NW (1990) A comparison of polypropylene mcsh, expanded polytetrafluoroethylene patch and polyglycolic acid mesh for the rcpair of experimental abdominal wall defects. Acta Chir Seand 156: 759–762

- 24. LeBlanc KA, Bellanger D, Rhynes KV 5th, Baker DG, Stout RW (2002) Tissue attachment strength of prosthetic meshes used in ventral and ineisional hernia repair. A study in the New Zealand White rabbit adhesion model. Surg Endose 16: 1542–1546
- Le Blank KA, Booth WV (1992) Laparoscopic repair of incisional abdominal hernia using expanded polytetraflouroethylene: preliminary findings. Surg Laparose Endose 3: 39–41
- Matthews BD, Pratt BL, Backus CL, Kereher KW, Heniford BT (2002) Comparison of adhesion formation to intra-abdominal mesh after laparoscopic adhesiolysis in the New Zealand White rabbit. Am Surg 68: 936–940
- Park A, Gagner M, Pomp A (1996) Laparoseopie repair of large ineisional hernia. Surg Laparose Endosc 6: 123–128
- Robbins SB, Pofahl WE, Gonzalez RP (2001) Laparoscopic ventral hernia repair reduces wound complications. Am Surg 67: 896–900
- Simmermacher RK, Sehakenraad JM, Bleichrodt RP (1994) Reherniation after repair of the abdominal wall with expanded polytetrafluoroethylene. J Am Coll Surg 178: 613–616
- Solar N, Verhaeghe P, Essomba A, Sevestre H, Stoppa R (1993) Treatment of postoperative incisional hernias by composite prosthesis (polyester–polyglactin 910). Clinical and experimental study. Ann Chir 47: 589–608
- Stoppa R, Louis D, Verhaeghe P, Henry X, Plachot JP (1987) Current treatment of postoperative eviscerations. Int Surg 72: 42–44
- Thomas DS, Phillips EH (2002) Current status of laparoscopic ventral hernia repair. Surg Endosc 16: 939–942
- 33. van 't Riet M, de van Vos Steenwijk PJ, Bonthuis F, Marquet RL, Steyerberg EW, Jeekel J, Bonjer HJ (2003) Prevention of adhesion to prosthetic mesh: comparison of different barriers using an ineisional hernia model. Ann Surg 237: 123–128
- Vrijland WW, Bonthuis F, Steyerberg EW, Marquet RL, Jeekel J, Bonjer HJ (2000) Peritoneal adhesions to prosthetic materials: choice of mesh for incisional hernia repair. Surg Endose 14: 960– 963
- Walker AP, Henderson J, Condon RE (1993) Double-layer prostheses for repair of abdominal wall defects in a rabbit model. Surg Res 55: 32–37

Surg Endosc (2006) 20: 824–829 DOI: 10.1007/s00464-005-0334-z

© Springer Science+Business Media, Inc. 2006

SUNUIDA/

and Other Interventional Techniques

# Robotic surgery and training: electromyographic correlates of robotic laparoscopic training

T. N. Judkins,<sup>1</sup> D. Oleynikov,<sup>2</sup> K. Narazaki,<sup>1</sup> N. Stergiou<sup>1</sup>

<sup>1</sup> University of Nebraska at Omaha, 6001 Dodge Street, Omaha, NE 68182, USA
 <sup>2</sup> University of Nebraska Medical Center, 983280 Nebraska Medical Center, Omaha, NE 68198-3280, USA

Received: 1 May 2005/Accepted: 25 October 2005/Online publication: 27 February 2006

#### Abstract

*Background:* Robotic laparoscopic surgery has been shown to decrease task completion time, reduce errors, and decrease training time, as compared with manual laparoscopic surgery. However, current literature has not addressed the physiologic effects, in particular muscle responses, to training with a robotic surgical system. The authors seek to determine the frequency response of electromyographic (EMG) signals of specific arm and hand muscles with training using the da Vinci Surgical System.

*Methods:* Seven right-handed medical students were trained in three tasks with the da Vinci Surgical System over 4 weeks. These subjects, along with eight control subjects, were tested before and after training. Electro-myographic (EMG) signals were collected from four arm and hand muscles during the testing sessions, and the median EMG frequency and bandwidth were computed.

*Results*: The median frequency and frequency bandwidth both were increased after training for two of the three tasks.

*Conclusion:* The results suggest that training reduces muscle fatigue as a result of faster and more deliberate movements. These changes occurred predominantly in muscles that were the dominant muscles for each task, whereas the more demanding task recruited more diverse motor units. An evaluation of the physiologic demands of robotic laparoscopic surgery using electromyography can provide us with a meaningful quantitative way to examine performance and skill acquisition.

Key words: da Vinci — Electromyography — Frequency analysis — Laparoscopy — Robotic surgery — Training Laparoscopy, a form of minimally invasive surgery, has revolutionized the treatment of abdominal pathologies. The benefits for patients include shorter recovery time with less pain [23], fewer adhesions [10], and better postoperative quality of life [14] than experienced with traditional open procedures. However, manual laparoscopy also has shown several limitations during surgery. These limitations include lack of depth perception, poor camera control, limited degrees of freedom at the instrument tips, and inverted hand–instrument movements [2, 3, 15]. Furthermore, these limitations lead to unnatural and painful surgical postures that result in fatigue for the surgeon [2, 3, 15].

The advent of robotic surgical systems, such as the da Vinci Surgical System (dVSS; Intuitive Surgical, Inc., Sunnyvale, CA, USA), have overcome some of the limitations associated with manual laparoscopy. The addition of three-dimensional visualization has provided depth perception [6] and increased dexterity [16, 17]. Wristlike articulations of the instruments also have been shown to improve surgeons' dexterity [16]. Coordinated hand–instrument movements have reduced the training time for robotic systems versus manual laparoscopy [5]. In addition, tremor abolition and motion scaling have been shown to enhance dexterity with the use of robotic systems, as compared with manual techniques [16].

Studies comparing robotic surgical systems with manual laparoscopy have examined the effectiveness of robotic procedures. Researchers have found that with the use of robotic systems, surgeons improve in dextcrity [16] and residents can be trained more quickly [5, 7, 12]. In addition, there are fewer errors, and the time required for task completion is reduced [11, 13, 16, 20, 22]. However, the current means of evaluating surgical performance and skill acquisition during training are limited to measurement of task completion time and the number of errors [11, 13, 16, 20, 22], or to subjective evaluation by an expert.

Correspondence to: N. Stergiou

To our knowledge, no studies have examined physiologic measures of surgeons during the performance of robotic surgical techniques. In manual laparoscopy, physiologie evaluations have been limited mainly to ergonomie measures [2, 3, 15]. These studies have found increased stress and fatigue associated with manual laparoscopy because of the surgeons' postures. Additionally, Quiek et al. [21] investigated electromyography during manual laparoscopy and found that both the task and the type of grasper used contributed to musele fatigue. This type of analysis has not yet been eonducted for robotic surgery.

Thus, it is not known whether robotic surgery leads to increased fatigue. Most importantly, it is not known whether training has an effect on fatigue. Electromyography can assist in the evaluation of physiologic muscular fatigue [1]. Specifically, frequency analysis of the electromyographic signals from the muscles involved has proved to be an effective method for measuring muscle fatigue and motor unit reeruitment during static force exertion [1]. Specifically, increased muscle fatigue is associated with a decreased median frequency of the power spectrum [1, 4]. In addition, increases in the frequency bandwidth of the power spectrum imply additional recruitment of motor units with varying conduction velocities [8, 9].

More recently, it has been shown that frequency analysis also can be applied to cyclic dynamic tasks to evaluate muscular fatigue [4]. Hypothetieally, training should decrease muscular exertion (i.e., median frequency) during surgery and thus improve the ability of the surgeon to operate. Therefore, this study sought to determine how muscle responses change during training with the dVSS.

#### Methods and materials

#### Subjects

For this study, 15 novice dVSS users consisting of first- and secondyear medical students (11 men and 4 women, ages  $25.3 \pm 2.6$  years) at the University of Nehraska Medical Center (UNMC) were recruited. All the participants were right-handed. Informed consent was obtained from each subject before participation in accordance with the Institution Review Board of UNMC.

#### Tasks

Subjects performed or practiced three tasks throughout this study: himanual carrying, needle passing, and suture tying. The bimanual carrying task required that two 15 × 2 mm ruhber pieces (one each with left and right graspers) be picked up simultaneously from 30 mm (diameter) metal caps and placed in two other metal caps 50 mm away (Fig. 1A). The caps were arranged in a square configuration such that the left graspers removed pieces from the top left cap and placed them in the hottom left cap. The right grasper removed pieces from the bottom right cap and placed them in the top left cap. The subject repeated the movement six times in succession. The needle-passing task required that a 26 mm surgical needle he passed through six holes in a latex tube (Fig. 1B). The subjects started from the proximal holes and proceeded in order to the distal holes. The suture tying task required that a  $150 \times 0.5$  mm surgical suture he passed through one of the holes in the latex tube and that three knots be tied using the intracorporeal knot (Fig. 1C). All three tasks were cyclic tasks designed to mimic



Fig. 1. Experimental tasks. A Bimanual carrying, B Needle passing. C Suture tying.

actual laparoscopic tasks that require significant himanual coordination.

#### Experimental protocol

Before performing the experiment, the subjects were randomly assigned to one of two groups: a training (7 subjects) or a control (8 subjects) group. The experiment was performed over a period of 4 weeks and consisted of one pretraining test (PRE), six training sessions, and one posttraining test (POST). The control subjects did not perform the six training sessions.



Fig. 2. Median frequency and bandwidth for idealized power spectrum. Median frequency is the frequency at half of the total power. Frequency handwidth is the difference between the maximum and minimum frequency when the power is greater than 1/2 the maximum power. In this figure, the maximum power is 1.

#### Pretraining test

During the first week, the subjects performed one pretraining test. At the beginning of the test, the subjects were given a verbal explanation of each task and allowed to practice the three tasks for 5 min at their own pace. During the practice, the subject could ask questions and receive further explanation and suggestions from the investigators. After the practice, the subjects performed one trial for each task while electromyographic signals were recorded.

#### Training sessions

Within 3 days after the pretraining test, the subject started the training sessions, which were scheduled on a per subject basis. Each training session lasted no more than 45 min, and each task was practiced three to four times. During practice, the subjects were allowed to ask questions and receive verbal explanation and suggestions from the investigators. At the end of the session, the subjects performed one trial for each task while electromyographic signals were recorded.

#### Posttraining test

During the last week and within 3 days after the sixth training session, a posttraining test was performed with the same procedures as in the pretraining session.

#### Electromyography

The electromyographic data were collected using DelSys Bagnoli (Delsys, Inc., Boston, MA, USA) surface electromyography (EMG) and sampled at 1,000 Hz through the PEAK Motus (Peak Performance Technologies, Englewood, CO, USA) data acquisition system. Surface electrodes were placed over the bellics of the following four muscles as described by Basmajian and Deluca [1]: flexor carpi radialis, extensor digitorum, biceps brachii, and triceps brachii. We chose the flexor carpi radialis as a primary wrist flexor muscle, the extensor digitorum as a primary wrist extensor muscle, the biceps brachii as a primary elbow flexor muscle, and the triceps brachii as a primary elbow extensor muscle, all of which are superficial and can be monitored by a surface EMG system. Although many other types of movements (e.g., flexion and extension of thumb, index, and middle fingers; forcarm pronation and supination), and thus many other muscles, are involved, we assumed that the contributions of these four muscles in



**Fig. 3.** EMG median frequency. **A** Bimanual carrying task. **B** Suture tying task. A diamond indicates a significant difference (p < 0.05) between PRE and POST. Error bars arc two standard errors.

the three tasks are considerably high, and thus that measuring the EMG activities of these muscles was important for the purpose of this study.

#### Data analysis

Frequency power spectrums for the signals of each muscle during each task were calculated using fast-Fourier transforms. The median frequency and bandwidth were calculated to measure the frequency response of each muscle during all three tasks for the PRE and POST conditions. The median frequency was calculated as

$$\int_0^{f_{\text{med}}} P(f) \, df = \int_{f_{\text{med}}}^{f_{\text{max}}} P(f) \, df$$

where  $f_{med}$  is the median frequency,  $f_{max}$  is the maximum frequency of the power spectrum (500 Hz), and P(f) is the power at a frequency f (Fig. 2).

Bandwidth was calculated by determining the difference between the minimum and maximum frequencies when the power exceeded one-half of the maximum power of the frequency spectrum (Fig. 2). All data were processed in MATLAB (The MathWorks, Inc., Concord, MA, USA).

#### Statistical analysis

A mixed two-factor (group by condition) analysis of variance with repeated measures on the condition factor was used to compare the group means of the median frequency and bandwidth for each muscle and for cach task. All statistical analyses were performed using SPSS 12.0 (SPSS, Chicago, IL, USA).

#### Results

#### Bimanual carrying

Significantly larger median frequencies were found after training for the flexor carpi radialis (p = 0.046), extensor digitorum (p = 0.033), and triceps brachii (p = 0.030) muscles in both groups (control and training; Fig. 3, left). The flexor carpi radialis had a median frequency of 136.6  $\pm$  7.3 Hz for PRE and 151.8  $\pm$  7.3 Hz for POST. The extensor digitorum had a median frequency of 89.6  $\pm$  5.2 Hz for PRE and 96.1  $\pm$  5.2 Hz for POST. The triceps brachii had a median frequency of 88.4  $\pm$  4.5 Hz for PRE and 98.7  $\pm$  4.5 Hz for POST. No significant differences were found in the median frequency between the two groups. No significant differences were found for the frequency bandwidth.





#### Needle passing

No significant differences were found for the median frequency across the two conditions for the two groups. A significant increase in frequency bandwidth was found after training for the extensor digitorum muscle (PRE, 79.9  $\pm$  10.7 Hz; POST, 104.0  $\pm$  10.7 Hz; p = 0.017) in both groups (Fig. 4, left). No significant differences were found between the two groups.

#### Suture tying

A significant increase in median frequency was found after training for the flexor carpi radialis muscle (PRE,  $135.3 \pm 6.3$  Hz; POST, 149.6  $\pm 6.3$  Hz; p = 0.021) in both groups (Fig. 3, right). A significant increase in frequency bandwidth also was found after training for the flexor carpi radialis muscle (PRE, 119.6  $\pm$  14.0 Hz; POST,  $150.9 \pm 14.0$  Hz; p = 0.023) in both groups (Fig. 4 right). No significant main effects (PRE/POST or group) were found for the frequency bandwidth. However, a significant interaction effect was found for the biceps brachii (p = 0.015; Fig. 5). For the control group, the bandwidth was found to be 84.7  $\pm$  15.0 Hz for PRE and 53.4  $\pm$  15.0 Hz for POST. For the training group, the bandwidth was found to be  $48.3 \pm 16.1 \text{ Hz}$ for PRE and 92.3  $\pm$  16.1 Hz for POST. This result indicated a differential response between the two groups for this task and for this muscle.

#### Discussion

Previous research has shown that robotic laparoscopic surgery is beneficial for both the patient and the surgeon. Patients recover faster with less pain [10, 14, 23], and surgeons can perform tasks faster [11, 13, 17–20, 22] and can be trained more easily [5, 7, 12] than with manual laparoscopy. However, the physiologic impacts of robotic surgery are not clearly understood.

In the current study, training with the dVSS generally increased the median frequency of the EMG power spectrum in the performance of bimanual tasks. This frequency shift has two likely causes. According to the available literature [11, 13, 17–20, 22], training results in shorter task completion times. Therefore, in the current study, each movement within a task occurred at a higher frequency, allowing completion of the task in a shorter time. These faster movements resulted in muscle bursts occurring at a higher frequency. In addition, findings have shown that a decrease in median frequency is associated with muscle fatigue during sustained contraction (for review, see [1]). Training may increase efficiency in the performance of a task, thereby reducing muscle fatigue. One or both of these factors resulted in the frequency shift.

Findings also showed that this frequency shift occurred independently of the subject's group (control or training). A possible explanation of this result is that the control group also had some training effects. Our previous pilot work showed that decreases in task completion time are possible even after one training session. Thus, it is not surprising that the control group also performed the tasks faster in just their second session (posttraining). Because EMG was not collected during the training sessions, this cannot be confirmed.

The frequency content of electromyograms also is directly related to the conduction velocities of recruited motor units [9]. It is well known that type 1 and type 2 muscle fibers have different conduction velocities [8]. Therefore, it can be inferred that a larger frequency bandwidth is indicative of a larger range of recruited motor units with different conduction velocities (i.e., different muscle fibers). We found that the EMG bandwidth had increased during the needle passing (extensor digitorum) and suture tying (flexor carpi radialis) tasks after training, as compared with pretraining. Furthermore, during the suture tying task, EMG bandwidth for the biceps brachii had increased after training for the training group, and had decreased after training for the control group, as compared with pretraining. These results suggest that subjects recruited a larger range of motor units with varying conduction velocities during training.

However, the bimanual carrying task showed no significant changes in EMG bandwidth. Because the bimanual carrying task requires more simplistic movements, as compared with the more complex the needle passing and suture tying tasks, the larger range of conduction velocities may be associated with more complex movements. It is possible that different motor units are recruited during different phases of the task depending on the precision needed for each phase. For example, looping the suture during the suture tying task may recruit different motor units than pulling the suture taut to finish the knot. Further analyses of the phases of the tasks are necessary to confirm this hypothesis and identify proper training tasks.

The findings also showed that when a significant difference in median frequency or bandwidth was found, it occurred in muscles that would be used predominantly for the given task. During the bimanual carrying task, the median frequency increased for the muscles used for grasping (flexor carpi radialis) and arm movements (biceps brachii and triceps brachii). Likewise, the median frequency for the flexor carpi radialis had increased after training during the suture-tying task, a grasping intensive task. Although care was taken to minimize electrode placement, future studies will monitor muscle 828



Fig. 5. Bandwidth interaction effects of biceps brachii during suture tying task. The control group decreased bandwidth POST-training while the training group increased bandwith POST-training. Error bars are two standard errors.

responses over a single session to confirm that changes in median frequency are attributable to reduced muscle fatigue and not other factors.

Bandwidth increases were found after training for the extensor digitorum during the needle passing task and for the flexor carpi radialis and biceps brachii during the suture tying task. For the needle passing task, a "hooking" motion is used to pass the needle through the latex tube, requiring more precise control of the fingers. Therefore, it is possible that different motor units may be recruited in the extensor digitorum to control the movement. The suture tying task requires fine control of grasping as well as looping of the suture for tying the knot. Therefore, bandwidth increases for the flexor carpi radialis and biceps brachii would be expected. It is worth mentioning that such detail in the examination of the different surgical tasks and the muscles involved will not be possible without the acquisition of physiologic measures, specifically electromyography. This verifies that future studies need to use our methodology to further establish learning and proficiency criteria for robotic surgical performance.

In conclusion, surgical training increased the median frequency response as a result of increased speed, reduced fatigue, or both during each task. These increases occurred predominantly in muscles that were the dominant muscles for each task. Additionally, the more demanding task recruited more diverse motor units, as shown by increases in bandwidth after training. It is cvident that an evaluation of the physiologic demands of robotic laparoscopic surgery using electromyography can provide us with a meaningful quantitative way to examine performance and skill acquisition. In our future work, we plan to include a group of expert surgeons and more realistic tasks (i.e., data collection during actual robotic laparoscopy) to further understanding of the physiologic demands of robotic surgery. We also propose that future studies should record EMG from both arms simultaneously for direct determination of bimanual coordination.

Acknowledgments. This research was funded by a grant awarded to Drs. Stergiou and Oleynikov by the Nebraska Research Initiative.

- Basmajian JV, De Luca CJ (1985) Muscles alive, their functions revealed by electromyography. Williams & Wilkins, Baltimore, MD
- Berguer R, Rab GT, Abu-Ghaida H, Alareon A, Chung J (1997) A comparison of surgeons' posture during laparoscopic and open surgical procedures. Surg Endose 11: 139–142
- Berguer R, Smith WD, Chung YH (2001) Performing laparoscopic surgery is significantly more stressful for the surgeon than open surgery. Surg Endosc 15: 1204–1207
- Bonato P, Roy SH, Knaflitz M, Luca CJD (2001) Time-frequency parameters of the surface myoelectric signal for assessing muscle fatigue during cyclic dynamic contractions. IEEE Trans Biomed Eng 48: 745–753
- Chang L, Satava RM, Pellegrini CA, Sinanan MN (2003) Robotic surgery: identifying the learning curve through objective measurment of skill. Surg Endose 17: 1744–1748
- D'Annibale AMD, Fiscon VMD, Trevisan PMD, Pozzobon MMD, Gianfreda VMD, Sovernigo GMD, Morpurgo EMD, Orsini CMD, Del Monte DMD (2004) The da Vinei rohot in right adrenalectomy: considerations on technique. Surg Laparosc Endosc Percutan Tech 14: 38–41
- De Ugarte DA, Etzioni DA, Gracia C, Atkinson JB (2003) Robolic surgery an resident training. Surg Endosc 17: 960–963
- Farina D, Fosei M, Merletti R (2002) Motor unit recruitment strategies investigated by surface EMG variables. J Appl Physiol 92: 235–247
- Farina D, Merletti R, Enoka RM (2004) The extraction of neural strategies from the surface EMG. J Appl Physiol 96: 1486–1495
- Gutt CN, Oniu T, Schemmer P, Mehrahi A, Buchler MW (2004) Fewer adhesions induced by laparoscopic surgery? Surg Endosc 18: 898–906
- Hashizume M, Shimada M, Tomikawa M, Ikeda Y, Takahashi I, Abe R, Koga F, Gotoh N, Konish K, Maehara S, Sugimaehi K (2002) Early experiences of endoscopic procedures in general surgery assisted by a computer-enhanced surgical system. Surg Endose 16: 1187–1191
- Hernandez JD, Bann SD, Munz Y, Moorthy K, Datta V, Martin S, Dosis A, Bello F, Darzi A, Rockall T (2004) Quali-

tative and quantitative analysis of the learning curve of a simulated surgical task on the da Vinci system. J Gastrointest Surg 18: 372–378

- Hubens G, Coveliers H, Balliu L, Ruppert M, Vaneerdeweg W (2003) A performance study comparing manual and robotic-assisted laparoscopic surgery using the da Vinei system. Surg Endose 17: 1595–1599
- 14. Korolija D, Sauerland S, Wood-Dauphinee S, Abbou CC, Eypasch E, Gareia Caballero M, Lumsden MA, Millat B. Monson JRT, Nilsson G, Pointner R, Schwenk W, Sbamiyeh A, Szold A. Targarona E, Ure B. Neugebauer E (2004) Evaluation of quality of life after laparoscopic surgery. Surg Endosc 18: 879–897
- McClure N, Gallagber AG, McGuigan J, Ritchie K, Sheehy NP (1997) An ergonomic analysis of the fulcrum effect in endoscopic skill acquisition. Gynaecol Endosc 6: 90
- Moorthy K, Munz Y, Dosis A, Hernandez J, Martin S. Bello F, Rockall T, Darzi A (2004) Dexterity enbancement with robotic surgery. Surg Endosc 18: 790–795
- Munz Y, Moorthy K, Dosis A, Hernandez J, Bann SD, Bello F, Martin S, Darzi A, Rockall T (2004) The benefits of stereoscopic

vision in robotic-assisted performance on bench models. Surg Endosc 18: 611-616

- Narazaki K, Oleynikov D, Stergiou N (in review) Robotic laparoscopy: objective assessment of proficiency with bimanual inanimate tasks. J Laparoendose Adv Surg Tech A
- Narazaki K, Oleynikov D, Stergiou N (2006) Robotic surgery training and performance: identifying objective variables for quantifying the extent of proficiency. Surg Endose 20: 96–103
- Prasad SM, Maniar HS, Soper NJ, Damiano RJ, Klingensmith ME (2002) The effect of robotic assistance on learning curves for basic laparoscopic skills. Am J Surg 183: 702–707
- Quiek NE, Gillete JC, Sbapiro R, Adrales GL, Gerlach D, Park AE (2003) The effect of using laparoscopic instruments on muscle activation patterns during minimally invasive surgery. Surg Endosc 17: 462–465
- Sarle R, Tewari A, Shrivastava A, Peabody J, Menon M (2004) Surgical robotics and laparoscopic training drills. J Endourol 18: 63–67
- Talamini MA, Stanfield CL, Chang DC, Wu AW (2004) The surgical recovery index. Surg Endose 18: 596–600

*Technique* 

Surg Endosc (2006) 20: 830–833 DOI: 10.1007/s00464-005-0450-9

© Springer Science+Business Media, Inc. 2006



and Other Interventianal Techniques

## Retroperitoneal hand-assisted laparoscopic surgery for endoscopic adrenalectomy

N. Sata, M. Shiozawa, A. Suzuki, K. Kurihara, J. Ohki, H. Nagai

Department of Surgery, Jichi Medical University, 3311-1 Yakushiji Shimotsuke Tochigi, 329-0498, Japan

Received: 23 June 2005/Accepted: 27 December 2005/Online publication: 16 March 2006

#### Abstract

Although hand-assisted laparoscopic surgery (HALS) is very common in various laparoscopic procedures, it is rarely used for retroperitoneal endoscopic adrenalectomy because of the small working area. The authors evaluate HALS in endoscopic adrenalectomy with respect to its use as a rescue procedure in complicated cases. In their department, 47 patients underwent endoscopic adrenalectomies between 1998 and 2004. Mainly because of complicated anatomy, three primary aldosteronism cases werc converted to retroperitoneal HALS. This involved making an additional 6 cm skin incision, into which the surgeon's left hand was inserted, with the palm used to create a sufficient visual field and working area. The fingers were used for tactile sensation and blunt resection. For these three cases, successful retroperitoneal HALS in endoscopic adrenalectomy resulted in no mortality or morbidity. These findings indicate that this procedure is a feasible technique for complicated benign adrenal tumor cases.

Key words: Endoscopic adrenalectomy — HALS — Retroperitoneal approach

Endoscopic adrenalectomy has been considered the standard surgical excision procedure for benign adrenal tumors during the past decade. Since the initial report of its use in 1992, this procedure has become a fcasible, minimally invasive method for benign adrenal tumors [5]. Transperitoneal and retroperitoneal approaches are the two principal laparoscopic routes to the adrenal gland. Although the transperitoneal approach is used more widely, the retroperitoneal approach offers particular advantages [1, 3, 4, 6, 12, 13, 18–20, 22].

Hand-assisted laparoscopic surgery (HALS) is an established alternative procedure in various types of laparoscopic surgery [9, 10, 14, 15, 21], including transperitoneal endoscopic adrenalectomy [2, 8, 16, 17]. However, HALS is rarely performed in retroperitoneal endo-

Table I	. Prof	ile of the	cases for	endosconic	adrenalee	tomy by year

	Transperitoneal approach (n)	Retroperitoneal approach ( <i>n</i> )	Conversion to open surgery (n)	Total
1998-2000	9	3	1	12
2001-2002	5	6		11
2003-2004	3	21		24
Total	17	30	1	47

scopic adrenalectomies because of the narrow working area.

In this report, we describe the details and distinct advantages of retroperitoneal endoscopic adrenalectomy using HALS. In particular, the use of HALS completely compensates for disadvantages otherwise encountered in retroperitoneal endoscopic adrenalectomy, such as disorientation, narrow working area, absence of tactile sensation, and technical limitations.

#### Patients

From 1998 to 2004, 47 patients (25 men and 22 women; mean age, 49.4 years) underwent endoscopic adrenalectomics in the Department of Surgery, Jiehi Medical University, Tochigi, Japan). The cases included 28 primary aldosteronisms, 11 Cushing's syndromes, 6 nonfunctioning adenomas, and 2 pheochromocytomas. The transperitoncal route was first applied to reach the adrenal gland as other abdominal organs assisted with orientation in the abdominal cavity. However, we sometimes encountered troublesome situations in the process of transperitoneal operation, especially in fatty cases or cases with a previous abdominal surgical procedure.

In 2000, we experienced a case of conversion to open surgery, which led to the gradual introduction of the retroperitoneal approach, which became the standard method after 2003 (Table 1). During the current study of 30 retroperitoneal endoscopic adrenalectomies, three were converted to HALS, mainly because of complicated anatomy (Table 2). The mean surgery time was 159 min, and the average blood loss was 138 ml in the 47 cases.

## Retroperitoneal hand-assisted laparoscopic adrenalectomy technique

Initially, all patients underwent a standard retroperitoneal endoscopic adrenalectomy using four trocars in the lateral bending position

Table 2. Cases converted to hand-assisted laparoseopic surgery (HALS) in endoscopic adrenalectomy

Site	Route	No. of eases	Cases eonverted to HALS	Mean operation time (min)	Average blood loss (ml)
Right	Transperitoneal	13	2	166	302
Retrope	Retroperitoneal	7	1	192	105
Left	Transperitoneal	4	0	203	228
	Retroperitoneal	23	2	148	34
Total		47	5	159	138



Fig. 1. a Placement of laparoseopic ports: (1) 12 mm port for videoscope, (2 and 3) 5 mm port for tissue manipulators and foreeps, (4) 2 mm port for fine forceps. b In the conversion to hand-assisted laparoscopic surgery (HALS), an additional 6 cm skin ineision (5) was made between the two left trocar incisions.

(Fig. 1a). The first 12 mm trocar for videoscope use was placed just beneath the 11th rih in the midaxillar line. After tissue expansion using a balloon dilator and pneumoretroperitoneum, three additional trocars (two 5 mm and one 2 mm) were inserted for tissue manipulation. Three such cases were converted to HALS.

The decision to convert depended on an assessment made 2 h after the commencement of surgery. The laparoscopic procedure was continued if it appeared that the tumor could be removed within the next hour. Otherwise, surgeons reverted to the HALS approach.

For HALS, a 6 cm skin incision was made between the two left trocar incisions, and the left hand of the surgeon was inserted in an air-



Fig. 2. Use of the surgeon's left fingers and palm. The thumb, ring finger, and little finger are used to hold the tumor; the forefinger and middle finger are used as blunt manipulators and to operate the forceps; and the palm is used as a working area and to create a visual field.

tight manner (Fig. 1b). The skin incision was kept as small as possible and extended to the lateral side if necessary for the convenience of the left hand maneuver. Afterward, only the 12 mm troear for the endoscope and the 5 mm troear for a manipulator were used for the HALS procedure. The operator stood at the right side of the patient for the right tumor, and at the left side for the left tumor.

The HALS approach involves two important factors. The movement and positioning of the surgeon's left hand is critical, and the volume of the retroperitoneal space is approximately that of a fist. A key feature of retroperitoneal HALS is that the thumb, ring finger, and little finger are used to hold the tumor, the forefinger and middle finger are used as blunt manipulators and foreeps, and the palm is the working area (Fig. 2). The position of the videoscope is adjusted for the use of two fingers to disseet connective tissue and to visualize vessels and the adrenal gland clearly. Small vessels are dissected using a harmonie scalpel, and the central vein and arteries are dissected using surgical elips (Fig. 3a and b).

The HALS technique was completed uneventfully, with no cases of mortality or morbidity. Dissection proceeded promptly and safely. All dissection points were clearly visualized, and tactile sensation resulted in reduction of the surgery time otherwise required. The additional surgery times required for HALS in the three cases were 47, 44, and 51 min, and the postoperative hospitalization times were 4, 6, and 8 days, respectively.

#### Discussion

The current report describes the use of retroperitoneal HALS in endoscopic adrenalectomy as a rescue procedure for technically or anatomically complicated eases. The HALS technique, a common and established procedure for various laparoscopic operations, provides significant advantages over the purely laparoscopic approach, such as safer and better manipulation, reduced



Fig. 3. Intraoperative views during retroperitoneal hand-assisted laparoscopic surgery (HALS) in endoscopic adrenalectomy. a Dissection of the central vein. b Dissection of the arteries. (1) adrenal adenoma, (2) central vein, (3) adrenal gland, (4) harmonic scalpel, (5) surgeon's thumb, (6) surgeon's middle finger, (7) surgeon's ring finger, (8) adrenal arteries. (9) surgical clip).

time and costs, and minimal additional overload. However, HALS is not commonly used for endoscopic adrenalectomy, especially for retroperitoneal approaches.

Bennett and Ray [2] first reported successful transperitoneal HALS in endoseopic adrenalectomy. They concluded that transperitoncal HALS is a feasible and less invasive alternative procedure for larger tumors and potentially malignant lesions. Although the abdominal cavity is large enough for placement of a hand port and insertion of a hand, the retroperitoneal space was not thought to be suitable for HALS because of its limited working area and visual field.

Retroperitoneal endoseopic adrenalectomy has several advantages over its transperitoneal eounterparts. It minimizes intraabdominal adhesions, subsequent bowel obstruction, and injury to the abdominal organs. Several studies have eoneluded that both methods are equally feasible for removal of adrenal tumors [13, 19, 20]. A very eautious view is that the major disadvantage of retroperitoneal endoscopic adrenalectomy is disorientation, because surgeons can have trouble locating the adrenal gland, which increases stress associated with the procedure.

We found that retroperitoneal HALS provided for easy orientation with taetile sensation, and thus eompletely compensated for the disadvantages otherwise associated with the retroperitoneal approach. For dissection within a small area, special left hand and finger techniques are required. With complete understanding of the technique, most surgeons can easily perform retroperitoneal HALS.

In the eurrent eases, HALS required approximately 50 additional minutes and did not affect the duration of the postoperative hospital stay. If cases requiring HALS were selected more quickly, operation time could be saved as well as overload to the patients. Intraoperative events, such as location of the kidney and the central vein as well as aeeidental puneture of the peritoneum, are possible factors that could predict the need for eonversion to HALS. Additionally, three-dimensional images of the adrenal gland and surrounding organs generated by preoperative multidetector raw computed tomography (CT) sean would work as a navigator for endoscopic adrenalectomy, possibly predicting the needs of HALS [21, 22].

The major advantage of laparoseopie adrenalectomy over open adrenalectomy is undoubtedly reduced postoperative pain and shorter hospitalization. In our department, a elinical pathway involving 4 days of postoperative hospitalization is applied to endoseopie adrenalectomy. Two of the eurrent patients were diseharged as scheduled, and the other patient was diseharged on postoperative day 8, mainly because of family and social reasons. In conclusion, retroperitoneal hand-assisted endoscopic adrenalectomy is feasible and can be used as an aid in the retroperitoneal approach, with intraoperative tactile localization of the adrenal gland. This procedure is especially applicable for anatomically complicated cases and for surgeons unfamiliar with the retroperitoneal route.

- Baba S, Iwamura M (2002) Retroperitoneal laparoscopic adrenalectomy. Biomed Pharmacother 56( Suppl 1): f13s-119s
- Bennett IC, Ray M (2002) Hand-assisted laparoscopic adrenalectomy: an alternative minimal invasive surgical technique for the adrenal gland. ANZ J Surg 72: 801–805
- Bonjer HJ, Sorm V, Berends FJ, Kazemier G, Steyerberg EW, de Herder WW, Bruining HA (2000) Endoscopic retroperitoneal adrenalectomy: lessons learned from 111 consecutive cases. Ann Surg 232: 796–803
- Chiu AW (2003) Laparoscopic retroperitoneal adrenalectomy: clinical experience with 120 consecutive cases. Asian J Surg 26: 139–144
- Gagner M, Lacroix A, Bolte E (1992) Laparoscopic adrenalectomy in Cushing's syndrome and pbeochromocytoma. N Engl J Med 329: 1033
- Giger U, Vonlantben R, Michel JM, Krahenbubl L (2004) Transand retroperitoneal endoscopic adrenalectomy: experience in 26 consecutive adrenalectomies. Dig Surg 21: 28–32
- Hurley ME, Herts BR, Remcr EM, Dylinski D, Gill IS (2003) Three-dimensional volume-rendered belical CT before laparoscopic adrenalectomy. Radiology 229: 581–586
- Kitaoka F, Yanaga K, Okudaira S, Tajima Y, Furui J, Kanematsu T (2005) Successful left adrenalectomy for metastatic hepatocellular carcinoma using hand-assisted laparoscopic surgery: report of a case. Surg Today 35: 172–174
- Klingler PJ, Hinder RA, Menke DM, Smith SL (1998) Handassisted laparoscopic distal pancreatectomy for pancreatic cystadenoma. Surg Laparosc Endosc 8: 180–184

- Kusminsky RE, Boland JP, Tiley EH, Deluca JA (1995) Handassisted laparoscopic splenectomy. Surg Laparosc Endosc 5: 463– 467
- Marvik R, Lango T, Tangen GA, Andersen JO, Kaspersen JH, Ystgaard B, Sjolie E, Fougner R, Fjosne HE, Nagelhus Hernes TA (2004) Laparoscopic navigation pointer for three-dimensional image-guided surgery. Surg Endosc 18: 1242–1248
- Munver R, Ilbeigi P (2005) Retroperitoneal laparoscopic adrenalectomy. Curr Urol Rep 6: 72–77
- Naya Y, Nagata M, Ichikawa T, Amakasu M, Omura M, Nisbikawa T, Yamaguchi K, Ito H (2002) Laparoscopic adrenalectomy: comparison of transperitoneal and retroperitoneal approaches. BJU Int 90: 199–204
- O'Reilly MJ, Saye WB, Mullins SG, Pinto SE, Falkner PT (1996) Technique of hand-assisted laparoscopic surgery. J Laparoendosc Surg 6: 239–244
- Ou H (1995) Laparoscopic-assisted minilaparatomy with colectomy. Dis Colon Rectum 38: 324–326
- Perrotti M, Ankem M, Peterson S, Geis WP (2003) Laparoscopic bilateral adrenalectomy with the use of a single midline hand-assist port. J Soc Lap Sur 7: 273–275
- Shen WT, Kebebew E, Clark OH, Duh QY (2004) Reasons for conversion from laparoscopic to open or hand-assisted adrenalectomy: review of 261 laparoscopic adrenalectomics from 1993 to 2003. World J Surg 28: 1176–1179
- Sung GT, Hsu TH, Gill IS (2001) Retroperitoneoscopic adrenalectomy: lateral approach. J Endourol 15: 505–511
- Suzuki K, Kageyama S, Hirano Y, Ushiyama T, Rajamahanty S, Fujita K (2001) Comparison of 3 surgical approaches to laparoscopic adrenalectomy: a nonrandomized, background matched analysis. J Urol 166: 437–443
- Takeda M (2000) Laparoscopic adrenalectomy: transperitoneal vs retroperitoneal approaches. Biomed Pharmacother 54(Suppl 1): 207s-210s
- Wolf JS Jr, Moon TD, Nakada SY (1997) Hand-assisted laparoscopic nephrectomy: technical considerations. Tech Urol 3: 123– 128
- Yagisawa T, Ito F, Isbikawa N. Matsuda K, Onitsuka S, Goya N, Toma H (2004) Retroperitoneoscopic adrenalectomy: lateral versus posterior approach. J Endourol 18: 661–664

### Letters to the editor

Surg Endosc (2006) 20: 834–835 DOI: 10.1007/s00464-005-0361-9

© Springer Science+Business Media, Inc. 2005



and Other Interventianal Techniques

## An alternative gallbladder extraction technique in laparoscopic cholecystectomy

Abstract. In this era of minimally invasive surgery, the challenge remains in finding techniques to reduce access trauma in terms of fewer and smaller size trocar ports. Our new described technique will allow a smaller subxiphoid port to be used to achieve extraction of the gallbladder without the need to change to a 5 mm laparoscope. We believe this method is easy to learn, safe and with no observable complications from our experience.

Key words: Gallbladder extraction — Laparoscopic cholecystectomy

We were very interested to read the recent article by Machado and Herman (Surg Endosc 2004; 18: 1289-1290) on the technique of removing the gallbladder during minilaparoscopic cholecystectomy [4]. We agree that it is a simple and cheap method of extraction using a self-made retrieval bag from a readily available surgical glove. We would like to ask the authors whether it is their usual practice to close the bag prior to delivery under direct vision? If not, should the operating surgeon be concerned with the possibility of spillage of infected bile or stones? We would like to describe our routine method of gallbladder extraction that is also simple to use and avoids the use of a 5-mm laparoscope. This method can be applied in laparoscopic cholecystectomy using the two- or three-trocars technique and can also be used in other laparoscopic procedures requiring specimen extraction, for example; laparosopic appendicectomy.

We perform laparoscopic cholecystectomy (LC) with the standard four-port method—three 5-mm right subcostal ports and one 10-mm subumbilical port. The method of gallbladder resection has been described previously. A manufactured extraction bag (Medical Technical Promotion, Dr. Karl-Storz, Tuttlingen, Germany) is used in cases where the gallbladder is severely inflamed, perforated (spontaneous or iatrogenic), or suspected to be malignant [2]. The use of an extraction bag is to reduce the amount of bile or stone spillage into the peritoneal cavity. It also serves to avoid the well-recognized port-site metastases in cases of malignancy [3].



Fig. 1. Closing string of the extraction bag is grasped with forceps introduced from the subxiphoid port.

The retrieval stage consists of first closing the extraction bag intracorporally by tightening the closing string. Then the forceps, inserted via the subxiphoid port, is used to grasp the end of this closing string (Fig. 1). While holding the closing string, the forceps can be advanced in a retrograde direction through the subumbilical port under direct vision by careful alignment (Fig. 2). It is usually easier for a single surgeon to perform this step as shown in the figure. The 10-mm subumbilical port is removed and the specimen can then be safely extracted with or without dilating the wound or extending the incision (Fig. 3).

This alignment technique can be applied to retrieve the gallbladder in uncomplicated elective cases where extraction bags are not necessary. Instead of grasping the closing string of the bag, a lengthened suture (at least 5 cm), used to ligate the cystic duct, can be used to advance through the subumbilical port.

In this era of minimally invasive surgery, the challenge remains in finding techniques to reduce access trauma in terms of fewer and smaller-size trocar ports. Our new described technique will allow a smaller subxiphoid port to be used to achieve extraction of the gallbladder without the need to change to a 5-mm laparoscope. We also recommend the use of a durable



Fig. 2. Alignment of forceps and laparoscope (external view, left). Forceps is advanced through the subumbilical port under direct vision (internal view, right).



Fig. 3. Subumbilical trocar removed (left). Extraction bag removed (right).

extraction bag, particularly for large stones, so as to allow various fragmentation techniques to be safely employed [1]. We believe this method is easy to learn, safe, and with no observable complications from our experience.

#### References

- Machado MA, Herman P (2004) A simple technique for removal of the gallbladder during microlaparoscopic cholecystectomy. Surg Endose 18: 1289–1290
- Kimura T, Goto H, Takeuchi Y, Yoshida M, Kobayashi T, Sakuramachi S, Harada Y (1996) Intraabdominal contamination after gallbladder perforation during laparoscopic cholecystectomy and its complications. Surg Endosc 10(9): 888–891

- Lundberg O (2000) Port site metastases after laparoscopic cholecystectomy. Eur J Surg Suppl 585: 27–30
- Bordelon BM, Hobday KA, Hunter JG (1992) Incision extension is the optimal method of difficult gallbladder extraction at laparoscopic cholecystectomy. Surg Endosc 6(5): 225–227

#### D. C. T. Wong C. N. Tang

Department of Surgery Pamela Youde Nethersole Eastern Hospital Chai Wan, Hong Kong SAR, China

Online publication: 24 November 2005

Surg Endosc (2006) 20: 836–837 DOI: 10.1007/s00464-005-0641-4

© Springer Science+Business Media, Inc. 2006



and Other Interventional Techniques

## A correlation between the shape of the epiglottis and obstructive sleep apnea

#### Abstract

The ineidence of Obstructive Sleep Apnea (OSA) is increasing with the rise in the prevalence of obesity in the population. Upon performing esophagogastroduodenoseopy (EGD) on more than 50 patients with BMI ranging from 21 to 63, we noticed an increase in the concavity of the posterior surface of the epiglottis in eorrelation with the increase in BM1. Since OSA is eaused by collapse of the airways, this same pressure seems to be responsible for the deformity of the epiglottis, which normally has a minimally concave posterior surface. Therefore the shape of the epiglottis reflects the degree of airway collapse and thus the severity of OSA. We recommend that patients with inereased concavity of the posterior epiglottal surface seen endoscopically should be tested for OSA.

Key words: Epiglottis — Coneavity — Obstructive Sleep Apnea

Obstructive sleep apnea (OSA) is a sleep disorder eharacterized by recurrent episodes of partial or complete closure of the upper airway during sleep [2]. This disease affects 2% to 4% of middle-aged men and 1% to 2% of middle-aged women in Western countries, with the majority of affected individuals remaining undiagnosed. In one survey, as many as 82% to 92% of people with symptoms of moderate to severe OSA were not diagnosed [1].

During sleep, muscle relaxation in the upper airways and diminished reflexes lead to partial airway collapse in all people. In normal individuals, this minor airway collapse has no major effect on the airflow. However, in patients with OSA, the airway collapse can be severe, leading to partial or complete obstruction [6]. In 80% of patients, this airway obstruction involves the retroglottal region. This has been confirmed by magnetic resonance imaging (MRI) and dynamic fiberoptic pharyngoscopy [4]. The compromise of airflow manifests as snoring, apneic episodes, and chronic nighttime hypoxemia, leading to significant complications including systemic and pulmonary hypertension, eor pulmonale, and many other pathologies.

Obstructive sleep apnea should be suspected in patients who exhibit loud snoring; episodes of apnea, choking, or gasping witnessed during sleep; hypertension; a neek eireumference of 17 inches or greater; a high body mass index (BM1); and laterally narrowed oropharynxes. The diagnosis of OSA can be established by polysomnography [5].

In an MR1 study of patients with OSA, it was found these patients have excess fat deposition, most often located anterolateral to the upper airway, as compared with control subjects [3]. The marked fat deposition in the neck tissues increases pressure on the airways, contributing to the airway collapse seen in patients with OSA [1].

#### Observation

Normally, the posterior surface of the epiglottis is slightly eoneave as seen from above [7]. When we performed esophagogastroduodenoseopy on patients with a variety of BM1s ranging from 21 to 63, we noticed a significant correlation between BMI and the degree of concavity of the posterior epiglottal surface. The posterior surface of the epiglottis in patients with normal BMI was minimally concave. However, the extent of concavity increased proportionally with the increase in BMI, to the point at which total closure of the epiglottis was observed in extreme cases (Fig. 1).

The degree of concavity can be estimated visually without difficulty. A numeric value grading the degree of concavity can be obtained by drawing lines cotangent to the posterior lateral surfaces of the epiglottis and then drawing perpendicular lines from the edge points of the epiglottis as shown in picture A. The angle produced by the intersection of these lines will be equal to the arc constituted by the epiglottis. This angle, which we named "alpha," was approximately 100° in a patient with a BMI of 28. The alpha was 360° in another patient with a BMI of 62.
Picture A Picture B

Fig. 1. Comparison between the epiglottis of a patient with a body mass index (BM1) of 28 (A) and that of a patient with a BM1 of 62 (B). The inferior side of the picture represents the posterior pharyngeal wall with the epiglottis anterior to it and the retroglottal space between them.

Conclusion

A positive correlation between high BMI and OSA is well known in the literature [1]. The pressure of the parapharyngeal fat pads and the chronic collapse of the retroglottal airway during sleep create the deformity of the epiglottis observed in patients with OSA. Thus, the shape of the epiglottis seen endoscopically reflects the degree of the chronic airway obstruction. On the basis of this finding, we recommend sleep studies for patients in whom an increased degree of epiglottis concavity is seen during esophagogastroduodenoscopy or nasolaryngoscopy.

#### References

- 1. Gibson GJ (2005) Obstructive sleep apnoea syndrome: underestimated and undertreated. Br Med Bull 72: 49-65 (print 2004)
- Kingman P (2000) Strohl: obstructive sleep apnea-hypopnea syndrome. 21st ed. Ceeil textbook of medicine, Goldman, p 462.
- Lin Z, Zhang H, Wang T, Li C, Bai Z (2000) The upper airway MRI of obstructive sleep apnea patients. Zhonghua Er Bi Yan Hou Ke Za Zhi 35: 51–54
- Lin Z, Zhang H, Wang T, Li C, Bai Z (2000) Upper airway occlusion sites of obstructive sleep apnea. Zhonghua Er Bi Yan Hou Ke Za Zhi 35: 150–152
- Olson EJ. Park JG, Morgenthaler TI (2005) Obstructive sleep apnea-hypopnea syndrome. Prim Care 32: 329–359
- Philip R. Westbrook MD (2002, June 25) An overview of obstructive sleep apnea: epidemiology, pathophysiology, clinical presentation, and treatment. UpToDate version

- Williams P (1989) Planchnology, respiratory system. In: Gray's anatomy. 37th ed. pp 1250–1252
- M. Gazayerli<sup>1</sup> W. Bleibel<sup>2</sup> A. Elhorr<sup>3</sup> E. Elakkary<sup>6</sup> D. Maxwell<sup>4</sup> R. Seifeldin<sup>5</sup>
- <sup>1</sup> North Oakland Medical Centers, Surgery Wayne State University
- United States
- <sup>2</sup> Caritas Carney Hospital, Internal Medicine
- Tufts University School of Medicine
- United States
- <sup>3</sup> North Oakland Medical, General Surgery
- Wayne State University
- United States
- <sup>4</sup> North Oakland Medical Centers, Pulmonary/Critical Care

Wayne State University

- United States
- <sup>5</sup> North Oakland Medical Centers, Family Medicine
- Wayne State University
- United States
- <sup>6</sup> North Oakland Medical Centers
- Wayne State University

United States

Online publication: 16 March 2006

## News and notices

Surg Endosc (2006) 20: 838–841 DOI: 10.1007/s00464-006-3032-6

© Springer Science+Business Media, Inc. 2006

## The Association of Endoscopic Surgeons of Great Britain and Ireland

The AESGBI office has moved from the Oaks Hospital Colchester. All correspondence should be addressed to:

Association of Endoscopic Surgeons of Great Britain and Ireland The Royal College of Surgeons 35/43 Lincoln's Inn Fields London WC2A 3PN, UK Tel: 0044 (0)171 973 0305 Fax: 0044 (0)171 430 9235 E-mail: jrahone@asghi.org.uk

#### **Courses at Colchester General Hospital**

Advanced laparoscopie courses take place in February and Oetober each year. They provide delegates with individual tuition including lectures, practical sessions and live operative demonstrations. The courses eover advanced techniques for gallstone disease and the operative sessions include laparoscopic hernia repair, hoth trans- and extra-peritoneal, fundoplication for hiatus hernia and colon resection or rectopexy.

Course Content:

Advanced techniques in laparoscopic cholecystectomy Detection of common bile duct stones/operative cholangiography Laparoscopic exploration of the common hile duct Simulator sessions including knot tying, suturing, cholangiography Case demonstrations as described above The course fee of £495 includes two nights hotel accommodation in-

eluding breakfast, course dinner, lunch and refreshments, course manual and videotape, all course materials and VAT.

For further information contact the course director:

Prof. Roger Motson Colchester General Hospital Turner Road Colchester, Essex, CO4 5JL, UK Tel: 0044 1206 742448 Fax: 0044 1206 742030 E-mail: sarah.wilby@essexrivers.nhs.uk

#### Courses at the Cuschieri Skills Centre

The Cuschieri Skills Centre, formerly The Surgical Skills Unit, provides high quality skills training aimed at improving surgical competence and execution related to particular specialties. The training covers predominantly, but not exclusively, endoscopic skills at three levels: basic, advanced, and procedure related. The Unit is also involved in research into the evaluation and assessment of psychomotor abilities of surgeons at all points in their career.

The Surgical Skills Unit was originally set up in 1992 hy Professor Sir Alfred Cuschieri, at the University of Dundee. In 1993 the unit was designated the MATTUS training unit for Scotland. Since that time it has expanded its remit to cover various surgical specialities continues to plan for new courses.

A variety of courses are regularly available: General Surgery, Laparoscopic Surgery Anaesthetics, General Practice Gynecology, Orthopedics, Otolaryngology, Pediatrics, Urology.

For further details, please contact the centre or visit our website.



and Other Interventional Techniques

Cuschieri Skills Centre Ninewalls Hospital & Medical School Dundee DDI 9SY, Scotland, United Kingdom Tel.: +44 (0) 1382 645857/496277 Fax: +44 (0) 1382 646042 www.dundee.ae.uk/surgicalskills

#### Ductal Calculi: The Laparoscopic Approach Practical Aspects of Laparoscopic Fundoplication Laparoscopic Colorectal Course Thoracoscopic Sympathectomy

Other specialist courses held in the unit include courses in anaesthetic techniques. arthroscopy, colonoscopy, endoscopy, gynecology, and otolaryngology.

For further information you can visit our weh site or contact:

Julie Struthers, Unit Co-ordinator Surgical Skills Unit Ninewells Hospital and Medical School Dundee DD1 9SY, Scotland, UK Tel: +44 1382 645857 Fax: +44 1382 646042 E-mail: j.e.a.struthers@dundee.ac.uk Web page: http://www.dundee.ac.uk/surgicalskills/

#### Courses at the Royal Adelaide Centre for Endoscopic Surgery

Basic and Advanced Laparoscopic Skills Courses are conducted hy the Royal Adelaide Centre for Endoscopic Surgery on a regular basis. The courses are limited to six places to maximize skill development and tuition. Basic courses are conducted over two days for trainees and surgeons seeking an introduction to laparoscopic cholecystectomy. Animal viscera in simulators is used to develop practical skills. Advanced courses are conducted over four days for surgeons already experienced in laparoscopic cholecystectomy who wish to undertake more advanced procedures. A wide range of procedures are ineluded, although practical sessions can be tailored to one or two procedures at the participants request. Practical skills are developed using training simulators and anaesthetised pigs.

Course fees: \$A300 (\$US225) for the basic course and \$A1,600 (\$US1,200) for the advanced course.

For further details and a broehure please contact:

Dr. D. 1. Watson or Professor G. G. Jamieson The Royal Adelaide Centre for Endoscopic Surgery Department of Surgery Royal Adelaide Hospital Adelaide, SA 5000, Australia Tel: +61 8 224 5516 Fax: +61 8 232 3471

#### Advanced Laparoscopic Suturing and Surgical Skills Courses

MOET Institute San Francisco, CA, USA Courses are offered year-round hy individual arrangement. The MOET Institute is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians and designates these CME activities for 20–40 eredit hours in Category 1 of the Physician's Recognition Award of the American Medical Association. These programs are also endorsed by the Society of Gastrointestinal Endoscopic Surgeons (SAGES).

For further information please contact:

Wanda Toy, Program Administrator Microsurgery & Operative Endoscopy Training (MOET) Institute 153 States Street San Francisco, CA 94114, USA Tel: (415) 626-3400 Fax: (415) 626-3444

## Fellowships in Minimally Invasive Thoracie and General Surgery

University of Pittsburgh Medical Center Pittsburgh, PA, USA

One-year fellowships in advanced Minimally Invasive Surgery in General and Thoracic Surgery are being offered at the University of Pittsburgh Medical Center. Requirements include completion of residence training programs in General or Thoracic Surgery. The fellowships will involve extensive clinical exposure as well as elinical and basic science research. These positions include a very competitive salary and travel allowance. Interested candidates are invited to send a letter of inquiry with a curriculum vitae to:

Philip R. Schauer, M.D.
Assistant Professor of Surgery and Co-Director for General Surgery James Luketieh, M.D.
Assistant Professor of Surgery and Co-Director for Thoracic Surgery The University of Pittsburgh Medical Center
C-800 Presbyterian University Hospital
200 Lothrop Street
Pittsburgh, PA 15213-3221, USA

#### Geisinger Medieal Center Fellowship in Minimally Invasive Surgery

A one-year fellowship in advanced minimally invasive general and thoracic surgery is available at Geisinger Medical Center. The fellowship will involve extensive clinical exposure to UGI, bariatric and thoracic minimally invasive surgery with emphasis on esophageal surgery for benign and malignant disease. Fellows will also participate in clinical outcomes research. Requirements include completion of residency training in general surgery or thoracic surgery. Competitive salary and benefits package including travel allowances are available. Interested candidates are invited to send a letter of inquiry and curriculum vitac to:

Anthony T. Petrick, MD Director, Minimally Invasive Surgery Geisinger Medical Center 100 North Academy Avenue Danville, PA 17821-2111. USA atpetrick@geisinger.edu

#### **Clinical Fellowship in Laparoseopie Surgery**

Mount Sinai School of Medicine, New York, NY, USA Program Director: Daniel Herron, M.D.

The Mount Sinai Clinical Fellowship in Laparoscopy is a 1-year position offered to two individuals each academic year. Fellows rotate through several clinical rotations, each one emphasizing a different aspect of minimally invasive surgery. During the bariatric rotation, the fellow will gain a broad exposure to laparoscopic gastric bypass, laparoscopic biliopancreatic diversion with duodenal switch and the laparoscopic adjustable gastric band. Additionally, the fellow will learn 2-stage approaches to supermorbid obesity and laparoscopic revisional surgery. In the GI rotation, the fellow joins one of the busiest laparoscopic foregut surgery groups in the East Coast, scrubbing on a large number of fundoplications, paraesophageal hernias, Heller myotomies, and gastroesophageal resections. This rotation also provides a strong experience in laparoscopic partial and subtotal colectomy as well as laparoscopic surgery for Crohn's disease. Solid organ surgery comprises another highlight of the fellowship, with two or more laparoscopic donor nephreetomies performed each week. Fellows will participate in several clinical research projects during the year and will be expected to present their results at a national meeting. Both the Fellowship Director and the Laparoscopic Division Chief have served on the Board of Governors of SAGES and strongly encourage Fellow participation in SAGES courses and committees. Fellows are provided with access to professional-quality video recording and editing equipment, and are formally trained in the use of this equipment to prepare academic video presentations. Numerous opportunities to write review articles and book chapters are available. Fellows will present in both General Surgery and Laparoscopie Surgery Grand Rounds. Fellows play an active role in the education of residents and medical students.

To apply, please visit www.misfellowshipcouncil.org.

#### Advanced Clinical Minimally Invasive Fellowship

Harvard Medical School Both Israel Deaconess Medical Center Boston, Massachusetts, USA Advanced Clinical Minimally Invasive Fellowship with emphasis on Bariatric Surgery, (One-year position begins July 2003). Interested candidates are invited to send curriculum vitac to:

Daniel B. Jones, M.D., F.A.C.S. Chief, Section of Minimally Invasive Surgery Harvard Medical School Beth Israel Deaconcess Medical Center Shapiro TCC 140 330 Brookline Avenue Boston, MA 02215, USA

#### Minimally Invasive Surgery Research Fellowships

Mount Sinai Minimally Invasive Surgery Center Mount Sinai School of Medicine, New York, NY, USA Center Director: Barry Salky, M.D.

Research fellowships are sponsored by the Mount Sinai MISC for qualified individuals who are interested in an in-residence experience working under the guidance of MISC faculty. The MISC research fellowship program is extremely competitive, and the Center receives numerous applications each year for only a few positions. Applications for research fellowships are accepted up to six months in advance of the expected start date.

For more information on this program (and all Mount Sinai Minimally Invasive Surgery Center activities), please visit www.mssm.edu/misc.

To apply for a research fellowship, contact:

Benjamin Green Manager Mount Sinai Minimally Invasive Surgery Center One Gustave L. Levy Place Box 1259 New York, NY 10029, USA E-mail: benjamin.green@mountsinai.org

#### East Bay (San Francisco Bay Area) Minimal Access Surgery Fellowship

A one-year fellowship starting July 2005 is being offered by the First Surgical Consultants Medical Group, at the Alta Bates Summit Medieal Center in Oakland/Berkeley, California, USA. The fellowship will involve extensive advanced laparoscopic experience as well as clinical research. Approximately 500–550 laparoscopic surgery procedures are performed annually, including bowel resection, hernia surgery, foregut surgery, and solid organ surgery. The fellow will also gain experience in bariatric surgery, use of intraoperative ultrasound, robotic surgery using Da Vinci Robot, as well as open (15–20%) surgery. First Surgical Consultants is an advanced laparoscopic surgery group, in private practice in the East Bay of the San Francisco bay area. The group includes five surgeons and two full-time physician assistants. Requirements for fellowship include completion of surgical residency and California License. Interested candidates are invited to contact:

Ajay Upadhyay, M.D. Fellowship Co-Director George Kazantsev, M.D. Fellowship Co-Director/Medical Director Bariatric Surgery First Surgical Consultants 365 Hawthorne Avenue, Suite #101 Oakland, CA 94609, USA Tel.: 510 465 5523 Fax: 510 832 6061 E-mail: Upadhyayak@pol.net

#### Intensive Courses of Laparoscopic Surgery 2006

Hospital Podlesi Center of Miniinvasive Surgery 739 61 Trinec, Czech Republic

Contact address: lada.bystronova@nempodlesi.cz, www.nempodlesi.cz

Head of the course: Stanislav Czudek, M.D., Ph.D. 24.–26.5.2006 4.–6.10.2006 1.–3.11.2006

#### 1st World Conference on Interventional Oncology

Cernobbio, Italy 12–16 June 2006

Scientific Organizer: Prof. Luigi Solbiati http://www.wcio2006.com/ Conference Sceretariat: Domm International s.r.l Via Rossini 1 20122 Milano, Italy Tel.: + 39 02 7779181 Fax: + 39 02 76000181

Contact: Stefania Sella, Lorella Filugello

#### 10th World Congress of Endoscopic Surgery hosted by the EAES and incorporating the 14th International Congress of the European

Association for Endoscopic Surgery (EAES)

Berlin, Germany 13–16 September 2006

#### "Techniques and Technologies for Endoscopic Surgery"

Congress President: Professor Gerhard F. Buess Congress Secretary: Professor Karl-Hermann Fuchs

#### Wednesday, 13 September 2006

Postgraduate courses:

Bariatric surgery, Laparoscopic treatment (TAPP-TEP) of inguinal hernia, GERD update 2006, Laparoscopic colorectal surgery, and TEM

#### Technology symposium:

State of the art systems and process analysis of today's OR, OR systems in practice, Economic constraints, Usability constrains, and OR systems of the future.

#### Opening ceremony: Thrusday-Saturday, 14-16 September 2006

## IFSES WORLD CONFERENCE SPECIAL TOPIC—LOW BUDGET SURGERY

Featuring a full day symposium focused on "low budget surgery," technologies for developing and emerging countries, surgical procedures, and structures for endoscopic surgery.

#### TECHNIQUES AND TECHNOLOGIES FOR ENDOSCOPIC SURGERY

Fcaturing a wide range of topics on the latest information and updates on pathology, surgical techniques, and advances in technologics in endoscopic surgery. Highlighted among these techniques and scientific aspects and technologies and scientific aspects topics will be:

#### **Education topics**

Education programmes in MAS around the world: Are training courses helpful in MAS? How can we take the danger out of the surgical learning curve?, and Simulation for training.

#### Deadlines:

Abstract submission: 1 April 2006 Early Registration: 15 June 2006

#### **Congress Secretariat:**

EAES Office P. O. Box 335 5500 AH Veldhoven The Netherlands Tel.: + 31 40 252 5288 Fax: + 31 252 3102 E-mail: info@eaes.org Website; www.eaes-eur.org

#### 4th Frankfurter Meeting "Obesity Surgery" Frankfurter Römer, Frankfurt, Germany

9-10 November 2006

Congress Organization: Chair: Rudolf A. Weiner, Frankfurt

Congress Secretary: Sylvia Weiner Krankenhaus Sachsenhausen Schulstr. 31 60594 Frankfurt Tel.: 0049-69-6605-1199 or 1146 Fax: 0049-6605-1510 E-mail: rweiner@khs-ffm.de

Scientific Program includes plenary sessions, conferences, free communications, and live-transmissions from New York and Frankfurt. Advances in all relevant disciplines, such as open and laparoscopic surgery, gastric banding, restriction and bypass, biliopancreatic diversion, gastric stimulation, endoscopy, new and experimental approaches, pre- and postoperative care, long-term results, and clinical, metabolic, and general assistance of morbidly obese surgical candidates will be reviewed.

Program and general information can be found at the website: www. obesity-academy.com

#### University Postgraduate Course in Laparoscopic Surgery

Victor Babes University of Medicine and Pharmacy Timisoara – Pius Branzeu Center for Laparoscopic Surgery and Microsurgery Swiss Association for Laparoscopic and Thoracoscopic Surgery Swiss National Science Foundation

Schedule 2006: October 11–13: Advanced course in digestive surgery Venue: Timisoara (Romania)

Course's language: English

Course's fees: 800 ECU (accommodations included), with the support of Swiss National Science Foundation

Accreditation: The course is accredited by Romanian College of Physicians and supported hy European Association of Endoscopic Surgery (25 credits)

The program of the course includes live demonstrations (held by European experts) and experimental laboratory training in live tissue (pig). All instructors are surgeons with experience in laparoscopie surgery.

For further information please visit our web site (www.umft.ro/ pius\_hranzeu\_center) or contact:

Dr Vareus Florian PB Center for Laparoscopic Surgery and Microsurgery Piata E. Murgu Nr. 2 RO 300041, TIMISOARA Tel. 0040724 7444 19, 0041 789 125 349 Email: vareusllorian@yahoo.com

#### Asia Pacific Hernia Congress

New Delhi, India 6–8 October 2006

The Asia Pacific Hernia Congress & Live Workshop - Second Congress of Asia Pacific Hernia Society in conjunction with Indian Association of Gastrointestinal Surgeons (IAGES) is scheduled to he held in New Delhi from 6–8 October 2006. The Congress is being organized by the Department of Minimal Access Surgery, Sir Ganga Ram Hospital, New Delhi (India)

Speciality sessions will include:

- Pathogenesis of hernia
- Actiology of recurrent hernias
- Polymers and Biomaterials: Advantages, short comings present and future
- · Biologieal Prosthesis: Indications and Expectations
- Industry Role: Progress towards better patient care
- Evolution in hernia repair: comparison of surgieal techniques
- Abdominal wall closure
- Hernia repairs (Conventional and Endoseopie)
- Inguinal
  - Ventral primary
  - Ventral incisional
  - Recurrent hernias

Live Surgical Workshop will include:

- Hiatus hernias
- · Diaphragmatic hernias
- Primary tissue repairs
- Mesh based groin hernia repairs
- Recurrent hernia repairs
- Endoscopic hernia repairs
- Laparoscopic IPOM repair for ventral / incisional hernias
- Laparoseopie Peritoneal flap repairs for ventral / incisional hernias
- Difficult hernias

Pre-Congress Master Class & Hands-on training in Endohernia Surgery on 5th October 2006.

For further inquiry, please contact the Congress Secretariat or visit our website www.aphs06.com Dr. Pradeep Chowbey - Organizing Chairman AYUSHMAN, Douhle Storey Market, R-Block, New Rajinder Nagar, New Delhi–110060 (India)

Tel.: +91-11-25868768 / 25748085 \* Fax.: 91-11-25748085 / 26519935 e-mail.: chowhey1@vsnl.com / info@aphs06.com wehsite: www.chowbev.com / www.aphs06.com

## **BUILD YOUR PRACTICE...** WHERE THE FUTURE OF MEDICINE LIVES.

Marshfield Clinic is seeking an experienced BC/BE BARIATRIC SURGEON to lead in the development of a Center for Excellence for Bariatric Surgery, Marshfield Clinic, renowned for its excellence in patient care, research and education, offers a unique mix of clinical practice and academics in a state of the art medical practice. We are seeking a skilled laparoscopic surgeon with experience managing a bariatric program.

Marshfield Clinic is one of the most respected multi-specialty groups in the country with over 700 physicians in 41 locations. Physicians enjoy a two year guaranteed salary and comprehensive benefit package including:

- 4 weeks of vacation
- Two weeks of CME time/\$5,500 CME allowance
- Health, Life, Dental, and Malpractice Insurance
- 401K and Retirement Plan
- Generous relocation allowance



To learn more, we encourage you to send your CV or inquire to: Beth Albee, Physician Recruitment, Marshfield Clinic, 1000 N. Oak Ave., Marshfield, WI MARSHFIELD 54449. Phone: 800-782-8581, ext. 19775; Fax #: 715-221-9779; E-mail: there the feture of madeins liver albee, beth@marshfieldclinic.org, Website; www.marshfieldclinic.org/recruit

Marshfield Clinic is an Affirmative Action/Equal Opportunity employer that values diversity. Minorities, females, individuals with disabilities and veterans are encouraged to apply. Sorry, not a health professional shortage area.



## Cleveland Clinic

Department of Colorectal Surgery at the Digestive Disease Center

## **3rd Annual** LAPAROSCOPIC COLORECTAL SURGERY COURSE &

27th Annual

## **TURNBULL SYMPOSIUM**

## SEPTEMBER 26-30, 2006

Intercontinental Hotel & MBNA Conference Center Cleveland Clinic Campus 
 Cleveland, Ohio

#### Featuring:

- Laparoscopic Cadaver Course
- Laparoscopic Congress with live surgery
- Turnbull Symposium with live surgery
- Procedure for Prolapsed Hemorrhoids and Pelvic Anatomy Courses

For more information call (216) 297-7330

or visit and register online www.clevelandclinicmeded.com/turnbull06







## **Surgical Symposium 2006** November 30<sup>th</sup> - December 1<sup>st</sup>

Present

## Westin New York at Times Square

Topics include colorectal and bariatric surgery, ventral and inquinal hernias, gastroesophageal reflux disease (GERD), surgical infections, vascular surgery, breast, endocrine and hepatobiliary surgery.

## **Course Directors:**

Armando E. Castro, M.D., George S. Ferzli, M.D., Michael Zenilman, M.D.

## **Invited Faculty:**

Tracey Arnell, M.D. + Robert Brolin, M.D. + Robert Cacchione, M.D. + Peter Faries, M.D. Charles Filipi, M.D., Robert Fitzgibbons, M.D., Giselle Hamad, M.D., William Inabnet, M.D. Kamal, Itani, M.D.+Jacques Marescaux, M.D.+Jeffrey Marks, M.D.+Adrian Park, M.D. Alfons Pomp, M.D., Jeffrey Ponsky, M.D., Bruce Ramshaw, M.D., Christine Ren, M.D. David Rivadeneira, M.D., Bruce Schirmer, M.D., Scott Shikora, M.D., Rache Simmons, M.D. Lee Swanstrom, M.D.+Larry Whelan, M.D.+Douglas Wong, M.D.

For more information please contact selectedtopics@lmcmc.com or call 718-270-1422

## springer.com



# Quick, what's the best way to keep up to date with new developments in your field?

As a researcher in the sciences, you know the dilemma. You need to stay current with new research, but you can't take time to wander the web in search of the latest publications.

## The answer is SpringerAlerts!

As an alerts subscriber, you get reliable "first word" on upcoming books and journal tables of contents. This service is fast, it's free, and best of all it's flexible.

## Customize your SpringerAlerts to deliver exactly the information you need

Choose your topic – select your discipline, sub-discipline and related topics from thousands of choices.

## springer.com/springeralerts

keep up TO DATE



011759



# SpringerLink

The premier online publishing service from Springer!

SpringerLink is a powerful central access point for researchers in science, technology and medicine. Today, our collection spans the universe of research – more than 1,200 peer reviewed journals and 2,500 volumes within our Online Book Series, plus a constantly expanding library of eBooks, Online Reference Works and Online Journal Archives.

## To browse journals visit springerlink.com

10101011011000010011011010001001101101



# **Digital Through And Through!**

IMAGE 1<sup>™</sup> – The New KARL STORZ Camera System



THE DIAMOND STANDARD

KARL STORZ GmbH & Co. KG, Mittelstraße 8, D-78532 Tuttlingen/Germany, Phone: +49 (0)7461 708-0, Fax: +49 (0)7461 708-105, E-Mail: info@karlstorz.de KARL STORZ Endoscopy America, Inc, 600 Corporate Pointe, Culver City, CA 90230-76 00, USA, Phone: +1/310/338-8100, Fax: +1/310/210 5527, E-Mail: Info@ksea.com KARL STORZ Endoscopia Latino-America, 815 N. W. 57 Av., Suite No. 480, Miami, FL 33126-2042, USA, Phone: +1/305/262-8980, Fax: +1/305/262-89 86, E-Mail: info@ksela.com KARL STORZ Endoscopy Canada Ltd., 2345 Argentia Road, Suite 100, Mississauga, Ontario L5N 8K4, Phone: +1/905/816-8100, Fax: +1/905/858-0933, E-Mail: info@karlstorz.ca www.karlstorz.com



after switching to the U-CLIP™ Approximation Device.

Whether you perform general, vascular or cardiac open or endoscopic surgical procedures, the self-closing U-CLIP device dramatically simplifies your suturing technique during tissue approximation by eliminating knottying with each individual placement.

For more information on the many applications for the U-CLIP device and how it can improve outcomes, please email rs.u-clip@medtronic.com





Caution: Federal law (USA) restricts this device to sale by or on the order of a physician. For a listing of indications, contraindications, precautions, and warnings, please refer to the Directions For Use. For more information contact your local Medtronic Cardiac Surgery Technologies Sales Representative or call Customer Service toll-free at 1-800-328-1357. ©Medtronic, Inc. 2005 UC200503839EN