AD _____

Award Number: W81XWH-07-1-0685

TITLE: BRCA1-Associated Protein BRCC36: A Novel Target for Breast Cancer Therapy

PRINCIPAL INVESTIGATOR: Xiaowei Chen, Ph.D.

CONTRACTING ORGANIZATION: Fox Chase Cancer Center Philadelphia, PA 19111

REPORT DATE: October 2008

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

✓ Approved for public release; distribution unlimited

Distribution limited to U.S. Government agencies only; report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

R	EPORT DOC		N PAGE		Form Approved OMB No. 0704-0188
Public reporting burden for this data needed, and completing a this burden to Department of D 4302. Respondents should be valid OMB control number. PL	collection of information is est ind reviewing this collection of lefense, Washington Headquar aware that notwithstanding an EASE DO NOT RETURN YOU	mated to average 1 hour per resp nformation. Send comments reg ters Services, Directorate for Info y other provision of law, no perso R FORM TO THE ABOVE ADD	ponse, including the time for revi arding this burden estimate or a normation Operations and Reports on shall be subject to any penalty RESS.	ewing instructions, s ny other aspect of thi s (0704-0188), 1215 s / for failing to comply	earching existing data sources, gathering and maintaining the s collection of information, including suggestions for reducing lefferson Davis Highway, Suite 1204, Arlington, VA 22202- with a collection of information if it does not display a currently
1. REPORT DATE		2. REPORT TYPE			B. DATES COVERED
4. TITLE AND SUBTIT	LE	Annual		5	ia. CONTRACT NUMBER
BRCA1-Associate	d Protein BRCC36	: A Novel Target for	Breast CancerTher	rapy t	56. GRANT NUMBER W81XWH-07-1-0685 56. PROGRAM ELEMENT NUMBER
6. AUTHOR(S)				Ę	d. PROJECT NUMBER
Xiaowei Chen, Ph.	D.			ť	e. TASK NUMBER
				ţ	if. WORK UNIT NUMBER
Email: Xiaowei.Cho	en@fccc.edu			\$	PERFORMING ORGANIZATION REPORT
Fox Chase Cance Philadelphia, Penr	r Center nsylvania 19111	AND ADDRESS(ES)			NUMBER
9. SPONSORING / MC U.S. Army Medica Fort Detrick. Marvl	NITORING AGENCY I Research and Ma and 21702-5012	NAME(S) AND ADDRES Iteriel Command	S(ES)	1	0. SPONSOR/MONITOR'S ACRONYM(S)
				1	1. SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION / A Approved for Public	VAILABILITY STATEI c Release; Distrib	MENT ution Unlimited			
13. SUPPLEMENTAR	YNOTES				
14. ABSTRACT					
Since tumor cells in ge may lead to a therapeu invasive breast cancers that abrogation of BRC protamine based siRN/ should improve the targ therapeutic target to er ImmunoPET imaging tr achieved.	neral are genomically u tic index in tumor cells s and that depletion of f C36 will sensitize breas A delivery system to sel geting of breast tumor c hance the efficacy of ra ials in breast cancer pa	nstable and have defects over "normal" cells. Prev BRCC36 sensitizes breas st tumors to the DNA-dan ectively deplete BRCC36 ells while reducing non-s adiation and chemotheran tients, clinical translation	s in DNA damage respor vious studies have demo st cancer cells to IR via ti nage based therapies. T 5 in breast tumor xenogra specific toxicity. The prop py which elicit DNA dam of successful preclinica	nses, it has been nstrated that BF he BRCA1 DNA Fo test this hypo afts. This cance posed studies w age. As the ant I results of our a	In proposed that targeting DNA repair pathways RCC36 is over-expressed in the vast majority of repair pathway. Therefore, we hypothesize thesis, we will utilize an antibody (anti-HER2)- er cell-specific or "smart" therapeutic approach ill clearly establish BRCC36 as a novel ibody is currently being developed for antibody-P/siRNA conjugates could be rapidly
BRCAL BRCC36	and siRNA ta	rget theranies			
16. SECURITY CLASS	SIFICATION OF:	- Jee energhtep	17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON
			OF ABSTRACT	OF PAGES	USAMRMC
а. керокт U	d. ABSTRACT U	C. THIS PAGE U	UU	61	196. IELEPHONE NUMBER (include area code)
			ı		Standard Form 298 (Rev. 8-98)

Table of Contents

Introduction	4
Body	5
Key Research Accomplishments	8
Reportable Outcomes	8
Conclusion	9
References	9
Appendices	10

INTRODUCTION

Breast cancer is the most common cancer affecting women, with an estimated lifetime risk of approximately 10% by 80 years of age. In the United States, it is estimated that approximately 182,450 new cases of female breast cancer will be diagnosed and greater than 40,000 breast cancer-related deaths will occur in 2008 (Jemal, et al., 2008). Approximately 13.2% of all American women (1 in 8) are expected to develop breast cancer sometime during their lifetime and 3.0% will subsequently die from the disease (Ries, et al., 2008). Despite advances in treatment and early detection, the breast cancer mortality rate among women in the United States decreased by only 2.2% per year between 1990 and 2002 (Jemal, et al., 2008). The BRCA1 gene (OMIM: 113705) is one of the most intensively studied breast cancer susceptibility genes and has a profound role in breast cancer etiology owing to its involvement in several important cellular processes. Importantly, estimates from previous studies (Collaborative Group on Hormonal Factors in Breast Cancer, 2001; Margolin, et al., 2006) indicate that family history is associated with 15% to 20% of breast cancer cases in the United States.

BRCA1-associated proteins as potential targets for Breast Cancer Therapies

Since BRCA1 was cloned in 1994 (Miki, et al., 1994), significant progress has been made in defining its biochemical and biological functions, as well as its role in breast and ovarian cancers (Chen, et al., In press; Narod and Foulkes, 2004). BRCA1 has been implicated in many cellular processes, including DNA repair, and protein ubiquitination (Chen, et al., In press). Because of the important role of BRCA1 in DNA repair, breast tumors with defective BRCA1 are believed to be more sensitive to DNA damage-based therapies. Nevertheless, defects in BRCA1 itself may not be the only reason for the loss of its activity nor the increased sensitivity of tumor cells to DNA damage-based agents. A number of studies have demonstrated that manipulation of BRCA1-associated proteins, such as RAD51, MRE11, and NBS1, can impact cellular sensitivity to ionizing radiation (IR) (Chen, et al., In press; Digweed, et al., 2002; Lio, et al., 2004; Nakanishi, et al., 2002; Russell, et al., 2003). BRCA1-associated proteins may, therefore, be considered as potential targets for breast cancer therapies. Despite a potentially significant role for BRCA1-associated protein complexes in modifying the activities of BRCA1, the total number of complexes and the identity and function of component proteins has yet to be fully elucidated. Thus, much of the scientific effort related to BRCA1 is currently directed at defining the biochemical functions of BRCA1 in association with these protein complexes.

BRCA1/2 containing complex (BRCC)

Using a combination of affinity purification of anti-FLAG and mass spectrometric sequencing, we have reported a novel multiprotein complex, termed BRCC (BRCA1/2 Containing Complex), which contains seven polypeptides including BRCA1, BRCA2, BARD1 and RAD51 (Dong, et al., 2003). In this multiprotein complex, one protein, referred to as BRCC36, has been found to be associated with BRCA1 and BRCA2. BRCC36 bears sequence homology to the JAMM domain family, and displays deubiquitinating (DUB) activities towards K63-linked ubiquitin substrates (Ambroggio, et al., 2004; Sobhian, et al., 2007). We have shown that cancer-causing truncations of BRCA1 abrogate the association of BRCC36 with BRCC and that BRCC36 expression is elevated in the majority of invasive breast tumors (Dong, et al., 2003). Importantly, we have also demonstrated that depletion of BRCC36 by small interfering RNAs (siRNAs) results in increased sensitivity in breast cancer cells to IR and disruption of IR-induced

BRCA1 phosphorylation and nuclear foci formation (Chen, et al., 2006). Based upon these findings, we believe that BRCC36 may be an ideal target for breast cancer therapy using the siRNA gene silencing approach.

Antibody-mediated siRNA delivery

Considerable research efforts have been focused on applying siRNA for human disease therapy, including cancer therapy. A novel method for *in vivo* delivery of siRNAs to specific cell types has been recently developed, and it takes advantages of the nucleic-acid binding properties of protamine as well as the specificity of fragment antibodies (Fab) (Sioud, 2006). This method shows that systemically administered siRNA can be targeted to cells that express a specific cell-surface receptor (Peer, et al., 2007; Song, et al., 2005). Compared to other siRNA delivery systems, antibody-based siRNA targeting provides many advantages (Sioud, 2006), including that (i) the siRNA is stable in the blood with a prolonged half-life; (ii) the siRNA can be transported across capillary endothelial walls; (iii) the siRNA can be specifically bound to the plasma membranes of target cells ("smart drug"); and (iv) the siRNAs can be efficiently delivered into the target cells through endocytosis.

BODY

Task 1: To express and purify anti-HER2 C6.5-protamine fusion protein.



Figure 1. Purification of C6.5protamine fusion proteins. Purification of C6.5-protamine fusion proteins was determined by HPLC (A. Up panel: C6.5; Low C6.5-protamine) panel: and Western blot analyses (B). (C) The levels of C6.5-protamine fusion protein were also examined by Western blot assay in protein samples extracted from soluble and insoluble cell fractions. Recombinant C6.5 protein was loaded as positive control (PC).

Expression of anti-HER2 C6.5db-P fusion protein:

The sequence of the truncated form of human protamine (amino acids 8-29: RSQSRSRYYRQRQRSRRRRRS) was first subcloned between the C-terminus of the anti-HER2 C6.5 and His-tag in the vector pCYN2. Anti-HER2 C6.5-P-His was then expressed in

TG1 E. coli, extracted in periplasmic extraction buffer (30mM Tris-HCl, 20% sucrose, 1mM EDTA). After centrifugation, anti-HER2 C6.5-P-His fusion protein in the soluble fraction was purified using the Ni-nitrilotriacetic acid (Ni-NTA) agarose followed by high-performance liquid chromatography (HPLC) size-exclusion chromatography over a Superdex 75 column (Amersham Pharmacia). HPLC and Western blot analyses showed that no trace of C6.5-P-His fusion protein was found in the flowthrough and elusion solution (Figure 1A and 1B). Further Western analysis showed that the abundant C6.5-protamine-His was detected in the insoluble fraction of extraction buffer. However, the C6.5-protamine-His fusion protein was not detected in the soluble fraction at all (Figure 1C). These results indicated that the C6.5-Protamine-His fusion protein was insoluble in current extraction buffer. As shown in Figure 2A and 2B, only strong denaturants such as, 6 M GuaHCl, 2% SDS, and 1% Triton X-100, were able to efficiently extract the fusion proteins C6.5-protamine. As 2% SDS and 1% TRitonX-100 may cause irreversible denaturation of the proteins, we chose 6 M GuaHCl to extract fusion proteins from insoluble section as described in a previous study (Li, et al., 2001). However, the yield of purified fusion protein was still too low to perform the siRNA delivery experiments (data not show). Based on the facts that protamine has strong binding capacity with the nucleotides, we have found that protamine fragment (a.a., 8-29) is a high charge molecular. As shown in Figure 3, the C6.5-protamine fusion protein has a charge value of +15.9 at pH7.0 in comparison to that C6.5 alone only has a positive charge of 3.7 at pH7.0. This high charge character of protamine fragment (a.a., 8-29) appears to be responsible for the insolubility of C6.5-protamine fusion protein.



Figure 2. Detection of C6.5protamine fusion proteins. Protein samples were first extracted from bacterial lysed with various reagents including 6 M GuaHCl, 2%SDS, or 1% Triton X- 100. The lysate from non-induced bacterial was used as negative control (NC), and recombinant C6.5 protein was loaded as positive control (PC). Protein samples were then fractionated in SDS-PAGE followed by Western blot analysis using antibodies against His tag (A) and protamine (B).

Conjugation of anti-Her2 antibody and protamine peptide

Due to the technical difficulties of purifying the anti-HER2 C6.5–protamine fusion protein, we have decided to use an alternative approach to first synthesize the protamine (a.a., 8-29)-Cys and then conjugate the protamine-Cys with cysteine-containing anti-HER2 antibodies. Because of the character of high charge in protamine peptide (a.a., 8-29), the scientists in Genscript took 3x tries to finally synthesize this peptide. We have first measured the baseline levels of free -SH group containing in cysteine using 5,5'-dithiobis-(2-nitrobenzoic acid) (Ellman's reagent; Pierce). As shown in **Figure 3A**, the -SH contents is much higher in the protamine and Herceptin, an anti HER2 antibody, but pretty low in anti-HER2-scFV3.9, which contains two cysteines in its sequence. Therefore, the protein of anti-HER2 scFv3.9 was first reduced by DTT to obtain monomer anti-HER2 sFv3.9-SH. In brief, 1M DTT was added to the scFv in buffer of 10 mM

HEPES and 150 mM NaCl, (pH 7.4) to a final concentration of 50 mM. After rotation at room temperature for 5–10 min, the protein was desalted on a Sephadex G-25 column (GE) (**Figure 3B**). To perform the conjugation, protamine (a.a., 8-29)-Cys was incubate with Herceptin or scFv3.9 at molecular ration of 10:1, respectively. The solution was then mixed by gentle rotation for 30 min at room temperature to produce HER2 antibody and Protamine conjugation.



Figure 3. Reduction of anti-HER2 scFv3.9. (**A**) Ellman's assay for the determination of free sulfhydryl groups. (**B**) Fractionation in desalting by Sephadex G-25.

Task 2: To determine if abrogation of BRCC36 by C6.5-P siRNA delivery can sensitize breast tumors to DNA damage-based therapies in mouse xenograft models.

BRCC36 siRNA delivery in HER2-positive cells via anti-HER2 antibodies and protamine conjugates

To examine if HER2 antibodies and protamine conjugates enable delivery of *BRCC36* siRNA specifically to the HER2-positive breast cancer cells, we have performed *in vitro* silencing studies in the SKBR-3 breast cancer cell line which constitutively expresses high levels of HER2 and BRCC36 (Chen, et al., 2006). In comparison to control cells without siRNA treatment, Western blot analysis revealed a ~50% decrease in BRCC36 levels in the cells which



Figure 4. Knocking-down BRCC36 expression by siRNA delivery via anti-HER2 antibody-protamine conjugates. For the siRNA delivery studies, SKBR-3 cells were plated at a density 5×10^3 cells/cm². After reaching 30% to 40% confluence, cells were transfected with BRCC36 siRNA using either oligofectamine or antibody/protamine conjugates in OPTI reduced serum medium. After 48 hours, cells were harvested, and Western blot analysis was performed to examine the protein level of BRCC36 in cell lysates. Protein loading level was evaluated by immunoblotting with anti-_β-actin antibody.

siRNAs were delivered by either lipid (i.e., Oligofectamine, Invitrogen) or the conjugates of Herceptin and protamine peptide transfected cells (**Figure 4**). However, the breast cancer cells added with the conjugates of anti-HER2 scFv3.9 and protamine peptide have no response to the BRCC36 siRNA treatment. Because of the low levels of free -SH group presented in anti-HER2 scFv3.9, we expect the conjugation between anti-HER2 scFv3.9 and protamine may be less effective. We are currently optimizing the condition for reduction of scFv in order to increase the level of free –SH in anti-HER2 scFv protein.

KEY RESEARCH ACCOMPLISHMENTS

• Subcloned the fragment of Protamine (a.a., 8-29) between the C-terminus of the anti-HER2 C6.5 and His-tag in the vector pCYN2.

• Expressed the anti-HER2 antibody (C6.5)- protamine fusion protein in TG1 E. coli.

• Demonstrated that anti-HER2 antibody (C6.5)-protamine fusion protein is a molecular which has very high positive charge and is not soluble in non-denature buffer.

• Synthesized protamine (a.a., 8-29) with additional cysteine, and performed conjugation between anti-Her2 antibodies and protamine peptide

• Demonstrated that siRNA delivered via the conjugates of Herceptin and protamine peptide enable knock-down of the level of BRCC36 in the HER2-positive breast cancer cells.

REPORTABLE OUTCOMES

Abstracts

1. **Chen, X**, Weaver, J, Bove, BA, Vanderveer, L, Miron, A, Daly, MB, Godwin. AK. Allelic imbalance in BRCA1 and BRCA2 gene expression is associated with an increased breast cancer risk. In: Era of Hope, Department of Defense (DoD) Breast Cancer Research Program Meeting, (AB# 32-11, poster presentation), 2008.

2. **Chen, X**, Amin, N, Godwin. AK. Abrogation of BRCC36 impairs IR-induced BRCA1 activation and sensitizes breast cancer cells to IR-induced apoptosis. In: Era of Hope, Department of Defense (DoD) Breast Cancer Research Program Meeting, (AB# 32-7, poster presentation), 2008.

3. **Chen, X**, Weaver, J, Bove, BA, Vanderveer, L, Miron, A, Daly, MB, Godwin, AK. Allelic imbalance in BRCA1 and BRCA2 gene expression is associated with an increased breast cancer risk. In: Annual meeting of American Association of Cancer Research, (AB# 1926, poster presentation), 2008.

Publications

1. **Chen, X**, Weaver, J, Bove, BA, Vanderveer, LA, Weil, SC, Miron, A, Daly, MB, Godwin, AK. Allelic Imbalance in BRCA1 and BRCA2 Gene Expression Is Associated with an Increased Breast Cancer Risk. Hum Mol Genet, 17: 1336-1348, 2008.

2. **Chen X,** Kistler JL, Godwin AK. BRCA1-associated proteins: novel targets for breast cancer radiation therapy. In: Columbus F., editor. Radiation therapy for breast cancer. Hauppauge, NY: Nova Science Publishers, Inc., in press.

CONCLUSIONS

Previous studies have demonstrated that BRCC36 is over-expressed in the vast majority of invasive breast cancers and that depletion of BRCC36 sensitizes breast cancer cells to IR via the BRCA1 DNA repair pathway. Therefore, we are examining if abrogation of BRCC36 will sensitize breast tumors to the DNA-damage based therapies. We have tested a cancer cell-specific or "smart" therapeutic approach utilizing the conjugation of anti-HER2 antibodies and protamine to deliver BRCC36 siRNA to HER2 positive breast cancer cells. Since tumor cells in general are genomically unstable and have defects in DNA damage responses, it has been proposed that targeting DNA repair pathways may lead to a therapeutic index in tumor cells over "normal" cells. This approach should lead to improving the targeting of breast tumor cells while reducing non-specific toxicity.

REFERENCES

- Collaborative Group on Hormonal Factors in Breast Cancer, 2001. Familial breast cancer: collaborative reanalysis of individual data from 52 epidemiological studies including 58,209 women with breast cancer and 101,986 women without the disease. Lancet 358(9291):1389-99.
- Ambroggio XI, Rees DC, Deshaies RJ. 2004. JAMM: a metalloprotease-like zinc site in the proteasome and signalosome. PLoS Biol 2(1):E2.
- Chen X, Arciero CA, Wang C, Broccoli D, Godwin AK. 2006. BRCC36 is essential for ionizing radiation-induced BRCA1 phosphorylation and nuclear foci formation. Cancer Res 66(10):5039-46.
- Chen X, Kistler JL, Godwin AK. In press. BRCA1-associated proteins: novel targets for breast cancer radiation therapy. In: Columbus F, editor. Radiation therapy for breast cancer. Hauppauge, NY: Nova Science Publishers, Inc.
- Digweed M, Demuth I, Rothe S, Scholz R, Jordan A, Grotzinger C, Schindler D, Grompe M, Sperling K. 2002. SV40 large T-antigen disturbs the formation of nuclear DNA-repair foci containing MRE11. Oncogene 21(32):4873-8.
- Dong Y, Hakimi MA, Chen X, Kumaraswamy E, Cooch NS, Godwin AK, Shiekhattar R. 2003. Regulation of BRCC, a holoenzyme complex containing BRCA1 and BRCA2, by a signalosome-like subunit and its role in DNA repair. Mol Cell 12(5):1087-99.

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T, Thun MJ. 2008. Cancer statistics, 2008. CA Cancer J Clin 58(2):71-96.
- Li X, Stuckert P, Bosch I, Marks JD, Marasco WA. 2001. Single-chain antibody-mediated gene delivery into ErbB2-positive human breast cancer cells. Cancer Gene Ther 8(8):555-65.
- Lio YC, Schild D, Brenneman MA, Redpath JL, Chen DJ. 2004. Human Rad51C deficiency destabilizes XRCC3, impairs recombination, and radiosensitizes S/G2-phase cells. J Biol Chem 279(40):42313-20.
- Margolin S, Johansson H, Rutqvist LE, Lindblom A, Fornander T. 2006. Family history, and impact on clinical presentation and prognosis, in a population-based breast cancer cohort from the Stockholm County. Fam Cancer 5(4):309-21.
- Miki Y, Swensen J, Shattuck-Eidens D, Futreal PA, Harshman K, Tavtigian S, Liu Q, Cochran C, Bennett LM, Ding W and others. 1994. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. Science 266(5182):66-71.
- Nakanishi K, Taniguchi T, Ranganathan V, New HV, Moreau LA, Stotsky M, Mathew CG, Kastan MB, Weaver DT, D'Andrea AD. 2002. Interaction of FANCD2 and NBS1 in the DNA damage response. Nat Cell Biol 4(12):913-20.
- Narod SA, Foulkes WD. 2004. BRCA1 and BRCA2: 1994 and beyond. Nat Rev Cancer 4(9):665-76.
- Peer D, Zhu P, Carman CV, Lieberman J, Shimaoka M. 2007. Selective gene silencing in activated leukocytes by targeting siRNAs to the integrin lymphocyte function-associated antigen-1. Proc Natl Acad Sci U S A 104(10):4095-100.
- Ries LAG, Melbert D, Krapcho M, Stinchcomb DG, Howlader N, Horner MJ, Mariotto A, Miller BA, Feuer EJ, Altekruse SF and others. 2008. SEER Cancer Statistics Review, 1975-2005. 2008 ed. Bethesda, MD: National Cancer Institute.
- Russell JS, Brady K, Burgan WE, Cerra MA, Oswald KA, Camphausen K, Tofilon PJ. 2003. Gleevec-mediated inhibition of Rad51 expression and enhancement of tumor cell radiosensitivity. Cancer Res 63(21):7377-83.
- Sioud M. 2006. RNAi Therapy: Antibodies guide the way. Gene Ther 13(3):194-5.
- Sobhian B, Shao G, Lilli DR, Culhane AC, Moreau LA, Xia B, Livingston DM, Greenberg RA. 2007. RAP80 targets BRCA1 to specific ubiquitin structures at DNA damage sites. Science 316(5828):1198-202.
- Song E, Zhu P, Lee SK, Chowdhury D, Kussman S, Dykxhoorn DM, Feng Y, Palliser D, Weiner DB, Shankar P and others. 2005. Antibody mediated in vivo delivery of small interfering RNAs via cell-surface receptors. Nat Biotechnol 23(6):709-17.

APPENDICES

1. **Chen, X,** Weaver, J., Bove, BA, Vanderveer, LA, Weil, SC, Miron, A, Daly, MB, Godwin, AK. Allelic Imbalance in BRCA1 and BRCA2 Gene Expression Is Associated with an Increased Breast Cancer Risk. Hum Mol Genet, 17: 1336-1348, 2008.

2. **Chen X,** Kistler JL, Godwin AK. BRCA1-associated proteins: novel targets for breast cancer radiation therapy. In: Columbus F., editor. Radiation therapy for breast cancer. Hauppauge, NY: Nova Science Publishers, Inc., in press.

Allelic imbalance in *BRCA1* and *BRCA2* gene expression is associated with an increased breast cancer risk

Xiaowei Chen¹, JoEllen Weaver¹, Betsy A. Bove¹, Lisa A. Vanderveer¹, Susan C. Weil³, Alexander Miron⁴, Mary B. Daly² and Andrew K. Godwin^{1,*}

¹Medical Science Division, ²Population Science Division, Fox Chase Cancer Center, Philadelphia, PA 19111-2497, USA, ³Morphotek Inc., Exton, PA 19341, USA and ⁴Department of Cancer Biology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA 02115, USA

Received October 1, 2007; Revised and Accepted January 16, 2008

The contribution of BRCA1 and BRCA2 to familial and non-familial forms of breast cancer has been difficult to accurately estimate because of the myriad of potential genetic and epigenetic mechanisms that can ultimately influence their expression and involvement in cellular activities. As one of these potential mechanisms, we investigated whether allelic imbalance (AI) of BRCA1 or BRCA2 expression was associated with an increased risk of developing breast cancer. By developing a quantitative approach utilizing allele-specific real-time PCR, we first evaluated AI caused by nonsense-mediated mRNA decay in patients with frameshift mutations in BRCA1 and BRCA2. We next measured AI for BRCA1 and BRCA2 in lymphocytes from three groups: familial breast cancer patients, non-familial breast cancer patients and age-matched cancer-free females. The AI ratios of BRCA1, but not BRCA2, in the lymphocytes from familial breast cancer patients were found to be significantly increased as compared to cancer-free women (BRCA1: 0.424 versus 0.211, P = 0.00001; BRCA2: 0.206 versus 0.172, P = 0.38). Similarly, the AI ratios were greater for BRCA1 and BRCA2 in the lymphocytes of non-familial breast cancer cases versus controls (BRCA1: 0.353, P = 0.002; BRCA2: 0.267, P = 0.03). Furthermore, the distribution of under-expressed alleles between cancer-free controls and familial cases was significantly different for both BRCA1 and BRCA2 gene expression (P < 0.02 and P < 0.02, respectively). In conclusion, we have found that AI affecting BRCA1 and to a lesser extent BRCA2 may contribute to both familial and non-familial forms of breast cancer.

INTRODUCTION

Breast cancer is the most common cancer affecting women, with a lifetime risk among females $\sim 10\%$ by the age of 80 years. In the USA, it has been reported that there will be approximately 180 510 new cases of breast cancer, and more than 40 910 breast cancer-related deaths in 2007 (1). Current estimates suggest that family history is associated with 10–20% of breast cancer (2,3). *BRCA1* (OMIM: 113705) and *BRCA2* (OMIM: 600185) are two of the most prominent breast cancer susceptibility genes and deleterious mutations in these two genes are estimated to account for about 15–30% of familial breast cancer (4–6).

Germline mutations affecting the coding region of *BRCA1* and *BRCA2* are thought to lead to expression of mutant proteins, which are either inactive or function as dominant negatives. However, these scenarios have not been supported by functional studies (7-9). In fact, *Brca1* and *Brca2* knockout mouse models have demonstrated that elimination of Brca1 or Brca2 proteins is sufficient for the development of mammary cancer (10,11). Previously, we have reported that mutant *BRCA1* mRNAs containing premature stop codons were eliminated or destabilized by nonsense-mediated mRNA decay (NMD) (12) and lead to a state of haploinsufficiency. As a result, the ratios between the expressions from the mutant alleles and the corresponding wild-type alleles were

^{*}To whom correspondence should be addressed at: Medical Science Division, Fox Chase Cancer Center, 333 Cottman Avenue, Philadelphia, PA 19111-2497, USA. Tel: +1 2157282205; Fax: +1 2157282741; Email: andrew.godwin@fccc.edu

© The Author 2008. Published by Oxford University Press. All rights reserved. For Permissions, please email: journals.permissions@oxfordjournals.org

significantly decreased, resulting in what was referred to as allelic imbalance (AI). AI of *BRCA1* or *BRCA2* expression could decrease the level of both transcripts and proteins and thus contribute to increased susceptibility of developing breast cancer.

There is growing evidence to support this concept. Epigenetic studies have shown that loss of BRCA1 expression due to promoter hypermethylation is associated with $\sim 10\%$ of sporadic cases of breast and ovarian cancer (13-18). However, screens to evaluate AI have not been applied in depth to study its potential role in the genesis of familial forms of these diseases. A previous study reported that 6 out of 13 human genes, including BRCA1 and p53, were expressed with significant difference between the two alleles, and this difference was transmitted by Mendelian inheritance (19). Furthermore, Yan et al. (20) observed that decreased expression of one of the adenomatous polyposis coli tumor suppressor gene (APC) alleles was associated with the development of familial adenomatous polyposis. Their studies also found that even more modest decreases in the expression of one APC allele could contribute to attenuated forms of polyposis (20). Based on these findings, we hypothesize that a subset of non-BRCA1/2 mutation carriers with a strong family history of breast cancer are at increased risk of developing this disease as a result of AI in BRCA1 and BRCA2 gene expression.

In the present study, we have developed a quantitative approach to measure the allele-specific expression of *BRCA1* and *BRCA2*. We compared *BRCA1/2* allelic variation in a cohort of *BRCA1/2* mutation-negative familial breast cancer patients, non-familial breast cancer patients and age-matched cancer-free volunteers. Since susceptibility to breast cancer is far from being fully understood, our study may help to further identify genetic factors which contribute to breast cancer susceptibility.

RESULTS

Development of a quantitative allelic imbalance assay

In order to determine if allele-specific real-time PCR is able to quantitatively measure the AI in BRCA1 and BRCA2 gene expression from the individual allele, RNAs were isolated from the blood lymphocytes of two individuals determined by genotype and sequence analysis to be homozygous for either BRCA1-c.4308T/T or BRCA1-c.4308C/C (Fig. 1A). This polymorphism was chosen since it is relatively common, based on NCBI dbSNP data. The samples were then reverse transcribed and the cDNAs were mixed at various ratios (8:1, 4:1, 2:1, 1:1, 1:2, 1:4 and 1:8) as described in the Materials and Methods section. BRCA1-c.4308T/T was detected by the VIC fluorescence signal and BRCA1-c.4308C/ C was detected by the FAM fluorescence signal. As shown in Figure 1B, with decreasing cDNA ratios of c.4308T to c.4308C, the VIC curve (detecting the c.4308T allele) shifted to the right with the increasing value of C_{T-c.4308T} (VIC), while the curve of FAM (detecting c.4308C allele) shifted to the left with the decreasing value of C_{T-c.4308C} (VIC). At the same time, the value of ΔC_T (C_{T-c.4308T} (VIC)⁻ C_{T-c.4308C (FAM)}) changed from the negative to the positive.

By the regression analysis, a linear relationship between Log₂ ratio of cDNAs c.4308T to c.4308C and ΔC_{T} was identified: Log_2 (c.4308T/C) = $-0.0877 + 1.57897 * \Delta C_T$ (P < (0.001) (Fig. 1C). The Pearson correlation coefficient (r) between Log₂ (c.4308T/c.4308C) and ΔC_T was 0.9798. To establish a similar standard curve for BRCA2 allelic expression, cDNAs from two individuals, who were either homozygous for BRCA2-c.3396A/A or BRCA2-c.3396G/G, were mixed at the following ratios: 8:1, 4:1, 2:1, 1:1, 1:2, 1:4 and 1:8 (c.3396A/A allele:c.3396G/G allele). BRCA2-c.3396A was detected by the VIC fluorescence signal and BRCA2-c.3396G was detected by the FAM fluorescence signal. As shown in Figure 1D, with decreasing ratios of c.3396A to c.3396G, the VIC curve (detecting c.3396A allele) shifted to the right while the FAM curve (detecting c.3396G allele) shifted to the left. After regression analysis, a linear relationship between Log₂ (c.3396A/ c.3396G) and ΔC_T was identified: Log₂ (c.3396A/G) = $0.11726 + 1.26458 * \Delta C_T (P < 0.001)$ (Fig. 1E). The Pearson correlation coefficient (r) between Log_2 (c.3396A/ G) and ΔC_T was 0.9868.

Detection of allelic imbalance caused by nonsense-mediated mRNA decay

To examine whether the allele-specific real-time PCR assay is able to detect AI of BRCA1 and BRCA2 gene expression in cell lines, we evaluated RNAs isolated from lymphoblastoid cell lines (LCLs) which were derived from deleterious mutation carriers heterozygous for BRCA1-c.3671ins4 or BRCA2-c.796delT. These frame-shift mutations create the premature stop codons, which are predicted to activate the NMD pathway and thus lead to decreased levels of mRNAs from the mutant alleles (12). As shown in Figure 2A and B, the ratios of BRCA1-c.4308T to -c.4308C between wild type and BRCA1c.3671ins4 heterozygous samples were 0.93 ± 0.04 and 2.07 ± 0.06 , respectively (P < 0.01). By subcloning and sequencing the individual transcripts, we found that the underexpressed allele contained both the BRCA1-c.3671ins4 mutation and the BRCA1-c.4308C polymorphism (detected by the FAM signal) (data not shown). To further examine if the loss of BRCA1-c.3671ins4 was associated with NMD, we treated the BRCA1-c.3671ins4 LCLs with puromycin, a translational inhibitor, 14 h prior to RNA isolation. The ratio of BRCA1-c.4308T to -c.4308C in BRCA1-c.3671ins4 heterozygous cells decreased \sim 30%, in comparison to the nontreatment group $(1.50 \pm 0.05 \text{ versus } 2.07 \pm 0.06, P < 0.01)$ (Fig. 2B). Our data indicated that treatment with puromycin was able to partially recover the AI caused by NMD. Significant AI was also observed for the BRCA2-c.796delT mutant allele. The ratios of BRCA2-c.3396G to -c.3396A between wild-type and BRCA2-c.796delT heterozygous samples were 0.98 ± 0.06 and 6.59 ± 1.31 , respectively (P < 0.01). After treating the BRCA2-c.796delT LCLs with puromycin, the ratio of BRCA2-c.3396G to -c.3396A in BRCA2-c.796delT heterozygous cells decreased \sim 31%, in comparison to the non-treatment group $(4.90 \pm 0.87 \text{ versus } 6.25 \pm 1.17)$ (Fig. 2C and D). Our results suggested that the loss of expression of BRCA1 or BRCA2 mutant alleles via NMD significantly contributed to the observed AI.





(A) Allele-specific real-time PCR amplification plot analyses of *BRCA1*-c.4308T (VIC) and -c.4308C (FAM) was performed in cDNAs generated by RT–PCR using RNAs from blood lymphocytes of two individuals homozygous for either the *BRCA1*-c.4308T/T or *BRCA1*-c.4308C/C. DNA sequencing chromatograms confirming the genotype are shown in the right panel. (B) Allele-specific real-time PCR amplification plot was analyzed in mixed cDNAs of *BRCA1*-c.4308T/T (detected by VIC) and *BRCA1*-c.4308C/C (detected by FAM) at the following ratios: 8:1, 4:1, 2:1, 1:1, 1:2, 1:4 and 1:8, respectively. (C) The standard curve for *BRCA1* allelic imbalance: Log₂ (c.4308T/C) = $-0.0877 + 1.57897 * \Delta C_T$. The Pearson correlation coefficient (*r*) between Log₂ (c.4308T/c.4308C) and ΔC_T was 0.9798 (Data expressed as Mean \pm SD, *n*=3; the mean value of ΔC_T for c.4308T/C=1 has been adjusted to zero). (D) Allele-specific real-time PCR amplification plot was analyzed in mixed cDNAs of *BRCA2*-c.3396A/A (detected by VIC) and *BRCA2*-c.3396A/G) = $0.11726 + 1.26458 * \Delta C_T$. The Pearson correlation coefficient (*r*) between Log₂ (c.3396A/G) = 1 has been adjusted to zero).



E BRCA2 Allelic Expression Standard Curve



Figure 1. continued

BRCA1 and *BRCA2* allelic imbalance is associated with breast cancer risk

To evaluate AI of BRCA1 and BRCA2 gene expression, genotype analysis of the two common polymorphisms, BRCA1-c.4308T/C and BRCA2-c.3396A/G, was performed on DNA samples isolated from fresh-frozen peripheral blood lymphocytes from 85 unrelated BRCA1/2 mutation-negative familial breast cancer carriers (median age at sample collection: 47), 112 non-familial breast cancer carriers (median age at sample collection: 52) and 102 age-matched cancer-free females (median age at sample collection: 51) (Table 1). From these analyses, 37 (43.5%), 48 (42.9%) and 41 (40.2%) of the samples evaluated were determined to be heterozygote for the BRCA1-c.4308T/C polymorphism for familial breast cancer patients, non-familial cancer patients and cancer-free controls, respectively (Table 1). Furthermore, 39 (45.9%), 44 (39.3%) and 36 (35.3%) of the samples above were found to be heterozygous for the BRCA2-c.3396A/G polymorphism (Table 1).

Since our initial validation studies were preformed using immortalized LCLs, we first compared AI in RNA isolated from 20 fresh-frozen lymphocytes versus 20 established Epstein-Barr Virus (EBV)-lines. No significant differences were detected between these two sample sets [BRCA1: 0.424 ± 0.129 versus 0.409 ± 0.127 (n = 11); BRCA2: 0.212 + 0.180 versus 0.225 + 0.209 (n = 10)]. However, to limit any AI variation potentially introduced by EBV transformation, all subsequent AI assays were performed using RNAs isolated from peripheral blood lymphocytes. Next, RNA isolated from BRCA1-c.4308T/C (n = 126) and BRCA2-c.3396A/G (n = 119) heterozygotes, including single heterozygotes and double heterozygotes, were evaluated for integrity and quantity. Those samples demonstrating high quality and the necessary quantities were used in the AI assay, as described in the Materials and Methods section.

To evaluate the AI, we used the absolute values of Log_2 (*BRCA1*-c.4308T/C) or Log_2 (*BRCA2*-c.3396A/c.3396G).



Figure 2. *BRCA1* and *BRCA2* allelic imbalance caused by NMD. (**A**) Allele-specific real-time PCR amplification plots of *BRCA1*-c.4308T (VIC) and -c.4308C (FAM) for non-template control, *BRCA1* wild-type lymphoblastoid cells (WT), *BRCA1* mutant (heterozygous *BRCA1*-c.3671ins4) lymphoblastoid cells without [PC (-)] or with [PC (+)] puromycin treatment. (**B**) Allelic expression ratios of *BRCA1*-c.4308T to *BRCA1*-c.4308C (a: versus WT; b: versus PC (+); *t*-test, *P* < 0.05). (**C**) Allele-specific real-time PCR amplification plots of *BRCA2*-c.3396A (VIC) and -c.3396G (FAM) for non-template control, *BRCA2* wild-type lymphoblastoid cells (WT), *BRCA2* mutant (heterozygous *BRCA2*-c.796delT) lymphoblastoid cells without [PC (-)] or with [PC (+)] puromycin treatment. (**D**) Allelic expression ratios of *BRCA2*-c.3396A (a: versus WT; b: versus PC (+); *t*-test, *P* < 0.05).



The mean value of Log₂ (c.4308T/C) of *BRCA1* in the lymphocytes from familial breast cancer carriers was found to be ~104% higher than that that in the lymphocytes from cancer-free controls $[0.424 \pm 0.157 \ (n=32)$ versus $0.211 \pm 0.169 \ (n=40), P=0.00001; t$ -test] (Table 2 and Fig. 3A and B). Log₂ of *BRCA1*-c.4308T/C in the lymphocytes from non-familial breast cancer carriers was 73% higher than that in cancer-free controls $[0.353 \pm 0.209 \ (n=32), P=0.002$ versus control] (Table 2 and Fig. 3A and C). In comparison, the mean value of Log₂ of *BRCA2*-c.3396A/G in the lymphocytes from familial breast cancer patients was moderately

higher (10%) than that in cancer-free controls $[0.206 \pm 0.180 \ (n=37)$ versus $0.172 \pm 0.123 \ (n=31)$, P=0.38; *t*-test] (Table 2 and Fig. 4A and B). A similar result (38% higher) was observed for Log₂ (c.3396A/G) of *BRCA2* in the lymphocytes of non-familial breast cancer carriers $[0.267 \pm 0.171 \ (n=26), P=0.03$ versus control] (Table 2 and Fig. 4A and C).

Interestingly, the distribution of under-expressed alleles of *BRCA1* and *BRCA2* was found to be significantly different between cancer-free control and familial breast carriers, but not between cancer-free control and non-familial breast carriers. As shown in Table 3 and Figure 3, under-expressed

Table 1. Characteristics of the study groups

Characters	Study groups		
	Familial	Non-familial	Cancer-free
Sample size	85	112	102
Age (median)			
At diagnosis	44	49	NA
At sample collection	47	52	51
Family history ^a			
2 or more	85	0	0
1	0	23	25
0	0	89	77
Genotypes			
BRCA1-c.4308T/C	37	48	41
BRCA2-c.3396A/G	39	44	36

^aNumber of first and/or second-degree relatives affected with either breast and/or ovarian cancer.

BRCA1-c.4308T (i.e. Log_2 [4308T/C] < 0) and *BRCA1*-c.4308C (i.e. Log_2 [4308T/C] > 0) alleles were found in ~53% (21 of 40) and ~47% (19 of 40) of cancer-free controls as compared to ~28% (9 of 32) and ~72% (23 of 32) of familial breast cancer carriers, respectively (P < 0.02). In addition, under-expressed *BRCA2*-c.3396A [i.e. Log_2 (3396A/G) < 0] and *BRCA2*-c.3396G [i.e. Log_2 (3396A/ G) > 0] alleles were found in ~45% (14 of 31) and ~55% (17 of 31) of cancer-free controls as compared to ~70% (26 of 37) and ~30% (11 of 37) of familial breast cancer carriers (P < 0.02), respectively (Table 3 and Fig. 4).

Inheritance effects of AI in BRCA1

A previous study has indicated that AI for several tumor suppressor genes could be transmitted by Mendelian inheritance (19). To test if the AI observed in our study may be inherited, we identified three affected women (i.e. probands) reporting a significant family history of breast and/or ovarian cancer for which we had blood from at least one of their sisters. Furthermore, each sister had to be heterozygous for the BRCA1-c.4308T/C polymorphism. As shown in family A (Table 4), sister Sis-02 displayed a similar AI pattern as compared to the proband, while sister Sis-01 displayed no AI. In the two other families, both the affected probands and their corresponding sisters showed AI (Table 4). We further performed a haplotype analysis to determine whether the alleles showing AI were shared between siblings. As shown in Table 4, sisters with the same AI phenotype shared the same haplotype with their affected sister. Importantly, sister Sis-01 in family A did not share the same haplotype. Her blood sample displayed no AI (0.007 + 0.147) for BRCA1 gene expressions whereas the AI was detected in her unaffected and affected sisters (Sis-02 and Proband, 0.382 + 0.176 and 0.375 + 0.06, respectively) (Table 4). The allele frequencies of the microsatellite markers used for haplotype construction are listed in Supplementary Material, Table S1.

DISCUSSION

In this study, we developed a quantitative AI assay to examine the expression difference between the alleles of *BRCA1* and

BRCA2 (Fig. 1). By performing this AI assay with specific primers and probes that target common single nucleotide polymorphisms in BRCA1 and in BRCA2, we were able to detect allelic imbalance associated with NMD in patients carrying frameshift mutations in BRCA1 and BRCA2 (Fig. 2). We next compared AI of BRCA1 and BRCA2 expression among three groups, familial breast cancer patients, non-familial breast cancer patients, and age-matched cancer-free females. AI ratios of BRCA1 in familial breast cancer cases were significantly higher than those from cancer-free controls (P =0.00001) (Table 2 and Fig. 3). Similar results were observed for AI ratios of BRCA1 in the lymphocytes from non-familial breast cancer patients (P=0.002). AI ratios of BRCA2 in familial or non-familial breast cancer cases were also higher than those from cancer-free controls (P = 0.38 or P = 0.03, respectively). However, the difference was not statistically significant in the ratios of mRNA expressed from the BRCA2 alleles found in familial breast cancer cases when compared to cancer-free controls (Table 2 and Fig. 4). In addition, the distribution of under-expressed alleles between cancer-free controls and familial cases was significantly different for both BRCA1 and BRCA2 gene expression (P < 0.02 and P <0.02, respectively) (Table 3). Furthermore, we have demonstrated that the AI patterns for BRCA1 expression, albeit in a small number of families, can be transmitted by Mendelian inheritance (Table 4). Although these findings are consistent with a previous study (19), future evaluations will benefit from evaluating AI in large families for evidence of disease segregation.

Several methods have been developed to evaluate allelespecific expression. The first method combines primer extension and capillary electrophoresis (19.21). The second approach utilizes microarray technology to measure allelespecific mRNA expression (22). Compared to the AI assay presented here, the method of primer extension plus capillary electrophoresis is also accurate but relatively time-consuming and expensive. The microarray approach provides a highthroughput and a powerful platform for the simultaneous analysis of large numbers of genes to analyze allele-specific gene expression, but it has less power to define the AI. Like the majority of allelic expression methods (23), our AI assay also requires a transcribed heterozygous variant in the individuals to be evaluated. In the present study, we targeted two common polymorphisms, BRCA1-c.4308T>C and BRCA2-c.3396A>G in the general population. Therefore, a substantial number of subjects homozygous for the polymorphisms had to be excluded. To overcome this limitation of population selection based on genotypes, other primers and probes will need to be developed to target other common polymorphisms in BRCA1 and/or BRCA2. In addition, our approach could easily be applied for studying AI in other cancer susceptibility genes, such as p53, APC, PTEN, etc.

In this study, we have demonstrated AI for both BRCA1 and BRCA2 in breast cancer populations. Interestingly, the increase of AI ratios in familial and non-familial breast cancer patients was more significant for BRCA1 than BRCA2. Loss of BRCA1 expression in breast cancer has been reported to be related to the pathogenesis of breast cancer (13–17). Loss of BRCA2 expression in cancers, in

Genes	Population	Sample number	AI $(Mean \pm SD)^a$	t-test (P-value) versus cancer-free
BRCA1	Cancer-free	40	0.211 ± 0.169	
	Familial	32	0.424 + 0.157	0.00001
	Non-familial	32	0.353 + 0.209	0.002
BRCA2	Cancer-free	31	0.172 + 0.123	
	Familial	37	0.206 + 0.180	0.38
	Non-familial	26	0.267 ± 0.171	0.03

Table 2. Allelic imbalance in BRCA1 and BRCA2 expression

^a To calculate the mean value of AI, all negative value of Log_2 (*BRCA1*-c.4308T/C) and Log_2 (*BRCA2*-c.3396A/c.3396G) in Figures 3 and 4 were changed to positive values.

contrast, is still controversial (24,25). These findings indicate that AI in *BRCA1* appears to be a more common event in breast cancer development than AI involving *BRCA2*. However, the mechanism(s) leading to the observed AI is for the most part unknown.

We have demonstrated that both *BRCA1* and *BRCA2* deleterious mutations can activate the NMD pathway and result in AI [Figure 2, and (12)]. However, all the familial breast cancer patients evaluated in the current study were determined to lack a mutation in *BRCA1* and *BRCA2* that would trigger NMD. Furthermore, we evaluated the *BRCA1* and *BRCA2* genes in the sporadic breast cancer patients and cancer-free controls demonstrating AI [i.e. allele expression ratio > 0.25 or < -0.25 (Figs 3 and 4)]. Again, no deleterious germline mutations were detected (data not shown). This is not entirely surprising given that germline mutations in *BRCA1* and *BRCA2* are rare in women affected with breast cancer without a strong family history of the disease (26–29).

Based on these observations, we conclude that NMD is not likely to be responsible for the observed AI in our casecontrol comparisons. Therefore, other mechanisms are likely to exist to account for the observed increased AI of BRCA1 and BRCA2 gene expression in female breast cancer patients. For example, the 5' and 3' non-coding regions of BRCA1 and BRCA2 are rarely evaluated through genetic testing, even though genetic alterations in these non-coding regions could be important in regulating BRCA1 and BRCA2 expression. For instance, genetic alterations within 5' DNA or the putative promoter regions are able to disrupt the binding of transcription factors to DNA regulatory elements and hence lead to the loss of allelic gene expression. Several studies have shown that large genomic deletions involving the BRCA1 promoter were associated with hereditary breast cancer (30-32). This concept is further supported by studies of Cowden syndrome (CS) showing that $\sim 10\%$ of CS-related PTEN mutations occur in the PTEN promoter and lead to a 50% reduction in PTEN expression (33,34). Also, allele-specific hypermethylation of the BRCA1 promoter region and decreased BRCA1 expression is associated with $\sim 10\%$ of sporadic breast cancer cases (18,30,35). Recent advances have identified a new pathway for gene regulation, i.e. via microRNAs (miRNAs) (36,37). These 21-22 nt RNA molecules are complementary to the 3'-UTR sequence of transcripts and mediate negative post-transcriptional regulation through RNA duplex formation (36,38). By performing in silico analyses in four BRCA1 SNPs and two BRCA2

SNPs (39), we have identified three rare *BRCA1* alleles (c.5628G, c.6273T, c.6924A) that could potentially create target sites for selected microRNAs (Supplementary Material, Table S2). Therefore, it is possible that altered mRNA targeting could contribute to AI of *BRCA1* gene expression in the absence of frameshift mutations. It will be important in future studies to determine the mechanisms that either disrupt transcription factors binding or alter miRNA binding, leading to constitutively decreased levels of *BRCA1* and *BRCA2* and an increased risk of developing breast cancer.

In summary, we have developed a quantitative approach to evaluate expression of BRCA1 and BRCA2 from individual alleles, and we have found that AI in BRCA1 and to a lesser extent BRCA2 is associated with increased breast cancer risk. Furthermore, we have demonstrated that the AI patterns for BRCA1 expression could be transmitted by Mendelian inheritance. Since susceptibility to breast cancer is far from being fully understood, our study suggests that alternate mechanisms, other than deleterious coding mutations, may contribute to breast cancer.

MATERIALS AND METHODS

Databases

RefSeqs (GenBank Accession No: NM_007295.2 and NM_000059.1) were used for *BRCA1* and *BRCA2* mRNA numbering, respectively. The A of ATG translation initiation codon is defined as position +1.

Subjects and genotype analysis

Three populations were used in this study, (i) *BRCA1/2* mutation-negative women reporting a personal and family history of breast cancer, i.e. familial; (ii) female breast cancer patients without a significant family history of disease, i.e. non-familial; and (iii) age-matched cancer-free female controls (Table 1). All participants were Caucasian women with European-American ancestry and were from the Delaware Valley, including the greater Philadelphia Metropolitan area in Pennsylvania. For family studies, eligible subjects were women with a personal and family history of cancer (at least two first and/or second-degree relatives affected with either breast and/or ovarian cancer) and were ascertained from the Family Risk Assessment Program (FRAP) at the Fox Chase Cancer Center (FCCC). All relevant institutional



Figure 3. *BRCA1* allelic expression ratios in cancer-free controls, familial and non-familial breast cancer patients. The AI assays were performed using specific primer and probe sets targeting *BRCA1*-c.4308T/C alleles. Log₂ ratios of *BRCA1*-c.4308T allele to -c.4308C allele expression were presented in cancer-free controls (**A**), familial (**B**) and non-familial breast cancer patients (**C**). (Data expressed as Mean \pm SD, n=3; the mean value of allelic expression ratios of total normal samples has been adjusted to zero).

review boards approved the study protocol and written informed consent was obtained from all participants. Genotype analyses of the two common polymorphisms, *BRCA1*-c.4308T/C and *BRCA2*-c.3396A/G were carried out

using ABI PRISM 7900HT Sequence Detection System and Assays-on-Demand SNP Genotyping products for fluorogenic polymerase chain reaction allelic discrimination (Applied Biosystems, Foster City, CA, USA).



Figure 4. *BRCA2* allelic expression ratios in cancer-free controls, familial and non-familial breast cancer patients. The AI assays were performed using specific primer and probe sets targeting *BRCA2*-c.3396A/G. Log₂ ratios of *BRCA2*-c.3396A allele to -c.3396G allele expression were presented in cancer-free controls (**A**), familial (**B**) and non-familial cancer patients (**C**). (Data expressed as Mean \pm SD, n = 3; the mean value of allelic expression ratios of total normal samples has been adjusted to zero).

Genes	Group	Under-expressed alleles		OR (95% CI)	P-value ^a
BRCA1		c.4308T, Log ₂ [4308T/C]<0	c.4308C, Log ₂ [4308T/C]>0		
	Cancer-free Controls	21	19		
	Familial	9	23	2.82 (1.05, 7.60)	0.02
	Non-Familial	16	16	1.11 (0.44, 2.80)	0.18
BRCA2		c.3396A, Log ₂ [3396A/G]<0	c3396G, Log ₂ [3396A/G]>0		
	Cancer-free Controls	14	17		
	Familial	26	11	0.35	0.02
	Non-familial	13	13	0.82	0.20

Table 3. Distribution of under-expressed alleles of BRCA1 and BRCA2

 $^{a}A~\chi^{2}$ test was used to assess the 2 by 2 tables.

Table 4. Allelic expression and haplotype analysis of BRCA1 in sisters from three breast cancer-prone families

Family	Members	Allelic expression [Log ₂ (<i>BRCA1</i> -c.4308T/C)]	Haplotypes D17S855-D17S1322-D17S1325
Family A	Proband ^a	0.375 + 0.060	145/155-121/121-193/193
5	Sis-01	0.007 + 0.147	145/153-121/121-195/193
	Sis-02	0.382 + 0.176	145/155-121/121-193/193
Family B	Proband ^b	0.477 + 0.070	145/151-121/124-193/193
5	Sis-01	0.232 + 0.214	145/151-121/124-193/193
Family C	Proband ^b	0.583 ± 0.243	145/153-121/127-189/189
-	Sis-01	0.522 ± 0.156	145/153-121/127-189/189

^aOvarian cancer carrier.

^bBreast cancer carrier.

Allelic imbalance assay

A 1.25 µl of the cDNA synthesized in the RT reaction was used in a real-time PCR reaction (25 µl total volume), performed with ABI PRISM 7900HT Sequence Detection System following methods recommended by the manufacturer. Optimal conditions were as follows: Step 1, 95°C for 10 min; Step 2, 92°C for 15 s, 60°C for 60 s with Optics; repeated for 40 cycles. The primer and probe sets used in real-time PCR reaction to detected BRCA1-c.4308T/C (rs1060915) and BRCA2-c.3396A/G (rs1801406) allelic expression were obtained from Applied Biosystem TaqMan® SNP Assay program (Assay ID: C.3178676 and C.7605673.1 for BRCA1 and BRCA2, respectively). Sequence information for primers and probes is available upon request. Each 96-well PCR plate included negative controls, positive controls and unknown samples. Real-time PCR data were analyzed with ABI SDS 2.2.2 software. In order to produce the BRCA1 allelic expression standard curve, cDNAs from the two samples with homozygous genotypes, BRCA1-c.4308T/T and BRCA2-c.4308C/C, were mixed as the following ratios: 8:1, 4:1, 2:1, 1:1, 1:2, 1:4 and 1:8 (c.4308T/T allele:c.4308C/T allele). For the same purpose, cDNAs from the two samples with homozygous genotypes, BRCA2-c.3396A/A and BRCA2-c.3396G/G, were mixed as the following ratios: 8:1, 4:1, 2:1, 1:1, 1:2, 1:4 and 1:8 (c.3396A/A allele:c.3396G/G allele).

The principles of quantitative real-time PCR provide the basis of this linear relation between Log_2 ratio and ΔC_T established in our approach to detect AI (40,41). Previous data have shown that AmpliTaq DNA polymerase cleaves the matched and well-hybridized probe and target sequences and produces

a fluorescent signal (42). In contrast, mismatches between a probe and target are expected to reduce the efficiency of probe hybridization, and AmpliTaq DNA polymerase is more likely to displace a mismatched probe without cleaving it, which does not produce a fluorescent signal.

Theoretically, Allele 1 gene copy number (detected by FAM):

$$Log_2[Allele - 1] = -A_1 * C_{T1} + B_1$$
 (1)

and Allele 2 gene copy number (detected by VIC):

$$Log_2[Allele - 2] = -A_2 * C_{T2} + B_2$$
 (2)

If the fluorescence probes have the same efficiency to hybridize with matched target sequence, that is, $A_1 = A_2 = A$, therefore,

$$Log_{2}[Allele - 1/2] = A*(C_{T2} - C_{T1}) + (B_{1} - B_{2})$$
 (3)

The function (3) was confirmed by two standard curves, Log_2 (c.4308T/C) = $-0.0877 + 1.57897 * \Delta C_T$ and Log_2 (c.3396A/G) = $0.11726 + 1.26458 * \Delta C_T$, set up by our experimental data (Fig. 1). Besides using function (3) to calculate the ratio of mRNA expression between the two alleles, function (1) and function (2) are able to be applied for examining the absolute value of each allele mRNA expression. However, the direct analysis of single allele expression is often complicated by the potential variations between individuals with different environmental or physiological background rather than genetic factors. Comparing the relative expression levels of two alleles of the same gene within the same biologic sample will help to minimize these variations.

Peripheral blood lymphocytes and LCLs

Lymphocytes were isolated from peripheral blood and stored at -150° C until needed. None of the blood samples from breast cancer patients were collected at the time of chemoor radiation therapy. In addition, a subset of cyropreserved lymphocytes from BRCA1 or BRCA2 mutation carriers (e.g. BRCA1-c.3671ins4 and BRCA2-c.796delT) or disease-free individuals were infected with EBV to establish immortal LCLs. LCLs were maintained in RPMI (GIBCO BRL) media supplemented with 20% fetal calf serum and antibiotics at 37°C, 5% CO₂ atmospheric condition and 95% humidity. The immortalized LCLs from cancer-free individuals that had been tested negative for mutations in BRCA1 and BRCA2 served as wild-type controls. To prevent potential degradation of unstable transcripts by NMD a translation inhibitor, puromycin (Sigma, St Louis, MO, USA) was added to the LCL cells as described in a previous study (12).

Subcloning the PCR product and sequence analysis

PCR fragments containing a common polymorphism and deleterious mutation were subcloned directly into pCR®4-TOPO vector (Invitrogen, Carlsbad, CA, USA). PCR was then performed to identify bacterial colonies containing appropriate inserts. Plasmid DNA was purified using QIAfilterTM Plasmid Maxi Kit (Qiagen Inc., Valencia, CA, USA) and the insert was sequenced using either the universal M13-primers or the primers for PCR reactions.

RNA isolation and reverse transcription (RT)

Total cellular RNAs were isolated from blood lymphocyte pellets using TRIzol reagent according to the protocols provided by the manufacturer (Invitrogen Corp., Carlsbad, CA, USA). Purified RNAs were further processed to remove any contaminating DNA (DNA-free kit, Ambion, Inc., Houston, TX, USA). After quantification with Bioanalyzer-2100 system using RNA 6000 Nano LabChip kits (Agilent Technologies, Palo Alto, CA, USA), 2 µg of total RNA from each sample was used as a template to be reverse-transcribed (RT) in a 20 µl reaction [containing 5 µM random hexamers, 500 μ M deoxynucleoside triphosphate mix, 1 × RT (reverse transcriptase) buffer, 5 mM MgCl₂, 1.5 units of RNase inhibitor and 7.5 units of MuLV reverse transcriptase]. All reagents were purchased from Applied Biosystems (Branchburg, NJ, USA). The RT reaction conditions were 10 min at 25°C, 1 h at 42°C and 5 min at 94°C.

Haplotype analysis

Haplotypes were constructed for *BRCA1* using three polymorphic microsatellite repeat markers located within (D17S855 and D17S1322) or adjacent (D17S1325) to the *BRCA1* locus. The sequences of the primer pairs were obtained from the Genome Database (http://www.gdb.org) and PCR reaction was carried out as previously reported (43,44). PCR products with fluorescent dye (HEX) labeled primer were mixed with Hi-Di Formamide and a fluorescent labeled internal size marker. The mixture was subjected to

electrophoreseis on an ABI 3100 Automated DNA Sequencer (Applied Biosystems, Foster City, CA, USA) and the data were analyzed by the GeneScan (Version 3.7) and GeneMapper (Version 4.0) software provided by the manufacturer.

Statistical analysis

Allele specific real-time PCR data were analyzed with ABI SDS software v2.2.2 (Applied Biosystems, Foster City, CA, USA). Statistical analysis was conducted using the SAS System (version 9) developed by the SAS Institute, Inc. (Cary, NC, USA). Student's *t*-test was employed for continuous data and results were presented as the mean \pm SD. We compared the distribution of under-expressed alleles in *BRCA1* or *BRCA2* between cases and controls using χ^2 95% confidence intervals (CI) and the difference in distribution of under-expressed alleles was estimated as odds ratios (OR). A value of *P* < 0.05 is considered significant.

SUPPLEMENTARY MATERIAL

Supplementary Material is available at HMG Online.

ACKNOWLEDGEMENTS

We would like to acknowledge Dr S. Litwin for help with statistical analyses and Ms K.A. Cattie with haplotype analysis. We also thank the numerous volunteers for providing blood samples for these studies.

Conflict of Interest statement. None of the authors has any conflict of interest.

FUNDING

This work is supported in part by Cheryl Herman and the Eileen Stein-Jacoby Fund, grants from the National Cancer Institute, i.e. P50 CA83638 (A.K.G.), U01 CA69631 (M.B.D.) and R25 CA057708 (S.W.); a fellowship from the Department of Defense, W81XWH-04-1-0573 (X.C.); and DOD grants, DAMD17-03-1-0707 (A.K.G.) and DAMD17-03-1-0312 (A.K.G.); and by an appropriation from the Commonwealth of Pennsylvania.

REFERENCES

- Jemal, A., Siegel, R., Ward, E., Murray, T., Xu, J. and Thun, M.J. (2007) Cancer statistics. *CA Cancer J. Clin.*, 57, 43–66.
- Collaborative Group on Hormonal Factors in Breast Cancer. (2001) Familial breast cancer: collaborative reanalysis of individual data from 52 epidemiological studies including 58,209 women with breast cancer and 101,986 women without the disease. *Lancet*, **358**, 1389–1399.
- Dite, G.S., Jenkins, M.A., Southey, M.C., Hocking, J.S., Giles, G.G., McCredie, M.R., Venter, D.J. and Hopper, J.L. (2003) Familial risks, early-onset breast cancer, and BRCA1 and BRCA2 germline mutations. *J. Natl Cancer Inst.*, 95, 448–457.
- Bove, B., Dunbrack, R.L., Jr and Godwin, A.K. (2002) In Pasqualini, J.R. (eds), *Breast Cancer: Prognosis, Treatment and Prevention*. Marcel Dekker Inc., pp. 555–624.
- King, M.C., Marks, J.H. and Mandell, J.B. (2003) Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science*, 302, 643–646.

- McClain, M.R., Palomaki, G.E., Nathanson, K.L. and Haddow, J.E. (2005) Adjusting the estimated proportion of breast cancer cases associated with BRCA1 and BRCA2 mutations: public health implications. *Genet. Med.*, 7, 28–33.
- Fan, S., Yuan, R., Ma, Y.X., Meng, Q., Goldberg, I.D. and Rosen, E.M. (2001) Mutant BRCA1 genes antagonize phenotype of wild-type BRCA1. *Oncogene*, 20, 8215–8235.
- Holt, J.T., Thompson, M.E., Szabo, C., Robinson-Benion, C., Arteaga, C.L., King, M.C. and Jensen, R.A. (1996) Growth retardation and tumour inhibition by BRCA1. *Nat. Genet.*, **12**, 298–302.
- Xia, F., Taghian, D.G., DeFrank, J.S., Zeng, Z.C., Willers, H., Iliakis, G. and Powell, S.N. (2001) Deficiency of human BRCA2 leads to impaired homologous recombination but maintains normal nonhomologous end joining. *Proc. Natl Acad. Sci. USA*, **98**, 8644–8649.
- Xu, X., Qiao, W., Linke, S.P., Cao, L., Li, W.M., Furth, P.A., Harris, C.C. and Deng, C.X. (2001) Genetic interactions between tumor suppressors Brca1 and p53 in apoptosis, cell cycle and tumorigenesis. *Nat. Genet.*, 28, 266–271.
- Jonkers, J., Meuwissen, R., van der Gulden, H., Peterse, H., van der Valk, M. and Berns, A. (2001) Synergistic tumor suppressor activity of BRCA2 and p53 in a conditional mouse model for breast cancer. *Nat. Genet.*, 29, 418–425.
- Chen, X., Truong, T.T., Weaver, J., Bove, B.A., Cattie, K., Armstrong, B.A., Daly, M.B. and Godwin, A.K. (2006) Intronic alterations in BRCA1 and BRCA2: effect on mRNA splicing fidelity and expression. *Hum. Mutat.*, 27, 427–435.
- Esteller, M., Silva, J.M., Dominguez, G., Bonilla, F., Matias-Guiu, X., Lerma, E., Bussaglia, E., Prat, J., Harkes, I.C., Repasky, E.A. *et al.* (2000) Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J. Natl Cancer Inst.*, **92**, 564–569.
- Lambie, H., Miremadi, A., Pinder, S.E., Bell, J.A., Wencyk, P., Paish, E.C., Macmillan, R.D. and Ellis, I.O. (2003) Prognostic significance of BRCA1 expression in sporadic breast carcinomas. *J. Pathol.*, 200, 207– 213.
- Russell, P.A., Pharoah, P.D., De Foy, K., Ramus, S.J., Symmonds, I., Wilson, A., Scott, I., Ponder, B.A. and Gayther, S.A. (2000) Frequent loss of BRCA1 mRNA and protein expression in sporadic ovarian cancers. *Int. J. Cancer*, 87, 317–321.
- Wang, C., Horiuchi, A., Imai, T., Ohira, S., Itoh, K., Nikaido, T., Katsuyama, Y. and Konishi, I. (2004) Expression of BRCA1 protein in benign, borderline, and malignant epithelial ovarian neoplasms and its relationship to methylation and allelic loss of the BRCA1 gene. *J. Pathol.*, 202, 215–223.
- Thrall, M., Gallion, H.H., Kryscio, R., Kapali, M., Armstrong, D.K. and DeLoia, J.A. (2006) BRCA1 expression in a large series of sporadic ovarian carcinomas: a Gynecologic Oncology Group study. *Int. J. Gynecol. Cancer*, 16 (Suppl. 1), 166–171.
- Baldwin, R.L., Nemeth, E., Tran, H., Shvartsman, H., Cass, I., Narod, S. and Karlan, B.Y. (2000) BRCA1 promoter region hypermethylation in ovarian carcinoma: a population-based study. *Cancer Res.*, 60, 5329– 5333.
- Yan, H., Yuan, W., Velculescu, V.E., Vogelstein, B. and Kinzler, K.W. (2002) Allelic variation in human gene expression. *Science*, **297**, 1143.
- Yan, H., Dobbie, Z., Gruber, S.B., Markowitz, S., Romans, K., Giardiello, F.M., Kinzler, K.W. and Vogelstein, B. (2002) Small changes in expression affect predisposition to tumorigenesis. *Nat. Genet.*, **30**, 25–26.
- Matyas, G., Giunta, C., Steinmann, B., Hossle, J.P. and Hellwig, R. (2002) Quantification of single nucleotide polymorphisms: a novel method that combines primer extension assay and capillary electrophoresis. *Hum. Mutat.*, 19, 58–68.
- Lo, H.S., Wang, Z., Hu, Y., Yang, H.H., Gere, S., Buetow, K.H. and Lee, M.P. (2003) Allelic variation in gene expression is common in the human genome. *Genome. Res.*, 13, 1855–1862.
- Buckland, P.R. (2004) Allele-specific gene expression differences in humans. *Hum. Mol. Genet.*, 13(Spec no. 2), R255–R260.
- Bieche, I., Nogues, C. and Lidereau, R. (1999) Overexpression of BRCA2 gene in sporadic breast tumours. *Oncogene*, 18, 5232–5238.
- Hilton, J.L., Geisler, J.P., Rathe, J.A., Hattermann-Zogg, M.A., DeYoung, B. and Buller, R.E. (2002) Inactivation of BRCA1 and BRCA2 in ovarian cancer. J. Natl Cancer Inst., 94, 1396–1406.

- Futreal, P.A., Liu, Q., Shattuck-Eidens, D., Cochran, C., Harshman, K., Tavtigian, S., Bennett, L.M., Haugen-Strano, A., Swensen, J., Miki, Y. *et al.* (1994) BRCA1 mutations in primary breast and ovarian carcinomas. *Science.* 266, 120–122.
- Lancaster, J.M., Wooster, R., Mangion, J., Phelan, C.M., Cochran, C., Gumbs, C., Seal, S., Barfoot, R., Collins, N., Bignell, G. *et al.* (1996) BRCA2 mutations in primary breast and ovarian cancers. *Nat. Genet.*, 13, 238–240.
- Newman, B., Mu, H., Butler, L.M., Millikan, R.C., Moorman, P.G. and King, M.C. (1998) Frequency of breast cancer attributable to BRCA1 in a population-based series of American women. *JAMA*, 279, 915–921.
- Southey, M.C., Tesoriero, A.A., Andersen, C.R., Jennings, K.M., Brown, S.M., Dite, G.S., Jenkins, M.A., Osborne, R.H., Maskiell, J.A., Porter, L. *et al.* (1999) BRCA1 mutations and other sequence variants in a population-based sample of Australian women with breast cancer. *Br. J. Cancer*, **79**, 34–39.
- Swensen, J., Hoffman, M., Skolnick, M.H. and Neuhausen, S.L. (1997) Identification of a 14 kb deletion involving the promoter region of BRCA1 in a breast cancer family. *Hum. Mol. Genet.*, 6, 1513–1517.
- 31. Smith, L.D., Tesoriero, A.A., Ramus, S.J., Dite, G., Royce, S.G., Giles, G.G., McCredie, M.R., Hopper, J.L. and Southey, M.C. (2007) BRCA1 promoter deletions in young women with breast cancer and a strong family history: a population-based study. *Eur. J. Cancer*, 43, 823–827.
- Brown, M.A., Lo, L.J., Catteau, A., Xu, C.F., Lindeman, G.J., Hodgson, S. and Solomon, E. (2002) Germline BRCA1 promoter deletions in UK and Australian familial breast cancer patients: identification of a novel deletion consistent with BRCA1:psiBRCA1 recombination. *Hum. Mutat.*, 19, 435–442.
- 33. Zhou, X.P., Waite, K.A., Pilarski, R., Hampel, H., Fernandez, M.J., Bos, C., Dasouki, M., Feldman, G.L., Greenberg, L.A., Ivanovich, J. *et al.* (2003) Germline PTEN promoter mutations and deletions in Cowden/ Bannayan-Riley-Ruvalcaba syndrome result in aberrant PTEN protein and dysregulation of the phosphoinositol-3-kinase/Akt pathway. *Am. J. Hum. Genet.*, **73**, 404–411.
- Pezzolesi, M.G., Zbuk, K.M., Waite, K.A. and Eng, C. (2007) Comparative genomic and functional analyses reveal a novel cis-acting PTEN regulatory element as a highly conserved functional E-box motif deleted in Cowden syndrome. *Hum. Mol. Genet.*, 16, 1058–1071.
- Signori, E., Bagni, C., Papa, S., Primerano, B., Rinaldi, M., Amaldi, F. and Fazio, V.M. (2001) A somatic mutation in the 5'-UTR of BRCA1 gene in sporadic breast cancer causes down-modulation of translation efficiency. *Oncogene*, **20**, 4596–4600.
- 36. Esquela-Kerscher, A. and Slack, F.J. (2006) Oncomirs-microRNAs with a role in cancer. *Nat. Rev. Cancer*, **6**, 259–269.
- Lai, E.C. (2002) Micro RNAs are complementary to 3'-UTR sequence motifs that mediate negative post-transcriptional regulation. *Nat. Genet.*, 30, 363–364.
- Wu, L., Fan, J. and Belasco, J.G. (2006) MicroRNAs direct rapid deadenylation of mRNA. Proc. Natl Acad. Sci. USA, 103, 4034–4039.
- Rusinov, V., Baev, V., Minkov, I.N. and Tabler, M. (2005) MicroInspector: a web tool for detection of miRNA binding sites in an RNA sequence. *Nucleic Acids Res.*, 33, W696–W700.
- Chiang, P.W., Song, W.J., Wu, K.Y., Korenberg, J.R., Fogel, E.J., Van Keuren, M.L., Lashkari, D. and Kurnit, D.M. (1996) Use of a fluorescent-PCR reaction to detect genomic sequence copy number and transcriptional abundance. *Genome Res.*, 6, 1013–1026.
- Heid, C.A., Stevens, J., Livak, K.J. and Williams, P.M. (1996) Real time quantitative PCR. *Genome Res.*, 6, 986–994.
- Livak, K.J., Flood, S.J., Marmaro, J., Giusti, W. and Deetz, K. (1995) Oligonucleotides with fluorescent dyes at opposite ends provide a quenched probe system useful for detecting PCR product and nucleic acid hybridization. *PCR Methods Appl.*, 4, 357–362.
- Weitzel, J.N., Lagos, V., Blazer, K.R., Nelson, R., Ricker, C., Herzog, J., McGuire, C. and Neuhausen, S. (2005) Prevalence of BRCA mutations and founder effect in high-risk Hispanic families. *Cancer Epidemiol. Biomarkers Prev.*, 14, 1666–1671.
- Rudkin, T.M., Hamel, N., Galvez, M., Hogervorst, F., Gille, J.J., Moller, P., Apold, J. and Foulkes, W.D. (2006) The frequent BRCA1 mutation 1135insA has multiple origins: a haplotype study in different populations. *BMC Med. Genet.*, 7, 15.

BRCA1-Associated Proteins: Novel Targets for Breast Cancer Radiation Therapy

Xiaowei Chen^a, Jamie L. Kistler, and Andrew K. Godwin

^aAuthor for correspondence:
Department of Medical Oncology
Fox Chase Cancer Center
333 Cottman Avenue
Philadelphia, PA 19111-2409, USA
Tel: 215-214-4286;
Fax: 215-728-2741;
E-mail: Xiaowei.Chen@fccc.edu

Abstract

Tumor cells, in general, are genomically unstable and have defects in DNA repair pathways, which subsequently hinder DNA damage responses. It has been proposed that therapeutic strategies specifically targeting DNA repair pathway proteins may lead to an increased therapeutic index in tumor cells versus normal cells. The BRCA1 pathways are known to play a critical role in DNA repair; thus, breast tumors with defects in proteins associated with the BRCA1 pathways are believed to be more sensitive to DNA damagebased therapies. BRCA1 can interact directly or indirectly with other tumor suppressors, DNA damage sensors, ubiquitin ligase partners, and signal transducers to form multisubunit protein complexes. These protein complexes are involved in a broad range of biological processes including DNA repair, cell cycle control, ubiquitination, and chromatin remodeling. Growing evidence suggests that mutation and/or aberrant expression of one or more key members of the BRCA1-associated multi-protein complexes may result in loss of normal BRCA1 activity and disruption of the BRCA1 pathways. These BRCA1-associated proteins are potential modifiers of BRCA1 functions and, therefore, potential targets for sensitizing breast cancer cells to radiation therapy.

Keywords: BRCA1, *breast cancer, radiation resistance, DNA repair, cell cycle, ubiquitination, chromatin remodeling.*

Introduction

Breast cancer is the most common cancer affecting women, with an estimated lifetime risk of approximately 10% by 80 years of age. In the United States, it is estimated that approximately 182,450 new cases of female breast cancer will be diagnosed and greater than 40,000 breast cancer-related deaths will occur in 2008 (Jemal *et al.*, 2008). Approximately 13.2% of all American women (1 in 8) are expected to develop breast cancer sometime during their lifetime and 3.0% will subsequently die from the disease (Ries *et al.*, 2008). Despite advances in treatment and early detection, the breast cancer mortality rate among women in the United States decreased by only 2.2% per year between 1990 and 2002 (Jemal *et al.*, 2008).

Importantly, estimates from previous studies (Collaborative Group on Hormonal Factors in Breast Cancer, 2001; Margolin *et al.*, 2006) indicate that family history is associated with 15% to 20% of breast cancer cases in the United States. The *BRCA1* gene (OMIM: 113705) is one of the most intensively studied breast cancer susceptibility genes and has a profound role in breast cancer etiology owing to its involvement in several important cellular processes. Deleterious mutations in *BRCA1* are thought to account for approximately 10% to 20 % of hereditary breast cancers (Bove *et al.*, 2002; King *et al.*, 2003; Walsh *et al.*, 2006). Among its many biological functions, the BRCA1 protein is involved in DNA repair. Because DNA repair pathways and associated proteins are targeted by radiation therapy, there is considerable interest in the development of novel therapeutic strategies to sensitize breast cancer patients with mutations in *BRCA1* to radiation therapy. This article will provide an overview of BRCA1 and its associated

proteins with a particular emphasis on their role in DNA repair, as well as summarize current paradigms for breast cancer treatment with a focus on the development of new strategies to exploit the role of BRCA1 associated proteins and improve the efficacy of breast cancer radiation therapy.

Radiation Therapy for Breast Cancer

Current treatment paradigms for breast cancer are complex and reflect the considerable heterogeneity of the disease (NCCN breast cancer treatment guidelines 2008). Treatment options for noninvasive breast cancers range from observation alone, to breast-conserving lumpectomy with or without breast radiation, to total mastectomy depending on the tumor biology and individual patient risk. Radiation therapy is also used in the adjuvant setting and in combination with lumpectomy for locoregional treatment of early stage invasive breast cancers. Systemic treatment for advanced or metastatic breast cancer includes chemotherapy, endocrine therapy, and newer types of targeted therapeutic agents (e.g., targeted monoclonal antibodies and tyrosine kinase inhibitors). Radiation therapy has been a treatment modality for breast cancer patients for more than 100 years and, over the last 3 decades, has become a critical component of successful treatment strategies for breast cancer. An increasing role for radiation therapy developed in the early 1970s, when Fletcher documented that radiation therapy was instrumental in decreasing local recurrences (Fletcher, 1972). In particular, supraclavicular metastases were reduced from 20% to 25% to only 1.3% to 3% with the addition of ionizing radiation (IR). Radiation therapy has also been utilized to treat patients with tumors that have undergone total mastectomy resulting in a reduction in local recurrences by greater

than two-thirds (Fletcher, 1972). This early work led to an expanded role for radiation therapy in breast cancer.

The emergence of radiation therapy to the forefront of modern breast cancer treatment lies in its application in breast conservation therapy. Current NCCN treatment guidelines support the preferred use of breast conservation therapy (i.e., lumpectomy with or without breast radiation) as a breast treatment for the majority of women with early stage breast cancers (i.e., ductal carcinoma in situ, stage I and II breast cancers). Evidence suggests that the addition of radiation therapy may significantly reduce recurrence in this patient population. Landmark studies on the necessity of radiation therapy in breast conservation therapy came from Fisher and colleagues, as a part of a clinical trial conducted by the National Surgical Adjuvant Breast and Bowel Project (NSABP; Protocol B-06) that showed lumpectomy with radiation therapy had much lower recurrence rates than lumpectomy alone (10% versus 35%; P < 0.001) after 12 years of follow-up (Fisher et al., 1995). This observation has been further validated by an extensive meta-analysis that supported improved local control with the addition of radiation therapy (Fisher et al., 2002; Veronesi et al., 2002). Recently, clinical research has examined the possible survival benefits of radiation therapy in breast cancer. The Early Breast Cancer Trialists' Collaborative Group (EBCTG) examined 78 trials involving more than 42,000 patients with breast cancer (Clarke et al., 2006). In the analyses of trials directly comparing patients receiving radiation therapy versus those not receiving radiation therapy, a clear reduction in local recurrences occurred in the radiotherapy group, including patients undergoing mastectomy or breast conservation

therapy (Clarke *et al.*, 2006). Interestingly, there was also a notable improvement in survival among patients treated with radiotherapy. In fact, patients receiving radiotherapy for their breast cancer had a nearly 6% reduction in their 15-year breast cancer mortality risk and a 4% to 5% reduction in overall mortality (Clarke et al., 2006). These findings support the contribution of radiotherapy to both the reduction of local recurrences and in 15-year overall mortality rates. Researchers have noted that breast cancer recurrences in the non-irradiated breast often occur within 3 years of initial diagnosis (Kurtz et al., 1989). In comparison, local recurrences in irradiated breast tissue occur much later, with the risk increasing with time (7% risk at 5 years, 14% risk at 10 years, and 20% risk at 20 years) (Smith et al., 2000). Thus, despite the benefits of radiation therapy in the treatment of breast cancer, patients continue to develop local recurrences in the targeted breast. The persistent recurrence of breast cancers following radiation therapy in multiple patient settings has prompted significant research efforts, particularly in understanding the etiology of radioresistant breast tumors and subsequent development of novel treatment paradigms to overcome this resistance.

Despite the benefits of radiation therapy in the treatment of breast cancer, patients continue to develop local recurrences in the targeted breast. Researchers have noted that breast cancer recurrences in the non-irradiated breast often occur within 3 years of initial diagnosis (Kurtz et al., 1989). In comparison, local recurrences in irradiated breast tissue occur much later, with the risk increasing with time (7% risk at 5 years, 14% risk at 10 years, and 20% risk at 20 years) (Smith et al., 2000). It is these recurrences that have spurred research into both breast cancer recurrences and the possibility of radio-resistant

breast tumors. One of the major radio-resistance mechanisms is related to a score of genes which are involved in the repair of DNA damage by radiation.

BRCA1 in Resistance to Breast Cancer Radiation Therapy

The clinical benefit of radiation therapy can be attributed to its mechanism of DNA damage and subsequent activation of apoptosis pathways. The damage caused by IR activates specific DNA damage cell cycle checkpoints, which leads to induction of various DNA repair pathways. The central component of these pathways is the ATM/CHEK2 kinase, which is activated upon DNA damage and subsequently phosphorylates multiple proteins, including BRCA1 (Canman et al., 1998; Cortez et al., 1999; Lee et al., 2000). In response to DNA damage induced by IR, BRCA1 is phosphorylated at specific tyrosine residues by ATM (the gene mutated in ataxia telangiectasia), CHEK2 (the human homologue of yeast checkpoint protein kinase [hCds1]), or by the ATM-related kinase, ATR (Cortez et al., 1999; Lee et al., 2000; Tibbetts et al., 1999). This phosphorylation, which occurs in a region containing clusters of serine-glutamine residues, has been shown to be functionally important using mouse models. In these studies, a mutated form of *BRCA1* lacking these phosphorylation sites failed to rescue radiation hypersensitivity when introduced into BRCA1-deficient cells (Cortez et al., 1999). In addition, phosphorylation by ATM/CHEK2 following DNA damage is critical for the recruitment of BRCA1 to both DNA repair and chromatin remodeling protein complexes (Zhong et al., 1999).

BRCA1 has been implicated in normal cellular processes, including DNA fidelity and damage repair, and has therefore been examined as having a possible role in the radioresistance of breast tumors. However, the specific role of BRCA1 in radioresistant breast cancer remains somewhat unclear. In vitro studies (Abbott et al., 1999; Foray et al., 1999; Mamon et al., 2003; Ruffner et al., 2001; Shen et al., 1998) have demonstrated an increased sensitivity to IR when *BRCA1* is mutated in human breast cancer cell lines. However, clinical observations in breast cancer patients fail to reliably support these in vitro findings (Baeyens et al., 2004; Garcia-Higuera et al., 2001; Leong et al., 2000). One study (Kirova et al., 2005) found that BRCA1 mutation carriers exhibited increased sensitivity to radiation therapy as assessed by the reduced rate of breast cancer recurrence following breast conserving treatment; however, Pierce and colleagues (Pierce et al., 2000) noted no significant differences in local recurrences between BRCA1 mutation carriers and patients with sporadic forms of breast cancer in a multicenter study. Two additional human studies (Baeyens et al., 2004; Leong et al., 2000) indicated that mutations in *BRCA1* may not account for clinical radiation hypersensitivity. These conflicting findings pose the question of whether BRCA1 mutations will indeed increase the sensitivities of tumor cells to the radiation-based therapies. Therefore, the role of BRCA1 and its influence on tumor cell sensitivity to radiation in vitro and in vivo will require further investigation.

Role of BRCA1 and Associated Proteins in Breast Cancer Etiology

Since its cloning and characterization in the mid-1990s (Miki *et al.*, 1994), BRCA1 has been implicated in many cellular processes including DNA repair, cell-cycle-checkpoint

control, protein ubiquitination, and chromatin remodeling. Although mutations in *BRCA1* are known to contribute to the development of hereditary breast and ovarian cancers, *BRCA1* mutations in sporadic breast cancers, which account for approximately 90% of all breast cancers, are surprisingly rare (Futreal *et al.*, 1994). In this aspect, various studies have indicated that loss of *BRCA1* expression through epigenetic mechanisms may contribute about 10% of sporadic breast cancer (Esteller *et al.*, 2000; Rio *et al.*, 1999; Yang *et al.*, 2001). In addition, accumulating evidence suggests that dysfunction of other genes, coding for proteins in pathways complementary to BRCA1, may be important in the pathogenesis of a significant proportion of sporadic, non-hereditary cancers. This hypothesis is supported by several lines of evidence, including phenotypic analyses of breast and ovarian tumors, as well as mechanistic studies of BRCA1-associated pathways (Farmer *et al.*, 2005; Jazaeri *et al.*, 2002).

BRCA1-Associated Proteins: Functional Modifiers of BRCA1 Activity

Due to its clinical significance, the *BRCA1* gene is one of the most intensively studied breast cancer susceptibility genes. The *BRCA1* gene encodes for a 220 kDa nuclear phosphoprotein that has been suggested to play a role in maintaining genomic stability and to act as a tumor suppressor. The BRCA1 protein interacts directly or indirectly with other tumor suppressor proteins (e.g., p53 and BRCA2), DNA damage sensors (e.g., RAD51, RAD50, MRE11 and NBS1), signal transducers (e.g., p21 and cyclin B), and ubiquitination proteins (e.g., BARD1, BRCC36, and RAP80) to form multi-subunit protein complexes (**Figure 1**), such as the BRCA1-associated genome surveillance complex (BASC) and the BRCA1 and BRCA2 containing complex (BRCC). Importantly, the proper formation of these multi-subunit protein complexes is critical in carrying out the multiple biological processes associated with BRCA1, including DNA repair, cell cycle control, chromatin remodeling, and ubiquitination.

The majority of BRCA1 functional studies have focused on its potential role in DNA damage responses. The implication that BRCA1 is a direct component of DNA damage response pathways comes from evidence of its interactions with BRCA2 and RAD51. The protein complex comprised of BRCA1, BRCA2, and RAD51 has been shown to activate DNA double-strand break (DSB) repair and to initiate homologous recombination, an observation which links the maintenance of genomic integrity to tumor suppression (Chen *et al.*, 1999). In addition, the BRCA1-associated MRE11-RAD50-NBS1 (MRN) complex has recently been demonstrated to activate CHEK2 downstream from ATM in response to replication-mediated DSBs (Takemura *et al.*, 2006). Disruption of any of these pathways may contribute to increased genomic instability and potentially sensitize cells to the effects of IR, specifically through the induction of cellular apoptosis.

The involvement of BRCA1 and its associated partners in normal DNA repair processes suggests that mutations in these tumor suppressor proteins would hinder DNA damage responses, predispose cells to additional accumulated mutations, and potentially contribute to subsequent malignant transformation. Importantly, compromised DNA repair mechanisms would also be expected to sensitize cells to the lethal effects of IR. Thus, while *BRCA1* mutations may play a profound role in breast cancer etiology, consequent disruption of normal DNA repair may actually be therapeutically exploited to

increase clinical radiation hypersensitivity in breast cancer patients who are *BRCA1* mutation carriers.

BRCA1 has also been shown to play a role in cell cycle control. For example, BRCA1 stimulates expression of the cyclin-dependent kinase (CDK) inhibitor, p21, and to inhibit cell-cycle progression into the S-phase (Somasundaram et al., 1997). In addition, research has shown that BRCA1 is not only essential for activating the CHEK1 kinase that regulates G2/M arrest induced by DNA damage, but also controls the expression, phosphorylation, and cellular localization of Cdc25C and Cdc2/cyclin B kinases (Yarden et al., 2002). Therefore, BRCA1 appears to be involved in regulating the onset of mitosis. Furthermore, a mouse study demonstrated that BRCA1 knockout mice, generated by removal of exon 11, have a defective G_2/M cell cycle checkpoint and extensive chromosomal abnormalities (Xu et al., 1999). It is also reported that elimination of one Tp53 allele (BRCA1 exon 11 - 77p53 + 7) rescued the embryonic lethality caused by the deletion of BRCA1 exon 11 and restored normal mammary gland development (Xu et al., 2001). However, most female mice homozygous for the *Brcal* exon 11 deletion and heterozygous for loss of the Tp53 gene developed mammary tumors within 6 to 12 months. Importantly, the resulting tumors lose the remaining Tp53 allele (Xu *et al.*, 2001). These findings indicated that the genetic interactions between Brca1 and p53 are associated with breast carcinogenesis.

BRCA1 and its associated protein have also been found to be involved in the process of chromatin remodeling. Wang and colleagues (Wang *et al.*, 2000) used

immunoprecipitation and mass spectrometry to identify a large multi-subunit protein complex referred to as BASC (BRCA1-associated genome surveillance complex), which is comprised of ATM, BLM, MSH2, MSH6, MLH1, the RAD50-MRE11-NBS1 complex, and the RFC1-RFC2-RFC4 complex. Confocal microscopy demonstrated that BRCA1, BLM, and the RAD50-MRE11-NBS1 complex co-localized to large nuclear foci, and BASC has subsequently been shown to be involved in chromatin remodeling at sites of double-strand DNA breaks (Wang *et al.*, 2000). In addition, BRCA1 directly interacts with the brahma-related gene 1 (BRG1) subunit of SW1/SNF-associated complex which has been demonstrated to be involved in chromatin-remodeling (Bochar *et al.*, 2000). This finding links chromatin remodeling processes to breast cancer. Furthermore, the BRCT domain of BRCA1 has been reported to be associated with the histone deacetylases, HDAC1 and HDAC2 (Yarden and Brody, 1999). Collectively, these findings may help explain the involvement of BRCA1 in multiple, seemingly unrelated processes such as transcription and DNA repair.

BRCA1 also interacts with a number of proteins and displays significant ubiquitin ligase activities. Importantly, deleterious mutations affecting the BRCA1 RING-finger domain, found in clinical specimens, abolish the ubiquitin ligase activity of BRCA1 (Ruffner *et al.*, 2001; Wu *et al.*, 1996). These findings support a relationship between the ligase activity of BRCA1 and the predisposition to breast cancer. Using a combination of affinity purification of anti-FLAG and mass spectrometric sequencing, a multiprotein protein complex, termed BRCC (<u>BRCA1/2 Containing Complex</u>), which contains seven polypeptides including BRCA1, BRCA2, BARD1 and RAD51, has been identified

(Dong et al., 2003). BRCC is an E3 ubiquitin ligase complex exhibiting activities in the E2-dependent ubiquitination of the tumor suppressor p53 (Dong et al., 2003). In this multiprotein complex, one protein, referred to as BRCC36, has been found to be directly interacted with BRCA1. Cancer-causing truncations of BRCA1 have been found to abrogate the association of BRCC36 with BRCC (Dong et al., 2003). We have also demonstrated that depletion of BRCC36 resulted in increased sensitivity in breast cancer cells to ionizing radiation (IR) and disruption of IR-induced BRCA1 phosphorylation and nuclear foci formation (Chen et al., 2006). Previous study has shown that a recombinant four-subunit BRCC complex containing BRCA1-BARD1-BRCC45-BRCC36 revealed an enhanced E3 ubiquitin ligase activity compared to that of BRCA1-BARD1 heterodimer (Dong et al., 2003). Furthermore, BRCC36 has recently been reported to also be present in a novel BRCA1-associated complex, BRCA1-BARD1-RAP80-Abraxas-BRCC36, which plays a role in recognizing DNA damage site (Wang *et al.*, 2007). These findings may suggest that the role of BRCC36 in DNA damage response could be dynamic and mediated by other protein partners (e.g., BRCC45, BRCC120, RAP80 or Abraxas) in the same complexes (Figure 2). In addition, BRCA1 has also been reported to interact with the RNA Pol II holoenzyme (Scully et al., 1997). Two recent reports have suggested that BRCA1 and BARD1 may be involved in the degradation of RNA polymerase II complex and siRNA-mediated knockdown of BRCA1 and BARD1 results in stabilization of RNAP II in the cells following UV exposure (Kleiman et al., 2005; Starita et al., 2005). These studies reported that BRCA1/BARD1 appears to initiate the degradation of stalled RNAP II and thus disrupts the coupled transcription by inhibiting RNA processing machinery in cells exposed to DNA damage. At present, the known substrates that are

polyubiquitinated by the BRCA1-BARD1 ubiquitin ligase are very limited and include RNA polymerase II, nucleophosmin/B23, and p53 (Dong *et al.*, 2003; Kleiman *et al.*, 2005; Sato *et al.*, 2004; Starita *et al.*, 2005).

BRCA1-associated Proteins as Potential Targets of Breast Cancer Therapies

In the last several decades, efforts have been made toward understanding the mechanism of response to both cytotoxic chemotherapy and radiation therapy in the treatment of breast cancer. Because tumor cells are typically genomically unstable with dysfunctional DNA damage responses, it has been proposed that targeting DNA repair pathways may lead to an increased therapeutic index in tumor cells versus normal cells. The involvement of BRCA1 and its associated partners in normal DNA repair processes suggests that mutations in these tumor suppressor proteins would hinder DNA damage responses, predispose cells to additional accumulated mutations, and potentially contribute to subsequent malignant transformation. Importantly, compromised DNA repair mechanisms would also be expected to sensitize cells to the lethal effects of IR. Thus, while *BRCA1* mutations may play a profound role in breast cancer etiology, consequent disruption of normal DNA repair may actually be therapeutically exploited to increase clinical radiation hypersensitivity in breast cancer patients who are *BRCA1* mutation carriers.

This speculation is supported by the recent development of the inhibitors of poly (ADPribose)-polymerase-1 (PARP). The PARP enzyme is involved in base excision repair which is critical pathway in the repair of DNA single-strand breaks (Ratnam and Low,

2007; Schreiber *et al.*, 2002). Farmer and colleagues have shown that defects in BRCA1 or BRCA2 profoundly sensitize cells to the inhibition of PARP enzymatic activity, resulting in chromosomal instability, cell cycle arrest, and subsequent apoptosis (Farmer *et al.*, 2005). PARP inhibitors are currently in clinical trials of patients with breast cancer or other malignancies who are *BRCA1* or *BRCA2* mutation carriers. Two phase I studies have shown that AZD2281 (AstraZeneca, UK), a potent orally active PARP inhibitor, is well tolerated and leads to significant PARP inhibition in patients carrying *BRCA1* and *BRCA2* mutations with breast or ovarian cancer (Fong *et al.*, 2008; Yap *et al.*, 2007). Importantly, clinical responses have been observed in all cohorts evaluated thus far, and future phase II studies are planned (Fong *et al.*, 2008; Yap *et al.*, 2007). Findings from these recent studies further suggest that the design of novel therapies, which inhibit components of particular DNA repair pathways, may provide effective and more tolerable therapeutic options for breast cancer patients with BRCA1 defects.

In vitro studies have demonstrated that breast cancer cells expressing mutated *BRCA1* have increased sensitivity to IR (Kennedy *et al.*, 2004; Powell, 2005). Notably, mutations in *BRCA1* itself may not be the only reason for loss of the encoded protein's activity. There is growing evidence that disruption of the BRCA1-associated multi-protein complexes, either through mutations or the aberrant expression of a key member(s) of these complexes, may result in loss of normal BRCA1 activity (Chen *et al.*, 2006; McCarthy *et al.*, 2003; Wang *et al.*, 2007; Wu *et al.*, 2007). In our own studies, we have tested the hypothesis that dysregulated expression (e.g., gain or loss) of protein(s) in BRCA1-associated pathways leads to a BRCA1 "null-like" phenotype and subsequent

DNA damage hypersensitivity in breast cancer cells (Chen *et al.*, 2006). As shown in **Figure 3**, BRCA1 and p53 are phosphorylated by the ATM kinase following IR. Depletion of the BRCA1-associated protein, BRCC36, prevents the phosphorylation of BRCA1 and disrupts BRCA1 nuclear foci formation following IR, an event that is associated with the induction of DNA repair. The proposed model illustrates that disruption of BRCA1 activation through depletion of BRCC36 may create an imbalance between the DNA repair and cell survival pathways and the apoptosis/cell death pathways following IR exposure. As a result, abrogation of BRCC36 sensitizes breast cancer cells to IR-induced apoptosis (Chen *et al.*, 2006).

This proposed mechanism is also supported by a number of studies that have demonstrated the impact of cellular resistance to IR upon manipulation of BRCA1associated proteins, such as RAD51, MRE11, and NBS1 (**Table 1**) (Billecke *et al.*, 2002; Boulton *et al.*, 2004; Chinnaiyan *et al.*, 2005; Digweed *et al.*, 2002; Garcia-Higuera *et al.*, 2001; Houghtaling *et al.*, 2005; Kim *et al.*, 2007; Lio *et al.*, 2004; Liu *et al.*, 2007; Nakanishi *et al.*, 2002; Russell *et al.*, 2003; Sobhian *et al.*, 2007; Wang *et al.*, 2007; Yan *et al.*, 2008). In addition, because multiple genetic hits are necessary for tumorigenesis, individuals that carry defects in DNA damage repair/response genes are particularly cancer prone, due to the genetic instability and hypermutability of their cells (Deng, 2006; Jasin, 2002). Therefore, these BRCA1-associated proteins are likely to be involved in tumorigenesis and are potential therapeutic targets.

Summary

Since BRCA1 was cloned a decade ago, significant progress has been made in defining its biochemical and biological functions, as well as its role in breast and ovarian cancers. BRCA1 has been implicated in many cellular processes, including DNA repair, and protein ubiquitination. Because of the important role of BRCA1 in DNA repair, breast tumors with defective BRCA1 are believed to be more sensitive to DNA damage-based therapies. Nevertheless, defects in BRCA1 itself may not be the only reason for the loss of its activity nor the increased sensitivity of tumor cells to DNA damage-based agents. A number of studies have demonstrated that manipulation of BRCA1-associated proteins, such as RAD51, MRE11, and NBS1, can impact cellular sensitivity to IR. BRCA1associated proteins may, therefore, be considered as potential targets for breast cancer therapies. Despite a potentially significant role for BRCA1-associated protein complexes in modifying the activities of BRCA1, the total number of complexes and the identity and function of component proteins has yet to be fully elucidated. Thus, much of the scientific effort related to BRCA1 is currently directed at defining the biochemical functions of BRCA1 in association with these protein complexes.

Acknowledgment

This work was supported in part by Cheryl Herman and the Eileen Stein-Jacoby Fund, a fellowship award from DOD BCRP, W81XWH-04-1-0573 (XC), and a DOD BCRP grant, W81XWH-07-1-0685 (XC).

Protein name	BRCA1 pathway affected	<i>BRCA1</i> gene manipulation approach	Increased resistance to IR	Increased sensitivity to IR
FANCD2	DNA repair	Defected		Garcia-Higuera et al, 2001 Houghtaling et al, 2005
NBS1	DNA repair	Defected	Nakanishi et al, 2002	
MRE11	DNA repair	Disrupted		Digweed et al, 2002
RAD51	DNA repair	Deficiency Blocking		Lio et al, 2004 Russell et al, 2003
		Overexpression	Vispe et al, 1998	
HDAC	Chromatin remodeling	Blocking		Chinnaivan et al, 2005
RB	Cell cycle checkpoint control	Decreasing Defected		Carlson et al, 2000 Billecke et al, 2002
BARD1	Ubiquitination	Depleted		Boulton et al, 2004
BRCC36	Ubiquitination	Depleted		Dong et al., 2003 Chen et al, 2006
RAP80	Ubiquitination	Depleted		Sobhian, et al, 2007 Yan et al, 2008
CCDC98/Abraxas	Ubiquitination	Deplete		Wang et al, 2007 Kim et al, 2007 Liu et al, 2007

Table 1. Radiation Sensitivity Studies Related to BRCA1-associated Proteins

IR, ionizing radiation.

Figure Legends

Figure 1. BRCA1-associated protein network. BRCA1 interacts with a number of proteins to form multi-subunit protein complexes, which are involved in DNA repair, cell cycle checkpoint control, protein ubiquitination, and chromatin remodeling.

Figure 2. BRCC36 in different BRCA1-associated protein complexes (BRCC or BRCA1 A Complex, respectively). Previous study has shown that BRCC36 potentiates the E3 ubiquitin ligase activity of BRCA1-BARD1 heterodimer. Recently, BRCC36 has been reported to also be present in a novel BRCA1-associated complex, BRCA1-BARD1-RAP80-Abraxas-BRCC36, which plays a role in recognizing DSB site.

Figure 3. A proposed model illustrating the role of BRCC36 in BRCA1-associated DNA repair pathway in response to ionizing radiation (IR). BRCA1 and p53 are phosphorylated by the ATM kinase following IR. The BRCA1 and p53 proteins are involved in DNA repair and apoptosis pathways, respectively. Depletion of the BRCA1associated protein, BRCC36, prevents the phosphorylation of BRCA1 and disrupts BRCA1 nuclear foci formation following IR, an event that is associated with the induction of DNA repair. The proposed model illustrates that disruption of BRCA1 activation through depletion of BRCC36 may create an imbalance between the DNA repair and cell survival pathways and the apoptosis/cell death pathways following IR exposure. As a result, abrogation of BRCC36 sensitizes breast cancer cells to IR-induced apoptosis.

REFERENCES

Collaborative Group on Hormonal Factors in Breast Cancer (2001). Familial breast cancer: collaborative reanalysis of individual data from 52 epidemiological studies including 58,209 women with breast cancer and 101,986 women without the disease. *Lancet* **358:** 1389-99.

NCCN breast cancer treatment guidelines (2008).

Abbott DW, Thompson ME, Robinson-Benion C, Tomlinson G, Jensen RA, Holt JT (1999). BRCA1 expression restores radiation resistance in BRCA1-defective cancer cells through enhancement of transcription-coupled DNA repair. *J Biol Chem* **274**: 18808-12.

Baeyens A, Thierens H, Claes K, Poppe B, de Ridder L, Vral A (2004). Chromosomal radiosensitivity in BRCA1 and BRCA2 mutation carriers. *Int J Radiat Biol* **80**: 745-56.

Billecke CA, Ljungman ME, McKay BC, Rehemtulla A, Taneja N, Ethier SP (2002). Lack of functional pRb results in attenuated recovery of mRNA synthesis and increased apoptosis following UV radiation in human breast cancer cells. *Oncogene* **21**: 4481-9.

Bochar DA, Wang L, Beniya H, Kinev A, Xue Y, Lane WS *et al* (2000). BRCA1 is associated with a human SWI/SNF-related complex: linking chromatin remodeling to breast cancer. *Cell* **102**: 257-65.

Boulton SJ, Martin JS, Polanowska J, Hill DE, Gartner A, Vidal M (2004).
BRCA1/BARD1 orthologs required for DNA repair in Caenorhabditis elegans. *Curr Biol* 14: 33-9.

Bove B, Dunbrack R, Godwin AK (2002). BRAC 1, BRAC2, and hereditary breast cancer. In: Pasqualini J (ed). *Breast Cancer: Prognosis, Treatment and Prevention*. Marcel Dekker Inc.: New York.

Canman CE, Lim DS, Cimprich KA, Taya Y, Tamai K, Sakaguchi K *et al* (1998). Activation of the ATM kinase by ionizing radiation and phosphorylation of p53. *Science* **281:** 1677-9.

Chen JJ, Silver D, Cantor S, Livingston DM, Scully R (1999). BRCA1, BRCA2, and Rad51 operate in a common DNA damage response pathway. *Cancer Res* **59**: 1752s-1756s.

Chen X, Arciero CA, Wang C, Broccoli D, Godwin AK (2006). BRCC36 is essential for ionizing radiation-induced BRCA1 phosphorylation and nuclear foci formation. *Cancer Res* **66**: 5039-46.

Chinnaiyan P, Vallabhaneni G, Armstrong E, Huang SM, Harari PM (2005). Modulation of radiation response by histone deacetylase inhibition. *Int J Radiat Oncol Biol Phys* **62**: 223-9.

Clarke M, Collins R, Darby S, Davies C, Elphinstone P, Evans E *et al* (2006). Effects of radiotherapy and of differences in the extent of surgery for early breast cancer on local recurrence and 15-year survival: an overview of the randomised trials. *Lancet* **366**: 2087-106.

Cortez D, Wang Y, Qin J, Elledge SJ (1999). Requirement of ATM-dependentphosphorylation of brca1 in the DNA damage response to double-strand breaks. *Science*286: 1162-6.

Deng CX (2006). BRCA1: cell cycle checkpoint, genetic instability, DNA damage response and cancer evolution. *Nucleic Acids Res* **34:** 1416-26.

Digweed M, Demuth I, Rothe S, Scholz R, Jordan A, Grotzinger C *et al* (2002). SV40 large T-antigen disturbs the formation of nuclear DNA-repair foci containing MRE11. *Oncogene* **21:** 4873-8.

Dong Y, Hakimi MA, Chen X, Kumaraswamy E, Cooch NS, Godwin AK *et al* (2003). Regulation of BRCC, a holoenzyme complex containing BRCA1 and BRCA2, by a signalosome-like subunit and its role in DNA repair. *Mol Cell* **12**: 1087-99. Esteller M, Silva JM, Dominguez G, Bonilla F, Matias-Guiu X, Lerma E *et al* (2000). Promoter hypermethylation and BRCA1 inactivation in sporadic breast and ovarian tumors. *J Natl Cancer Inst* **92:** 564-9.

Farmer H, McCabe N, Lord CJ, Tutt AN, Johnson DA, Richardson TB *et al* (2005).Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 434: 917-21.

Fisher B, Anderson S, Bryant J, Margolese RG, Deutsch M, Fisher ER *et al* (2002). Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med* **347:** 1233-41.

Fisher B, Anderson S, Redmond CK, Wolmark N, Wickerham DL, Cronin WM (1995). Reanalysis and results after 12 years of follow-up in a randomized clinical trial comparing total mastectomy with lumpectomy with or without irradiation in the treatment of breast cancer. *N Engl J Med* **333**: 1456-61.

Fletcher GH (1972). Local results of irradiation in the primary management of localized breast cancer. *Cancer* **29**: 545-51.

Fong PC, Boss DS, Carden CP, Roelvink M, De Greve J, Gourley CM *et al* (2008). AZD2281 (KU-0059436), a PARP (poly ADP-ribose polymerase) inhibitor with single agent anticancer activity in patients with BRCA deficient ovarian cancer: Results from a phase I study. *J Clin Oncol* **26:** A5510.

Foray N, Randrianarison V, Marot D, Perricaudet M, Lenoir G, Feunteun J (1999). Gamma-rays-induced death of human cells carrying mutations of BRCA1 or BRCA2. *Oncogene* **18:** 7334-42.

Futreal PA, Liu Q, Shattuck-Eidens D, Cochran C, Harshman K, Tavtigian S *et al* (1994).BRCA1 mutations in primary breast and ovarian carcinomas. *Science* 266: 120-2.

Garcia-Higuera I, Taniguchi T, Ganesan S, Meyn MS, Timmers C, Hejna J *et al* (2001).Interaction of the Fanconi anemia proteins and BRCA1 in a common pathway. *Mol Cell*7: 249-62.

Houghtaling S, Newell A, Akkari Y, Taniguchi T, Olson S, Grompe M (2005). Fancd2 functions in a double strand break repair pathway that is distinct from non-homologous end joining. *Hum Mol Genet*.

Jasin M (2002). Homologous repair of DNA damage and tumorigenesis: the BRCA connection. *Oncogene* **21**: 8981-93.

Jazaeri AA, Yee CJ, Sotiriou C, Brantley KR, Boyd J, Liu ET (2002). Gene expression profiles of BRCA1-linked, BRCA2-linked, and sporadic ovarian cancers. *J Natl Cancer Inst* **94**: 990-1000.

Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T *et al* (2008). Cancer statistics, 2008. *CA Cancer J Clin* **58**: 71-96.

Kennedy RD, Quinn JE, Mullan PB, Johnston PG, Harkin DP (2004). The role of BRCA1 in the cellular response to chemotherapy. *J Natl Cancer Inst* **96:** 1659-68.

Kim H, Huang J, Chen J (2007). CCDC98 is a BRCA1-BRCT domain-binding protein involved in the DNA damage response. *Nat Struct Mol Biol* **14:** 710-5.

King MC, Marks JH, Mandell JB (2003). Breast and ovarian cancer risks due to inherited mutations in BRCA1 and BRCA2. *Science* **302**: 643-6.

Kirova YM, Stoppa-Lyonnet D, Savignoni A, Sigal-Zafrani B, Fabre N, Fourquet A (2005). Risk of breast cancer recurrence and contralateral breast cancer in relation to BRCA1 and BRCA2 mutation status following breast-conserving surgery and radiotherapy. *Eur J Cancer*.

Kleiman FE, Wu-Baer F, Fonseca D, Kaneko S, Baer R, Manley JL (2005). BRCA1/BARD1 inhibition of mRNA 3' processing involves targeted degradation of RNA polymerase II. *Genes Dev* **19:** 1227-37.

Kurtz JM, Amalric R, Brandone H, Ayme Y, Jacquemier J, Pietra JC *et al* (1989). Local recurrence after breast-conserving surgery and radiotherapy. Frequency, time course, and prognosis. *Cancer* **63**: 1912-7.

Lee JS, Collins KM, Brown AL, Lee CH, Chung JH (2000). hCds1-mediated phosphorylation of BRCA1 regulates the DNA damage response. *Nature* **404**: 201-4.

Leong T, Whitty J, Keilar M, Mifsud S, Ramsay J, Birrell G *et al* (2000). Mutation analysis of BRCA1 and BRCA2 cancer predisposition genes in radiation hypersensitive cancer patients. *Int J Radiat Oncol Biol Phys* **48**: 959-65.

Lio YC, Schild D, Brenneman MA, Redpath JL, Chen DJ (2004). Human Rad51C deficiency destabilizes XRCC3, impairs recombination, and radiosensitizes S/G2-phase cells. *J Biol Chem* **279**: 42313-20.

Liu Z, Wu J, Yu X (2007). CCDC98 targets BRCA1 to DNA damage sites. *Nat Struct Mol Biol* **14:** 716-20.

Mamon HJ, Dahlberg W, Azzam EI, Nagasawa H, Muto MG, Little JB (2003). Differing effects of breast cancer 1, early onset (BRCA1) and ataxia-telangiectasia mutated (ATM) mutations on cellular responses to ionizing radiation. *Int J Radiat Biol* **79:** 817-29.

Margolin S, Johansson H, Rutqvist LE, Lindblom A, Fornander T (2006). Family history, and impact on clinical presentation and prognosis, in a population-based breast cancer cohort from the Stockholm County. *Fam Cancer* **5**: 309-21.

McCarthy EE, Celebi JT, Baer R, Ludwig T (2003). Loss of Bard1, the heterodimeric partner of the Brca1 tumor suppressor, results in early embryonic lethality and chromosomal instability. *Mol Cell Biol* **23**: 5056-63.

Miki Y, Swensen J, Shattuck-Eidens D, Futreal PA, Harshman K, Tavtigian S *et al* (1994). A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* **266**: 66-71.

Nakanishi K, Taniguchi T, Ranganathan V, New HV, Moreau LA, Stotsky M *et al* (2002). Interaction of FANCD2 and NBS1 in the DNA damage response. *Nat Cell Biol* **4:** 913-20.

Pierce LJ, Strawderman M, Narod SA, Oliviotto I, Eisen A, Dawson L *et al* (2000). Effect of radiotherapy after breast-conserving treatment in women with breast cancer and germline BRCA1/2 mutations. *J Clin Oncol* **18:** 3360-9. Powell SN (2005). The roles of BRCA1 and BRCA2 in the cellular response to ionizing radiation. *Radiat Res* **163:** 699-700.

Ratnam K, Low JA (2007). Current development of clinical inhibitors of poly(ADPribose) polymerase in oncology. *Clin Cancer Res* **13**: 1383-8.

Ries LAG, Melbert D, Krapcho M, Stinchcomb DG, Howlader N, Horner MJ *et al.* (2008). National Cancer Institute: Bethesda, MD.

Rio PG, Maurizis JC, Peffault de Latour M, Bignon YJ, Bernard-Gallon DJ (1999). Quantification of BRCA1 protein in sporadic breast carcinoma with or without loss of heterozygosity of the BRCA1 gene. *Int J Cancer* **80**: 823-6.

Ruffner H, Joazeiro CA, Hemmati D, Hunter T, Verma IM (2001). Cancer-predisposing mutations within the RING domain of BRCA1: loss of ubiquitin protein ligase activity and protection from radiation hypersensitivity. *Proc Natl Acad Sci U S A* **98**: 5134-9.

Russell JS, Brady K, Burgan WE, Cerra MA, Oswald KA, Camphausen K *et al* (2003). Gleevec-mediated inhibition of Rad51 expression and enhancement of tumor cell radiosensitivity. *Cancer Res* **63**: 7377-83. Sato K, Hayami R, Wu W, Nishikawa T, Nishikawa H, Okuda Y *et al* (2004). Nucleophosmin/B23 is a candidate substrate for the BRCA1-BARD1 ubiquitin ligase. *J Biol Chem* **279:** 30919-22.

Schreiber V, Ame JC, Dolle P, Schultz I, Rinaldi B, Fraulob V *et al* (2002). Poly(ADP-ribose) polymerase-2 (PARP-2) is required for efficient base excision DNA repair in association with PARP-1 and XRCC1. *J Biol Chem* **277**: 23028-36.

Scully R, Anderson SF, Chao DM, Wei W, Ye L, Young RA *et al* (1997). BRCA1 is a component of the RNA polymerase II holoenzyme. *Proc Natl Acad Sci U S A* **94:** 5605-10.

Shen SX, Weaver Z, Xu X, Li C, Weinstein M, Chen L *et al* (1998). A targeted disruption of the murine Brca1 gene causes gamma-irradiation hypersensitivity and genetic instability. *Oncogene* **17:** 3115-24.

Smith TE, Lee D, Turner BC, Carter D, Haffty BG (2000). True recurrence vs. new primary ipsilateral breast tumor relapse: an analysis of clinical and pathologic differences and their implications in natural history, prognoses, and therapeutic management. *Int J Radiat Oncol Biol Phys* **48**: 1281-9.

Sobhian B, Shao G, Lilli DR, Culhane AC, Moreau LA, Xia B *et al* (2007). RAP80 targets BRCA1 to specific ubiquitin structures at DNA damage sites. *Science* **316**: 1198-202.

Somasundaram K, Zhang H, Zeng YX, Houvras Y, Peng Y, Wu GS *et al* (1997). Arrest of the cell cycle by the tumour-suppressor BRCA1 requires the CDK-inhibitor p21WAF1/CiP1. *Nature* **389**: 187-90.

Starita LM, Horwitz AA, Keogh MC, Ishioka C, Parvin JD, Chiba N (2005).
BRCA1/BARD1 ubiquitinate phosphorylated RNA polymerase II. *J Biol Chem* 280: 24498-505.

Takemura H, Rao VA, Sordet O, Furuta T, Miao ZH, Meng L *et al* (2006). Defective Mre11-dependent activation of Chk2 by ataxia telangiectasia mutated in colorectal carcinoma cells in response to replication-dependent DNA double strand breaks. *J Biol Chem* **281**: 30814-23.

Tibbetts RS, Brumbaugh KM, Williams JM, Sarkaria JN, Cliby WA, Shieh SY *et al* (1999). A role for ATR in the DNA damage-induced phosphorylation of p53. *Genes Dev* **13:** 152-7.

Veronesi U, Cascinelli N, Mariani L, Greco M, Saccozzi R, Luini A *et al* (2002). Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med* **347**: 1227-32.

Walsh T, Casadei S, Coats KH, Swisher E, Stray SM, Higgins J *et al* (2006). Spectrum of mutations in BRCA1, BRCA2, CHEK2, and TP53 in families at high risk of breast cancer. *Jama* **295**: 1379-88.

Wang B, Matsuoka S, Ballif BA, Zhang D, Smogorzewska A, Gygi SP *et al* (2007). Abraxas and RAP80 form a BRCA1 protein complex required for the DNA damage response. *Science* **316**: 1194-8.

Wang Y, Cortez D, Yazdi P, Neff N, Elledge SJ, Qin J (2000). BASC, a super complex of BRCA1-associated proteins involved in the recognition and repair of aberrant DNA structures. *Genes Dev* **14**: 927-39.

Wu LC, Wang ZW, Tsan JT, Spillman MA, Phung A, Xu XL *et al* (1996). Identification of a RING protein that can interact in vivo with the BRCA1 gene product. *Nat Genet* **14**: 430-40.

Wu W, Nishikawa H, Hayami R, Sato K, Honda A, Aratani S *et al* (2007). BRCA1 ubiquitinates RPB8 in response to DNA damage. *Cancer Res* **67:** 951-8.

Xu X, Qiao W, Linke SP, Cao L, Li WM, Furth PA *et al* (2001). Genetic interactions between tumor suppressors Brca1 and p53 in apoptosis, cell cycle and tumorigenesis. *Nat Genet* **28**: 266-71.

Xu X, Weaver Z, Linke SP, Li C, Gotay J, Wang XW *et al* (1999). Centrosome amplification and a defective G2-M cell cycle checkpoint induce genetic instability in BRCA1 exon 11 isoform-deficient cells. *Mol Cell* **3**: 389-95.

Yan J, Yang XP, Kim YS, Jetten AM (2008). RAP80 responds to DNA damage induced by both ionizing radiation and UV irradiation and is phosphorylated at Ser 205. *Cancer Res* **68**: 4269-76.

Yang Q, Sakurai T, Mori I, Yoshimura G, Nakamura M, Nakamura Y *et al* (2001). Prognostic significance of BRCA1 expression in Japanese sporadic breast carcinomas. *Cancer* **92:** 54-60.

Yap TA, Boss DS, Fong PC, Roelvink M, Tutt A, Carmichael J *et al* (2007). First in human phase I pharmacokinetic (PK) and pharmacodynamic (PD) study of KU-0059436 (Ku), a small molecule inhibitor of poly ADP-ribose polymerase (PARP) in cancer patients (p), including BRCA1/2 mutation carriers. *J Clin Oncol* **25:** A3529.

Yarden RI, Brody LC (1999). BRCA1 interacts with components of the histone deacetylase complex. *Proc Natl Acad Sci U S A* **96:** 4983-8.

Yarden RI, Pardo-Reoyo S, Sgagias M, Cowan KH, Brody LC (2002). BRCA1 regulates the G2/M checkpoint by activating Chk1 kinase upon DNA damage. *Nat Genet* **30**: 285-9.

Zhong Q, Chen CF, Li S, Chen Y, Wang CC, Xiao J *et al* (1999). Association of BRCA1 with the hRad50-hMre11-p95 complex and the DNA damage response. *Science* **285**: 747-50.

Figure 1







Figure 3.



Affiliations

- Xiaowei Chen Department of Medical Oncology Fox Chase Cancer Center 333 Cottman Avenue Philadelphia, PA 19111-2409, USA Tel: +1 215 214 4286 Fax: +1 215 728 2741 Xiaowei.Chen@fccc.edu
- Jamie L. Kistler^a Department of Medical Oncology Fox Chase Cancer Center 333 Cottman Avenue Philadelphia, PA 19111-2409, USA
- Andrew K. Godwin Department of Medical Oncology Fox Chase Cancer Center 333 Cottman Avenue Philadelphia, PA 19111-2409, USA

^aCurrently employed with MedErgy Marketing, Yardley, PA.