

REPORT DOCUMENTATION PAGE

*Form Approved
OMB No. 0704-0188*

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 11/13/2009	2. REPORT TYPE Final Performance Technical Report	3. DATES COVERED (From - To) 21 February 2005 – 30 September 2009
--	---	---

4. TITLE AND SUBTITLE Diapycnal Mixing in a Coastal Regime	5a. CONTRACT NUMBER
	5b. GRANT NUMBER N00014-05-1-0362
	5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S) Michael C. Gregg	5d. PROJECT NUMBER
	5e. TASK NUMBER
	5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Applied Physics Laboratory, University of Washington 1013 N.E. 40th Street Seattle, WA 98105	8. PERFORMING ORGANIZATION REPORT NUMBER
---	---

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Scott Harper, ONR 322 Office of Naval Research 875 North Randolph Street Arlington, VA 22203-1995	10. SPONSOR/MONITOR'S ACRONYM(S)
	11. SPONSOR/MONITOR'S REPORT NUMBER(S)

12. DISTRIBUTION/AVAILABILITY STATEMENT
UU

13. SUPPLEMENTARY NOTES

14. ABSTRACT
As part of ONR's AESOP DRI, during August 2006 M. Gregg's research team surveyed turbulent mixing across Monterey Bay, sequentially profiling along a set of tracks and repeating each sequence for a half-daily tidal cycle, 12.4 hours. In addition, a 300 kHz Workhorse ADCP was installed to monitor currents at one place. The observations revealed an unexpectedly rich array of processes within the bay and on the adjacent upper shelf.

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 1	19a. NAME OF RESPONSIBLE PERSON Michael C. Gregg
a. REPORT UU	b. ABSTRACT UU	c. THIS PAGE UU			19b. TELEPHONE NUMBER (Include area code) 206-543-1353

Reset

Final Technical Report

Diapycnal Mixing in a Coastal Regime

Michael C. Gregg, P.I.
Applied Physics Laboratory, University of Washington
1013 NE 40th St.
Seattle, WA 98105-6698
Phone: 206-543-1353 Fax: 206-543-6785
Email: gregg@apl.washington.edu

Performance Period: 21 February 2005 – 30 September 2009

Grant # N00014-05-1-0362

As part of ONR's AESOP DRI, during August 2006 M. Gregg's research team surveyed turbulent mixing across Monterey Bay, sequentially profiling along a set of tracks and repeating each sequence for a half-daily tidal cycle, 12.4 hours. In addition, a 300 kHz Workhorse ADCP was installed to monitor currents at one place. The observations revealed an unexpectedly rich array of processes within the bay and on the adjacent upper shelf.

Six tracks were run with Advanced Microstructure Profilers (AMP), all across the shelf break because the AMP can profile to 1100 m (Fig. 1). Nineteen tracks were run with Modular Microstructure Profilers (MMPs) that carry a pumped Sea-Bird CTD. In addition, a 300 kHz Workhorse ADCP was set on the bottom while we were in the bay.

The 300 kHz Workhorse revealed a steady mean flow into the bay superimposed on barotropic tides with maximum amplitudes of ± 0.07 m s⁻¹. Analysis shows a strongly modal structure to the baroclinic tidal currents, with the dominant modes changing during the two-weekly tidal cycle. At times, shear in the modes dominated mixing in MMP runs over the Workhorse.

The first group of MMP profiles in the bay discovered vertical columns of intense acoustic backscatter. These were visible in the ship's 12 kHz echo sounder as well as our 120 kHz and 208 kHz BioSonics acoustics backscatter systems. From the sixth run along the track an MMP sampled one of the features, which we subsequently learned were large aggregations of fish, probably anchovies. As seen from the dissipation profiles, ϵ within this fish aggregation were several decades larger than in adjacent profiles outside the aggregation. These were the largest dissipation rates measured in the bay. Gregg worked with John Horne, UW School of Fisheries,

to begin an assessment of the biological and physical nature of the aggregations and the mixing they produce. A manuscript was submitted to the Journal of Physical Oceanography in Spring 2009, which has now been published. Gregg and Horne (2009) provided the first observations of turbulence within aggregations of fish, demonstrating that, in at least some cases, mixing efficiency is much lower than assumed by estimates that biomixing may be a major mixing process.

In addition to the large effect of the tides on mixing, we also found solitons, both as trains and individual upward pulses. We plan a separate paper about them as well as about the nature of tides in the bay, as well as one summarizing mixing across the bay and comparing it with that on other shelves, looking for common themes and parameterizations.

PUBLICATION

Gregg, M.C. and J.K. Horne, Turbulence, acoustic backscatter and pelagic nekton in Monterey Bay, *J. Phys. Oceanogr.*, 39, 1097-1114, 2009.