

**A REDESCRIPTION OF THE HOLOTYPE MALE OF AEDES
(STEGOMYIA) TONGAE EDWARDS WITH A NOTE
ON TWO TOPOTYPIC FEMALES
(DIPTERA: CULICIDAE)**

YIAU-MIN HUANG

Reprinted from
PROCEEDINGS OF THE ENTOMOLOGICAL SOCIETY OF WASHINGTON
Vol. 74, No. 3, September 1972
pp. 338-342
Made in the United States of America

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE SEP 1972	2. REPORT TYPE	3. DATES COVERED 00-00-1972 to 00-00-1972			
4. TITLE AND SUBTITLE A Redescription of the Holotype Male of Aedes (Stegomyia) Tongae Edwards with a Note on Two Topotypic Females		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Smithsonian Institution, Medical Entomology Project, Washington, DC, 20560		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	6	

**A REDESCRIPTION OF THE HOLOTYPE MALE OF AEDES
(STEGOMYIA) TONGAE EDWARDS WITH A NOTE
ON TWO TOPOTYPIC FEMALES**

(DIPTERA: CULICIDAE)^{1,2}

YIAU-MIN HUANG, *Southeast Asia Mosquito Project, Department of
Entomology, Smithsonian Institution, Washington, D.C. 20560*

ABSTRACT—The holotype male of *Aedes (Stegomyia) tongae* Edwards from Ha'apai, Tonga, South Pacific is redescribed and illustrated. Two topotypic females of this species of Buxton & Hopkins' expedition to Ha'apai, Tonga, South Pacific, 1925 are also described here.

The name *Aedes tongae* Edwards 1926 has caused much confusion. At least two species were being mistaken for *tongae* Edwards as reported by Belkin (1962:476) and Ramalingam and Belkin (1965:2).

According to Ramalingam and Belkin (1965:2-3), the description and figures by Belkin (1962:475-476, 349-350) for "*tongae*" are actually those of another species which they named *tabu*. Thus, no detailed description and figures of true *tongae* are available at the present.

Through the kindness of Dr. P. F. Mattingly, I have had the opportunity to examine the holotype male of *tongae* Edwards and the two females which bear the same data as the holotype. I take advantage of this opportunity to give a full description of this holotype male and the two topotypic females so that the identity of Edwards' *tongae* should no longer remain in doubt.


Species of the *scutellaris* group in the Tonga area are being investigated further under a SEAMP-WHO project and the full results will be published later. Due to the highly variable nature of the group in both adult and immature characteristics, a large number of progeny rearings from many different localities are required to provide sound criteria for establishing the true identity of the species.

¹ This work was supported by Research Contract No. DA-49-193-MD-2672 from the U.S. Army Medical Research and Development Command, Office of the Surgeon General.

² Immediate publication secured by full payment of page charges—Editor.

→

Fig. 1. *Aedes (Stegomyia) tongae* Edwards, holotype ♂: A, dorsal aspect; B, lateral aspect of abdomen; C, lateral aspect of thorax; D, anterior surface of legs.



Holotype male
Aedes (Stegomyia) tongae Edwards

Richard M. Moulton

***Aedes (Stegomyia) tongae* Edwards**

(Figs. 1, 2)

Aedes (Stegomyia) variegatus var. *tongae* Edwards, 1926, Bull. Ent. Res. 17:103 (♂ *, ♀). Type locality: Ha'apai, Tonga Islands.

Type male, in fair condition (Pres. by Imp. Bur. Ent. 1926. 62) with associated terminalia on a slide, Ha'apai, Tonga, South Pacific, 26-II-1925 (P. A. Buxton and G. H. Hopkins collectors). Deposited in the British Museum (Natural History), London.

Head.—Proboscis dark scaled, with a few pale scales on ventral side, slightly longer than fore femur; palpus dark, with a white basal band on segments 2, 3; rest of palpus missing; antenna plumose, shorter than proboscis; clypeus bare; torus covered with white scales except on dorsal side; decumbent scales of vertex all broad and flat; erect forked scales dark, not numerous, restricted to occiput; vertex with a median stripe of broad white scales, with broad dark ones on each side interrupted by a lateral stripe of broad white scales followed by a patch of white broad ones ventrally. *Thorax*. Scutum with narrow dark scales and a prominent median longitudinal stripe of similar white ones, median stripe narrows slightly posteriorly and forks at beginning of prescutellar space, prescutellar line with yellowish scales; posterior dorsocentral yellowish lines do not reach to middle of scutum; supraalar line of broad white scales present; acrostichal bristles absent; dorsocentral bristles present; scutellum with broad white scales on all lobes and with a few broad dark ones at apex of mid lobe; anterior pronotum with broad white scales; posterior pronotum with narrow dark scales on upper portion and with broad white scales on lower portion forming a white stripe instead of a white patch; paratergite with broad white scales; postspiracular area without scales; subspiracular area without scales; patches of broad white scales on propleuron, on upper and lower sternopleuron and on upper and lower mesepimeron; lower mesepimeral scale patch of medium size and narrowly connected to upper mesepimeral scale patch; lower mesepimeron without bristles; metameron bare. *Wing*. With dark scales on all veins except for a minute basal spot of white scales on costa; first forked cell 1.5 times as long as its stem. *Halter*. With dark scales. *Leg*. Coxae with patches of white scales; knee-spots present on all femora; fore and mid femora dark anteriorly, paler posteriorly; hind femur anteriorly with a broad white longitudinal stripe which widens towards base and is separated from apical white knee-spot; fore and mid tibiae dark anteriorly, paler posteriorly; hind tibia dark; fore tarsus with basal white bands on tarsomeres 1, 2; mid tarsus missing; hind tarsus with basal white bands on tarsomeres 1-4, ratio of length of white band to total length of tarsomere is $\frac{1}{3}$, $\frac{1}{3}$, $\frac{2}{5}$ and $\frac{1}{2}$; tarsomere 5 all white; fore leg with tarsal claws unequal, larger one toothed, smaller one simple; hind leg with tarsal claws equal, simple. *Abdomen*. Abdominal segment I with white scales on laterotergite; tergum II with a small basal median spot and with lateral white spots; terga III-V each with a complete sub-basal pale yellowish band connected to lateral white spots. *Terminalia*. Basimere 3.5 times as long as wide; its scales restricted to dorsolateral, lateral and ventral areas; with a patch of hairs on the basomesal area of dorsal surface; mesal surface membranous; claspette simple, slender, sternal and tergal sides parallel, rounded apically, with 6 modified setae in a row on apical $\frac{1}{3}$ of sternal side; lateral surface with hairs extending basad




Fig.2

Aedes (Stegomyia) tongae Edwards

Fig. 2. *Aedes (Stegomyia) tongae* Edwards, holotype ♂, tergal aspect of terminalia with claspette enlarged.

to about level of modified setae; apex tergally with hairs about $\frac{1}{2}$ as long as entire lobe length; distimere simple, elongate, as long as basimere, slightly swollen near tip; with a spiniform process and a few hairs near apex; aedeagus with a distinct sclerotized lateral toothed plate on each side; paraprocts without teeth; cercal setae absent; ninth tergum with middle rounded and with a hairy lobe on each side.

FEMALE. Based on 2 topotypic females, with same data as type male. Deposited in British Museum. Essentially as in male, differing in the following respects: palpus $\frac{1}{5}$ of proboscis, with white scales on apical half. Wing with first forked cell about 2 times as long as its stem. Mid tarsus which is absent in the male, with basal white bands on tarsomeres 1, 2; fore and mid legs with tarsal claws equal, simple. Abdominal tergum II dark dorsally with lateral white spots only; terga III-VI each with a complete or incomplete sub-basal pale yellowish band and with lateral white spots which are turned dorsomesally and connected to sub-basal pale yellowish bands; tergum VII with lateral white spots only or with a small basal median spot as well; segment VIII completely retracted.

ACKNOWLEDGMENTS

I am grateful to Dr. Botha de Meillon for the helpful assistance in connection with this paper and for critical review of the manuscript. I extend my thanks to Mr. Vichai Malikul of the Southeast Asia Mosquito Project for his help in making the drawings. I also wish to express my gratitude to Dr. P. F. Mattingly, Department of Entomology, British Museum (Natural History), for the loan of the specimens described above.

REFERENCES

- BELKIN, J. N. 1962. The mosquitoes of the South Pacific. Univ. Calif. Press, Berkeley, 2 vols., 608 and 412 pp.
- EDWARDS, F. W. 1926. Mosquito notes.-VI. Bull. Ent. Res. 17:101-131.
- RAMALINGAM, S. and J. N. BELKIN. 1965. Mosquito studies (Diptera, Culicidae). III. Two new *Aedes* from Tonga and Samoa. Contr. Amer. Ent. Inst. 1(4):1-10.