

# Growth and Phenomenology of Phytoplankton Thins Layers in the Gulf of Maine

Mary Jane Perry  
School of Marine Sciences and  
Ira C. Darling Marine Center  
University of Maine  
193 Clark's Cove Road  
Walpole ME 04573 - 3307  
phone: (207) 563-3146 ext. 245 fax: (207) 563-3119 email: [perrymj@maine.edu](mailto:perrymj@maine.edu)

David W. Townsend  
School of Marine Sciences  
5706 Aubert Hall, Rm 341  
University of Maine  
Orono, ME 04469  
phone: (207) 581-4367 fax: (207) 581- 4388 e-mail: [davidt@maine.edu](mailto:davidt@maine.edu)

Award Number: N000140410633  
<http://optics.dmc.maine.edu/slocum/nemo/>

## LONG-TERM GOALS

Our overarching long-term goal is to understand what controls phytoplankton distribution, optical properties and production in the coastal ocean. Our project-specific goals are to understand the mechanisms responsible for the creation, maintenance and demise of subsurface phytoplankton layers in the Gulf of Maine; to decrease uncertainty in interpretation of chlorophyll *a* fluorescence as an estimator of phytoplankton biomass by better understanding daytime fluorescence quenching; and to determine the vertical distribution of toxigenic species of *Alexandrium* in the Gulf of Maine.

## OBJECTIVES

The specific objectives of the past year's work were: 1) to continue to analyze data from two cruises in 2005 and 2006 in the Gulf of Maine to better understand how the subsurface distributions of phytoplankton and suspended particles are controlled by light, nitrate and density structure over a broad range of hydrographic conditions; 2) to improve analysis of nitrate concentrations derived from *in situ* UV absorption spectra with ISUS; 3) to continue to analyze data from optically-instrumented gliders deployed in Gulf of Maine; 4) to confirm diel and vertical patterns in phytoplankton UV absorption coefficients; 5) to analyze patterns in photoinhibition of variable fluorescence and *in situ* chlorophyll *a* fluorescence; and 6) to continue to analyze distributions of *Alexandrium* species in the Gulf of Maine.

## APPROACH

In the 2005 field program we successfully located phytoplankton thin layers in the Gulf of Maine, but we were not able to stay within a single layer. In the 2006 field year we broadened out approach to a more general study of the vertical structure of subsurface phytoplankton layers in the Gulf of Maine,

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>2006</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Growth and Phenomenology of Phytoplankton Thins Layers in the Gulf of Maine</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>School of Marine Sciences and Ira C. Darling Marine Center University of Maine 193 Clark's Cove Road Walpole ME 04573 - 3307</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>7</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			


and carried out a survey program in order to encounter a wide range of optical and hydrographic conditions within a relatively restricted geographic area. On the order of 150 profiles were taken of temperature, salinity, chlorophyll *a* and CDOM fluorescence, beam *c*, optical backscattering, dissolved oxygen, and PAR; approximately 100 profiles included continuous nitrate measurements (Satlantic ISUS). At a number of depths and stations, discrete water samples were collected for chlorophyll, phytoplankton absorption, variable fluorescence, nutrients and samples of *Alexandrium spp.* distributions; in addition *Alexandrium* samples were collected with a vertical pumping system. At the time of the 2006 cruise we also deployed a Slocum glider instrumented with a SBE CTD, Aanderaa optode, and two WET Labs Eco pucks (chlorophyll *a* and CDOM fluorescence; 3 wavelengths of backscattering).

## WORK COMPLETED

In the past year we have focused on analyzing data from the 2006 cruise in the Gulf of Maine, completing fluorescence quenching experiments, confirming diel and vertical patterns in phytoplankton UV absorption coefficients, testing a Slocum glider with Satlantic radiometers, and completing analysis of distributions of *Alexandrium* species in the Gulf of Maine. In 2007 additional fieldwork was carried out from a small boat in Wilkinson Basin in the Gulf of Maine and in the Damariscotta River Estuary to continue studies of phytoplankton absorption, variable fluorescence, and fluorescence quenching. A brief glider mission was carried out in 2007 with a Slocum glider instrumented with Satlantic radiometers.


## RESULTS

One major focus has been analysis of data from the 2006 cruise in the Gulf of Maine. Although we found relatively few “thin” layers, most of the summer water masses were highly stratified and almost all had well-defined subsurface phytoplankton layers. The only exception was a very well-mixed water mass in the Eastern Maine Coastal Current north of the Penobscot River discharge. We used the method of Twardowki et al. (2001) to characterize particle scattering as organic or mineral; high ratios of optical backscattering to beam *c* are indicative of minerals, while low ratios are characteristic of organic particles. In Figure 1 we use relative counts to display the bimodal distribution of the backscattering ratio. Most high ratios were found at depth, as expected for waters dominated by




**Figure 1. Relative ratios of optical backscattering-to-beam *c* in the Gulf of Maine.**

resuspended sediments, and most low ratios were found in near-surface waters dominated by organic particles, with the exception of the Eastern Maine Coastal Current (Fig. 2). Ratios of fluorescence-to-beam  $c$  could then be used in near-surface waters dominated by organic particulate to assess phytoplankton photoadapted state – fluorescence quenched, high-light or low-light adapted. In 60% of all well-defined peaks, the beam  $c$  maxima was within  $\pm 0.5$  m of the fluorescence maxima. Subsurface oxygen maxima were typically observed close to the beam  $c$  maxima, indicating that phytoplankton in the subsurface layers were actively growing.




**Figure 2. Distribution of relative ratios of optical backscattering-to-beam  $c$  in the Gulf of Maine.**

During the 2006 cruise, we successfully collected a number of continuous profiles of nitrate with the ISUS and analyzed over 1,000 wet chemistry nitrate samples for its calibration. This year, in collaboration with Ken Johnson of MBARI, we were able to tune ISUS calibrations (Fig. 3), thereby improving the accuracy of the nitrate profiles. We are using profile data to analyze vertical gradients in phytoplankton and nitrate to understand the role of nitrate flux in controlling the magnitude and maintenance of the subsurface layers.




**Figure 3. Distribution of relative ratios of optical backscattering-to-beam  $c$  in the Gulf of Maine.**

Data from previous years were analyzed and new experiments were completed to study chlorophyll *a* fluorescence quenching and diel variability in absorption by MMA-like pigments. Both daytime fluorescence quenching with in situ fluorometers and variable fluorescence (Fv/Fm) with Satlantic FIRE show similar patterns in response to ambient light, i.e., decreases with increases in light (Fig. 4).




**Figure 4. Variable fluorescence (Fv/Fm) as a function of irradiance.**

These results suggest that variable fluorescence measurements on autonomous platforms would be a useful tool for determining if chlorophyll is quenched and therefore help reduce biases in estimates of phytoplankton biomass from fluorescence. The time course of recovery of fluorescence and variable fluorescence shows that a small amount of light is essential for photosystem recovery (Fig. 5).




**Figure 5. Light recovery of variable fluorescence (Fv/Fm); lack of recovery in complete darkness.**


Last year strong diel patterns in UV-absorbing pigments were observed during a diel study in Wilkinson Basin in the Gulf of Maine. Absorption coefficients at 320 nm increased in the morning and then decreased throughout the afternoon, with the magnitude of absorption decreasing with depth (Figs. 6, 7). The diel plasticity of the UV absorption was unexpected and not previously reported in field studies. Observations this summer confirmed the phenomenon (Fig. 8). A Slocum glider instrumented with Satlantic radiometers was tested in the Gulf of Maine (Fig. 9). Analyzes of *Alexandrium* distributions show higher concentrations downstream of the Eastern Maine Coastal Current (Fig. 10).





*Figure 6. Absorption spectra of phytoplankton from 0 m, Wilkinson Basin, Gulf of Maine.*




*Figure 7. Absorption spectra of phytoplankton from 25 m, Wilkinson Basin, Gulf of Maine.*



**Figure 8. Ratio of absorption 330 to 676 nm five depths, Wilkinson Basin, Gulf of Maine; solid circles from 2006 and stars from 2007.**



**Figure 9. Irradiance spectra ( $E_d$ ) from Slocum glider deployed in Gulf of Maine.**



*Figure 10. Distribution of Alexandrium spp. in June 2006.*

## IMPACT/APPLICATIONS

A better understanding of the interaction of hydrography, nutrients, and light in controlling the subsurface distribution of phytoplankton is important to the Navy in its goal of predicting and understanding how biota affect the optical properties of operational importance to the Navy.

## RELATED PROJECTS

The underwater glider was acquired under award N000140510412 to Perry, Townsend and colleagues, entitled "Acquisition of Underwater Gliders for Autonomous Sampling in the Gulf of Maine".

## REFERENCES

Twardowski, M.S., E. Boss, J.B. Macdonald, W.S. Pegau, A.H. Barnard, and J.R.V. Zaneveld. 2001. A model for estimating bulk refractive index from the optical backscattering ratio and the implications for understanding particle composition in Case I and Case II waters. *J. Geophys. Res.*, 106(C7), 14,129-14,142.