



NSF/ONR Workshop on Data Assimilation in Ocean Research

LOOPS/Poseidon: A Distributed System for Real-Time Interdisciplinary Ocean Forecasting with Adaptive Modeling and Sampling

P.F.J. Lermusiaux, C. Evangelinos, P.J. Haley Jr, W.G. Leslie, N.M. Patrikalakis, A.R. Robinson, R. Tian

PIs: N.M. Patrikalakis, J.J. McCarthy, A.R. Robinson, H. Schmidt Scientists: C. Evangelinos, P.J. Haley Jr., P.F.J. Lermusiaux, R. Tian http://czms.mit.edu/poseidon





Report Documentation Page					Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.						
1. REPORT DATE SEP 2003		2. REPORT TYPE		3. DATES COVE 00-00-2003	ered 3 to 00-00-2003	
4. TITLE AND SUBTITLE					5a. CONTRACT NUMBER	
LOOPS/Poseidon: A Distributed System for Real-Time Interdisciplinary					5b. GRANT NUMBER	
Ocean Forecasting with Adaptive Modeling and Sampling					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER		
					5e. TASK NUMBER	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Massachusetts Institute of Technology,Laboratory for Information and Decision System,Cambridge,MA,02139					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited						
13. SUPPLEMENTARY NOTES NSF/ONR Workshop on Data Assimilation in Ocean Research, 9-11 Sep 2003, Williamsburg, VA						
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFIC		17. LIMITATION OF	18. NUMBER	19a. NAME OF		
a. REPORT unclassified	ь. ABSTRACT unclassified	c. THIS PAGE unclassified	ABSTRACT Same as Report (SAR)	OF PAGES 32	RESPONSIBLE PERSON	

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std Z39-18

Ocean Science and Data Assimilation



- SV: STATE VARIABLE
- P: PARAMETER
- OBSERVED
 MEASURED
- U: UNOBSERVED OR UNMEASURED
- E: ERROR
- ###: DYNAMICAL LINKAGES

- Field and remote observations
- Models:
 - Dynamical
 - Measurement
 - Error
- Assimilation schemes
- Sampling strategies
- State and parameter estimates
- Uncertainty estimates
- A Dynamic Data-Driven Application System (DDDAS)



LOOPS/Poseidon Adaptive Interdisciplinary Ocean Forecasting in a Distributed Computing Environment



- Research coupling Physical and Biological Oceanography with Ocean Acoustics.
- More effective Real-Time Ocean Forecasting for Naval and Maritime Operations, Pollution Control, Fisheries Management, Scientific Data Acquisition, etc.
- MIT OE (IT, Acoustics) and Harvard DEAS (Ocean Physics-Biology-Acoustics).

Key points

- Web interface
- Remote visualization
- Metadata for code and data
- Metadata/Ontology editors
- Legacy application support
- Grid computing infrastructure
- Transparent data access
- Data assimilation (ESSE, OI)
- Interdisciplinary interactions
- Adaptive modeling
- Adaptive sampling
- Feature Extraction
- Prototype for community-use



Physical-Biological-Acoustical Oceanography with HOPS



- Primitive Equation (PE) physical dynamics model
- Multiple biological models
- Interfaces to acoustical models
- Adaptable to different domains
- Nested-domains parallelism
- Software: F77-matlab-C
- I/O: NetCDF, stdin

Error Subspace Statistical Estimation (ESSE)



- Uncertainty forecasts (with dynamic error subspace, error learning)
- Ensemble-based (with nonlinear and stochastic model)
- Multivariate, non-homogeneous and non-isotropic DA
- Consistent DA and adaptive sampling schemes
- Software: not tied to any model, but specifics currently tailored to HOPS

IT Design Motivations

- <u>Real-time</u> predictions of **interdisciplinary** ocean fields and *uncertainties*
 - Data Assimilation (DA) using ESSE is currently ensemble-based and thus ideal for high throughput distributed computing
 - Interdisciplinary interactions and multiscale/nested simulations ideal for parallel computing
- Develop autonomous adaptive models for physics & biology
 - Adaptive parameter values, model structures and state variables
 - Error metrics and criteria for adaptation
- Towards automated, distributed management of observed and modeled data
 - Consistent use of metadata helps provide transparent data management, including quality control
 - Forecasting workflow is being automated, including DA
- Web access from lightweight clients eases operational use and system control
- Interactive visualizations for better understanding and decision-making

Software Strategies

- Exploit parallelism (especially throughput) opportunities
- Maximize performance, facilitate users, but limited changes
 - -For new generalized adaptive biological model: MPI coding
 - -For existing software: automate file I/O based workflows
 - -Work to the maximum extent possible at the binary level
 - -Metadata for software use (and installation) in XML
- Use Grid technologies
 - -For user: compute and data access solutions
 - -Drive forecasting, visualization workflows on the Grid
 - -Present results to user's web browser

GRID COMPUTING - MIDDLEWARE



Interdisciplinary Data Assimilation (DA)

- Is in its infancy, but can contribute significantly to understanding physical-acoustical-biogeochemical processes, including quantitative development of fundamental models
- Required for interdisciplinary ocean field prediction and parameter estimation
- Model-model, data-data and data-model compatibilities are essential
- Care must be exercised in understanding, modeling and controlling errors and in performing sensitivity analyses to establish robustness of results
- Dedicated interdisciplinary research needed

Coupled Physical-Acoustical Filtering via ESSE

Coupled assimilation of sound-speed and TL data for a joint estimate of soundspeed and TL fields

•"Truth" ocean

"truth" provides

natural data

75m depth



Coupled Physical-Biogeochemical Smoothing via ESSE



Coupled Physical-Biogeochemical DA via ESSE (continued)



e) Difference betweenESSE smoothingestimate on Aug. 25 andnowcast on Aug. 25

f) Forecast for Sep. 2,starting from ESSEsmoothing estimate onAug. 25

(g): as d), but for Chl-a at 20 m depth

(h): RMS differences between Chl-a data on Sep. 2 and the field estimates at these datapoints as a function of depth (specifically, "RMS-error" for persistence, dynamical forecast and ESSE filtering estimate)





Interdisciplinary Adaptive Sampling

- Use forecasts and their uncertainties to alter the observational system in space (locations/paths) and time (frequencies) for physics, biology and acoustics.
- Locate regions of interest, based on:
 - Uncertainty values (error variance, higher moments, pdf's)
 - Interesting physical/biological/acoustical phenomena (feature extraction, Multi-Scale Energy and Vorticiy analysis)
 - Maintain synoptic accuracy
- Plan observations under operational, time and cost constraints to maximize information content (e.g. minimize uncertainty at final time or over the observation period).

Integrated Ocean Observing and Prediction Systems

Platforms, sensors and integrative models

HOPS/ESSE- AOSN-II Accomplishments

- 23 sets of real-time nowcasts and forecasts of temperature, salinity and velocity released from 4 August to 3 September
- 10 sets of real-time ESSE forecasts issued over same period: total of 4323 ensemble members (stochastic model, BCs and forcings)
- Adaptive sampling recommendations suggested on a routine basis
- Web: http://www.deas.harvard.edu/~leslie/AOSNII/index.html for daily distribution of forecasts, scientific analyses, data analyses, special products and control-room presentations
- Assimilated ship (Pt. Sur, Martin, Pt. Lobos), glider (WHOI and Scripps) and aircraft SST data, within 24 hours of appearance on data server (after quality control)
- Forecasts forced by 3km and hourly COAMPS flux predictions

Real-time Adaptive Sampling – Pt. Lobos

- Large uncertainty forecast on 26 Aug. related to predicted meander of the coastal current which advected warm and fresh waters towards Monterey Bay Peninsula.
- Position and strength of meander were very uncertain (e.g. T and S error St. Dev., based on 450 2-day fcsts).
- Different ensemble members showed that the meander could be very weak (almost not present) or further north than in the central forecast
- Sampling plan designed to investigate position and strength of meander and region of high forecast uncertainty.





Surf. Temperature Fct 872 24 18.6 18.3 17.9 37°13 17.5 17.1 87° N 16.8 16.4 38° 48 16.0 15.6 15.3 88*36 14.9 14.5 36° 24 14.1 13.8 26712 13.4 13.0 12.6 -98° M 12.3 11.9 369 48 Min= 1.10486+01) 1.530602+0 6.00 Day Forecast : 28 Aug 2003

Salinity Error Fct



ESSE field and error modes forecast for August 28 (all at 10m)









Error Covariance Forecast for 28 August



Real-time Adaptive Coupled Models



- Different Types of Adaptive Couplings:
 - •Adaptive physical model drives multiple biological models (biology hypothesis testing)
 - •Adaptive physical model and adaptive biological model proceed in parallel, with some independent adaptation
- •Implementation
 - •For performance and scientific reasons, both modes are being implemented using message passing for parallel execution
 - •Mixed language programming (using C function pointers and wrappers for functional choices)

Generalized Adaptable Biological Model



A Priori Biological Model



Example: Use P data to select parameterisations of Z grazing

Table 1. Parameterization of grazing on multiple types of prey with passive selection (g_{max} : maximum grazing rate; K: Half-saturation constant (but saturation constant in Eq. 1); P_0 threshold below which grazing is zero; p_i : preference coefficient; ?, a, ? : constant).

Function	References		
(1) Rectilinear $g_{i} = \begin{cases} g_{\max} \frac{p_{i}P_{i}}{K}, & \text{for } R \leq K \\ g_{\max}, & \text{for } R > K \end{cases}, R = \sum_{i=1}^{n} p_{i}P_{i}$	Armstrong, 1994		
(2) Ivlev function for each prey type: $g_i = g_{\max}(1 - e^{-\alpha_i P_i})$	Leonard et al., 1999		
(3) Ivlev function with interference between prey types: $g_i = g_{\max} \left(1 - e^{-aR}\right) \frac{p_i P_i}{R}$, with $R = \sum_{i=1}^n p_i P_i$	Hofmann and Ambler, 1988		
(4) Mechanistic disc function: $g_i = g_{\max} \frac{a_i N_i}{1 + \sum_{j=1}^n a_j \tau_j N_j}$	Murdoch and Oaten, 1975; Holt, 1983; Gismervik and Anderson, 1997; Strom and Loukos, 1998		
(5) Michaelis Menten Function: $g_i = g_{\max} \frac{p_i P_i}{K + \sum_{j=1}^{n} p_j P_j}$	Murdoch, 1973; Real, 1977; Moloney and Field, 1991; Verity, 1991; Gismervik and Anderson, 1997; Strom and Loukos, 1998		
(6) Threshold MM function: $g_{i} = g_{\max} \left(\frac{R - P_{0}}{K + R - P_{0}} \right) \frac{p_{i}P_{i}}{R}, \text{ with } R = \sum_{i=1}^{n} p_{i}P_{i}$	Evans, 1988; Lancelot et al., 2000		
(7) Modified MM function: $g_i = g_{\max} \frac{p_i P_i}{1 + \sum_{i=1}^{n} p_i P_i}$	Verity, 1991; Fasham et al. (1999) and Tian et al. (2001)		

Table 2. Parameterization of grazing on multiple types of prey with active switching selection (g_{max} : maximum grazing rate; K: Half-saturation constant; P_0 threshold below which grazing is zero; p_i : preference coefficient; α , a, τ : constant).

Function	References		
(1) Switching MM predation: $g_{i} = g_{\max} \frac{p_{i}P_{i}^{2}}{K\sum_{j=1}^{n} p_{j}P_{j} + \sum_{j=1}^{n} p_{j}P_{j}^{2}}$	Fasham et al., 1990; Strom and Loukos, 1998; Pitchford and Brindley, 1999; Spitz et al., 2001		
(2) Mechanistic disc switching predation: $g_{i} = g_{\max} \frac{b_{i}N_{i}^{2}}{(1 + c_{i}N_{i})(1 + \sum_{j=1}^{n} \frac{b_{j}h_{j}N_{j}^{2}}{1 + c_{j}N_{j}^{2}})}$	Chesson, 1983		
(3) Generalized switching function: $g_{i} = g_{\max} a_{i} \frac{(p_{i}P_{i})^{m}}{\sum_{i=1}^{n} (p_{i}P_{i})^{m}}$	Tansky, 1978; Teramoto, 1979; Matsuda et al., 1986		
(4) Generalized switching function: $g_{i} = g_{\max} \frac{(p_{i}P_{i})^{m}}{\left(\sum_{i=1}^{n} (p_{i}P_{i})\right)^{m}}$	Vance, 1978		
(5) Generalized switching MM function: $g_{i} = g_{\max} \frac{(p_{i}P_{i})^{m}}{1 + \sum_{i=1}^{n} (p_{i}P_{i})^{m}}$	Gismervik and Andersen (1997)		
(6) Generalized switching MM function: $g_{i} = g_{\max} \frac{\left(p_{i}(P_{i} - P_{0i})\right)^{m}}{1 + \sum_{i=1}^{n} \left(p_{i}(P_{i} - P_{0i})\right)^{m}}$	This work		

Distributed/Grid Computing, Forecasting and Data assimilation with Legacy codes

- Distributed technologies (Sun Grid Engine) with web portal front-end ready to be tested with ESSE and HOPS
- Partial parallelism within ESSE easy because open-source routines (Sun Lapack) were used from the start
- HOPS, ESSE and acoustics codes: Fortran-matlab legacies
 - Relatively complex codes and makefile options
 - Hundreds of build and runtime parameters
- For other (future) codes, source code might not be available
- Classic encapsulation techniques that compartmentalize the code into subroutines, called from wrappers require constant reworking
- Thus: we chose to encapsulate at the binary level, with generic approach, so as to handle new codes with limited/no rewriting

Metadata for handling legacy software

- Hierarchical structure for describing code (can also handle binary-only case)
- Basic assumptions about codes thus encapsulated:
 - No independent GUI, all runtime control from the command line and input/stdin files
 - All build-time parameterization done by altering the makefile and selecting values (parameters) in include-files
- Datatypes and relevant ranges for each parameter checked to ensure validity



XML Encapsulation for Legacy Binaries

- Descriptions of I/O files, runtime parameters, stdin and command line arguments, makefile parameters, requirements and conflicts for options, invocation mechanisms are needed:
 - Essentially a computer readable install and user guide
 - XML description provides software <u>use</u> and <u>build</u> metadata
 - Design of appropriate hierarchical XML Schemas (evolutionary)
 - Simulation datafile metadata are also usable (e.g. NcML for NetCDF)
 - Provides the constraints for generation of workflows (file I/O based)
- Binaries can be built on demand from generated makefiles
- Developers need to keep XML description up-to-date with their code (incremental effort) without switching to more elaborate approaches
- Concept is generally applicable, directly useful with other ocean models

Java-Based GUI for Legacy Binaries

- Prototype GUI, accepts generic set of description files and generates user interface for building and running the binary. Implemented as an applet.
- Validates user choices, generates relevant scripts
- Integral part of the Grid-portal for LOOPS/Poseidon, it can be re-implemented in a more server-centric way (JSP etc.)
- Future directions for enhancement include:
 - Workflow composition: Employing the descriptions of the binaries and their input/output files as constraints. We are currently using predefined workflows.
 - Context mediation: When dataflow endpoints mismatch

GUI: validity checking

✓ AppletApplication	_ = ×					
<u>F</u> ile <u>T</u> ools <u>H</u> elp						
PARAMETERS INDEX						
Information Edit Value						
Name: DTUV						
Type: integer						
Value: 900 s	V Edit Variable					
Information Edit Value	Name: MIXVEL					
Name: DTSF	Type: enumerated					
Type: integer	Value: 4					
Value: 900 s	4 not in set {1;2;3}					
Information Edit Value						
Set 3	acceptableValues: 1;2;3 [List of acceptable values, separated by semicolon¢]					
Name: MIXVEL	OK Cancel					
Type: enumerated						
Value: 1						
Information Edit Value						
MIXVEL momentum equation horizontal mixing scheme:						
[2] Laplacian at constant depths.						
[3] Laplacian along terrain following levels.						
Parsing without schema-validation is succuessful.						

Interactive Visualization and Targeting of pdf's 0.28 2.76





Advanced Visualization and Interactive Systems Lab: A. Love, W. Shen, A. Pang

Interactive Visualization and Targeting of pdfs (cont.)





0.83

0,83

0.83



Status:

Search 2D CPs

Search 2D CP Candidates

Quit

Reset 2D CPs

CONCLUSIONS: Present and Future

- Advanced systems for adaptive sampling and adaptive modeling in a distributed computing environment
- Web interface, Remote visualization, Metadata for code and data, XML-based encapsulation of software, Grid computing infrastructure (SunGridEngine)
- Interdisciplinary data assimilation should contribute significantly to understanding, especially to the quantitative development of fundamental/simplified coupled models
- More interdisciplinary research and education needed: mathematics, computer science, physical-biogeochemicalacoustical ocean science, atmospheric science, earth science and complex system science
- Short-term impacts likely overestimated, long-term effects likely under-estimated

Feature Extraction for Adaptive Sampling

- Developing automated procedures to identify physical features of interest in the flow: upwelling, eddies & gyres, jets/fronts etc.
- Procedure can be based on a threshold for a derived quantity or a more complicated set of rules.
- Graphical output (in conjunction with uncertainty information) helps the user plan sampling patterns and vehicle paths.

