



The Millennium Cohort: a 21-Year Contribution to the Understanding of Military and Veterans' Health

Second Annual Trauma Stress Disorders Conference
Natcher Auditorium, NIH

December 10, 2009

MOMRP
Science to Soldier

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 10 DEC 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE The Millennium Cohort: a 21-Year Contribution to the Understanding of Military and Veterans' Health				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center, DoD Center for Deployment Health Research, San Diego, CA, 92106-3521				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at The Second Annual Trauma Spectrum Disorders Conference: A Scientific Conference on the Impact of Military Service on Families and Caregivers, 10 December 2009, Bethesda, MD					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 24	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Deployment Health Research



Lessons learned from 1991 Gulf War




The Origins of the Millennium Cohort Study



- IOM recommended coordinated prospective cohort study of service members
 - Capitalize on new DoD surveillance and health care data
 - Data sources that were not available at the time of the Gulf War
- For the first time, measure the impact of deployment prospectively

Section 743 of the FY1999 Strom Thurmond Act authorized the Secretary of Defense to establish a... ***longitudinal study*** to evaluate data on the health conditions of members of the Armed Forces upon their return from deployment.



Background




- **The Millennium Cohort Study is a longitudinal study designed to evaluate long-term subjective health and chronic diagnosed health problems, especially in relationship to exposures of military concern and deployments**
- **All services, active duty Reserve/National Guard**
- **Participants are re-surveyed at 3-year intervals including after service through 2022**
- **New accession cohorts were added in FY2004 and FY2007**
- **New accessions planned for FY2010, including a family component**

Basic Methodology



- Survey refined based on focus group testing, pilot study, and expert review
- Questionnaire leverages standard instruments (PHQ, PCL, SF-36V, others)
- Includes measures of physical health, behavioral health, mental health
- Includes exposure questions, and other metrics (deployment, sleep, etc.)
- Participants respond via traditional paper, or over secure website





Current Status

2001: Study launched

77,047 enrolled in Panel 1 (Wave 1)

2004: Panel 1 follow-up and Panel 2 enrollment initiated

86,131 enrolled / followed-up

2007: Panel 1 and 2 follow-up; Panel 3 enrollment


~ 115,000 enrolled / followed-up

2010: Panel 1, 2, and 3 follow-up; Panel 4 enrollment

Follow-up on > 150,000

- **> 70% with at least 1 follow-up**
- **~ 50% deployed in support of operations in Iraq and Afghanistan**
- **~ 20% have left military service**
- **> 30 peer-reviewed publications**
- **> 150 scientific presentations with many awards**

DoD and VA Data Sources





PTSD and Depression

- **PTSD Checklist-Civilian Version (PCL-C)²**
 - 17-item self-report measure (Likert 1 to 5)
 - PTSD if moderate or above level of at least one intrusion symptom, three avoidance symptoms, and two hyperarousal symptoms
 - And a score of 50 or more (range 17 to 85)

- **PRIME-MD Patient Health Questionnaire (PHQ)¹**
 - Psychosocial assessment based on scores of several health concepts
 - Major depressive syndrome (9 items)
 - Panic syndrome (15 items)
 - Other anxiety syndrome (6 items)
 - Eating disorders (4 items; binge and bulimia nervosa)

Has your doctor or other health professional EVER told you that you have any of the following conditions?

...

PTSD

Depression

...

(Spitzer, 1994; Spitzer, 1999; Spitzer, 2000)¹, (Weathers, 1993; Blanchard, 1996)²

Baseline PTSD Prevalence



- **At baseline, the weighted prevalence of PTSD was 3.6%**
 - **1.2% reported PTSD diagnosis without current symptoms**
 - **2.0% had PTSD symptoms without reported diagnosis**
 - **0.4% reported PTSD diagnosis with symptoms**

- **Those with PTSD at baseline were more likely to be:**
 - **Women**
 - **Less educated**
 - **Never married or divorced**
 - **Current smokers**
 - **Problem alcohol drinkers**

Baseline PTSD Prevalence



	Millennium Cohort N = 74,947	PTSD diagnosis without current symptoms N = 951	No PTSD diagnosis with current PTSD symptoms N = 1,487	PTSD diagnosis with current PTSD symptoms N = 284
Functional health				
MCS, weighted mean (95% CI)	53.0 (52.9, 53.1)	48.5 (47.8, 49.3)	27.4 (26.8, 28.1)	26.0 (24.5, 27.4)
PCS, weighted mean (95% CI)	53.4 (53.2, 53.6)	50.0 (49.3, 50.7)	48.0 (47.3, 48.7)	43.2 (41.6, 44.9)

- **Those with PTSD symptoms (2.4%)**
 - Significantly less favorable PCS and MCS scores
- **Those with reported PTSD diagnosis without symptoms (1.2%)**
 - Lower scores but closer to overall Cohort means



New-Onset PTSD

New-onset PTSD symptoms or diagnosis, over the approximate 3 year period between baseline and follow-up, was identified in:

7.6% - 8.7% of those who deployed with combat

1.4% - 2.1% of those who deployed without combat

2.3% - 3.0% of those who did not deploy

Smith TC, Ryan MAK, Wingard DL, Slymen DJ, Sallis JF, Kritz-Silverstein D, for the Millennium Cohort Study Team. New onset and persistent symptoms of posttraumatic stress disorder self-reported after deployment and combat exposures: prospective population-based US military cohort study. *British Medical Journal*. 2008 Feb;336(7640):366-71.



Persistent PTSD

Persistent PTSD symptoms or diagnosis, over the approximate 3 years between baseline and follow-up, was identified in:

43.5% - 47.9% of those who deployed with combat

22.4% - 26.2% of those who deployed without combat

45.9% - 47.6% of those who did not deploy



Photo source: <http://www.defenselink.mil/multimedia>

Prior Assault and New-Onset PTSD



- **New-onset PTSD symptoms or diagnosis among combat deployers was identified in:**

Women


- * **21.7% of women who reported prior assault**
- * **10.1% of women who did not report prior assault**

Men

- * **12.4% of men who reported prior assault**
- * **5.9% of men who did not report prior assault**

- **In contrast to hypotheses that survival from trauma represents or confers resilience, these findings suggest vulnerability to combat stress and PTSD among survivors of prior assault**

Functional Health and PTSD Symptoms



LeardMann CA, Smith TC, Smith B, Wells TS, Ryan MAK, for the Millennium Cohort Study Team. Baseline self-reported functional health predicts vulnerability to posttraumatic stress disorder following combat deployment: prospective US military cohort study. *British Medical Journal*. 2009; 338:b1273.

Positive and Negative Health Factors Associated with PTSD Symptoms



- **Vigorous physical activity was associated with decreased odds of new-onset and persistent PTSD symptoms**
- **Light/moderate exercise at follow-up was also associated with decreased odds of new-onset PTSD symptoms**
- **Those with new-onset PTSD symptoms experienced a 3-4 lb weight gain over those with no PTSD symptoms**
- **New-onset and persistent PTSD symptoms in nondeployed was associated with weight gain over those with resolved or no PTSD symptoms**
- **Herbal therapy and megavitamin use significantly more prevalent among participants with PTSD symptoms**

PTSD or Any Mental Health Diagnosis Among Those with PTSD Symptoms



PTSD Symptoms*	Dx code from time of symptom at		
	1 year (%)	3 years (%)	5 years (%)
2001-2003 N = 1876			
PTSD dx (ICD-9-CM code 309.81)	2.3	4.8	6.7
Any mental health dx (codes 290-319x)	26.4	41.7	49.4

*PTSD sensitive criteria among Panel 1 baseline participants

- A relatively small percentage received a PTSD-specific diagnosis through their care in the military health system within 5 years
- Nearly half received any mental health diagnosis within 5 years
- Among those with PTSD symptoms and subsequent PTSD diagnosis code
 - ~ 2/3 had persistent PTSD symptoms at 3-year follow-up
 - ~ 1/3 had resolved PTSD symptoms at 3-year follow-up

Alcohol Problems



- **Factors most strongly predictive of alcohol problems included younger age, smoking, and prior alcohol problems**
- **Reserve/Guard with reported combat exposures at increased odds of heavy weekly drinking, binge drinking, and alcohol-related problems**
- **Results consistent between men and women**
- **Results consistent across service branches**

Smoking



- **Deployment associated with an increase in smoking**
 - Increase predominantly due to smoking reuptake rather than smoking initiation
 - Among past smokers, deployment with combat, deploying multiple times, and deployment >9 months increased risk of smoking reuptake
 - Among baseline smokers, deployment not associated with increased amount of smoking





New-Onset Depression

- **New-onset depression symptoms:**
 - **5.7% of men and 15.7% of women who deployed with combat**
 - **2.3% of men and 5.1% of women who deployed without combat**
 - **3.9% of men and 7.7% of women who did not deploy**
 - **Deployed men and women who reported combat exposures had a significantly increased risk for depression compared with nondeployed service members**
 - **In contrast, deployed men and women who did not report combat exposures were at significantly lower risk for depression than nondeployed men and women**

Limitations of Presented Studies



- **Millennium Cohort is a random sample of the military population that may not be representative of all military personnel or those who deploy**
- **Those who were ill may have declined or participated in follow-up at different levels biasing new-onset estimates**
- **Self-reported exposure assessment is limited and not specific to deployment**
- **Use of a mental health screen as a surrogate for clinician diagnosis is imperfect**

Strengths of Presented Studies



- **71% follow-up**
- **Prospective investigations of specific characteristics and select populations possible from a large population-based sample of all services and components**
- **Preliminary findings investigating follow-up non-response did not find significant differences in measures of effect for PTSD after adjusting for non-response**
- **Self-reported mental health symptoms may be a better representation of symptom prevalence than diagnoses in medical databases**

Conclusions



- **Combat exposures, rather than deployment itself, significantly affect onset of mental health symptoms, problem alcohol drinking, and cigarette smoking post-deployment**
- **Significant amount of newly reported smoking and problem drinking associated with newly reported mental health symptoms post-combat deployment**
- **Specific populations including those with poor mental and/or physical health, and prior stressful life events could be targeted for PTSD prevention programs**

Acknowledgments



Millennium Cohort Study Team, San Diego

Melissa Bagnell, MPH; Ava Conlin, DO, MPH; Gina Creaven, MBA; James Davies; Lacy Farnell; Nisara Granado, MPH, PhD; Gia Gumbs, MPH; Lesley Henry; Jaime Horton; Isabel Jacobson, MPH; Kelly Jones; Cynthia LeardMann, MPH; Travis Leleu; Gordon Lynch; Jamie McGrew; Amanda Pietrucha, MPH; Teresa Powell, MS; Donald Sandweiss, MD; Amber Seelig, MPH; Beverly Sheppard; Besa Smith, MPH, PhD; Katherine Snell; Steven Speigle; Kari Welch, MA; Martin White, MPH; James Whitmer; and Charlene Wong, MPH

Millennium Cohort Co-Investigators

Dr. Tyler Smith, Dr. Paul Amoroso, Dr. Edward Boyko, Dr. Gary Gackstetter, Dr. Gregory Gray, Dr. Tomoko Hooper, Dr. James Riddle, Dr. Margaret Ryan, and Dr. Timothy Wells

Scientific Steering and Advisory Committee Members

Dr. Elizabeth Barrett-Conner; Dr. Dan Blazer; Dr. Harold Koenig; Mr. Michael O'Rourke; Dr. Larry Palinkas; Dr. Michael Peddecord; Mr. Joe Sharpe; Dr. Marie Swanson

Additional Collaborators and Supporters

Scott Seggerman, Management Information Division, DMDC; COL Carl Castro, Dr. Dennis Goodes, Mr. Mark Bither, MAJ Bonilla, Dr. Joan Hall, and Dr. Kate Nassauer, MOMRP; COL Karl Friedl and Dr. Richard Satava, MRMC; Dr. Charles Hoge, WRAIR; Dr. Susan Proctor and Dr. Kristin Heaton, USARIEM; COL Charles Engel, Dr. Kristie Gore, Dr. Michael Freed, and Dr. Xian Liu WRAMC; Dr. Kathryn M Magruder, Univ of South Carolina; Dr. William Schlenger, Abt Associates; Dr. John Fairbank, Duke University; Dr. Charles Marmar, UCSF; Dr. Seth Eisen, VA; Dr. Roger Gibson, USUHS; Dr. Deborah Wingard, Dr. Donna Kritz-Silverstein, and Dr. Thomas Patterson, UCSD; Dr. Caroline Macera, Dr. James Sallis, and Dr. Donald Sylmen, SDSU; Dr. Pam Keel, Univ of Iowa; COL Gaston Bathalon, USARIEM; Dr. Alyson Littman, Seattle ERIC; Anna Bukowinski, Sydney Lee and Carter Sevick, DoD BIHR; LtCol Chris Phillips, RAP; Dr. Robin Harris, Dr. Eyal Shahar, Dr. Grant Skrepnek, and Dr. Stephen Coons, Univ of Arizona; Dr. Sarah Fortuna, AFRL; CAPT Robert Koffman, BUMED; and professionals from the Institute for Systems Biology

We are indebted to the Millennium Cohort Study members for their continued participation!



MILLENNIUM COHORT STUDY



MOMRP
Science to Soldier