

Process Modeling of Large Amplitude Internal Waves in the South China Sea

Kraig Winters
University of California San Diego
Scripps Institution of Oceanography
9500 Gilman Drive
La Jolla CA 92093-0209

phone: (858) 822-3420 fax: (858) 534-0300 email: kraig@coast.ucsd.edu

Award Number: N00014-05-1-0573

<http://cfcd.ucsd.edu/>

LONG-TERM GOALS

My long-term objectives are to develop and employ numerical techniques to enhance our physical understanding of the life-cycle of energetic oceanic internal waves as well as our ability to predict the spatial and temporal variability of the induced flow fields. This goal encompasses the generation, long-range propagation and ultimately, the decay of such waves, either by shoaling or through energy transfer to other waves or modes of motion.

OBJECTIVES

My immediate objectives are twofold: (i) to assess and quantify the numerical requirements for simulation of NLIW evolution, including intensification and degeneration into packets, as they propagate in the nearly inviscid limit over distances comparable to the South China Sea basin and (ii) to validate the approach of combining numerical methods of high accuracy with an immersed boundary method to simulate the generation of internal waves produced by tidal flow over steep isolated sills as occur for example in the Luzon Strait.

APPROACH

To address item (i), simplified experiments were run in which wave generation was idealized using a localized near-surface body force. Waves excited at tidal frequency at energy levels comparable to observations in Luzon Strait and the SCS basin. The waves were then allowed to propagate freely across the deep basin. Given the approximate scales for such waves in the SCS, a scaling analysis of the governing equations, showing a near balance between nonlinear steepening and non-hydrostatic dispersion, was used as a basis for estimating the maximum allowable numerical errors in the various terms in the equations of motion to avoid contamination over the time required for such waves to traverse the basin. Simulations were then carried out using different numerical differentiation schemes with varying formal levels of accuracy on grids of varying resolution in an effort to discern the grid requirements for the different numerical approaches. To address (ii), a series of idealized experiments in which a time-periodic flow of uniformly stratified fluid was forced over an isolated obstacle were carried out. The primary experimental parameters were the degree of nonlinearity of the forcing (measured by the ration of the particle excursion length to the half-width of the sill) and the slope criticality parameter, i.e. whether the downward branches of free waves at the forcing frequency point into the topography (subcritical topography) or into the fluid (supercritical topography). These

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2008		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Process Modeling Of Large Amplitude Internal Waves In The South China Sea				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Scripps Institution of Oceanography, University of California San Diego, 9500 Gilman Drive, La Jolla, CA, 92093-0209				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT My long-term objectives are to develop and employ numerical techniques to enhance our physical understanding of the life-cycle of energetic oceanic internal waves as well as our ability to predict the spatial and temporal variability of the induced flow fields. This goal encompasses the generation, long-range propagation and ultimately, the decay of such waves, either by shoaling or through energy transfer to other waves or modes of motion.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

simulations were designed to complement and carefully compared with PIV results from the laboratory experiments of Peacock, Echeverri and Flynn at MIT (also under ONR NLIWI support).

WORK COMPLETED

Both sets of experiments have been completed and analyzed. Manuscripts have been written and are under internal review. The first will be submitted to Ocean Modeling and the second to the Journal of Fluid Mechanics. A new set of experiments has been initiated which look at the wave generation process in detail for more realistic stratification (thin, highly stratified surface layer overlying weaker stratification below) at much higher values of the nonlinearity parameter. I am collaborating with Larry Army at SIO on these experiments.

RESULTS

Accuracy requirements were quantified for simulation of nonlinear wave packets over basin scales. Whether or not such waves are represented numerically in a physically reasonable manner is shown to be highly scale-dependent and intrinsically coupled to the discretization methods employed as well as the time and space scales of interest. For ocean basin-scale propagation, as considered here, physically meaningful resolution of these waves demands not only a high degree of overall numerical accuracy but also a careful consideration of the behavior of the explicit but a-physical dissipation operator employed. The introduction of such an operator is an unfortunate necessity. One would like to capture the dynamics at all discrete scales in the inviscid limit but cannot in the face of significant nonlinearities that transfer energy to small scales. At best, we can hope for a calculation in which the scales of interest are treated in a demonstrably nearly adiabatic limit while confining the effects of the dissipation operator, over the time scales of interest, to spatial scales that exist on the given numerical grid but must be considered unresolved in the context of the physics. The experiments for different numerical methods at different grid resolution show the feasibility of achieving these goals using different approaches.

From a technical perspective, good agreement was demonstrated between experimentally measured flow fields using PIV and numerically generated fields combining high-order numerical schemes with an immersed boundary approach. The fields were decomposed into vertical modes and radiation fluxes computed. Results from both methods were compared with the linear theoretical predictions of Petrelis, Llewellyn-Smith and Young (2006).

The more nonlinear, nonuniform stratification experiments are still in the early stages but we are simulating flows very reminiscent of the Knight Inlet observations in which packets of steep waves propagate upstream from the sill prior to the reversal of the tidal currents.

IMPACT/APPLICATIONS

Potential impacts include an objective framework to discuss and assess the performance of different classes of numerical models for the generation and long-range propagation of NLIWs.

RELATED PROJECTS

Internal wave generation in the laboratory (NLIWI), Prof. Thomas Peacock. Described above.

PUBLICATIONS

Density Stratification, Turbulence, but How Much Mixing? G.N. Ivey, K. B. Winters, and J. R. Koseff, Annual Review of Fluid Mechanics Vol. 40: 169-184 (Volume publication date January 2008) (doi:10.1146/annurev.fluid.39.050905.110314) [published, refereed]

Growth of Inertia-Gravity Waves in Sheared Inertial Currents, Winters, K. B. (2008), J. Fluid Mech. 601 , 85 – 100
[published, refereed]