AD_____

Award Number: W81XWH-08-1-0464

TITLE: Generation of Breast Cancer Stem Cells by the EMT

PRINCIPAL INVESTIGATOR: Robert A. Weinberg, Ph.D.

CONTRACTING ORGANIZATION: Whitehead Institute for Biomedical Research, Cambridge, MA 02142

REPORT DATE: October 2010

TYPE OF REPORT: Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

X Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
Public reporting burden for this data needed, and completing a this burden to Department of D 4302. Respondents should be valid OMB control number PI	collection of information is estir ind reviewing this collection of ir efense, Washington Headquart aware that notwithstanding any CASE DO NOT PETURN YOU	nated to average 1 hour per resp formation. Send comments rega ers Services, Directorate for Infor other provision of Iaw, no person FORM	onse, including the time for revie arding this burden estimate or an mation Operations and Reports 1 shall be subject to any penalty f	wing instructions, searc y other aspect of this co (0704-0188), 1215 Jeffe for failing to comply with	hing existing data sources, gathering and maintaining the illection of information, including suggestions for reducing rson Davis Highway, Suite 1204, Arlington, VA 22202- a collection of information if it does not display a currently	
1. REPORT DATE (DD	D-MM-YYYY)	2. REPORT TYPE		3. D	ATES COVERED (From - To)	
4. TITLE AND SUBTIT	LE	final Report		5a.	CONTRACT NUMBER	
Generation of	Breast Cancer	Stem Cells by	the EMT			
			5b. W 8	GRANT NUMBER		
				5c.	PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d.	PROJECT NUMBER	
				5e.	TASK NUMBER	
Robert A. Weinberg, Ph.D.						
					WORK UNIT NUMBER	
weinberg(<i>a</i>)wi.mit.e	du ANIZATION NAME(S)	AND ADDRESS(ES)		8. P	ERFORMING ORGANIZATION REPORT	
				N	IUMBER	
Whitehead Institute for Biomedical Research						
Cambridge, MA 02142						
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research Materiel Command				10.	SPONSOR/MONITOR'S ACRONYM(S)	
Fort Detrick, Maryland				11.	SPONSOR/MONITOR'S REPORT	
21702-5012					NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT						
Approved for public release; distribution unlimited						
13. SUPPLEMENTARY NOTES						
14 APSTRACT						
The epithelial-mesenchymal transition plays an important role in the progression of benign human breast cancier cells into						
highly malignant carcinoma derivatives. Recently, we have fo und that in addition to promoting the invasive and motile						
phenotypes of cancer cells, the EMT also confers on cancer cells certain stem-cell properties. However, it rem ained unclear						
whether such EMT products are actually stem cells or only cells having many of the properties of stem cells. In this study, we						
provided evi dence that the EMT indeed produces bona fide mammary stem cells, which are able to reconstitute entire						
mammary ductal trees in vivo. We also identified key EMI-associated transcription factors that act as master regulators of						
maintaining pre-existing stem cells						
15. SUBJECT TERMS						
			/ _			
16. SECURITY CLASSIFICATION OF:U			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC	
a. REPORT	b. ABSTRACT	c. THIS PAGE	ບບ		19b. TELEPHONE NUMBER (include area	
U	U	U		7	code)	

Generation of Breast Cancer Stem Cells by the EMT Robert A. Weinberg, Ph.D.

Table of Contents

<u>Page</u>

Introduction1	
Body1	
Key Research Accomplishments	3
Reportable Outcomes	3
Conclusion	3
References	
Appendices	-

Introduction

The progression of be nign hum an breast cancer cel ls into highly m alignant derivatives is accompanied by the acquisition of m esenchymal cell ch aracteristics, motility, and invasivenes s. These multifaceted changes in cell pheno types are in volved in the transd ifferentiation program term ed the epithelial-mesenchymal transition (EMT). Recently, we discovered that in a ddition to conferring cell-biological traits of high-grade m alignancy, the EMT program also coverts m ore differentiated m ammary epithelial cells into derivatives ha ving many of the attributes of epit helial stem cells, a conversion that operates in both imm ortalized and in transformed, tum origenic mammary epithelial cells. Ho wever, it remained unclear whether. The outcom e of t hese experiments holds im portant im plications for the mechanisms controlling the form ation of normal mammary epithelial stem cells as w ell as the formation of breast cancer stem cells. In this study period, we have gained substan tial insight into the identities of the master regulators of mammary stem cells.

Body

Our previous work demonstrated that differentiated mammary epithelial cells (MECs) that have undergone an epithelial-mesenchymal transition (EMT) acquire many properties of mammary stem cells, including the expression of stem-cell-surface markers and the ability to form mammospheres. In this study, we posed the questions of whether forced passage through an EMT can indeed form normal mammary gland stem cells (SCs), which can reconstitute an entire mammary ductal tree upon being transplanted into a cleared mammary fat pad, and whether the EMT-inducing transcription factors can act as master regulators of mammary SCs.

In the previous progress report cycle, we found that mammary stem cells specifically express the EMT-inducing Slug transcription factor (TF) relative to luminal progenitor cells or fully differentiated luminal cells (~90 fold higher in mammary SCs). This suggested to us that Slug was likely to be an excellent candidate for the regulators of the mammary SC state. We therefore asked whether induction of an EMT by Slug in mammary epithelial cells could generate mammary SCs. To test this, we transiently expressed Slug in MECs for 10 - 12 days using tetracycline-inducible lentiviral constructs, and subjected the cells to the in vivo gland-reconstitution assay. We found that cells having undergone an EMT induced by Slug had 40 - to 100-fold higher mammary gland-reconstituting activity. This experiment showed that induction of an EMT indeed leads to generation of bona fide mammary stem cells. However, because this experiment was done with MECs containing pre-existing SCs, we could not rule out the possibility that Slug may expand pre-existing SCs, rather than converting non-stem cells into stem cells.

In the past year or so, we focused on convincingly demonstrating that the EMT is able to convert non-stem cells in to mammary SCs. First, we isolated either luminal progenitor cells or fully differentiated luminal cells that are devoid of mammary stem cells by FACS on the basis of CD49f and CD61. We then tested whether Slug could generate SCs in each nonstem cell population. We infected the cells with the tetracycline-inducible Slug construct or control vector and induced Slug expression for 5 days in adherent culture, and then stopped the doxycycline treatment and subjected the cells to an in vitro 3D Matrigel culture system in order



Figure 1: Luminal progenitor cells (top) or fully differentiated luminal cells (bottom) were infected with control or Slug vectors. After a 5-day treatment of doxycycline, cells were subjected to 3D matrigel culture to assay for mammary stem-cell activity.

1

to assay mammary SC activity. In this type of assay, mammary stem cells can form complex 3D ductallobular organoids, As shown in Figure 1, transient expression of Slug efficiently induced mammary stem cells from luminal progenitor cells, however it failed to induce mammary stem cells from fully differentiated luminal cells. This experiment demonstrated that induction of an EMT by Slug can indeed generate stem cells from non-stem cells; however it could only do so in the progenitor cells. We reasoned that relative to luminal progenitor cells, the fully differentiated luminal cells may lack certain factor(s) that can cooperate with Slug to induce mammary SC state.

We therefore gathered a dozen or so TFs that have been associated with stem cell biology in general or factors that have been shown to act together with EMT-inducing TFs during developmental processes. We co-expressed each individual factor with Slug in fully differentiated luminal cells in order to screen for factors that can act together with Slug to induce stem-cell activities in the fully differentiated cells. We found that co-expression of Sox9 with Slug can robustly induce stem-cell activity in the fully differentiated cells (Figure 2), while expression of either Slug or Sox9 alone failed to induce mammary SC. We analyzed the expression pattern of Sox9 in different cell types of



mammary epithelium using a strain of Sox9-GFP reporter mice. We found that Sox9 is highly expressed in the luminal progenitor cells and a subset of basal cell population that contains stem cells. The fact that only luminal progenitors but not fully differentiated luminal cells express Sox9 explains why Slug expression can generate mammary SCs only in luminal progenitor cells but not in fully differentiated cells.

We further asked whether Sox9 and Slug indeed convert fully differentiated luminal cells to mammary SCs that can regenerate an entire mammary ductal tree. We infected the fully differentiated luminal cells with tetracyclineinducible Slug and Sox9 and induced the expression of constructs by doxycycline for 5 days in adherent culture, and then transplanted the cells in a 1:1 mixture with RFPexpressing control MECs in cleared mammary fat pads. As anticipated, the control vectorexpressing luminal cells had no reconstitution activity at all. In stark contrast, the



Figure 3: Slug and Sox9 induce bona fide mammary stem cells. Left panels: represent images of cleared fat pads injected with indicated cell types; middle panels: flow cytometry analysis of GFP and RFP cells in transplanted fat pads; right graph: the ratio of GFP vs RFP cells in fat pads transplanted by each cell type.

Weinberg, Robert A. (BC073843)

Slug/Sox9-expressing luminal cells reconstituted the fat pads robustly (Figure 3).

The above experiments demonstrated that the EMT in collaboration with Sox9 can convert even the fully differentiated MECs – cells at the bottom of differentiation hierarchy – back into a bona fide stem-cell state. We further asked whether Slug and Sox9 are required for maintaining the pre-existing stem cells in the mammary epithelial cell population. We knocked down the expression of Slug or Sox9 in unfractionated MECs using lentivirus vector-expressed shRNAs and then gauged mammary stem-cell activity of these cells using both in vitro or in vivo assays. In both assays, knocking down of either genes leads to severe to complete inhibition of mammary stem-cell activity (13-fold decrease to complete inhibition). These data demonstrated that EMT-inducing factor Slug and Sox9 act as master regulators of mammary stem-cell state. They are both sufficient in inducing mammary SCs from non-SCs and necessary for maintaining pre-existing SCs.

Key Research Accomplishments

- The induction of an EMT has now been demons trated to convert a normal murine m ammary epithelial c ells into m ammary epithelial s tem cells that can, following orthotop ic implantation, generate an entire m ammary ductal tree. This o bservation offers us for the first tim e a definitive proof of the connection between EMT and entran ce into a stem -cell state. The EMT-inducing transcription factor Slug acts as master regulator for the stem-cell state.
- Genetic screens have identified S ox9 as an important co-factor for Slug-mediated stem-cell induction and maintenance.

Reportable Outcomes

This supported the following publications:

-Valastyan, S., Reinhardt, F., Benaich, N., Calogrias, D., Sza', A.M., Wang, Z.C., Brock, J.E., Richardson, A.L., and Weinberg. R.A. (2009) A pleiotropically acting MiroRNA, miR-31, inhibits breast cancer metastasis. Cell, 137: 1032-1046. PMCID: PMC2766609

-Valastyan, S., Benaich, N., Chang, A., Reinhardt, F., and Weinberg, R.A. (2009) Concomitant repression of three target genes can explain the impact of a microRNA on metastasis. Genes and Development, 23:1-6 PMCID: PMC2779763

-Valastyan S, Chang A, Benaich N, Reinhardt F, Weinberg RA.(2010) Concurrent suppression of integrin alpha5, radixin, and RhoA phenocopies the effects of miR-31 on metastasis. Cancer Res. 2010 Jun 15;70(12):5147-54. Epub 2010 Jun 8. PMCID: PMC2891350

Conclusion

One major implication of the reported work derives from experiments reported herein in which we have succeeded in greatly increasing the proportion of normal mammary epithelial stem cells in a population of differentiated mammary epithelial cells, doing so by expressing EMT-inducing transcription factors in these cells transiently. Since the EMT can otherwise be induced by contextual signals that have been and are being identified, this suggests that a protocol can be developed involving exposure of differentiated epithelial cells to a mixture to EMT-inducing signals that can substantially increase the proportion of normal epithelial stem cells in these cultures. This holds, in turn, important implications for future modalities of regenerative therapy designed to restore damaged epithelial tissues through implantation of normal, syngeneic epithelial stem cells.

A second major outcome of this work is that we have now started to understand the core transcription regulatory network of mammary stem cells by identifying the master regulators of the

Weinberg, Robert A. (BC073843)

mammary stem-cell state. This core stem-cell transcription network may also play important role in determining the cancer stem cell state, which is being studied intensively by us now.