

This article was downloaded by:

On: 18 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Energetic Materials

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713770432>

Synthesis of Insensitive 5,5'-Azotetrazolate Salts

Kirstin F. Warner^a; Richard H. Granholm^a

^a Indian Head Division, Naval Surface Warfare Center, Indian Head, Maryland

Online publication date: 10 January 2011

To cite this Article Warner, Kirstin F. and Granholm, Richard H.(2011) 'Synthesis of Insensitive 5,5'-Azotetrazolate Salts', *Journal of Energetic Materials*, 29: 1, 1 – 6

To link to this Article: DOI: 10.1080/07370652.2010.490255

URL: <http://dx.doi.org/10.1080/07370652.2010.490255>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE JAN 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Synthesis of Insensitive 5,5'-Azotetrazolate Salts				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Surface Warfare Center, Indian Head Division, 4104 Evans Way, Suite 102, Indian Head, MD, 20640				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Synthesis of Insensitive 5,5'-Azotetrazolate Salts

KIRSTIN F. WARNER and
RICHARD H. GRANHOLM

Indian Head Division, Naval Surface Warfare Center,
Indian Head, Maryland

Salts of 5,5'-azotetrazolate with 5-aminotetrazole and 2,4,6-triamino-s-triazine cations were synthesized as potential ingredients for gun propellants. These salts were characterized by nuclear magnetic resonance (NMR) and infrared spectroscopy. Thermal properties were studied by differential scanning calorimetry (DSC). Their reactions to mechanical stimuli were also determined by small-scale sensitivity testing, small-scale shock reactivity testing, and small-scale internal blast testing.


Keywords: 5,5'-azotetrazolate salts, energetic salts, explosives, high nitrogen

Introduction

In recent years, there has been a great deal of interest in high nitrogen compounds [1–8]. A series of tests was performed to gain insight into their thermal stability, handling, and sensitivity and the explosive characteristics of these salts. These salts are potential ingredients for gun propellants and gas generators.

This article not subject to US copyright law.


Address correspondence to Kirstin F. Warner, Indian Head Division, Naval Surface Warfare Center, 4104 Evans Way, Suite 102, Indian Head, MD 20640. E-mail: kirstin.warner@navy.mil



Scheme 1. Synthesis of 5,5'-azotetrazolate.

This article addresses the synthesis and insensitivity of 5,5'-azotetrazolate salts with 5-aminotetrazole and 2,4,6-triamino-*s*-triazine cations.

5-Aminotetrazole monohydrate and 2,4,6-triamino-*s*-triazine are commercially available and were used as received. The corresponding hydrochloride salt was easily prepared by reacting



Scheme 2. Synthesis of 2,4,6-triamino-*s*-triazinium 5,5'-azotetrazolate.

5-aminotetrazole monohydrate or 2,4,6-triamino-s-triazine with 37% hydrochloric acid in methanol as shown in Schemes 1 and 2 [9]. The corresponding hydrochloride salt was allowed to react with sodium 5,5'-azotetrazolate in an aqueous solution to afford 5-aminotetrazolium 5,5'-azotetrazolate (**1**) and 2,4,6-triamino-s-triazinium 5,5'-azotetrazolate (**2**), respectively.

Results and Discussion

Differential scanning calorimetry (DSC) measurements were performed at 5°C/min in a nitrogen atmosphere at a flow rate of 50 mL/min. 5-Aminotetrazolium 5,5'-azotetrazolate (**1**) began to decompose at 134.3°C with a peak exotherm at 170°C. 2,4,6-Triamino-s-triazinium 5,5'-azotetrazolate (**2**) began to decompose at 248.3°C with a peak exotherm at 270°C. As shown in Table 1, the small-scale sensitivity data of (**1**) and (**2**) were in the low to medium sensitivity range for impact, friction, and electrostatic discharge compared to RDX with medium sensitivity.

Explosive behavior is summarized in Table 2. The small-scale shock reactivity test (SSRT) produces a dent, with its depth correlating with reactivity [10]. 5-Aminotetrazolium 5,5'-azotetrazolate produced a dent depth of 2.5 mm and an internal blast peak quasi-static overpressure (P_{QS}) of 15 psig,

Table 1
Small-scale sensitivity data

Material	ERL impact 50% hgt (cm)	ABL friction 20 TIL (psig)	ABL ESD 20 TIL (joules)	DSC onset/ peak (°C)
5-aminotetrazolium 5,5'-azotetrazolate	178	180	0.326	134.3/270
2,4,6-triamino-s- triazinium 5,5'- azotetrazolate	100	≤560	≤0.095	248.3/270
Reference: RDX	21	135	0.165	212/235

Table 2
Small-scale shock reactivity and internal blast data

Material	Dent depth (mm)	P _{QS} (psig)
5-aminotetrazolium 5,5'-azotetrazolate	2.5	15
2,4,6-triamino-s-triazinium 5,5'-azotetrazolate	2.2	13
TAGzT	4	18
Detonator (PETN RP-80)	2.2	18

and 2,4,6-triamino-s-triazinium 5,5'-azotetrazolate salt produced a dent depth of 2.2 mm and an internal blast peak quasi-static overpressure (P_{QS}) of 13 psig. These two test compounds produced dents about the same as the blank test (detonator alone), indicating that the compounds have little or no potential to detonate. Triaminoguanidinium 5,5'-azotetrazolate (TAGzT) was used as a standard that had a dent depth of 4.0 mm and an internal blast peak quasi-static overpressure (P_{QS}) of 18 psig. TAGzT does show shock reactivity and has a potential to detonate. Internal blast peak quasi-static overpressure (P_{qs}) generally correlates with total energy release. However, in the case of high-nitrogen propellants, their intrinsic lower flame temperatures combined with heat loss in a small-scale test may prevent efficient burning of combustible product species. The low internal blast pressures for the two test compounds compared to TAGzT suggest lower reactivity.

Experimental

CAUTION: *Although we experienced no difficulty handling these salts, care should be taken when handling these energetic materials.*

General Procedure

Sodium 5,5'-azotetrazole pentahydrate was prepared following the Thiele method [1]. It was added to a magnetically stirred

solution of the respective chloride in water. After 2 hr at room temperature, the solution was concentrated under vacuum to afford a solid.

5-Aminotetrazolium 5,5'-azotetrazolate (1). Yield 90%. DSC: $T_{\text{Dec.}} = 170^{\circ}\text{C}$, $T_{\text{Onset}} = 134.3^{\circ}\text{C}$ (274 J/g). IR (cm^{-1}): 3,580 (m), 1,650 (vs), 1,588 (vs), 1,400 (s), 1,265 (m), 1,195 (m), 1,192 (m), 1,148 (m), 1,071 (s), 1,050 (s), 779 (s), 739 (s), 733 (s), 680 (m). ^1H NMR (DMSO): δ 6.4, 3.2; ^{13}C NMR (DMSO): δ 174, 157. Density: 1.73 g/cm³.

2,4,6-Triamino-s-triazinium 5,5'-azotetrazolate (2). Yield 90%. DSC: $T_{\text{Dec.}} = 270^{\circ}\text{C}$, $T_{\text{Onset}} = 248.3^{\circ}\text{C}$ (682 J/g). IR (cm^{-1}): 3,389 (m), 3,285 (m), 3,149 (m), 1,650 (vs), 1,662 (vs), 1,643 (vs), 1,606 (s), 1,559 (vs), 1,512 (m), 1,483 (m), 1,406 (m), 1,390 (m), 1,334 (m), 1,261 (m), 1,205 (m), 1,179 (m), 1,092 (m), 1,063 (m), 1,023 (m), 972 (m), 871 (m), 785 (m), 741 (m), 728 (m), 679 (m). ^{13}C NMR (DMSO): δ 163, 157. Density: 1.70 g/cm³.

Conclusions

Salts of 5,5'-azotetrazolate dianion with 5-aminotetrazolate and 2,4,6-triamino-s-triazine cations were synthesized and studied. These salts showed good thermal stability and low to medium sensitivity to impact, friction, and electrostatic discharge. 5-Aminotetrazolium 5,5'-azotetrazolate had a dent depth of 2.5 mm, which was slightly more than an inert response, and 2,4,6-triamino-s-triazinium 5,5'-azotetrazolate had a dent depth of 2.2 mm, which was inert, producing a dent equivalent to the detonator.

Acknowledgments

Financial support was provided by IHD, NSWC CORE Program. Dr. Daniel N. Sorensen provided the DSC data. Daniel L. Remmers provided small-scale safety data. Sodium 5,5'-azotetrazolate salt was provided by Jerry Salan. Dr. Christine Michienzi was a source of helpful discussions.

References

- [1] Thiele, J. 1898. Synthesis of 5,5'-azotetrazolate. *Liebigs Annalen*, 303: 57–75.
- [2] Reddy, G. O. and A. K. Chatterjee. 1983. A thermal study of salts of azotetrazole. *Thermochemica Acta*, 66: 231–244.
- [3] Koldobskii, G. I. and V. A. Ostrovskii. 1994. Tetrazole. *Russian Chemical Reviews*, 63(10): 797–814.
- [4] Hammerl, A., G. Holl, T. M. Klapotke, P. Mayer, H. Noth, H. Piotrowski, and M. Warchhold. 2002. Salts of 5,5'-azotetrazole. *European Journal of Inorganic Chemistry*, 2002(4): 834–845.
- [5] Silvabalan, R., M. B. Talawar, N. Senthilkumar, B. Kavitha, and S. N. Asthana. 2004. Studies of azotetrazolate based high nitrogen content high energy materials potential additives for rocket propellants. *Journal of Thermal Analysis and Calorimetry*, 78: 781–792.
- [6] Klapotke, T. M. 2007. New nitrogen rich high explosives. *Structure and Bonding*, 125: 85–121.
- [7] Klapötke, T. M. and C. Miró Sabaté. 2008. Nitrogen-rich tetrazolium azotetrazolate salts: A new family of insensitive energetic materials. *Chemistry of Materials*, 20(5): 1750–1763.
- [8] Klapötke, T. M. and C. Miró Sabaté. 2009. New energetic compounds based on the nitrogen-rich 5,5'-azotetrazolate anion ($[\text{C}_2\text{N}_{10}]^{2-}$). *New Journal of Chemistry*, 33(7): 1605–1617.
- [9] Marchewka, M. K. 2002. Infrared and Raman spectra of melamine hemihydrate. *Materials Science and Engineering*, B95: 214–221.
- [10] Granholm, R. H. and H. W. Sandusky. 2006. Small-scale shock reactivity and internal blast test. In *Shock Compression of Condensed Matter-2005*, M. D. Furnish, M. Elert, T. P. Russel, and C. T. White (eds.), *American Institute of Physics Conference Proceedings*, 845, Pt II: 1257–1260.