

Coastal Benthic Optical Properties of Coral Environments: ROV/AUV Imaging

Kendall L. Carder

Marine Science Department, University of South Florida
St. Petersburg, FL 33701-5016

Phone: (813) 553-1148 Fax: (813) 553-3918 email: kcarder@monty.marine.usf.edu

David K. Costello

Marine Science Department, University of South Florida
St. Petersburg, FL 33701-5016

Phone: (813) 553-3953 Fax: (813) 553-3918 email: dkc@monty.marine.usf.edu

Award Number: N00014-97-1-0006

LONG-TERM GOALS

The deconvolution, quantification, and interpretation of the various components of water-leaving radiance in shallow coastal waters with emphasis on coral environments are the long-term goals of the project.

OBJECTIVES

In this project, objectives include the development of instrumentation and models to measure and predict the contribution of bottom reflectance to upwelling radiance in coastal waters. An underlying objective, then, is the development of the methodologies required to remotely classify bottom types in varying water depths. Intrinsic in this effort are the quantification of the optical properties of the water column and the need to perform rigorous data calibration/validation while working toward optical closure and to address the inherent problems of scale between *in situ* and remotely sensed data.

An collateral objective is to create a world-wide-web repository for data from the CoBOP project to enable investigators to efficiently explore the large and diverse data set.

APPROACH

In the first funding year of the CoBOP project (FY96), transects over coral bottoms in the Dry Tortugas were laid out and mapped by divers and by the Fluorescence Imaging Laser Line Scanner (FILLS). Instrumentation aboard our Remotely Operated Vehicle (ROV) and Autonomous Underwater Vehicle (AUV) platforms were used to determine the color and intensity of bottom elements from different altitudes (Costello et al., 1997). The goal was to correct imagery for path radiance and attenuation, providing bottom albedo estimates for the dominant bottom types/features, to image bottom fluorescence, and to measure the vertical spectral structure of the upwelling and downwelling light fields. The analyses required rigorous validation/calibration efforts and modeling efforts. Simultaneously, effort was expended toward developing simple, relatively low-cost methods that could exploit gross bottom reflectance signatures to yield useful data.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 1998	2. REPORT TYPE	3. DATES COVERED 00-00-1998 to 00-00-1998			
4. TITLE AND SUBTITLE Coastal Benthic Optical Properties of Coral Environments: ROV/AUV Imaging		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of South Florida, Department of Marine Science, St. Petersburg, FL, 33701-5016		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

In this past funding year, a CoBOP field campaign was conducted from the Caribbean Marine Research Center (CMRC) located on Lee Stocking Island, Exuma Islands, Bahamas. Our responsibilities in the 1998 fieldwork were similar to those of the Dry Tortugas exercises with the addition of extensive atmospheric measurements and the inclusion of sediment, grassbed, and mixed bottom types in addition to coral environments.

WORK COMPLETED

Work during this period has consisted of reduction and modeling of CoBOP'96 and TWIST2 (Sombrero Reef, FL Keys) data, the participation in the CoBOP'98 field campaign, and the construction of a data repository to serve the CoBOP project.

General efforts:

- Significant effort has been expended toward building a JAVA-based, data archive site for the CoBOP project team for interactive data exploration on the world wide web. Sample data sets have been obtained, a searchable heading format has been designed, and a prototype structure has been built which allows limited interactive plotting of geophysical data from a location of interest within the study area (Hou et al., 1998b).
-
- A rigorous check of the geometric collection performance (cosine response) of USF-developed, 512-channel in situ irradiometers and of a LICOR-1800 has been performed (English et. al. 1998).
-
- Methods have been presented to estimate scattering by large (>0.1 mm) particles (Hou et. al. 1997) and a forward-angle scattering correction for transmissometers determined by the large-particle size distribution (Hou et. al. 1998a).
-
- A method has been described which utilized color video taken from the OV-II AUV to calculate the percent live sea-bottom cover (Renadette et. al. 1997, 1998).
- A method has been developed to model and utilize downwelling irradiance spectra affected by wave focusing (Costello et. al. 1998a).
-
- A method to vicariously calibrate aircraft and satellite spectrometers using cloud shadows also corrects for atmospheric effects is applicable to NRL's Ocean Imaging Spectrometer (Reinersman et al., 1998).

CoBOP'98 field efforts

- 490 Hyperspectral (512-channel) downwelling irradiance and upwelling irradiance spectra were acquired during 25 vertical profiles performed using the ROSEBUD ROV.
-
- CTD, c-488, c-660, FL-685 and Bb-830 data were acquired during 30 vertical profiles performed using the ROSEBUD ROV launched from the R/V Subchaser.
-

- 6-channel, intensified bottom-albedo-video as well as NTSC-color-video bottom imagery were also obtained during each ROV deployment.
-
- 47 JSD drop-package casts were performed from the R/V Suncoaster. The JSD recorded CTD, ac-9, 0.2 micron-filtered ac-9, 488-nm attenuation, 830 nm backscatter, 6-channel HS-6 backscatter, Flashpak CDOM fluorescence, and chlorophyll fluorescence.
-
- Sea surface Rrs, filter pad (pigment, detrital, and CDOM) absorption (44 stations), HPLC (37 stations), and fluorometric chlorophyll (44 stations) were obtained.
-
- Color video bottom imagery was obtained during mapping transects using the R/V Subchaser stern-mounted bottom camera. The imagery is overlaid with GPS position and time. The Subchaser ship computer simultaneously logs water depth and surface temperature, boat speed and heading, and data from the boat meteorological instrumentation.

RESULTS

- Water-Raman scattering and chlorophyll a fluorescence are extremely significant components of the upwelling light field at depths > 2 m and wavelengths > 520 nm over coralline environments and cannot be ignored in evaluating bottom-reflected (actively or passively) radiance.
- Solar-stimulated fluorescence at 685 nm from sediments due to benthic diatoms is ubiquitous on the Florida shelf and is apparent in bottom imagery from depths of 7 m to > 20 m but < 30 m during the summer.
- Animals (e.g. sponges) and man-made objects are readily apparent by their dark contrast with the bright, red, bottom fluorescence from benthic diatoms, coral symbionts, and macrophytes.
- Range to various components in an image greatly affects the 685-fluorescence signal since the e-folding depth through water is 2.5 m. Correction for range is critical for image interpretation (see Carder and Costello: N00014-96-1-5013)
- Wave focusing has a very significant spectral impact on the instantaneous downwelling light field on clear days providing red-rich irradiance in focal zones and blue-rich irradiance in divergence zones. Coral and vegetation fluorescence are spectrally and temporally dependent on the incident light field, and fluorescence “spill-over” may occur when photosynthetic reaction centers are full during wave focusing events but not for steady-state conditions providing the same time-averaged photon quality and quantity. Field measurements of IOPs and AOPs have allowed spectral model closure calculations to simulate the instantaneous spectral light field measurements. Increases in aerosols decrease these fluctuations due to wave focusing, stabilizing the light field (Costello et al., 1998a). In clear, shallow coastal waters, upwelling radiance can be significantly affected by bottom reflection of focused downwelling irradiance (Costello et al., 1998b).

IMPACT/APPLICATIONS

In our analysis of our hyperspectral light profile data sets, it soon became apparent that wave focusing and Raman scattering introduced significant complexity to models of the submarine light field. Moreover, modeling efforts toward addressing the complexities using Smith and Baker's (1981) water absorption numbers were generally fruitless. Use of Pope and Fry (1997) numbers, modeled skylight and sunlight fields (Gregg and Carder, 1990) and focus/defocus of the sunlight contribution provided modeled light fields consistent with measurements. Fluorescence and primary production models need to consider the nonlinear aspects of this light field versus traditional time-averaged methods.

The differences in the 685 nm fluorescence yields, suggested by our observations, ranged from high for hard-bodied coral, medium for branching coral, to low for benthic diatoms. These differences suggest possible automatic classification schemes if adequate range information is available. Certainly, non-vegetative bottom features such as animals and man-made objects are sharply discernible when viewed at 685 nm, and path radiance due to backscattering does not reduce image contrast for fluorescence-dominated scenes.

Aircraft and space sensor calibrations change with vibration, temperature, and time. Recalibration is critical since a small change in calibration can introduce large errors. For example, the atmosphere contributes 90% (or more) of the radiance received at a space-borne sensor. Just a 2% shift in calibration would, then, result in a 20% error in the calculated water-leaving radiance. Reinersman et al. (1998) provide a vicarious method to re-calibrate a sensor that simply depends on finding small, compact clouds in a scene.

TRANSITIONS

The hyperspectral data acquired during this project has aided other projects (see RELATED PROJECTS section) in efforts toward the development and validation of algorithms for remote sensing of water constituents and bathymetry in coastal waters. Additionally, the Real-time Ocean Bottom Optical Topographer (ROBOT) module, an AUV-borne, laser-line-based, instrument system which was first deployed during the CoBOP'98 field campaign will participate in Navy-sponsored underwater mine detection exercises in December, 1998.

RELATED PROJECTS

As part of the CoBOP Directed Research Initiative, this project is synergistic with numerous other CoBOP investigations and several multi-discipline investigations are underway. This project also provides significant data to and benefits from important instrumentation developed under "Optical Variability and Bottom Classification in Turbid Water" (ONR CODE 3220M).

Other collaborative projects:

ONR - *Shelf Erosion and Sedimentation Processes: Quantification and Prediction of Their Effects on the Optical Properties of Coastal Waters.* PIs - K.L. Carder, D.K. Costello (USF)

- ONR - *An AUV-based investigation of the role of nutrient variability in the predictive modeling of physical process in the littoral ocean.* PIs - K.A. Fanning and J.J. Walsh (USF).
- ONR - *A simulation analysis of the time-dependent roles of phytoplankton and CDOM in effecting the 3-dimensional structure of inherent optical properties on the West Florida shelf.* PI - J.J. Walsh (USF)
- ONR - *A Multi-Disciplinary Investigation of the Nature and Predictability of Sediment Resuspension in Shallow Water: Its Effect on Water Column and bottom Optical Properties.* PIs - A.C. Hine, D.P. Howd, D. Mallinson, D. Naar (USF), D. Wilson (USGS)
- NRL - *Hyperspectral Modeling of Harmful Algal Blooms on the West Florida Shelf.* PI - W. P. Bissett (Florida Environmental Research Institute)
- NASA Project - Hyperspectral characterization of gelbstoff for application to remote sensing of carbon cycling in coastal regions. PIs - P.Coble and C.Castillo (USF)**
- NASA (EOS) - *High spectral resolution MODIS algorithms for ocean chlorophyll in case II waters.* PI - K. Carder
- NASA (SIMBIOS) - *Stray Light and Atmospheric Adjacency Effects for Large-FOV Ocean-Viewing Space Sensor.* PI - K. Carder
- NOAA - *Ocean Color Algorithm Evaluation for Remote Sensing of Coastal and Estuarine Waters.* PIs - R. Stumpf, P. Testor, J. Pennock, C. Tomas, R. Arnone, and K. Carder

REFERENCES

- Carder, K.L., R. Chen, and S. Hawes. Nitrogen-Depletion Temperatures for use in Parameterization of MODIS Algorithms. Abstract Submitted to 1998 AGU Ocean Sciences Meeting. 1997.
- Costello, D.K., K.L. Carder, and J.S. Patch. 1998a. Methods for Utilizing Hyperspectral In-situ Light Profiles in the Presence of Wave Focusing and the Absence of Above-water Measurements. EOS AGU/ASLO.
- Costello D.K., K. L. Carder, W. Hou, T.G. Peacock, and J.E. Ivey. 1998b. Hyperspectral Measurements of Upwelling Radiance During CoBOP: the Role of Bottom Albedo and Solar Stimulated Fluorescence. Ocean Optics XIV. Kailua-Kona.
- Costello D.K. and K. L. Carder. 1997. In situ optical data collected aboard unmanned underwater vehicles in coastal water. ASLO 97. Santa Fe.
- English D.C., R.G. Steward, D.K. Costello, and K.L. Carder. Characterizations and Techniques needed for the use of Spectral Radiometers to Collect Plane Irradiance Measurements. Abstract Submitted to 1998 AGU Ocean Sciences Meeting.
- Gregg, W.W. and K.L. Carder, 1990. A simple solar irradiance model for cloudless maritime atmospheres. *Limnol. Oceanogr.*, **35**(8), 1657-1675.
- Hou, W., K.L. Carder, and D.K. Costello. 1998a. Forward-Scattering Corrections for Transmissometers Using In Situ Particle-Size Measurements EOS AGU/ASLO.

- Hou, W. K.L. Carder, D.K. Costello, D.C. English, J.E. Ivey, and C. Mazel. 1998b. Database Structure of the CoBOP project with Visual Inspection via WWW. Ocean Optics XIV. Kailua-Kona.
- Hou, W. 1997. Characteristics of Large Particles and their Effects on the Submarine Light Field. Ph.D. dissertation, University of South Florida.
- Lee, Z.P., K.L. Carder, C. Mobley, R.G. Steward, and J. S. Patch. 1998a. Hyperspectral remote sensing for shallow waters: 1. A semi-analytical model. *Appl. Opt.*, 37(27), 6329 - 6338.
- Lee, Z.P., K.L. Carder, C. Mobley, R.G. Steward, and J. S. Patch. 1998b. Hyperspectral remote sensing for shallow waters: 2. Deriving Bottom Depths and Water Properties by Optimization. *Appl. Opt.*, submitted.
- Lee, Z. P., M. R. Zhang, K. L. Carder, and L. O. Hall. 1998d. A neural network approach to deriving optical properties and depths of shallow waters. Ocean Optics XIV. Kailua-Kona.
- Lee, Z.P., K.L. Carder, C. Mobley, R.G. Steward, and J.S. Patch. 1998e. Deriving Optical Properties and Water Depth of Shallow Waters by Inversion of a Remote-sensing Reflectance Model. EOS AGU/ASLO.
- Lee, Z.P., K.L. Carder, R.G. Stewart, T.G. Peacock, C.O. Davis, and J.L. Mueller. 1996. Remote-Sensing Reflectance and Inherent Optical Properties of Oceanic Waters Derived from Above-Water Measurements. Ocean Optics XIII. SPIE Vol. 2963, 160-166.
- Lee, Z.P., K.L. Carder, R.G. Stewart, T.G. Peacock, C.O. Davis and J.S. Patch. 1998c. An empirical algorithm for light absorption by ocean water based on color. *JGR*, accepted.
- Peacock, T.G., K.L. Carder, and R. Waterbury. 1997. A Dual-Path, Long-Path, Absorption, Spectrometer for Obtaining Absorption Coefficients of Dissolved Organic Matter (DOM). Presented at ASLO 1997, Santa Fe, New Mexico, 10-14 February 1997.
- Patch, J.S., K.L. Carder, Z.P. Lee, and R.G. Steward. Effects of Colored Dissolved Organic Matter (CDOM) and Pigment Packaging on Remote Sensing Reflectance Algorithms in the Southeastern Bering Sea. Abstract Submitted to 1998 AGU Ocean Sciences Meeting.
- Pratt, P., K.L. Carder, and D.K. Costello. 1997. Remote Sensing Reflectance Algorithms Developed to Correct Underwater Coral Imagery for the Effects of Optical Thickness to Assist in Benthic Classification. Presented at the Fourth International Conference on Remotes Sensing for Marine and Coastal Environments, Orlando, Florida, 17-19 March 1997.
- Reinersman, P.N., K.L. Carder, and F.I. Chen. 1998. Satellite-sensor calibration verification with the cloud-shadow method. *Appl. Opt.* 37(24); 5541-5549.
- Renadette, L.A., K.L. Carder, D.K. Costello, W. Hou, and D.C. English. 1998. Characterization of Bottom Albedo Using Landsat TM Imagery. EOS AGU/ASLO.
- Renadette, L.A., K.L. Carder, D.K. Costello, and W. Hou. 1997. AUV Data: Interpretation in Terms of Aircraft and Satellite Imagery. ASLO 1997, Santa Fe.
- Smith, R.C. and K. Baker, 1981. Optical properties of the clearest natural waters. *Appl. Opt.*, 20(2), 177-184.

World Wide Web site: <http://iceman.marine.usf.edu>