

Microsporidiosis

*Ann Cali, Ronald C. Neafie and
Peter M. Takvorian*

Introduction

Definition

Microsporidiosis is infection by eukaryotic unicellular protists of the phylum Microsporidia.¹ They are considered most closely related to the fungi,² but customarily are discussed among the protozoa. Several genera of microsporidia have been identified in human infections: *Nosema*,^{3,4,5} *Brachiola*,^{6,7} *Vittaforma*,⁸ *Pleistophora*,^{9,10} *Trachipleistophora*,¹¹ *Enterocytozoon*,^{12,13} *Encephalitozoon*,¹⁴⁻¹⁶ *Septata*,¹⁷ and *Anncaliia*.¹⁸ All microsporidia are obligate intracellular parasites, but pathologic changes vary with genus and species. In humans, infection may be latent or subclinical until the immune system is suppressed.¹⁹⁻²¹ Microsporidia are a significant opportunistic pathogen in patients with AIDS. Clinical features vary with the location and extent of infection. Microsporidia may infect virtually any tissue or organ of the body,²²⁻²⁵ including muscle,²⁶⁻²⁸ intestine,^{13,17,29,30} gallbladder,³¹ liver,^{14,31} kidneys,^{28,32,33} eyes,^{16,34-38} brain,^{28,39} lungs,⁴ skin, and nasal sinuses.^{33,40,41} Intestinal microsporidiosis is most common, occurring in 30% to 50% of AIDS patients with chronic diarrhea.⁴²⁻⁴⁴ Untreated intestinal, renal, cerebral, hepatic and disseminated infections are usually fatal.

Synonyms

The phylum Microsporidia^{1,45,46} was also known as phylum Microspora,^{47,48} formerly subphylum Microspora.⁴⁹

Encephalitozoon and *Nosema* were considered synonymous^{50,51} until 1970, when life cycle features demonstrated that they are separate genera.⁵² *Nosema corneum* is now *Vittaforma corneae*,⁸ a new genus having been established based on the ultrastructure of this organism's developmental stages and inconsistencies with any established genus. *Septata intestinalis* is also known as *Encephalitozoon intestinalis*.⁵³ *Nosema algerae* was reclassified as *Brachiola algerae*,⁵⁴ and is now *Anncaliia algerae*.¹⁸ *Nosema connori* was *Brachiola connori*,⁵⁴ and is now *Anncaliia connori*.¹⁸ Microsporidia with diagnostic spores but no identifiable developing stages are collectively called *Microsporidium*.^{47,48}

The terms nosematosis and encephalitozoonosis are occasionally used to describe microsporidiosis caused by *Nosema* sp or *Encephalitozoon* sp.

Epidemiology

Microsporidia are ubiquitous in animals (primarily insects, fish, and mammals) in developed and undeveloped, tropical and temperate regions of the world.^{28,47,48,55} Human infections have been reported in Africa,⁵⁶ Australia,⁵⁷ Singapore,⁵⁸ Europe,¹² and the United States.^{3,19,59,60} Between 1924 and 1985, less than a dozen cases of human microsporidiosis were reported.⁶¹ Since 1985, thousands of cases have been documented, primarily in AIDS patients⁶² but also in immunocompetent,^{63,64} immunosuppressed⁶⁵ and organ

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE JUN 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Microsporidiosis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Rutgers University, Dept. of Biological Sciences, 195 University Avenue, Boyden Hall, Newark, NJ, 07102				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADA545141. Chapter 14 from e-book, Topics on the Pathology of Protozoan and Invasive Arthropod Diseases.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 24	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			


**Figure 14.1**

Diagram of the internal structure of a microsporidian spore. The spore coat has an outer electron-dense region called the exospore (Ex) and an inner thicker electron-lucent region, the endospore (En). A unit membrane (P) separates the spore coat from the spore contents. The extrusion apparatus, anchoring disk (A), polar tubule (Pt), lamellar polaroplast (Lp), spongiform polaroplast (Sp) and tubular polaroplast (Tp), dominates the spore contents and is diagnostic for microsporidian identification. The posterior vacuole (Pv) is a membrane-bound vesicle which sometimes contains a “membrane whirl,” a “glomerular-like” structure, flocculent material, or some combination of these structures. The spore cytoplasm is dense and contains ribosomes (R) in a tightly coiled helical array. The nucleation may consist of a single nucleus or a pair of abutted nuclei, a diplokaryon (Dn). The size of the spore depends on the particular species and can vary from less than 1 μm to more than 10 μm. The number of polar tubule coils also varies from a few to 30 or more, again depending on the species observed.

transplant patients.^{66,67} With the aid of molecular technology, surveys of humans with no clinical signs of infection have been identified as positive for microsporidia.⁶³ Additionally, animal reservoir hosts have been identified for the four most common microsporidian organisms found in human infections.⁶⁸

Infectious Agents

Microsporidia are named for the small, resistant spore stage characteristic of this group. The phylum contains more than 170 genera and approximately 1300 species.^{28, 69} These diverse eukaryotes have relic mitochondria⁷⁰ no centrioles, contain 70S ribosomes, similar to prokaryotes, and have the smallest genome of any eukaryote thus far reported.⁷¹ Their ribosomal sequences are more similar to bacteria than to eukaryotes.⁷² However, their histochemically identifiable Golgi organelles indicate that they may be parasitically evolved

**Figure 14.2**

Electron micrograph of *Anncaliia* (*Brachiola*, *Nosema*) *connori* spore in adrenal gland showing the coiled polar filament (arrow) and two nuclei. x30,000

degenerate protists,^{73,74} microtubule gene data^{75,76} as well as several enzyme processes, place them closest to the fungi.²

Spores infecting humans are 1 to 5 μm in length and ~1 μm in width. All microsporidian spores contain a single long coiled structure called the polar filament, a unique structure attached at the anterior end by a large, mushroom-shaped anchoring disk.⁷⁷ Electron microscopy reveals this structure coiled around the single- or double-nucleated sporoplasm inside the thick, resistant and refractile spore coat (Figs 14.1 & 14.2). The straight, anterior portion of the polar filament is surrounded by a polaroplast which may be lamellar, tubular, or both. Light microscopy reveals a PAS-positive granule in the anterior end of the spore (Fig 14.3).

While the spore structures themselves are characteristic of microsporidia, the number of spores produced in sporogony, the manner in which they are produced, and the host-parasite interface (Table 14.1) vary among genera of microsporidia. Host-parasite interface may involve: 1) direct contact with host-cell cytoplasm, 2) indirect contact with host-cell cytoplasm by production of a parasite-secreted envelope (sporophorous vesicle, SPOV), 3) indirect contact by production of a parasite-induced, host-produced envelope (parasitophorous vacuole), or 4) indirect contact by a host-produced envelope (parasitophorous vacuole)⁵ and parasite-produced secretions.^{28,61,78} These characteristics, along with morphologic features of developmental stages, nucleation, site of infection, and serologic and molecular features, help to identify the genus and species.

Table 14.1. Interfacial Relationships of the Microsporidia

Type I. Direct contact	<p>The parasite plasmalemma is in direct contact with the host-cell cytoplasm (hyaloplasm).</p> <p>e.g. <i>Nosema</i>, <i>Brachiola</i>, <i>Anncaliia</i> and <i>Enterocytozoon</i>.</p>
Type II. Indirect contact by host produced isolation.	<p>A. Parasitophorous vacuole-host formed single membrane surrounding the developing parasite cell cluster. This is present during both the proliferative phase and the sporogonic phase, however the parasite relationship to it changes. The developing parasite cells maintain a very close relationship with this envelope until they develop the thickened sporont plasmalemma, then they appear loose within the vacuole.</p> <p>e.g. <i>Encephalitozoon cuniculi</i>.</p> <p>B. Host endoplasmic reticulum (ER) double membrane, surrounds parasite cells throughout development. In the proliferative phase the host ER double membranes follows the plasmalemma of the dividing cells so that no obvious vacuole is formed. In sporogony, the host ER does not divide with the sporonts and instead forms a double membraned parasitophorous vacuole surrounding the cluster of organisms formed in sporogony.</p> <p>e.g. <i>Vittaforma</i> sp, <i>Endoreticulatus</i> sp.</p>
Type III. Indirect contact by parasite produced isolation:	<p>A. Parasite secreted envelope, surrounds parasite cells throughout development. It becomes a sporophorous vesicle (SPOV) in sporogony when the parasite plasmalemma pulls away from the secreted envelope and then the plasmalemma thickens.</p> <p>e.g. <i>Pleistophora</i> sp.</p> <p>B. The parasite develops in direct contact with the host cell cytoplasm during early development but then a parasite formed membrane isolates the sporonts from host cytoplasmic contact.</p> <p>e.g. <i>Vairimorpha</i> sp.</p>
Type IV. Indirect contact by host and parasite produced isolation.	<p>A. Host ER closely abutts the parasite plasmalemma in merogony. Then the parasite produces “blisters” arising from the plasmalemma to form the interfacial envelope. Thus SPOV is formed in sporogony. It may also contain tubules.</p> <p>e.g. <i>Loma</i> sp, and <i>Glugea</i> sp.</p> <p>B. The host and parasite contribute to the formation of a thick interfacial envelope that surrounds all stages of parasite cells.</p> <p>e.g. <i>Trachipleistophora</i> sp.</p> <p>C. Host formed parasitophorous vacuole surrounds the parasite cluster and parasite secreted material surrounds each parasite cell inside the parasitophorous vacuole.</p> <p>e.g. <i>Septata</i> sp.</p>




Figure 14.3
Section of appendix showing *Anncaliia connori* spores in muscularis. The anterior end of the spore has a PAS positive granule (arrows).
PAS x1,260

Life Cycle and Transmission

The life cycles of microsporidia infecting humans have 3 phases: infective, proliferative, and sporogonic (Fig 14.4). The infective phase begins when a spore has developed fully in a host cell. Whether it remains within the host or passes to the environment through sputum, urine, or feces, the spore must receive the proper stimulus (pH or specific ions) in order to trigger the eversion of the polar filament resulting in the formation of a long polar tubule that emerges with sufficient speed/force to penetrate a new host cell (Fig 14.5).⁷⁹ A sporoplasm passes through the tubule and is deposited in a new host cell in less than a second, initiating a new infection.⁸⁰ Some spores are autoinfective and will evert their polar tubules, releasing sporoplasms within the same host, establishing additional sites of infection.

The proliferative phase begins when a sporoplasm grows and divides producing many organisms within the host-cell cytoplasm. The beginning of the sporogonic phase, sporogony, is usually signaled by one or more changes: secretions deposited on the surface membrane now called a thickened plasmalemma and/or formation of an isolating envelope (SPOV). Sporonts divide one or more times then become sporoblasts, which metamorphose into spores.

Environmental sources of human exposure to microsporidia are not known definitively, but since many human infections are intestinal and cause severe diarrhea, fecal-oral transmission is likely. In disseminating infections of the renal system, spores are passed in urine, providing another possible source of exposure. A strong correlation between soil exposure and microsporidial keratitis in HIV negative patients has been reported.⁵⁸ Municipal drinking water chlorination does not inactivate spores, so this is a possible reservoir for microsporidia as well as spores on food con-




Figure 14.4
Life cycle. Diagrammatic illustration of a typical developmental cycle of the microsporidia. The three regions represent the three phases of a microsporidian life cycle. Phase I is the Infective/Environmental phase, the extracellular phase of the cycle. It contains the mature spores in the environment. Under appropriate conditions, the spore is activated (e.g. if the spore is ingested by an appropriate host, it is activated by the gut environment) and triggered to evert its polar filament which becomes a hollow tubule. If the polar tubule pierces a susceptible host cell and injects the sporoplasm into it, phase II begins. Phase II is the proliferative phase. This is the first phase of intracellular development. During the proliferative part of the microsporidian life cycle, organisms are usually in direct contact with the host cell cytoplasm and they increase in number. The transition to phase III, the sporogonic phase represents the organisms' commitment to spore formation. In many life cycles this is morphologically indicated by parasite secretions through the plasmalemma producing the "thickened" membrane. The number of cell divisions that follow varies, depending on the genus in question, and results in spore production.



Figure 14.5
Anncaliia algerae. Spore in germination medium viewed by phase contrast microscopy. Sporoplasm (arrowhead) still connected to its polar tubule (PT) and spore shell (SS).

sumed raw (Fig 14.6).^{81,82} The use of molecular tools has also led to the identification of human-infecting microsporidia in fecal samples from cats, dogs, primates, cattle, and birds.⁸³

Families: Nosematidae & Tublinosematidae

There are over 100 species of *Nosema*, most of which are parasitic in insects. A few *Nosema* species have been described from human ocular infections,^{22,58} *N. oculorum*,




Figure 14.6
The food-water connection between microsporidia and human infection.

Nosema sp.^{29,84} and *N. corneum*.^{13,17,29,30,85} The genus *Vittaforma* was created for the organism originally named *Nosema corneum*, after Silveira and Canning suggested that the diplokaryotic arrangement of nuclei was the only characteristic indicative of the genus *Nosema*. *Vittaforma corneae*,⁸ as it is now known, is polysporoblastic and thus will probably be omitted from family Nosematidae as new molecular sequence data become known. The *Nosema* sp of Curry⁸⁴ is accepted in the *Nosema* genus because of the presence and/or absence of features eliminating other possibilities. *Nosema* sp of Ashton³⁴ lacked definitive generic information and was moved to the “group” Microsporidium with a species designation of *M. celonensis*⁵⁵ as was *Nosema* sp of Pinnolis⁴ to *M. africanum*.⁵⁵ More recently Loh conducted a four year study that confirmed 124 cases of microsporidial keratitis. While the pathology and therapy were described, the generic designations of the organisms were not demonstrated.⁵⁸

Three *Brachiola* spp have been described from human infections and recently moved to a new family, based on molecular and morphologic features.¹⁸

The new family, Tublinosematidae, was established in 2005 for organisms with many similarities to Nosematidae but with different morphological and mo-

lecular features.⁸⁶ While the family was established for microsporidia infecting insects, the human as well as the insect infecting *Brachiola* species were subsequently placed here because of the sequence similarity of *B. algerae* and *Anncaliia meligethi*.^{18,87} These organisms possess a thickened plasmalemma throughout development and an elaborate surface tubular network that extends into the host cytoplasm. The genus *Anncaliia* was added to this family (2006) and its sequence data so closely resembled *Brachiola algerae* that the genus *Brachiola* became a synonym to the previously described genus *Anncaliia*.¹⁸ However, we retain the genus *Brachiola* in this family, for its type species, *vesicularum*, because of its unique protoplasmic branches that extend into the host cytoplasm, similar to fungal hyphae, not demonstrated in any other microsporidia.⁶

Morphologic Description

Nosema and *Anncaliia* (*Brachiola*) spores are approximately 4 µm in length and have paired abutted nuclei (diplokarya) in every stage of development. The parasite plasmalemma is in direct contact with the host-cell cytoplasm. If there are several nuclear divisions before cytokinesis




Figure 14.7
Microsporidia infection in patient's eye. Central stroma of cornea is necrotic and surrounded by acute inflammatory cells. H&E x210

takes place, proliferative cells may become elongated. After many cell divisions, large clusters of individual diplokaryotic cells develop. Sporonts produce two sporoblast cells that develop into two spores, all in direct contact with the host-cell cytoplasm.^{28,52}

Nosema sp form a thickened plasmalemma only in the sporogonic phase. However, all developmental stages of *Anncaliia* (*Brachiola*) sp form a thickened plasmalemma; these stages additionally form elaborate vesiculotubular appendages on the plasmalemmal surface.⁸¹ *Brachiola vesicularum* uniquely forms protoplasmic extensions on elongated cells during the proliferative phase and produces spores that contain one to three rows (usually two) of polar filament coils.⁶

Clinical and Pathologic Features

Two *Nosema* infections, subsequently changed to *Microsporidium celonensis* and *M. africanum* have been reported in HIV-negative patients with infections of the corneal stroma that led to perforation or blindness, followed by enucleation.^{3,4,34,35,85,88} Histologic examination of corneal sections from one patient revealed organisms consistent with microsporidia (Fig 14.7). The central stroma was necrotic and surrounded by acute inflammatory cells. Immediately above Descemet's membrane were abundant refractile spores measuring 3.5 μ m by 1.5 μ m, some free and some in macrophages (Fig 14.8). No organisms were found in the exudate. In both infections, the genus *Nosema* was suggested but no specific identification could be made.^{4,34,89} Two other microsporidia-associated infections of the corneal stroma were reported in the early 1990s in otherwise healthy individuals.^{3,88} The first patient was a 39-year-old man from Ohio with a corneal ulcer. Examination of Gram-stained biopsy tissue by electron microscopy revealed microsporidia. Spores were 5 μ m by 3 μ m, binucleated, and




Figure 14.8
Refractile microsporidia (arrow) at edge of cornea next to the acute inflammatory cells. H&E x400




Figure 14.9
Nosema ocularum. High-power electron micrograph of spore showing 9–12 polar tube cross sections. Bar = 1.0 μ m

contained 9 to 12 polar filament coils (Fig 14.9). The parasite was subsequently named *Nosema ocularum*.^{3,35} The second patient was a 45-year-old man from South Carolina with no history of prior trauma to the infected eye.⁸⁸ Biopsy of the stroma revealed a microsporidium parasite that was isolated and grown in cell culture. The spores were 3.7 μ m by 1 μ m and contained 5 to 6 polar filament coils. This organism, originally named *Nosema corneum*,^{85,88} is now known as *Vittaforma corneae*.⁸

From 1990 to the present, hundreds of microsporidian keratoconjunctivitis cases have been reported, in both HIV negative and positive patients, however, only a few reports have identified the organisms beyond being microsporidia. The pathologic changes, however, have been described from the survey of 124 microsporidia positive patients (134 eyes).⁵⁸ “Common features were follicular papillary conjunctivitis and coarse punctuate epithelial lesions in three patterns—diffuse, peripheral, and paracentral—evolving




Figure 14.10

Anncaliia connori (4 μ m x 2 μ m) in bowel of 4-month-old child: Gram-positive microsporidia in smooth muscle of ileum, Brown and Hopps (B&H) x2180

into nummular keratitis before resolution”...(99 percent of the cases were resolved with topical fluoroquinolone monotherapy). “New clinical features were diffuse endotheliitis (19.4%) with corneal edema and limbitis.”⁵⁸

The first well documented human infection with a microsporidium was in a 4-month-old male with thymic aplasia, severe diarrhea, and malabsorption (Fig 14.10).^{7,89,90} At autopsy, *Anncaliia* (*Brachiola*, *Nosema*) *connori* spores measuring 4 μ m by 2 μ m were found in the small and large bowel; no other infectious agent was discovered.⁷ Infection had disseminated to the lungs, stomach, kidneys, adrenal glands, myocardium, liver, and diaphragm (Figs 14.11 to 14.18).⁸⁹

A *Brachiola* sp infection was reported in a 31-year-old male AIDS patient who had pain and progressive muscular weakness of the lower extremities of five months duration. It was named *B. vesicularum*.⁶ The tissue biopsy contained spores measuring 2.5 μ m to 2.9 μ m by 1.9 μ m to 2 μ m. Intramuscular infection and cytolysis were observed by light microscopy. Electron microscopy revealed diplokaryotic microsporidia in all stages of development (Figs 14.19 & 14.20). An elaborate array of vesiculotubular appendages emanated from the plasmalemmal surface of most stages (Fig 14.21), and spores presented with variable polar filament arrangements (Fig 14.22). Additionally, protoplasmic extensions develop on some proliferative stages (Figs 14.23 & 14.24) that are unique to this microsporidium. There was loss of muscle striation in surrounding areas.

Anncaliia (*Brachiola*, *Nosema*) *algerae*, a parasite of mosquitoes, has been cultured in mammalian cells at 29° to 37°C^{54,91} and documented by PCR




Figure 14.11

Silvered *Anncaliia connori* in wall of ileum of patient in Fig 14.10. 90 minute Grocott methenamine silver (GMS) x1300




Figure 14.12

Anncaliia connori (4 μ m x 2 μ m) in kidney of patient in Figure 14.10. H&E x1800




Figure 14.13

Anncaliia connori in adrenal gland of patient in Figure 14.10. H&E x1475




Figure 14.14

Anncaliia connori in adrenal gland of patient in Figure 14.10. Warthin-Starry (WS) x1860




Figure 14.15

Anncaliia connori in myocardium of patient in Figure 14.10. H&E x450




Figure 14.16

Anncaliia connori in myocardium of patient in Figure 14.10. WS x2260




Figure 14.17

Anncaliia connori in myocardium of patient in Figure 14.10. Ziehl-Neelsen (ZN) x1440




Figure 14.18

Anncaliia connori in liver of patient in Figure 14.10. Giemsa x1910

techniques in ocular and dermal infections of immunocompetent patients.³⁸ It has also been documented in deep tissue infections in skeletal muscle from a patient with rheumatoid arthritis being treated with immunosuppressive drugs (Figs 14.24, 14.25a & 14.25b). The infection progressed until the patient died.^{65,92} Infection in both epithelial and connective tissue cells of the false vocal chord area (Figs 14.26 a-e & 14.27) was documented in a terminal cancer patient.⁹³

Family: Pleistophoridae

This family was named from the fish parasite *Pleistophora typicalis* (Gurley 1893). Since that time, approximately two dozen species had been described, all of them in fish⁵⁵ until 1985 when a human muscle infection with *Pleistophora* sp was identified.⁹ This infection from an immunodeficient, but HIV negative male from the USA, was subsequently studied from the original tissue blocks and named *Pleistophora ronneaei*.^{10,92} Two additional *Pleistophora* sp cases were identified in HIV+ males, one in Australia⁹⁴ and one from Spain.⁹⁵ It was not until 1996, that yet another genus, *Trachipleistophora*, was established for human infection from this family of microsporidia.¹¹ The first species, *T. hominis* was described from muscle and eye infections in an AIDS patient from Australia.^{11,27} A second species, *T. anthropoptera*,⁹⁶ was described from two AIDS patients.⁹⁷ This species disseminated to multiple organs including the brain, heart, kidney, pancreas, thyroid, parathyroid, liver, bone marrow, lymph nodes, and spleen. The most heavily infected cells were epithelia, cardiac myocytes, and astrocytes.²⁴

Morphologic Description

A thick, parasite-secreted, envelope is produced on the surface of proliferative cells. It separates from the surface and becomes an SPOV in sporogony. *Pleistophora ronneaei* and *Trachipleistophora* sp develop within the host-cell cytoplasm in this type of parasite produced vesicle. Nuclei are single, even when there are many nuclei (not diplokaryotic) within a plasmodial proliferative cell. In *P. ronneaei*, proliferative and sporogonic plasmodia divide by multiple fragmentation of large cells. *Pleistophora* is polysporous; 16 to over 100




Figure 14.19

Brachiola vesicularum. Electron microscopic overview of an infected muscle cell. Developmental stages of the parasite appear as clusters surrounded by the striated muscle cell filaments (M). The electron-density variations of the parasite cells are readily observable, represent different stages of parasite development. The most dense stages are sporoblasts (Sb) and spores (S). Diplokaryotic nucleation (Nu) is apparent in the majority of parasitic cells. Some dividing cells contain two diplokarya (arrows) x4,300.




Figure 14.20

Brachiola vesicularum. A proliferative cell in advanced stage of cytokinesis (broad arrowheads), the spindle plaques (arrows) are still present. The cytoplasm of this proliferative cell is more densely granular and endoplasmic reticulum (er) is more abundant indicating that this cell is also advanced in the parasite developmental cycle. Note the presence of myofilaments (F) and of vesiculotubular material (T).






Figure 14.21


Brachiola vesicularum. A late proliferative/sporont cell completing cytokinesis. The plasmalemmal invagination is almost complete and the diplokaryotic nucleus (Nu) pairs appear to be in interphase. Note the presence of vesiculotubular material (T).

**Figure 14.22 a,b,c**

Brachiola vesicularum a. Mature spore containing a fully developed electron-lucent endospore coat (average thickness 90 nm to 100 nm). The exospore (62 nm) surface has several vesiculotubular structures (T) on it. Note the presence of nine polar filament (Pf) cross sections arranged in two rows. Ribosomes (R) appear in a spiral-like array forming rows around the nuclear area (Nu). x 41,600; b. Spore containing ten polar filament (Pf) cross sections clustered into three rows. (endospore coat = 82 nm, exospore coat = 63 nm) x 36,800; c. Section through a spore revealing the presence of the anterior anchoring disc complex (A) of the polar filament (Pf) and the manubroid (Mpf) portion of it. The cross sections of the polar filament coils arranged in a single row is visible. Multiple rows of ribosomes (R) are also present. Note the presence of vesiculotubular material (T). x26,000.

**Figure 14.23 a,b**


Brachiola vesicularum. Proliferative cells with protoplasmic extensions. a. A very elongated parasite cell (7.1 μ m in length and varies between 1.0 μ m and 1.2 μ m wide) with a vesiculotubular "cap" complex (TvC) at one end and a scalloped thick plasmalemmal surface which contains several channels (arrowheads). Additionally, this cell possesses several protoplasmic extensions (E) of varying lengths, projecting from the cell surface (broad arrows). At the ends of these protoplasmic extensions are vesiculotubular (T) structures with the electron-dense fibrous coating, similar to those previously illustrated on the typical proliferative cells. Note the presence of vesiculotubular (T) structures and myofilaments (F) in the host cytoplasm. x19,300. b. A portion of a parasite cell protoplasmic extension complex measuring 4.80 μ m in length and between 0.5 μ m to less than 0.3 μ m in width. A number of branches of varying lengths, have formed from the cell surface and project into the host cytoplasm. These projections also end in vesiculotubular (T) structures. The surface of the protoplasmic extensions also have some scalloping and shallow indentations present. In the lower third of the figure is a section of a parasite (P) cell with several vesiculotubular (T) structures attached to the cell surface. x 27,000.

**Figure 14.24**



Annaliia algerae in muscle after the patient presented with myositis and muscle pain showing multiple organisms in the muscle fibers (arrows) with associated cell lysis but little or no inflammation. H&E x100.

spores may be produced from sporonts encased in the parasite-secreted SPOV envelope (Figs 14.28 & 14.29).

In the two species of *Trachipleistophora*, *T. hominis*^{11,98} and *T. anthropophthera*⁹⁶, proliferative cells have 2 to 4 nuclei and divide by binary fission. In the sporogony phase, division into sporoblasts is effected by repeated binary fission, producing 2 to 32 spores within the SPOVs, no plasmodial stages are produced (Figs 14.30 to 14.34). Spores of *T. hominis* are approximately 4 by 2 μ m (Fig 14.35). *Trachipleistophora anthropophthera* is dimorphic,

**Figure 14.25 a,b**

Annaliia algerae in muscle from patient in Figure 14.24. a. A diplokaryon, a thickened plasmalemma (arrows), and vesiculotubular extensions are evident in proliferative forms. Nu denotes nucleus. x14,000; b. A mature *A. algerae* spore with a single row of nine polar filaments (arrowheads) in cross section. Mature spores in the biopsy specimen had only single rows of 8 to 12 polar filaments in cross section. x18,300








Figure 14.28

Pleistophora ronneafiei. Developmental stages (A,B,C,D) in human skeletal muscle. Electron micrograph demonstrates the location of the infection, abutting bundles of actin and myosin filaments arranged as normal, functional contractile unit of host muscle (HM) and containing a host cell nucleus (HN). The early stages of parasitic development are each surrounded by parasite-secreted dense material. The vesicle wall is most elaborate in proliferative development as illustrated between proliferative cells A and B. The proliferative cells A and B are probably sister cells, as in early proliferative development, the cells divide with the secretions. In sporogony, multiple parasite cells (cluster D) may be found within each sporophorous vesicle, which becomes more homogeneously dense in sporogony making the elongated oval spores obvious.



a



b

Figure 14.29 a,b

Pleistophora ronneafiei.

Sporoblasts of *Pleistophora* in human skeletal muscle. After sporogonial plasmotomy is completed, the sporophorous vesicles are filled with many uninucleated cells, the sporoblasts. a. Sporophorous vesicle containing early sporoblasts (SB) abutted to a proliferative cell (P) and the lack of projections on the vesicle walls (straight arrows) where it abuts the host cytoplasm (HM). b. Late sporoblast or early spore contains a single nucleus (N), the developing polar filament (PF), the Golgi (G), and the beginning of spore wall (SW) thickening.

in sporogony, 2 types of SPOVs and spores are formed (Figs 14.36). One type of SPOV contains thick-walled spores (approximately 8, measuring 3.7 by 2.0 μm), each containing 9 polar filament coils. The other type contains 2 thin-walled spores with 3 to 5 polar filament coils and measuring 2.2 μm to 5 μm by 1.8 μm to 2.0 μm .⁹⁶

Clinical and Pathologic Features

Pleistophora sp primarily infect the muscles of marine and freshwater fish, but three *Pleistophora* infections in human skeletal muscle have been reported.^{9,26,94,95} The first, in 1985, was in a 20-year-old immunocompromised man from Florida.^{9,99,100} Over a 7-month period, the patient experienced

progressive wasting and generalized muscle weakness leading to contractures. Biopsies of skeletal muscle contained large clusters of organisms visible by light microscopy with H&E (Fig 14.37), acid-fast (Fig 14.38) and Giemsa stains (Fig 14.39). Electron microscopy of the same biopsy tissue revealed the diagnostic features of *Pleistophora*.⁹ Developmental stages from this case were subsequently described (Fig 14.28 & 14.29) and the parasite named *P. ronneafiei*.¹⁰ The two additional cases of myositis caused by *Pleistophora* in patients with AIDS, in 1993 and in 1996 have not been compared.^{94,95}

Trachipleistophora hominis was described from an AIDS patient in Australia.^{11,27} The infection was primarily (Fig 14.30) muscular¹⁰¹ but organisms were also found in corne-




Figure 14.30
Trachipleistophora hominis. Spores and larger brown spore precursors forming masses within skeletal muscle fibers. Free spores are visible in the adjacent connective tissues (arrow), and discrete early aggregates of spore precursors are visible in fibers (arrowhead). Warthin-Starry x 400.




Figure 14.31
Trachipleistophora hominis. Corneal scraping showing sporophorous vesicles containing spores and spore precursors in epithelial cells (arrows). Note the dispersed spores in the background showing the posterior vacuole (arrowhead). Modified trichrome x1000.




Figure 14.32
Trachipleistophora hominis. Multiple sporophorous vesicles closely abutting within a skeletal muscle cell (open arrow) with a single meront (arrow). Note the skeletal muscle nucleus (arrowhead). x1,600.




Figure 14.33
Trachipleistophora hominis. Two adjacent binucleate meronts with thick outer coats (arrow). x3,000.




Figure 14.34
Trachipleistophora hominis. Sporonts undergoing division in a skeletal muscle cell (arrows). x3,000.





Figure 14.35
Trachipleistophora hominis. Spore with anchoring disc (arrow), straight portion of polar tube (open short arrow) extending through the polaroplast (open arrow) posterior vacuole, and tangential sections through coils of tube (arrowhead). x12,000.



← **Figure 14.36**
Trachipleistophora anthropophthera. Type I mature spore's polar filament has thicker and inward displaced thinner posterior coils. Bar = 0.5 μm.




Figure 14.37
Pleistophora ronneaei in skeletal muscle showing large clusters of organisms: H&E x560




Figure 14.38
Pleistophora ronneaei in skeletal muscle showing large clusters of organisms. ZN x490




Figure 14.39
Pleistophora ronneaei in skeletal muscle showing large clusters of organisms. Giemsa x520





Figure 14.40
(A) Gross picture of a brain from a patient with *Trachipleistophora anthropophthera* infection demonstrating multiple necrotic lesions in the gray matter. (B) Light microscopic section from the brain shown in panel A, demonstrating spores in astrocytes and other cells. GMS x400

al epithelium (Fig 14.31) and nasopharyngeal washings. In experimentally inoculated athymic mice, infection spread to tissue of the bladder and large intestine.¹¹

Trachipleistophora anthropophthera was identified in 2 AIDS patients in the United States. Both had disseminated infections involving the heart, kidneys, and brain (Figs 14.40a & 14.40b), manifesting in seizures and impaired cognition, suggestive of toxoplasmosis.^{24,28,96,97}

Family: Enterocytozoonidae

Enterocytozoon was the first genus of microsporidia created for a human infection.^{12,30} It has subsequently been found in pigs and cattle.¹⁰² This microsporidium has many unique developmental features¹³ and researchers have had only limited success at growing them in culture.

Morphologic Description

Enterocytozoon organisms develop in direct contact with the host-cell cytoplasm. As nuclei multiply, plasmodia enlarge. *Enterocytozoon* forms two unique organelles: electron-lucent inclusions and electron-dense disks. They both form in a multinucleate plasmodial cell in direct contact with the host cell cytoplasm (Fig 14.41).¹³ Electron-lucent inclusions appear very early in the development of the proliferative plasmodia, increase in size and number as the plasmodia grow, and are present throughout the life cycle. Electron-dense disks form at the surface of the electron-lucent inclusions, often in small stacks similar to a stack

of red blood cells. Plasmodial cells containing these disks have many rounded nuclei. The disks eventually fuse and form the spores' polar filament. Finding several polar filaments within a multinucleate plasmodium is diagnostic for *Enterocytozoon* (Fig 14.42).¹³ The presence of these organelles and the development of multiple polar tubules within a multinucleate parasite cell are all unique features of the developmental cycle of *Enterocytozoon*. The plasmodium divides by multiple fission; producing a dozen or more sporoblasts which mature into spores, all in direct contact with the host-cell cytoplasm. Spores are 1.3 μm by 0.8 μm and contain a single nucleated sporoplasm surrounded by approximately 6 polar tubule coils, arranged in a double row (Fig 14.43).^{13,24}

Clinical and Pathologic Features

Enterocytozoon bienersi is one of the most frequently reported microsporidial infection in humans. Incidence in AIDS patients is approximately 7% in Africa,⁵⁶ 20% to 30% in the United States and Australia,^{43,57,59} and 50% in France.⁴² The organism infects the apical portion of enterocytes of the small bowel (Figs 14.44 & 14.45). Endoscopically, infection appears as a slight flattening of the epithelium.¹⁰³ Histologically, the only visible pathologic feature is villus atrophy due to more rapid desquamation of infected enterocytes.^{12,13,24,30,43,59} The parasite can disseminate to the epithelial lining of the common bile duct, gallbladder epithelium, and biliary and respiratory tracts.^{19,31,59,104} *Enterocytozoon bienersi* most commonly infects male AIDS patients caus-



Figure 14.41
Enterocytozoon bienewsi. Sporogonial plasmodium containing at least 12 nuclei (N) in a single plane of section. The round dense nuclei are each associated with electron dense disc complexes (arrows) and electron lucent inclusions (*). Electron dense discs fuse into arcs forming polar tube coils. Despite the advanced maturation and organelle separation associated with each nucleus, there is not yet any evidence of cytokinesis or plasmalemmal thickening. x25,600. Bar = 1 μ m.




Figure 14.42
Enterocytozoon bienewsi. Late sporogonial plasmodium with advanced stages of polar tubule formation. Fused electron dense discs are seen in coiled (single arrow), stacked (double arrows) and cross sectional (triple arrows) profiles throughout the cytoplasm. Anterior anchoring discs (A) and associated polaroplast membranes appear at this stage even though individual sporoblast membranes have not yet developed. The electron lucent inclusions are seen in elongated (E) and cross section (C) views. x33,231. Bar = 1 μ m. Insert: Connection and arrangement of various structures developing in the plasmodium. Umbrella-shaped anchoring disc (A) and associated polaroplast membranes (P) attached to the manubroid portion of the polar tube (M) which connects with arcs formed by the coiled region of the developing polar tube. This complex of polar tube and associated structures surrounds a single nucleus (N). x28,000. Bar = 1 μ m.

ing malabsorption resulting in chronic diarrhea,^{12,13,19,30,105,106} and has subsequently been associated with AIDS-related sclerosing cholangitis.^{31,104} As awareness of microsporidial infection has grown, *E. bienewsi* has also been reported

in HIV-negative³³ and female HIV-positive¹⁰⁷ patients with and without diarrhea,¹⁰⁸ and in tracheal,³² bronchial,¹⁰⁹ and nasal^{110,111} epithelium. *Enterocytozoon bienewsi* can also be diagnosed from stool samples. The spores can be visualized by modified trichrome¹¹² (Figs 14.46).




Figure 14.43
Spore of *Enterocytozoon bienewsi* demonstrating the characteristic six turns of the polar tubule, which are organized into two tiers of three turns each and which are out of register by 45°. x83,000

Family: *Encephalitozoonidae*

Encephalitozoon cuniculi was first discovered in 1924 in rabbits.¹¹³ It was placed in the microsporidia in 1964 as a junior synonym in the genus *Nosema*.⁵¹ In 1971, it was reclassified as a genus of *Microsporidia*⁵² and described from rabbits, mice and hamsters.¹¹⁴ The family was established in 1989.¹¹⁵ This parasite has subsequently been reported from over 30 different mammalian hosts⁶¹ and the first human infection with *E. cuniculi* was reported in 1987.¹⁴

From 1989 to 1991, six cases of microsporidian keratoconjunctivitis were reported in patients with AIDS, four from New York, one from Texas, and one from Ohio.^{3, 16, 35, 36, 116, 117} All had conjunctivitis, blurred vision, and photophobia. By 1999 over 20 cases were characterized, reported and reviewed.²² Organisms were observed in corneal epithelial cell scrapings examined by light and electron microscopy.^{16, 118, 119} The organisms were morphologically similar to *E. cuniculi*, but a clearly defined parasitophorous vacuole surrounding the organisms was not always visible.¹⁶ Didier et al. reported that the organism was morphologically the same, but serologically different from *E. cuniculi* and named it *Encephalitozoon hellem*.¹⁵

First reported in 1991, *Septata* was the second microsporidial genus created for a human infection.^{17, 29} It was placed in this family as a new genus, based on similarity of some of its morphological features with the type species while presenting some unique features including intestinal infection. Infection with *S. intestinalis* has been reported in the United




Figure 14.44
Enterocytozoon bienewsi infects the apical portion of enterocytes of the duodenum: Gram-positive spores (1.5 µm x 1.0 µm) B&H x1010




Figure 14.45
Enterocytozoon bienewsi infects the apical portion of enterocytes of the duodenum containing acid-fast spores ZN x800




Figure 14.46
Enterocytozoon bienewsi in feces. Modified trichrome x700

States,^{17, 29, 59, 120} Europe,^{23, 59, 60} and Australia.^{57, 59, 121, 122} Subsequent molecular data suggested a closer relationship between the two genera and the species: *S. intestinalis* was moved into the genus *Encephalitozoon*.⁵³




Figure 14.47

Encephalitozoon hellem. Epithelial cells with large vacuolated area containing oval-shaped organisms and amorphous material. The oval bodies vary in appearance, reflecting different stages of development. X9,100




Figure 14.48

Encephalitozoon hellem. Developing stages of *E. hellem* growing in direct contact with host cell cytoplasm of RK-13 cell line. Four early sporonts (SP) with "thickening membranes" (arrows) are present and the elongated sporont contains at least two nuclei (Nu). Note host rough endoplasmic reticulum (er) in close proximity to sporonts. x23,000

Morphologic Description

The genus *Encephalitozoon* is characterized by a phagosome-like parasitophorous vacuole that surrounds developing parasites and isolates them from the host-cell cytoplasm. Developing stages (Fig 14.47) may contain one or more nuclei but the nuclei are not attached to each other in diplokaryotic arrangement. Multinucleate cells are long and narrow, not plasmodial (Fig 14.48).¹¹⁴ Proliferative cells usually abut the vacuole membrane and break free of it in sporogony. Sporonts form a thickened plasmalemma on individual cells. Within the parasitophorous vacuole, each sporont elongates, divides, and produces spores. Spores are 1 μm to 1.5 μm by 0.5 μm and contain a sporoplasm with a single nucleus and approximately 6 polar tubule coils (Fig 14.49), arranged in a single row.^{114,123} *Encephalitozoon hellem* is morphologically the same as *E. cuniculi* except that the parasitophorous vacuole is not always present.¹²⁴

Encephalitozoon (Septata) intestinalis is characterized by parasite-secreted material surrounding the developing stages and spores (Fig 14.50) inside the parasitophorous vacuole. Proliferative and sporogonic stages have 1 to 4 nuclei. Cells are rounded when single-nucleate and elongated when they contain 2 to 4 nuclei. In sporogony, the plasmalemma thickens and elongated sporonts are produced. Each sporont di-

vides, producing up to 4 single-nucleate sporoblasts. After a final cell division, sporoblasts develop the polar filament complex and metamorphose into spores. *Encephalitozoon intestinalis* cells develop in clusters. During early development, these clusters appear tightly packed. During sporogony and spore formation, some cells condense, leaving a space between individual developing forms. When this happens, the parasite-secreted fibrillar matrix surrounding the different stages is apparent. Early and late forms develop asynchronously, with parasite secretions surrounding individual cells and a parasitophorous vacuole surrounding the cluster (Fig 14.50). Spores are 2 μm by 1.2 μm, with a single nucleus and 4 to 7 (usually 5) polar tubule coils arranged in a single row.^{17,29}

Clinical and Pathologic Features

Encephalitozoon cuniculi is probably the most studied microsporidian. Primarily a parasite of animals, it has been reported in over 30 mammalian hosts.^{55,61,114} In 1984, Bergquist et al reported serologic evidence of *E. cuniculi* in AIDS patients with a history of travel to the tropics.^{125,126} In 1987, Terada et al. found *E. cuniculi* organisms in the liver of an AIDS patient with hepatitis.^{14,127}




Figure 14.49

Encephalitozoon hellem. Two spores are cut in longitudinal (upper) and cross section. Both have a thick spore coat including thin dense exospores (EX), thick electron-lucent endospore (EN), and the thin inner spore coat membrane (M). In the organism above, a single, large, rounded nucleus (N) is centrally located. Peripheral to and on each side of the nucleus are seven to eight round dense bodies that are cross sections of the polar tubule (PT) coils. In the organism below, the polar tubule is coiled parallel to the plane of the section and appears as a single dense ring situated adjacent to the inner aspect of the spore coat. x32,500.



Figure 14.50

Encephalitozoon (Septata) intestinalis. Deposition of material results in a uniformly thick plasmodium surrounding the sporont cells. These sporonts continue to secrete the fibrillar lamina. The sporont (ST) is a tetranucleate (n) elongated cell in the process of cytokinesis (arrowhead). This cluster of parasite cells also contains many mature electron-dense spores, proliferative cells (P), and a dense fibrillar lamina separating the individual parasite cells. x9,000

Since then, *E. cuniculi* has been demonstrated in peritoneal, cerebral, and disseminated infections in AIDS patients (Figs 14.51 & 14.52).^{39,128-130}

In cell culture, *E. hellem* may develop with or without a surrounding vacuole (Fig 14.48). *Encephalitozoon hellem*, originally reported from eye infections, has since been associated with disseminated infection and infection of the sinuses,⁴⁰ nasal tissue,¹³¹ tongue, respiratory system (Figs 14.53 & 14.54),^{132,133} kidneys (Fig 14.55) and male genital tract (Figs 14.56 & 14.57).^{109,134,135} Schwartz et al¹³² used immunofluorescence and H&E to demonstrate the organisms in tissues (Figs 14.58).

Encephalitozoon intestinalis is an intestinal epithelial cell parasite, causing diarrhea, malabsorption, and wasting. Because it can also infect macrophage,¹²⁰ fibroblastic, and endothelial cells, it can infect the lamina propria below the enterocytes and disseminate to other parts of the body (Figs 14.59 to 14.61).¹⁰³ Infections have been reported in the liver, renal system, colon,¹²¹ gallbladder, lungs, sinuses,^{121,136} and conjunctiva.^{17,20, 32,37,57,137} It has been misdiagnosed as *En-*

terocytozoon bienewsi in the intestine and as *Encephalitozoon hellem* in the eye, sinuses, and urine.^{59,138}

Although size, nucleation, and number of polar tubule coils are similar in *Encephalitozoon intestinalis* and *Enterocytozoon bienewsi*, the organisms can be distinguished by the arrangement of polar filaments in the spore stage. In *Enterocytozoon bienewsi*, the polar filament forms a double row of coils; in *Encephalitozoon intestinalis*, they form a single row. Another diagnostic feature is observable in tissue sections. Spores and developing stages of *Encephalitozoon intestinalis* are within individual chambers inside a vacuole in the host-cell cytoplasm. *Enterocytozoon bienewsi* development is in direct contact with the host-cell cytoplasm (Fig 14.53).^{17,20,120}

Diagnosis

Identifying the spore stage containing the polar filament is diagnostic for microsporidia. Spores can be seen by light microscopy using stains¹³⁹ such as Giemsa,^{100,140-142} Ziehl-




Figure 14.51
Encephalitozoon cuniculi (arrows) spores in bronchiole epithelium. B&B x650.




Figure 14.52
Encephalitozoon cuniculi acid-fast spores (arrows) in bronchiole epithelium. (Not all spores are acid-fast) ZN x875




Figure 14.53
Encephalitozoon hellem spores (arrows) in the trachea: Some acid-fast spores ZN x620




Figure 14.54
Encephalitozoon hellem spores in the trachea: Silvered spores in mucosa, GMS x800

Neelsen,⁸⁹ Brown & Brenn,¹⁰³ Brown & Hopps, Weber's trichrome,¹¹² Warthin-Starry,^{57,122,143} and fluorescence.^{60, 119,132,144} Hematoxylin-eosin is not particularly useful for identifying microsporidia. Spores are birefringent and can be seen with polarized light (Fig 14.8). A PAS-positive granule in the anterior end of the spore is diagnostic for larger microsporidia such as *Nosema*, *Anncaliia* and *Pleistophora* (Fig 14.3), but trying to identify the granule in the smaller, more common human-infecting microsporidia, *Encephalitozoon* and *Enterocytozoon*, is impractical.

Touch preps and smears can be made from biopsy materials, eye scrapings, tissue specimens (Fig 14.31) and aspirates stained with Giemsa or Gram's stain. Spores can be identified in urine, fecal samples (Fig 14.46), or duodenal fluid by light, fluorescence,^{132,144} or electron microscopy.^{3,61,132} Where available, immunologic technology (such as ELISA,¹⁴⁵ Western blot,^{15,146} monoclonal and polyclonal an-




Figure 14.55
Cluster of *Encephalitozoon hellem* spores (arrows) in kidney. H&E x850




Figure 14.56
Cluster of *Encephalitozoon hellem* spores in kidney. Black bodies are Gram-positive spores. B&H x875




Figure 14.57
Encephalitozoon hellem spores in bladder exudates. B&H x650




Figure 14.58
Tubular epithelium of the kidney infected with *Encephalitozoon hellem*. Immunofluorescence staining reveals brightly staining clusters of intra-epithelial spores (arrow). x750




Figure 14.59
Encephalitozoon (Septata) intestinalis cells in tightly packed clusters in skin: Silvered 3 μ m x 1 μ m spores. GMS x790

tibody,^{147,148} fluorescent antibody,^{132,135,144,149}) and PCR^{150,151} techniques are significant diagnostic aids.^{28,63,152-156}

Electron microscopy is the preferred method for identifying microsporidia.^{19,55,61} It can be used to identify the presence of a polar filament in the spore. Additionally the polar filament arrangement and number of coil cross sections may aid in identification of organisms. Example: a double row of polar filament coils distinguishes spores of *Enterocytozoon bieneusi* from those of *Encephalitozoon intestinalis* in fecal or other samples. In tissue biopsies, microsporidial families can be identified by the host/parasite interface (Table 14.1). For example: there is no separation from host cytoplasm (Nosematidae, Enterocytozoonidae) by vacuole (Encephalitozoonidae) or SPOV (Pleistophoridae).

For histologic examination of tissue sections, glutaraldehyde fixation and plastic embedding are highly recommen-

ded, even if electron microscopy is unavailable. Parasites are often masked by the host tissue in 6 μ m paraffin-embedded preparations. One-micrometer-thick plastic sections, heat-fixed to a glass slide and stained with toluidine blue, provide better visualization of microsporidia, even by light microscopy, because of the thinness of the section (Fig 14.36). Plastic embedding provides a permanent tissue specimen that can be re-evaluated by electron microscopy if the need or opportunity arises. Additionally, paraffin-embedded tissue can be deparaffinized and reprocessed for plastic embedding.¹⁰

The most common microsporidia that infect humans are able to disseminate. Therefore, if they are found in a single location, other locations to which they are known to disseminate should be examined/tested.



Figure 14.60
Nodular Cutaneous Microsporidiosis of the leg caused by *Encephalitozoon intestinalis* in an HIV positive patient




Figure 14.61
Encephalitozoon intestinalis cells in skin from patient in Figure 14.60 demonstrating gram-positive spores with bipolar staining and central band. B&H x900

Treatment

Several drugs have been used to treat microsporidiosis in humans, with varying results. Fumagillin, a compound that inhibits microsporidiosis in honeybees, inhibits replication of *Encephalitozoon cuniculi* in human tissue culture.¹⁵⁷ Although not recommended or approved for internal use in humans, it has been used topically to treat *Encephalitozoon hellem* infections of the eye with some success.¹⁵⁸ Topical fluoroquinolone monotherapy resulted in resolution of 99% of 124 cases of keratitis.⁵⁸

Albendazole is the most effective treatment for microsporidiosis in humans. The usual regimen is 400 mg administered orally twice a day for 4 to 6 weeks. Albendazole was first used against *Enterocytozoon bienewisi* infection in HIV-positive patients with AIDS, chronic diarrhea, and wasting. After 4 weeks of treatment with albendazole, all patients regained continence and either gained weight or stopped losing weight. Within a month after stopping treatment, most patients had a recurrence of symptoms but responded well to further albendazole treatment.¹⁵⁹

Albendazole is more effective against *Encephalitozoon intestinalis* than against *Enterocytozoon bienewisi*.^{137,160} Parasites are cleared from the urine and intestinal tract within 6 weeks, with no recurrence after treatment. Biopsy reveals normal enterocyte morphology with only disintegrating spore remnants in macrophages.¹⁶⁰ Albendazole is also effective against other *Encephalitozoon* sp.^{128,161-164} and *Brachiola* sp.,⁶ clearing infections with no recurrence. It has proven effective against *Vittaforma corneae* in culture.¹⁶⁵⁻¹⁶⁷ The effects of albendazole, fumagillin, and TNP-470 on microsporidian replication in culture has also been studied.¹⁶⁸

References

1. Sprague V, Becnel JJ, Hazard EI. Taxonomy of phylum Microspora. *Crit Rev Microbiol.* 1992;18:285-395.
2. Weiss LM. The First United Workshop on Microsporidia from Invertebrate and Vertebrate Hosts. *Folia Parasitol (Praha).* 2005;52:1-7.
3. Bryan R, Cali A, Owen R, Spencer H. Microsporidia: newly recognized opportunistic pathogens in patients with AIDS. In: Sun T, ed. *Progress in Clinical Parasitology*. Vol 2. Blue Bell, PA: Field and Wood, Inc Pub; 1990:1-26.
4. Pinnolis M, Egbert PR, Font RL, Winter FC. Nosematosis of the cornea. Case report, including electron microscopic studies. *Arch Ophthalmol.* 1981;99:1044-1047.
5. Cali A. Comparison of the biology and pathology of microsporidia from different host groups. In: Samson RA, Vlak JM, Peters D, eds. *Fundamental and Applied Aspects of Invertebrate Pathology*. 1986:356-359. Netherlands, Fourth International Colloquium of Invertebrate Pathology; 1986:356-359.
6. Cali A, Takvorian PM, Lewin S, et al. *Brachiola vesicularum*, n. g., n. sp., a new microsporidium associated with AIDS and myositis. *J Eukaryot Microbiol.* 1998;45:240-251.
7. Margileth AM, Strano AJ, Chandra R, Neafie R, Blum M, McCully RM. Disseminated nosematosis in an immunologically compromised infant. *Arch Pathol.* 1973;95:145-150.
8. Silveira H, Canning EU. *Vittaforma corneae* n. comb. for the human microsporidium *Nosema corneum* Shadduck, Meccoli, Davis and Font, 1990, based on its ultrastructure in the liver of experimentally infected athymic mice. *J Eukaryot Microbiol.* 1995;42:158-165.
9. Ledford DK, Overman MD, Gonzalvo A, Cali A, Mester SW, Lockey RF. Microsporidiosis myositis in a patient with the acquired immunodeficiency syndrome. *Ann Intern Med.* 1985;102:628-630.
10. Cali A, Takvorian PM. Ultrastructure and development of *Pleistophora ronneafiei* n. sp., a Microsporidium (Protista) in the skeletal muscle of an immune-compromised individual. *Journal of Eukaryotic Microbiology.* 2003;50:77-85.
11. Hollister WS, Canning EU, Weidner E, Field AS, Kench J, Marriott DJ. Development and ultrastructure of *Trachipleistophora hominis* n.g., n.sp. after in vitro isolation from an AIDS patient and inoculation into athymic mice. *Parasitology.* 1996;112:143-154.
12. Desportes I, Le Charpentier Y, Galian A, et al. Occurrence of a new microsporidian: *Enterocytozoon bienewisi* n. g., n. sp., in the enterocytes of a human patient with AIDS. *J Protozool.* 1985;32:250-254.
13. Cali A, Owen RL. Intracellular development of *Enterocytozoon*, a unique microsporidian found in the intestine of AIDS patients. *J Protozool.* 1990;37:145-155.

14. Terada S, Reddy KR, Jeffers LJ, Cali A, Schiff ER. Microsporidian hepatitis in the acquired immunodeficiency syndrome. *Ann Intern Med.* 1987;107:61-62.
15. Didier ES, Didier PJ, Friedberg DN, et al. Isolation and characterization of a new human microsporidian, *Encephalitozoon hellem* (n. sp.), from three AIDS patients with keratoconjunctivitis. *J Infect Dis.* 1991;163:617-621.
16. Cali A, Meisler DM, Rutherford I, et al. Corneal microsporidiosis in a patient with AIDS. *Am J Trop Med Hyg.* 1991;44:463-468.
17. Cali A, Kotler DP, Orenstein JM. *Septata intestinalis*, n. g., n. Sp., an intestinal microsporidian associated with chronic diarrhea and dissemination in AIDS patients. *J Eukaryot Microbiol.* 1993;40:101-112.
18. Franzen C, Nasonova ES, Schölermerich J, Issi IV. Transfer of the members of the Genus *Brachiola* (Microsporidia) to the Genus *Anncalia* based on ultrastructural and molecular Data. *J Eukaryot Microbiol.* 2006;53:26-35.
19. Weber R, Bryan RT, Schwartz DA, Owen RL. Human microsporidial infections. *Clin Microbiol Rev.* 1994;7:426-461.
20. Cali A. Cytological and taxonomical comparison of two intestinal disseminating microsporidiosis. *AIDS.* 1993;7(suppl 3):S12-S16.
21. Sak B, Kasicková D, Kvác M, Kvetonová D, Ditrich O. Microsporidia in exotic birds: intermittent spore excretion of *Encephalitozoon* spp. in naturally infected budgerigars (*Melipittacus undulatus*). *Vet Parasitol.* 2010;168:196-200.
22. Friedberg DN, Ritterband DC. Ocular microsporidiosis. In: Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999:293-314.
23. Gunnarsson G, Hurlbut D, DeGirolami PC, Federman M, Wanke C. Multiorgan microsporidiosis: report of five cases and review. *Clin Infect Dis.* 1995;21:37-44.
24. Kotler DP, Orenstein JM. Clinical syndromes associated with microsporidiosis. In: Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999:258-292.
25. Orenstein JM, Gaetz HP, Yachnis AT, Frankel SS, Mertens RB, Didier ES. Disseminated microsporidiosis in AIDS: are any organs spared? *AIDS.* 1997;11:385-386.
26. Cali A, Takvorian PM, Keohane E, Weiss LM. Opportunistic microsporidian infections associated with myositis. *J Eukaryot Microbiol.* 1997;44:86S.
27. Field AS, Marriott DJ, Milliken ST, et al. Myositis associated with a newly described microsporidian, *Trachipleistophora hominis*, in a patient with AIDS. *J Clin Microbiol.* 1996;34:2803-2811.
28. Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999.
29. Cali A, Orenstein JM, Kotler DP, Owen RL. A comparison of two microsporidian parasites in enterocytes of AIDS patients with chronic diarrhea. *J Protozool.* 1991;38:96S-98S.
30. Modigliani R, Bories C, Le Charpentier Y, et al. Diarrhea and malabsorption in acquired immune deficiency syndrome: a study of four cases with special emphasis on opportunistic protozoan infestations. *Gut.* 1985;26:179-187.
31. Pol S, Romana CA, Richard SR, et al. *Enterocytozoon bienersi* infection in acquired immunodeficiency syndrome-related sclerosing cholangitis. *Gastroenterology.* 1992;102:1778-1781.
32. Orenstein JM. Pathology of intestinal microsporidiosis in HIV-infection [abstract]. *AIDS: Workshop on Intestinal Microsporidiosis in HIV Infection.* 1993;7(suppl 3).
33. Sandfort J, Hannemann A, Gelderblom H, Stark K, Owen RL, Ruf B. *Enterocytozoon bienersi* infection in an immunocompetent patient who had acute diarrhea and who was not infected with the human immunodeficiency virus. *Clin Infect Dis.* 1994;19:514-516.
34. Ashton N, Wirasinha PA. Encephalitozoonosis (nosematosis) of the cornea. *Br J Ophthalmol.* 1973;57:669-674.
35. Cali A, Meisler DM, Lowder CY, et al. Corneal microsporidiosis: characterization and identification. *J Protozool.* 1991;38:215S-217S.
36. Didier ES, Shadduck JA, Didier PJ, Millichamp N, Vossbrinck CR. Studies on ocular microsporidia. *J Protozool.* 1991;38:635-638.
37. Lowder CY, McMahon JT, Meisler DM, et al. Microsporidian keratoconjunctivitis caused by *Septata intestinalis* in a patient with acquired immunodeficiency syndrome. *Am J Ophthalmol.* 1996;121:715-717.
38. Visvesvara GS, Bellomo M, Moura H, et al. Isolation of *Nosema algerae* from the cornea of an immunocompetent patient. *J Eukaryot Microbiol.* 1999;46:10S.
39. Weber R, Bryan RT. Microsporidial infections in immunodeficient and immunocompetent patients. *Clin Infect Dis.* 1994;19:517-521.
40. Lacey CJ, Clarke AM, Fraser P, Metcalfe T, Bonsor G, Curry A. Chronic microsporidian infection of the nasal mucosae, sinuses, and conjunctivae in HIV disease. *Genitourin Med.* 1992;68:179-181.
41. Schwartz DA, Cali A, Visvesvara GS, Roseberger D, Hewan-Lowe KO, Bryan RT. A nasal microsporidian with unusual features from a patient with AIDS. *Abstracts of International Conference on AIDS.* 1993;1:384.
42. Molina JM, Sarfati C, Beauvais B, et al. Intestinal microsporidiosis in human immunodeficiency virus-infected patients with chronic unexplained diarrhea: prevalence and clinical and biological features. *J Infect Dis.* 1993;167:217-221.
43. Orenstein JM, Chiang J, Steinberg W, Smith PD, Rotterdam H, Kotler DP. Intestinal microsporidiosis as a cause of diarrhea in human immunodeficiency virus-infected patients: a report of 20 cases. *Hum Pathol.* 1990;21:475-481.
44. Verre J, Marriot D, Hing M, Field A, Harkness J. Evaluation of light microscopic detection of microsporidia spores in feces from HIV-infected patients [abstract]. *AIDS: Workshop on Intestinal Microsporidiosis in HIV Infection.* 1993;7(suppl 3).
45. Issi IV. Microsporidia as a phylum of parasitic protozoa. *Protozoology (Leningrad).* 1986;10:1-136.
46. Sprague V, Becnel JJ. Note on the name-author-date combination for the taxon *Microsporidies Balbiani*, 1882, when ranked as a phylum. *J Invertebr Pathol.* 1998;71:91-94.
47. Sprague V, Vavra J. Biology of the Microsporidia. In: Bulla LA, Cheng TC, eds. *Comparative Pathobiology*. Vol 1. New York, NY: Plenum Press; 1977:1-371.
48. Sprague V, Vavra J. Systematics of the Microsporidia. In: Bulla LA, Cheng TC, eds. *Comparative Pathobiology*. Vol 2. New York, NY: Plenum Press; 1977:1-510.
49. Levine ND, Corliss JO, Cox FE, et al. A newly revised classification of the protozoa. *J Protozool.* 1980;27:37-58.
50. Lainson R, Garnham PC, Killick-Kendrick R, Bird RG. Nosematosis, a microsporidian infection of rodents and other animals, including man. *Br Med J.* 1964;5407:470-472.
51. Weiser J. On the taxonomic position of the genus *Encephalitozoon Levaditi*, Nicolau and Schoen, 1923 (Protozoa: Microsporidia). *Parasitology.* 1964;54:749-751.
52. Cali A. Morphogenesis in the genus *Nosema*. Proceedings of the IVth International Colloquium on Insect Pathology. College Park, Md: 1970;431-438.
53. Hartskeerl R., Van Gool T, Schuitema AR, Didier ES, Terpstra WJ. Genetic and immunological characterization of the microsporidian *Septata intestinalis* Cali, Kotler and Orenstein, 1993: reclassification to *Encephalitozoon intestinalis*. *Parasitology.* 1995;110:277-285.
54. Lowman PM, Takvorian PM, Cali A. The effects of elevated temperatures and various time-temperature combinations on the development of *Brachiola (Nosema) algerae* N. Comb. in mammalian cell culture. *J Eukaryot Microbiol.* 2000;47:221-234.
55. Canning EU, Lom J, Dykova I. *The Microsporidia of Vertebrates*. London, England: Academic Press; 1986.
56. Lucas SB, Papadaki L, Conlon C, Sewankambo N, Goodgame R, Serwadda D. Diagnosis of intestinal microsporidiosis in patients with AIDS [letter]. *J Clin Pathol.* 1989;42:885-890.
57. Field AS, Canning EU, Hing MC, Verre J, Marriott DJ. Microsporidia in HIV-infected patients in Sydney, Australia: a report of 37 cases, a new diagnostic technique and the light microscopy and ultrastructure of a disseminated species. *AIDS: Workshop on Intestinal Microsporidiosis in HIV Infection* 1993;7(suppl 3):S27-S33.
58. Loh RS., Chan CM, Ti SE, Lim L, Chan KS, Tan DT. Emerging prevalence of microsporidial keratitis in Singapore: epidemiology, clinical features, and management. *Ophthalmology.* 2009;116:2348-2353.
59. Desportes-Livage I, Levy JP. Intestinal microsporidia and HIV Infection. *AIDS 1993; 7(Suppl 3): S1-S2*. Editorial, Workshop on Intestinal Microsporidia in HIV Infection, 15-16 December 1992, Paris, France
60. van Gool T, Snijders F, Reiss P, et al. Diagnosis of intestinal and disseminated microsporidian infections in patients with HIV by a new rapid fluorescence technique. *J Clin Pathol.* 1993;46:694-699.
61. Cali A, Owen RL. Microsporidiosis. In: Balows A, Hausler WJ Jr, Ohashi M, and Turano A, eds. *The Laboratory Diagnosis of Infectious Diseases: Principles and Practice: Vol I: Bacterial, Mycotic, and Parasitic Diseases*. New York: Springer-Verlag; 1988:928-949.
62. Schwartz DA, Sobottka I, Leitch GJ, Cali A, Visvesvara GS. Pathology of microsporidiosis: emerging parasitic infections in patients with acquired immunodeficiency syndrome. *Arch Pathol Lab Med.* 1996;120:173-188.
63. Sak B, Brady D, Pelikanova M, et al. Unapparent microsporidian infection among immunocompetent humans in the Czech Republic. *J Clin Microbiol.* 2010;PMID: 21191056
64. Microsporidian keratoconjunctivitis in patients with AIDS. *MMWR Morb Mortal Wkly Rep.* 1990;39:188-189.

65. Coyle C, Weiss LM, Rhodes LV, et al. Fatal myositis due to the microsporidian *Brachiola algerae*, a mosquito pathogen. *N Engl J Med*. 2004;351:42-47.
66. Talabani H, Sarfati C, Pillebout E, van Gool T, Derouin F, Menotti J. Disseminated infection with a new genovar of *Encephalitozoon cuniculi* in a renal transplant recipient. *J Clin Microbiol*. 2010;48:2651-2653.
67. Lanterrier F, Boutboul D, Menotti J, et al. Microsporidiosis in solid organ transplant recipients: two *Enterocytozoon bienewsi* cases and review. *Transpl Infect Dis*. 2009;11:83-88.
68. Mathis A, Weber R, Deplazes P. Zoonotic potential of the microsporidia. *Clin Microbiol Rev*. 2005;18:423-445.
69. Cali A, Takvorian PM. Microsporidia. In: *Handbook of Protozoa*, 2nd ed. Margulis L, Chapman M, eds. Jones and Bartlett, Boston. In press. 2011.
70. Williams B, Hirt R, Lucocq JM, Embley TM. A mitochondrial remnant in the microsporidian *Trachipleistophora hominis*. *Nature*. 2002;418:865-869.
71. Katinka MD, Duprat S, Cornillot E, et al. Genome sequence and gene compaction of the eukaryote parasite *Encephalitozoon cuniculi*. *Nature*. 2001;414:450-453.
72. Vossbrinck CR, Maddox JV, Friedman S, Debrunner-Vossbrinck BA, Woese CR. Ribosomal RNA sequence suggests microsporidia are extremely ancient eukaryotes. *Nature*. 1987;326:411-414.
73. Takvorian PM, Cali A. Enzyme histochemical identification of the Golgi apparatus in the microsporidian, *Glugea stephani*. *J Eukaryot Microbiol*. 1994;41:63S-64S.
74. Takvorian PM, Cali A. Polar tube formation and nucleoside diphosphatase activity in the microsporidian, *Glugea stephani*. *J Eukaryot Microbiol*. 1996;43:102S-103S.
75. Edlind T, Katiyar S, Visvesvara G, Li J. Evolutionary origins of Microsporidia and basis for benzimidazole sensitivity: an update. *J Eukaryot Microbiol*. 1996;43:109S.
76. Edlind TD, Li J, Visvesvara GS, Vodkin MH, McLaughlin GL, Katiyar SK. Phylogenetic analysis of beta-tubulin sequences from amitochondrial protozoa. *Mol Phylogenet Evol*. 1996;5:359-367.
77. Cali A. General microsporidian features and recent findings on AIDS isolates. *J Protozool*. 1991;38:625-630.
78. Cali A, Takvorian PM. Developmental morphology and life cycles of the Microsporidia. In: Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999:85-128.
79. Cali A, Takvorian PM. *Brachiola algerae* sporoplasms. *J Eukaryot Microbiol*. 2001;Suppl:81S-82S.
80. Lom J, Vavra J. The mode of sporoplasm extrusion in microsporidian spores. *Acta Protozool*. 1963;1:81-89.
81. Cali A, Takvorian PM. The Microsporidia: pathology in man and occurrence in nature. *Southeast Asian J Trop Med Pub Health*. 2004;35:58-64.
82. Enriquez FJ, Taren D, Cruzlopez A, Muramoto M, Palting JD, Cruz P. Prevalence of intestinal encephalitozoonosis in Mexico. *Clin Infect Dis*. 1998;26:1227-1229.
83. Lobo ML, Teles A, da Cunha MB, et al. Microsporidia detection in stools from pets and animals from the zoo in Portugal: a preliminary study. *J Eukaryot Microbiol*. 2003;50 (Suppl):581-582.
84. Curry A, Hardeep SM, Sumedh D, Canning EU, Wagner BE. A case of bilateral microsporidial keratitis from Bangladesh: infection by insect parasite from the genus *Nosema*. *J Med Microbiol*. 2007;56:1250-1252.
85. Shadduck JA, Meccoli RA, Davis R, Font RL. Isolation of a microsporidian from a human patient. *J Infect Dis*. 1990;162:773-776.
86. Franzen C, Fischer S, Schroeder J, Scholmerich J, Schneuwly S. Morphological and molecular investigations of *Tubulinosema ratisbonensis* gen. nov., sp. nov. (Microsporidia: Tubulinosematidae fam. nov.), a parasite infecting a laboratory colony of *Drosophila melanogaster* (Diptera: Drosophilidae). *J Eukaryot Microbiol*. 2005;52:141-152.
87. Cali A, Weiss LM, Takvorian PM. An Analysis of the microsporidian genus *Brachiola*, with comparisons of human and insect isolates of *B. algerae*. *J Eukaryot Microbiol*. 2004;51:678-685.
88. Davis RM, Font RL, Keisler MS, Shadduck JA. Corneal microsporidiosis. A case report including ultrastructural observations. *Ophthalmology*. 1990;97:953-957.
89. Strano AJ, Cali A, Neafie R. Microsporidiosis. In: Binford CH, Connor DH, eds. *Pathology of Tropical and Extraordinary Diseases*. Vol 1. Washington, DC: AFIP; 1976:336-339.
90. Sprague V. *Nosema connori* n. sp., a microsporidian parasite of man. *Trans Am Microsc Soc*. 1974;93:400-403.
91. Trammer T, Chioralia G, Maier WA, Seitz HM. In vitro replication of *Nosema algerae* (Microsporidia), a parasite of anopheline mosquitoes, in human cells above 36 degrees C. *J Eukaryot Microbiol*. 1999;46:464-468.
92. Cali A, Weiss LM, Takvorian PM. A review of the development of two types of human skeletal muscle infections from microsporidia associated with pathology in invertebrates and cold-blooded vertebrates. *Folia Parasitologica (Praha)*. 2005;52:51-61.
93. Cali A, Neafie R, Weiss LM, et al. Human vocal cord infection with the Microsporidium *Anncalia algerae*. *J Eukaryot Microbiol*. 2010;57:562-567.
94. Chupp GL, Alroy J, Adelman LS, Breen JC, Skolnik PR. Myositis due to *Pleistophora* (Microsporidia) in a patient with AIDS. *Clin Infect Dis*. 1993;16:15-21.
95. Grau A, Valls ME, Williams JE, Ellis DS, Muntane MJ, Nadal C. Myositis caused by *Pleistophora* in a patient with AIDS [in Spanish]. *Med Clin (Barc)*. 1996;107:779-781.
96. Vavra J, Yachnis AT, Shadduck JA, Orenstein JM. Microsporidia of the genus *Trachipleistophora*—causative agents of human microsporidiosis: description of *Trachipleistophora anthropophthera* n. sp. (Protozoa: Microsporidia). *J Eukaryot Microbiol*. 1998;45:273-283.
97. Yachnis AT, Berg J, Martinez-Salazar A, et al. Disseminated microsporidiosis especially infecting the brain, heart, and kidneys. Report of a newly recognized pansporoblastic species in two symptomatic AIDS patients. *Am J Clin Pathol*. 1996;106:535-543.
98. Weidner E, Canning EU, Hollister WS. The plaque matrix (PQM) and tubules at the surface of intramuscular parasite, *Trachipleistophora hominis*. *J Eukaryot Microbiol*. 1997;44:359-365.
99. Macher AM, Neafie R, Angritt P, Tuur SM. Microsporidial myositis and the acquired immunodeficiency syndrome (AIDS): a four-year follow-up [letter]. *Ann Intern Med*. 1988;109:343.
100. Macher AM, Neafie R, Angritt P, Tuur SM, Nelson RP, Cali A. AIDS. Case for diagnosis series, 1988. *Mil Med*. 1988;153:M41-M48.
101. Curry A, Beeching NJ, Gilbert JD, Scott G, Rowland PL, Currie BJ. *Trachipleistophora hominis* infection in the myocardium and skeletal muscle of a patient with AIDS. *J Infect*. 2005;51:e139-e144.
102. Rinder H, Thomschke A, Dengie B, Gothe R, Loscher T, Zahler M. Close genotypic relationship between *Enterocytozoon bienewsi* from humans and pigs and first detection in cattle. *J Parasitol*. 2000;86:185-188.
103. Orenstein JM, Lew E, Poles MA, Dieterich D. The endoscopic brush cytology specimen in the diagnosis of intestinal microsporidiosis. *AIDS*. 1995;9:1199-1201.
104. McWhinney PH, Nathwani D, Green ST, Boyd JF, Forrest JA. Microsporidiosis detected in association with AIDS-related sclerosing cholangitis. *AIDS*. 1991;5:1394-1395.
105. Canning EU, Hollister WS. *Enterocytozoon bienewsi* (Microspora): prevalence and pathogenicity in AIDS patients. *Trans R Soc Trop Med Hyg*. 1990;84:181-186.
106. Orenstein JM, Friedberg DN, Cali A, Kotler DP. Microsporidiosis in AIDS: an update of enteritis caused by *E. bienewsi* and a second species, and *E. cuniculi* as a cause of keratoconjunctivitis. *Abstracts of Sixth International Conference on AIDS*. 1990:Th.B.522.
107. Sun T, Kaplan MH, Teichberg S, Weissman G, Smilari T, Urmacher C. Intestinal microsporidiosis: report of five cases. *Ann Clin Lab Sci*. 1994;24:521-532.
108. Rabeneck L, Gyorkey F, Genta RM, Gyorkey P, Foote LW, Risser JM. The role of Microsporidia in the pathogenesis of HIV-related chronic diarrhea. *Ann Intern Med*. 1993;119:895-899.
109. Weber R, Kuster H, Visvesvara GS, Bryan RT, Schwartz DA, Luthy R. Disseminated microsporidiosis due to *Encephalitozoon hellem*: pulmonary colonization, microhematuria, and mild conjunctivitis in a patient with AIDS. *Clin Infect Dis*. 1993;17:415-419.
110. Eeftink Schattenkerk JK, van Gool T, Schot LS, van den Bergh Weerman MA, Schot LJ, Dankert J. Chronic rhinosinusitis in HIV-infected patients with microsporidiosis. *AIDS: Workshop on Intestinal Microsporidiosis in HIV Infection*. 1993;7(suppl 3):S36-S37.
111. Hartskeerl RA, Schuitema AR, van Gool T, Terpstra WJ. Genetic evidence for the occurrence of extra-intestinal *Enterocytozoon bienewsi* infections. *Nucleic Acids Res*. 1993;21:4150.
112. Weber R, Bryan RT, Owen RL, Wilcox CM, Gorelkin L, Visvesvara GS. Improved light-microscopical detection of microsporidia spores in stool and duodenal aspirates. The Enteric Opportunistic Infections Working Group. *N Engl J Med*. 1992;326:161-166.

113. Levaditi C, Nicolau S, Schoen R. Etiologic agent of epizootic encephalitis in rabbits (*Encephalitozoon cuniculi*) [in French]. *Comp Rend Soc Biol*. 1923;89:984-986.
114. Pakes SP, Shadduck JA, Cali A. Fine structure of *Encephalitozoon cuniculi* from rabbits, mice and hamsters. *J Protozool*. 1975;22:481-488.
115. Voronin VN. Macrotaxonomy of the Microsporidia phylum. *Parazitologiya*. 2001;35:35-44.
116. Friedberg DN, Stenson SM, Orenstein JM, Tierno PM, Charles NC. Microsporidial keratoconjunctivitis in acquired immunodeficiency syndrome. *Arch Ophthalmol*. 1990;108:504-508.
117. Lowder CY, Meisler DM, McMahon JT, Longworth DL, Rutherford I. Microsporidia infection of the cornea in a man seropositive for human immunodeficiency virus. *Am J Ophthalmol*. 1990;109:242-244.
118. Desser SS, Hong H, Yang YJ. Ultrastructure of the development of a species of *Encephalitozoon* cultured from the eye of an AIDS patient. *Parasitol Res*. 1992;78:677-683.
119. Schwartz DA, Visvesvara GS, Dieneshouse MC, et al. Pathologic features and immunofluorescent antibody demonstration of ocular microsporidiosis (*Encephalitozoon hellem*) in seven patients with acquired immunodeficiency syndrome. *Am J Ophthalmol*. 1993;115:285-292.
120. Orenstein JM, Tenner M, Cali A, Kotler DP. A microsporidian previously undescribed in humans, infecting enterocytes and macrophages, and associated with diarrhea in an acquired immunodeficiency syndrome patient. *Hum Pathol*. 1992;23:722-728.
121. Dore GJ, Marriott DJ, Hing MC, Harkness JL, Field AS. Disseminated microsporidiosis due to *Septata intestinalis* in nine patients infected with the human immunodeficiency virus: response to therapy with albendazole. *Clin Infect Dis*. 1995;21:70-76.
122. Fields AS, Hing MC, Milliken ST, Marriott DJ. Microsporidia in the small intestine of HIV-infected patients. A new diagnostic technique and a new species. *Med J Aust*. 1993;158:390-394.
123. Sprague V, Vernick SH. The ultrastructure of *Encephalitozoon cuniculi* (Microsporidia, Nosematidae) and its taxonomic significance. *J Protozool*. 1971;18:560-569.
124. Cali A, Weiss LM, Takvorian PM. Microsporidian taxonomy and the status of *Septata intestinalis*. *J Eukaryot Microbiol*. 1996;43:106S-107S.
125. Bergquist NR, Morfeldt-Mansson L, Pehrson PO, Petrini B, Wasserman J. Antibody against *Encephalitozoon cuniculi* in Swedish homosexual men. *Scand J Infect Dis*. 1984;16:389-391.
126. Bergquist NR, Stintzing G, Smedman L, Waller T, Andersson T. Diagnosis of encephalitozoonosis in man by serological tests. *Br Med J (Clin Res Ed)*. 1984;288:902.
127. Gordon SC, Reddy KR, Gould EE, et al. The spectrum of liver disease in the acquired immunodeficiency syndrome. *J Hepatol*. 1986;2:475-484.
128. De Groote MA, Visvesvara G, Wilson ML, et al. Polymerase chain reaction and culture confirmation of disseminated *Encephalitozoon cuniculi* in a patient with AIDS: successful therapy with albendazole. *J Infect Dis*. 1995;171:1375-1378.
129. Weber R, Deplazes P, Flepp M, et al. Cerebral microsporidiosis due to *Encephalitozoon cuniculi* in a patient with human immunodeficiency virus infection. *N Engl J Med*. 1997;336:474-478.
130. Zender HO, Arrigoni E, Eckert J, Kapanci Y. A case of *Encephalitozoon cuniculi* peritonitis in a patient with AIDS. *Am J Clin Pathol*. 1989;92:352-356.
131. Canning EU, Curry A, Lacey CJ, Fenwick JD. Ultrastructure of *Encephalitozoon* sp. infecting the conjunctival, corneal and nasal epithelia of a patient with AIDS. *Eur J Protistol*. 1992;28:226-237.
132. Schwartz DA, Bryan RT, Hewan-Lowe KO, et al. Disseminated microsporidiosis (*Encephalitozoon hellem*) and acquired immunodeficiency syndrome: autopsy evidence for respiratory acquisition. *Arch Pathol Lab Med*. 1992;116:660-668.
133. Schwartz DA, Visvesvara GS, Leith GJ, et al. Pathology of symptomatic microsporidial (*Encephalitozoon hellem*) bronchiolitis in the acquired immunodeficiency syndrome: a new respiratory pathogen diagnosed from lung biopsy, bronchoalveolar lavage, sputum, and tissue culture. *Hum Pathol*. 1993;24:937-943.
134. Schwartz DA, Visvesvara GS, Weber R, Bryan RT. Male genital tract microsporidiosis and AIDS: prostatic abscess due to *Encephalitozoon hellem*. *J Eukaryot Microbiol*. 1994;41:61S.
135. Visvesvara GS, Leith GJ, Moura H, Wallace S, Weber R, Bryan RT. Culture, electron microscopy, and immunoblot studies on a microsporidian parasite isolated from the urine of a patient with AIDS. *J Protozool*. 1991;38:105S-111S.
136. Molina JM, Oksenhendler E, Beauvais B, et al. Disseminated microsporidiosis due to *Septata intestinalis* in patients with AIDS: clinical features and response to albendazole therapy. *J Infect Dis*. 1995;171:245-249.
137. Orenstein JM, Dieterich DT, Kotler DP. Systemic dissemination by a newly recognized intestinal microsporidia species in AIDS. *AIDS*. 1992;6:1143-1150.
138. Coyle CM, Wittner M, Kotler DP, et al. Prevalence of microsporidiosis due to *Enterocytozoon bienewsi* and *Encephalitozoon (Septata) intestinalis* among patients with AIDS-related diarrhea: determination by polymerase chain reaction to the microsporidian small-subunit rRNA gene. *Clin Infect Dis*. 1996;23:1002-1006.
139. Luna LG. *Manual of Histologic Staining Methods of the Armed Forces Institute of Pathology*. New York, NY: McGraw-Hill; 1968.
140. Rijpstra AC, Canning EU, Van Ketel RJ, Eeftink Schattenkerk JK, Laarman JJ. Use of light microscopy to diagnose small-intestinal microsporidiosis in patients with AIDS. *J Infect Dis*. 1988;157:827-831.
141. Simon D, Weiss LM, Tanowitz HB, Cali A, Jones J, Wittner M. Light microscopic diagnosis of human microsporidiosis and variable response to ocreotide. *Gastroenterology*. 1991;100:271-273.
142. Weber R, Schwartz DA, Deplazes P. Laboratory diagnosis of microsporidiosis. In: Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999:315-362.
143. Field AS, Marriott DJ, Hing MC. The Warthin-Starry stain in the diagnosis of small intestinal microsporidiosis in HIV-infected patients. *Folia Parasitol (Praha)*. 1993;40:261-266.
144. van Gool T, Canning EU, Dankert J. An improved practical and sensitive technique for the detection of microsporidian spores in stool samples. *Trans R Soc Trop Med Hyg*. 1994;88:189-190.
145. Hollister WS, Canning EU. An enzyme-linked immunosorbent assay (ELISA) for detection of antibodies to *Encephalitozoon cuniculi* and its use in determination of infections in man. *Parasitology*. 1987;94:209-219.
146. Weiss LM, Cali A, Levee E, et al. Diagnosis of *Encephalitozoon cuniculi* infection by western blot and the use of cross-reactive antigens for the possible detection of microsporidiosis in humans. *Am J Trop Med Hyg*. 1992;47:456-462.
147. Aldras AM, Orenstein JM, Kotler DP, Shadduck JA, Didier ES. Detection of microsporidia by indirect immunofluorescence antibody test using polyclonal and monoclonal antibodies. *J Clin Microbiol*. 1994;32:608-612.
148. Keohane E, Takvorian PM, Cali A, Tanowitz HB, Wittner M, Weiss LM. The identification and characterization of a polar tube reactive monoclonal antibody. *J Eukaryot Microbiol*. 1994;41:48S.
149. Alfa Cisse O, Ouattara A, Thellier M, et al. Evaluation of an immunofluorescent-antibody test using monoclonal antibodies directed against *Enterocytozoon bienewsi* and *Encephalitozoon intestinalis* for diagnosis of intestinal microsporidiosis in Bamako (Mali). *J Clin Microbiol*. 2002;40:1715-1718.
150. Garcia L. Laboratory identification of microsporidia. *Journal of Clinical Microbiology*. 2002;40:1892-1901.
151. Weiss LM, Zhu X, Cali A, Tanowitz HB, Wittner M. Polymerase chain reaction identification of *Microsporidia* using ribosomal RNA. *AIDS: Workshop on Intestinal Microsporidiosis in HIV Infection*. 1993;7(suppl 3):S62-S63.
152. Weiss LM, Zhu X, Cali A, Tanowitz HB, Wittner M. Utility of microsporidian rRNA in diagnosis and phylogeny: a review. *Folia Parasitol (Praha)*. 1994;41:81-90.
153. Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM. Nucleotide sequence of the small ribosomal RNA of *Encephalitozoon cuniculi*. *Nucleic Acids Res*. 1993;21:1315.
154. Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM. Nucleotide sequence of the small subunit rRNA of *Septata intestinalis*. *Nucleic Acids Res*. 1993;21:4846.
155. Zhu X, Wittner M, Tanowitz HB, Cali A, Weiss LM. Ribosomal RNA sequences of *Enterocytozoon bienewsi*, *Septata intestinalis* and *Ameson michaelis*: phylogenetic construction and structural correspondence. *J Eukaryot Microbiol*. 1994;41:204-209.
156. Zhu X, Wittner M, Tanowitz HB, Kotler D, Cali A, Weiss LM. Small subunit rRNA sequence of *Enterocytozoon bienewsi* and its potential diagnostic role with the use of the polymerase chain reaction. *J Infect Dis*. 1993;168:1570-1575.
157. Shadduck JA. Effect of fumagillin on in vitro multiplication of *Encephalitozoon cuniculi*. *J Protozool*. 1980;27:202-208.
158. Yee RW, Tio FO, Martinez JA, Held KS, Shadduck JA, Didier ES. Resolution of microsporidian epithelial keratopathy in a patient with AIDS. *Ophthalmology*. 1991;98:196-201.

159. Blanshard C, Ellis DS, Tovey DG, Dowell S, Gazzard BG. Treatment of intestinal microsporidiosis with albendazole in patients with AIDS. *AIDS*. 1992;6:311-313.
160. Orenstein JM, Dieterich DT, Lew EA, Kotler DP. Albendazole as a treatment for intestinal and disseminated microsporidiosis due to *Septata intestinalis* in AIDS patients: a report of four patients. *AIDS: Workshop on Intestinal Microsporidiosis in HIV Infection*. 1993;7(suppl 3):S40-S42.
161. Koudela B, Lom J, Vitovec J, Kucerova Z, Ditrich O, Travnicek J. In vivo efficacy of albendazole against *Encephalitozoon cuniculi* in SCID mice. *J Eukaryot Microbiol*. 1994;41:49S-50S.
162. Lecuit M, Oksenhendler E, Sarfati C. Use of albendazole for disseminated microsporidian infection in a patient with AIDS. *Clin Infect Dis*. 1994;19:332-333.
163. Weiss LM, Michalakakis E, Coyle CM, Tanowitz HB, Wittner M. The in vitro activity of albendazole against *Encephalitozoon cuniculi*. *J Eukaryot Microbiol*. 1994;41:65S.
164. Aarons EJ, Woodrow D, Hollister WS, Canning EU, Francis N, Gazzard BG. Reversible renal failure caused by a microsporidian infection. *AIDS*. 1994;8:1119-1121.
165. Didier ES, Maddy JA, Kwong CD, Green LC, Snowden KF, Shadduck JA. Screening of compounds for antimicrosporidian activity in vitro. *Folia Parasitol (Praha)*. 1998;45:129-139.
166. Silveira H, Canning EU. In vitro cultivation of the human microsporidium *Vittaforma corneae*: development and effect of albendazole. *Folia Parasitol (Praha)*. 1995;42:241-250.
167. Franzen C, Salzberger B. Analysis of the tubulin gene from *Vittaforma corneae* suggests benzimidazole resistance. *Antimicrob Agents Chemother*. 2008;52:790-793.
168. Didier ES. Effects of albendazole, fumagillin, and TNP-470 on microsporidian replication in vitro. *Antimicrob Agents Chemother*. 1997;41:1541-1546.

Acknowledgements

The following figures are used by permission. "Copyright © American Society for Microbiology."

Figures 14.1, 14.4.

Cali A, Takvorian PM. Developmental morphology and life cycles of the Microsporidia. In: Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999:85-128.

Figures 14.30 to 14.35.

Field AS, Marriott DJ, Milliken ST, et al. Myositis associated with a newly described Microsporidian, *Trachipleistophora hominis*, in a patient with AIDS. *J Clin Microbiol*. 1996;34:2803-2811.

Licensed to the Armed Forces Institute of Pathology. "Copyright © American Society for Microbiology."

Figures 14.40a,b and 14.43.

Kotler DP, Orenstein JM. Clinical syndromes associated with microsporidiosis. In: Wittner M, Weiss LM, eds. *The Microsporidia and Microsporidiosis*. Washington, DC: ASM Press; 1999:258-292.

The following figures are used by permission
"Copyright © Wiley Publications."

Figure 14.5.

Cali A, Takvorian PM. *Brachiola algerae* sporoplasms. *J Eukaryot Microbiol* 2001;Suppl:81S-82S.

Figures 14.19 to 14.23.

Cali A, Takvarian PM, Lewin S, et al. *Brachiola vesicularum*, n. g., n. sp., a new microsporidium associated with AIDS and myositis. *J Eukaryot Microbiol*. 1998;45:240-251.

Figures 14.28, 14.29a,b.

Cali A, Takvorian PM. Ultrastructure and development of *Pleistophora ronneqfiei* n. sp, a Microsporidium (Protista) in the skeletal muscle of an immune-compromised individual. *J Eukaryot Microbiol*. 2003;50:77-85.

Figure 14.36

Licensed to the Armed Forces Institute of Pathology.

Vavra J, Yachnis AT, Shadduck JA, Orenstein JM. Microsporidia of the genus *Trachipleistophora*-causative agents of human microsporidiosis: description of *Trachipleistophora anthropophthera* n. sp. (Protozoa: Microsporidia). *J Eukaryot Microbiol*. 1998;45:273-283.

Fig 14.41, 14.42.

Cali A, Owen RL. Intracellular development of *Enterocytozoon*, a unique microsporidian found in the intestine of AIDS patients. *J Protozool*. 1990;37:145:155.

Figure 14.48.

Cali A, Takvorian PM, Lewin S, et al. Identification of a new Nosema-like microsporidian associated with myositis in an AIDS patient. *J Eukaryot Microbiol*. 1996;43:108S.

Figure 14.50

Cali A, Kotler DP, Orenstein JM. *Septata intestinalis*, n. g, n. sp, an intestinal microsporidian associated with chronic diarrhea and dissemination in AIDS patients. *J Eukaryot Microbiol*. 1993;40:101-112.

The following figures are used by permission,
"Massachusetts Medical Society © 2010."

Figure 14.24, 14.25a,b

Coyle C, Weiss LM, Rhodes LV, et al. Fatal myositis due to the microsporidian *Brachiola algerae*, a mosquito pathogen. *N Engl J Med*. 2004;351:42-47.

The following figures are used by permission,
"Copyright © American Journal of Tropical
Medicine and Hygiene."

Figures 14.47, 14.49

Cali A, Meisler DM, Rutherford I, et al. Corneal microsporidiosis in a patient with AIDS. *Am J Trop Med Hyg*. 1991;44:463-468.

The following figure used by permission.

Figure 14.58

Compliments of Govinda S. Visvesvara, Ph.D. Centers for Disease Control and Prevention.