

The Spectrum of Generalized Petersen Graphs

RALUCCA GERA and PANTELIMON STĂNICĂ*

Department of Applied Mathematics

Naval Postgraduate School

Monterey, CA 93943, USA

{rgera,pstanica}@nps.edu

September 5, 2010

*Also associated to the Institute of Mathematics of the Romanian Academy, Bucharest, Romania.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 05 SEP 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE The Spectrum of Generalized Petersen Graphs				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Applied Mathematics, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Australasian Journal of Combinatorics 49 (2011), 39-45					
14. ABSTRACT In this paper, we completely describe the spectrum of the generalized Petersen graph $P(n; k)$, thus adding to the classes of graphs whose spectrum is completely known.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Corresponding author:

Pantelimon Stănică

¹

Department of Applied Mathematics

Naval Postgraduate School

Monterey, CA 93943, USA

pstanica@nps.edu

¹Also associated to the Institute of Mathematics of the Romanian Academy, Bucharest, Romania.

Abstract

In this paper, we completely describe the spectrum of the generalized Petersen graph $P(n, k)$, thus adding to the classes of graphs whose spectrum is completely known.

1 Introduction and motivation

Let $G = (V(G), E(G))$ be a simple graph. The spectrum of a graph G is the multiset of eigenvalues of the adjacency matrix. The graph spectrum is an important tool one can use to find information about the physical properties of a network, such as robustness, diameter, connectivity [3]. In this research we completely describe the spectrum for the class of graphs, defined below.

The *generalized Petersen graph* (GPG) $P(n, k)$ has vertices, respectively, edges given by

$$\begin{aligned} V(P(n, k)) &= \{a_i, b_i, 0 \leq i \leq n-1\}, \\ E(P(n, k)) &= \{a_i a_{i+1}, a_i b_i, b_i b_{i+k} \mid 0 \leq i \leq n-1\}, \end{aligned}$$

where the subscripts are expressed as integers modulo n ($n \geq 5$), and k is the “skip”. Note that $k \leq \lfloor \frac{n-1}{2} \rfloor$, because of the obvious isomorphism $P(n, k) \cong P(n, n-k)$. Let $A(n, k)$ (respectively, $B(n, k)$) be the subgraph of $P(n, k)$ consisting of the vertices $\{a_i \mid 0 \leq i \leq n-1\}$ (respectively, $\{b_i \mid 0 \leq i \leq n-1\}$) and edges $\{a_i a_{i+1} \mid 0 \leq i \leq n-1\}$ (respectively, $\{b_i b_{i+k} \mid 0 \leq i \leq n-1\}$). We will call $A(n, k)$ (respectively, $B(n, k)$) the *outer* (respectively, *inner*) subgraph of $P(n, k)$. We display in Figure 1 the graph $P(12, 3)$.

For other graph theoretical terminology the reader could refer to [7].

2 Eigenvalues of $P(n, k)$

In this section we find our description for the spectrum of generalized Petersen graphs $P(n, k)$. We denote the adjacency matrix of the GPG $P(n, k)$

AMS Mathematics Subject Classification: 05C70, 05C12, 05C50, 15A42.

Keywords: Generalized Petersen Graphs; Eigenvalues; Spectrum.




Figure 1: The Generalized Petersen Graph $P(12, 3)$

by $A(P(n, k))$. Let $\lambda_0 = 3 > \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{2n-1}$ be the sequence of eigenvalues of $P(n, k)$.

We call an $n \times n$ matrix *circulant*, and denote it by $\text{circ}(a_1, a_2, \dots, a_n)$ if it is of the form

$$\text{circ}(a_1, a_2, \dots, a_n) = \begin{pmatrix} a_1 & a_2 & a_3 & \dots & a_n \\ a_n & a_1 & a_2 & \dots & a_{n-1} \\ \vdots & & & & \vdots \\ a_2 & a_3 & a_4 & \dots & a_1 \end{pmatrix}.$$

Lemma 2.1 *The $(2n) \times (2n)$ adjacency matrix of the GPG $P(n, k)$ has the block form*

$$A(P(n, k)) = \begin{pmatrix} C_k^n & I_n \\ I_n & C^n \end{pmatrix},$$

where I_n is the $n \times n$ identity matrix, C^n, C_k^n are circulant matrices, with $C^n = \text{circ}(0, 1, 0, 0, \dots, 0, 1)$ and $C_k^n = \text{circ}(\underbrace{0, \dots, 0}_{k \text{ times}}, 1, 0, \dots, 0, 1, \underbrace{0, \dots, 0}_{k-1 \text{ times}})$ being the adjacency matrix for $A(n, k)$ and $B(n, k)$, respectively. Thus, C^n

is the adjacency matrix of a cycle graph on n vertices \mathcal{C}_n , respectively, C_k^n is the union of d cycle graphs $\mathcal{C}_{n/d}$ on n/d vertices, where $d = \gcd(n, k)$.

Proof. The outer subgraph (whose adjacency matrix is C^n) of $P(n, k)$ is the cycle graph \mathcal{C}_n and the inner subgraph (whose adjacency matrix is C_k^n) has d connected components each isomorphic to $\mathcal{C}_{n/d}$. Also, the adjacency matrix (which depends on the labeling) has the claimed form where the labels used on the outer subgraph are consecutively $1, 2, \dots, n$, and on the inner subgraph the adjacent labels are $i, i + k, i + 2k, \dots$ (where $i + sk$ is understood as $1 + (i - 1 + sk) \pmod{n}$). Note that b_0 is adjacent to vertex b_k in the subgraph and to vertex labeled b_{n-k} in $B(n, k)$, and so $C_k^n = \text{circ}(\overbrace{0, \dots, 0}^{k \text{ times}}, 1, 0, \dots, 0, 1, \overbrace{0, \dots, 0}^{k-1 \text{ times}})$ \square

We recall the Chebyshev's polynomial of the first kind [5], defined by the identity $T_n(\cos \theta) = \cos(n\theta)$, with the generating function $\sum_{n=0}^{\infty} T_n(x)t^n = \frac{1-xt}{1-2xt+t^2}$. We now present the eigenvectors and eigenvalues for \mathcal{C}_n (see [2, p. 53 and pp. 72–73]). Let \mathbf{v}^t denote the transpose of \mathbf{v} .

Lemma 2.2 *The eigenvalues of the cycle graph \mathcal{C}_n on n vertices are*

$$\alpha_j = 2 \cos \left(\frac{2\pi j}{n} \right)$$

with a corresponding eigenvector

$$\mathbf{v}_j = (1, \zeta^j, \zeta^{2j}, \dots, \zeta^{(n-1)j})^t,$$

$0 \leq j \leq n - 1$. The characteristic polynomial of the cycle \mathcal{C}_n is $2T_n(x/2) - 2$ where T_n is the Chebyshev's polynomial of the first kind.

Corollary 2.3 *The eigenvalues corresponding to the circulant C in the adjacency matrix $A(P(n, k))$ are $\alpha_j = 2 \cos \left(\frac{2\pi j}{n} \right)$ ($0 \leq j \leq n - 1$), and the eigenvalues corresponding to C_k are $\beta_j = 2 \cos \left(\frac{2\pi jk}{n} \right)$ ($0 \leq j \leq n - 1$).*

We now state our main theorem which adds to the class of graphs whose spectrum is now known.

Theorem 2.4 *The eigenvalues of $P(n, k)$, say $\delta_{2j}, \delta_{2j+1}$, are all roots of the quadratic equation*

$$\delta^2 - (\alpha_j + \beta_j)\delta + \alpha_j\beta_j - 1 = 0, \tag{1}$$

where $\alpha_j = 2 \cos \left(\frac{2\pi j}{n} \right)$, $\beta_j = 2 \cos \left(\frac{2\pi jk}{n} \right) = 2T_k(\alpha_j/2)$ ($0 \leq j \leq n - 1$) are the eigenvalues of C , respectively C_k .

Proof. We first consider the case of $d = \gcd(n, k) = 1$. Since $d = 1$, then C_k is the adjacency matrix of a cycle graph isomorphic to \mathcal{C}_n , and so it is similar to C , that is, there exists a permutation matrix P , such that $P^{-1}C_kP = C$. This implies that the two matrices will have the same eigenvalues and eigenvectors. Then α_j, β_j are eigenvalues corresponding to the same eigenvector, say $\mathbf{v}_j = (1, \zeta_n^j, \dots, \zeta_n^{(n-1)j})^t$. We are looking for an eigenvector for $A(P(n, k))$ of the form $\mathbf{w}_j = (a_j\mathbf{v}_j, \mathbf{v}_j)^t$, where a_j will be determined later. If two distinct values for a_j are to be found, for any $0 \leq j \leq n-1$, then we are done with our search for the eigenvectors/eigenvalues.

With this value for \mathbf{w}_j , we need δ (dependent on j) such that

$$\begin{pmatrix} C_k & I_n \\ I_n & C \end{pmatrix} \begin{pmatrix} a_j\mathbf{v}_j \\ \mathbf{v}_j \end{pmatrix} = \delta \begin{pmatrix} a_j\mathbf{v}_j \\ \mathbf{v}_j \end{pmatrix}$$

and so, we get the system

$$\begin{cases} a_j C_k \mathbf{v}_j + \mathbf{v}_j = \delta a_j \mathbf{v}_j \\ a_j \mathbf{v}_j + C \mathbf{v}_j = \delta \mathbf{v}_j \end{cases} \iff \begin{cases} a_j \beta_j \mathbf{v}_j + \mathbf{v}_j = \delta a_j \mathbf{v}_j \\ a_j \mathbf{v}_j + \alpha_j \mathbf{v}_j = \delta \mathbf{v}_j \end{cases}$$

which implies

$$\begin{cases} a_j(\delta - \beta_j)\mathbf{v}_j = \mathbf{v}_j \\ (\delta - \alpha_j)\mathbf{v}_j = a_j\mathbf{v}_j \end{cases}$$

and so, $(\delta - \beta_j)(\delta - \alpha_j) = 1$, which renders the claim for this case, that is, δ must satisfy the equation $\delta^2 - (\alpha_j + \beta_j)\delta + \alpha_j\beta_j - 1 = 0$.

The case of $d > 1$ is treated similarly. The eigenvectors \mathbf{w}_j must have the form $\mathbf{w}_j = (a_1\mathbf{v}'_j, a_2\mathbf{v}'_j, \dots, a_d\mathbf{v}'_j, \mathbf{v}_j)$, with \mathbf{v}_j as before and $\mathbf{v}'_j = (1, \zeta_n^j, \dots, \zeta_n^{(n'-1)j})^t$, $n' = n/d$, for some appropriate multipliers a_i . A similar system to the one for $d = 1$ case will be obtained and, interestingly enough, the same polynomial whose roots are the eigenvalues λ_i will be found. The theorem is proved. \square

Using the quadratic formula in (1) and simplifying we get the following corollary.

Corollary 2.5 *The eigenvalues of $P(n, k)$ are given by*

$$\cos\left(\frac{2\pi j}{n}\right) + \cos\left(\frac{2\pi jk}{n}\right) \pm \sqrt{\left(\cos\left(\frac{2\pi j}{n}\right) - \cos\left(\frac{2\pi jk}{n}\right)\right)^2 + 1}, \quad 0 \leq j \leq n-1.$$

The largest eigenvalue of $P(n, k)$, $\lambda_0 = 3$, is one of the two values obtained for $j = 0$ in the previous corollary. It is known (see [2, Thm. 3.11]) that if a graph is bipartite, then its spectrum is symmetric with respect to 0. In our case, we have the following result.

Corollary 2.6 *If n is even and k is odd, then the eigenvalues of the bipartite graph $P(n, k)$ are given by ± 3 and*

$$\begin{aligned} & \cos(2j\pi/n) + \cos(2jk\pi/n) \pm \sqrt{(\cos(2j\pi/n) - \cos(2jk\pi/n))^2 + 1} \\ & - \cos(2j\pi/n) + (-1)^k \cos(2jk\pi/n) \mp \sqrt{(\cos(2j\pi/n) + (-1)^k \cos(2jk\pi/n))^2 + 1}, \end{aligned}$$

for $0 \leq j < n/2$.

3 Bounds on the eigenvalues of $P(n, 2)$

In the previous section we found the complete set of eigenvalues of $P(n, k)$ under no restrictions on n and k . Here, we would like to find some bounds on some eigenvalues. Eigenvalue interlacing techniques (see the great survey by Haemers [4] on the topic) will not work easily since there is no visible connection between the various $P(n, k)$, and moreover, the technique is not sensitive enough for our purpose. We shall use a different method.

Here, we will take $k = 2$ and consider $P(n, 2)$ (this includes the case of the classical Petersen graph $P(5, 2)$). Since the second Chebyshev polynomial of the first kind is $T_2(x) = 2x^2 - 1$, we immediately obtain the following:

Theorem 3.1 *The eigenvalues of $P(n, 2)$ are (for $0 \leq j \leq n - 1$)*

$$2 \cos^2(2j\pi/n) + \cos(2j\pi/n) - 1 \pm \sqrt{(2 \cos^2(2j\pi/n) - \cos(2j\pi/n) - 1)^2 + 1}.$$

To find good bounds on the eigenvalues in this case, we look for the extreme points of the two functions

$$f_{\pm}(x) = 2x^2 + x - 1 \pm \sqrt{(2x^2 - x - 1)^2 + 1}, \quad (2)$$

in the interval $-1 \leq x \leq 1$. Certainly, we cannot expect exact or even tight results, in general, since the sequence $\frac{2j\pi}{n}$, $0 \leq j < n - 1$, is finite and therefore, $\cos\left(\frac{2j\pi}{n}\right)$ is not dense in this interval. However, we will have lower

and upper bounds, which is what we are interested in. Since any differentiable function in a compact domain attains its extreme points at either the critical points or on the boundary, we proceed by studying first the functions' critical points:

$$f'_{\pm}(x) = 4x + 1 \pm \frac{(2x^2 - x - 1)(4x - 1)}{\sqrt{(2x^2 - x - 1)^2 + 1}} = 0,$$

has solutions (computed by Mathematica) at $x_1 \sim -0.41100$ (for f_+) and $x_2 \sim -0.65041, x_3 \sim -0.04610$ (for f_-). The values of the corresponding f_{\pm} at these critical points are

$$\begin{aligned} f_+(-0.41100) &= -0.04210\dots \\ f_-(-0.65041) &= -1.92081\dots \\ f_-(-0.04610) &= -2.42092\dots \end{aligned}$$

Further, we look at the values of f_{\pm} at $|x| = 1$. Thus, $f_+(1) = 3, f_+(-1) = \sqrt{5}$, and $f_-(1) = 1, f_-(-1) = -\sqrt{5}$. Certainly, the maximum value is 3, and the minimum value is approximately -2.42092. We sketch in Figure 2 the two functions f_{\pm} , to visualize our analysis from above:




Figure 2: The top function is f_+ and the bottom function is f_-

Every value of f_+ is above every value of f_- , and so the minimum is attained by f_- and the maximum is attained by f_+ . Furthermore, we see

that the second largest eigenvalue of $P(n, 2)$ is

$$\begin{aligned} \lambda_1 &= f_+ \left(\cos \left(\frac{2\pi}{n} \right) \right) \\ &= \cos \left(\frac{2\pi}{n} \right) + \cos \left(\frac{4\pi}{n} \right) + \sqrt{4 \left(2 \cos \left(\frac{2\pi}{n} \right) + 1 \right)^2 \sin^4 \left(\frac{\pi}{n} \right) + 1}, \end{aligned} \quad (3)$$

which increases as n increases (shown simply by using Calculus techniques). For instance, for $3 \leq n \leq 20$ the sequence $\lambda_1 = \lambda_1(n)$ is

$$0, 0.41421, 1., 1.41421, 1.71083, 1.93185, 2.10199, 2.23607, 2.34356, 2.43091, \\ 2.50268, 2.56224, 2.61211, 2.65421, 2.69002, 2.7207, 2.74716, 2.77011.$$

Since $\lim_{n \rightarrow \infty} \cos \left(\frac{2\pi}{n} \right) = 1$, we obtain the next result.

Theorem 3.2 *The eigenvalues of $P(n, 2)$ are*

$$\lambda_0 = 3 > \lambda_1 \geq \dots \geq \lambda_{2n-1} \geq -2.42092.$$

Moreover, the second largest eigenvalue satisfies $\lim_{n \rightarrow \infty} \lambda_1(n) = 3$.

Acknowledgement. The authors are grateful to the referee whose suggestions have led to an improvement in the presentation of this paper. During the preparation of this paper, the authors were partially supported by a RIP grant from NPS.

4 Further comments

All of our results for $P(n, 2)$ can be certainly extended to $P(n, 3)$, $P(n, 4)$, etc., but to find sensitive bounds on eigenvalues for arbitrary GPG $P(n, k)$ does not seem to be easy, since the sequence of the involved Chebyshev's polynomials of the first kind does not have a "controllable" behavior in $|x| \leq 1$.

Also, it would be interesting to investigate the number of and distinct values among the eigenvalues of $P(n, k)$, and that is presumably doable. We suspect that the methods of this paper can be also applied to the I -graphs of [1] or the *supergeneralized* Petersen graphs of [6].

References

- [1] M. Boben, T. Pisanski, and A. Žitnik, I -graphs and the corresponding configurations, *J. Combinatorial Designs* **13:6** (2005), 406–424.
- [2] D.M. Cvetkovic, M. Doob, and H. Sachs, *Spectra of Graphs*, Academic Press, 1980.
- [3] E.R. van Dam and R.E. Kooij, The minimal spectral radius of graphs with a given diameter, *Linear Algebra Appl.* **423** (2007), 408–419.
- [4] W.H. Haemers, Interlacing eigenvalues and graphs, *Linear Algebra Appl.* **226–228** (1995), 593–616.
- [5] T.J. Rivlin, *Chebyshev Polynomials*, New York: Wiley, 1990.
- [6] M.L. Saražin, W. Paccob, and A. Previtali, Generalizing the generalized Petersen graphs, *Discrete Math.* **307** (2007), 534–543.
- [7] D. West, *Introduction to graph theory*, 2nd Edition, Prentice Hall, Upper Saddle River, NJ, 2001.