

2010 Annual Report

PI: Anireddy S.N. Reddy

Organization: Colorado State University

ONR Award Number: N000140810470

Award Title: Metabolic Engineering of Plants to Produce Precursors (Phloroglucinol and 1,2,4-butanetriol) of Energetic Materials

A. Scientific and Technical Objectives

Current chemical-based synthesis of energetic materials uses toxic ingredients and produces many environmentally undesirable wastes. In addition, chemical-based production is not sustainable because many of the chemicals used in making these precursors are petroleum-based. An alternative strategy is to produce precursors of energetic materials using a bio-based approach. Recently, significant progress has been made in identifying the genes and enzymes in microbes that can produce precursors of energetic materials. This knowledge can now be exploited to use plants for producing precursors of energetic materials in a sustainable manner. Plants offer many advantages in producing energetic ingredients. These include sustainability as they are renewable sources, likely to be highly cost-effective as they use light energy and CO₂ to produce sugars and other organic compounds that can be converted into desirable carbon compounds. The focus of our project is to use bacterial enzymes to engineer plant metabolism to produce precursors of energetic materials. The specific objectives of this proposal are: 1) Metabolic engineering of plants to produce high levels of phloroglucinol by introducing a bacterial gene that converts malonyl Co-A to phloroglucinol into plants. 2) Metabolic engineering of plants to produce butanetriol. This will be accomplished by introducing bacterial genes involved in butanetriol synthesis from xylose and arabinose. 3) To develop regeneration and robust stable transformation technologies for *Miscanthus* to introduce and express genes involved in synthesis of energetic materials.

B. Approach

Our approach is to express bacterial genes in plants to produce phloroglucinol and 1,2,4-butanetriol. Most plants naturally produce the precursors of these chemicals but not the enzymes to convert them to the precursors of energetic materials. However, bacteria have the enzymes that convert these precursors into end products. We will clone these bacterial genes, transfer them into plant transformation vectors under constitutive promoters and generate transgenic lines. The genes to be introduced are: xylose dehydrogenase, xylonate dehydratase, arabinose dehydrogenase, arabinonate dehydratase, benzoylformate decarboxylase, dehydrogenase and *PhlD*. Because of many available resources for *Arabidopsis*, we are using this plant as a model system to test this concept. Once proven, these constructs will be introduced into *Miscanthus*, a non-food crop that is known to grow well on marginal soils with minimum or no inputs. Transformation of plants will be done using *Agrobacterium*-based binary vectors using floral-dip method (in case of *Arabidopsis*) and regeneration of plants in tissue culture after co-cultivation with *Agrobacterium* or after particle bombardment (in case of *Miscanthus*). Most of our initial work will be performed in *Arabidopsis* and the best gene constructs will be transferred to *Miscanthus*, a grass species.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2010		2. REPORT TYPE Annual		3. DATES COVERED -	
4. TITLE AND SUBTITLE Metabolic Engineering of Plants to Produce Precursors (Phloroglucinol and 1,2,4-butanetriol) of Energetic Materials				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Colorado State University				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 8	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

We will be using a sterile hybrid of *Miscanthus* for these studies. Propagation of this variety is done through rhizomes.

C. Concise Accomplishments

We cloned and sequence verified and/or synthesized all genes needed for this work. These include *PhlD*, xylose dehydrogenase, xylonate dehydratase, arabinose dehydrogenase, L-arabinonate dehydratase and benzoylformate decarboxylase) that are necessary to produce these chemicals in plants. We then cloned all eleven genes (bacterial and synthetic) into plant transformation vectors, each with a different constitutive promoter. Arabidopsis transgenic lines were generated with *PhlD* for phloroglucinol production and xylose and arabinose pathway genes for the production of D- butanetriol and L-butanetriol, respectively. We obtained several independent transgenic plants expressing these constructs. Expression of all introduced genes was verified in T1 lines. We are in the process of obtaining homozygous lines of all these transgenic lines. Arabidopsis plants transformed with the *PhlD* gene are being analyzed for phloroglucinol production. Once we establish that we can produce these chemicals in Arabidopsis, these genes will be then introduced into *Miscanthus*, a non-food crop that is known to grow well on marginal soils with minimum or no inputs thus providing a renewable and less costly resource for the production of precursors of energetic material. Towards this goal, we are developing transformation methods for *Miscanthus*. We have done over three hundred hormonal combinations to optimize callus induction from *Miscanthus* organs. Of all the organs tested, roots from tissue culture grown plants and inflorescence yielded callus induction with 80-90% efficiency. We were able to regenerate whole plants from inflorescence calli with over 90% efficiency. We are now using the inflorescence calli to develop transformation methods for *Miscanthus* using *Agrobacterium* and gene gun.

D. Expanded Accomplishments

Objective 1: Metabolic engineering of plants to produce high levels of phloroglucinol by introducing a bacterial gene that converts malonyl Co-A to phloroglucinol into plants.

We cloned *PhlD* from *Pseudomonas fluorescens* and introduced into two entry vectors Impact vector 1.1 and pSAT4A. The entry vectors contain promoters and terminators for expression of the genes in plants. The promoter, gene, terminator cassette was digested from the entry vectors and cloned into binary vectors pRCS2-nptii and pBin. Both binary vectors were used to transform *Agrobacterium* and those transformants were then used to transform Arabidopsis plants. Seeds were collected from the transformed plants and were grown on kanamycin containing media. Transformants were obtained from the *PhlD* pRCS2-nptii construct (See Figure 1) and plants were transferred to soil (See Figure 2). Tissue from the leaves of transformants was analyzed for expression of the mRNA for *PhlD*. Expression was confirmed in different lines (Figure 3). These plants are selfed to generate homozygous lines. We recently obtained homozygous lines, which we are in the process of analyzing for the presence of phloroglucinol. In addition, we have synthesized *PhlD* gene to optimize for codon usage in Arabidopsis and generated transgenic lines with synthetic *PhlD*. Several lines expressing synthetic *PhlD* are identified by RT-PCR (see Figure 4).




Figure 1. Left: Selection of transformants from PhlD transformation. Right: Transformants obtained with PhlD driven by 35S promoter are grown on kanamycin plates.




Figure 2. Transgenic Arabidopsis lines carrying PhlD gene driven by 35S promoter.




Figure 3. Expression of *PhlD* in transgenic lines. **Top:** RT-PCR analysis showing the levels of *PhlD* transcript in four transgenic lines and wild type (WT). C, positive control. **Bottom:** Northern blot. Numbers 1 to 15 are transgenic lines in which *PhlD* is driven by 35S promoter, B, transgenic line in which *PhlD* is driven by rubisco promoter. Wt, wild type.




Figure 4. Expression of synthetic *PhlD* in transgenic lines. RT-PCR analysis showing the levels of *PhlD* transcript in twelve transgenic lines (indicated with numbers) and wild type (W).

Objective 2: Metabolic engineering of plants to produce butanetriol. This will be accomplished by introducing bacterial genes involved in butanetriol synthesis from xylose and arabinose.

Xylose dehydrogenase (XDG) was cloned from *Calobacter crescentus* and cloned into pSAT3A, xylonate dehydratase (XDT) was cloned from *E. coli* and cloned into pSAT4A, benzoylformate decarboxylase (BFD) was cloned from *Pseudomonas putida* and cloned into pSAT6A. Arabinose dehydrogenase (AD) and arabinonate dehydratase (ADT) were constructed synthetically based on the genes from *Burkholderia vietnamiensis* G4 and *Pseudomonas fluorescens* and were cloned into pSAT3A and pSAT4A, respectively. Xylose dehydrogenase (XDG), XDT, and BFD were also synthesized. The synthetic genes have altered codons that are optimized for expression in *Arabidopsis*. The pSAT vectors have the promoters and terminators for expression of the genes in plants.

To construct the binary vectors with XDG XDT, BFD (bacterial or synthetic) and AD, ADT, BFD (synthetic genes) each gene was cloned sequentially into a promoter vector and then moved the promoter with the gene into a plant transformation vector and verified by PCR. Verification of all constructs in plant transformation vectors is shown in Figures 5 to 7. We have transformed plants with all three constructs. T₀ seeds for xylose pathway with bacterial genes are obtained. For xylose pathway with synthetic genes as well as arabinose pathway with synthetic genes, transgenic lines were obtained and expression of introduced genes at the mRNA level was verified using RT-PCR (see Figures 8 and 9)

3



2

Clone +Con
G T B G T B




Figure 5. Left: Schematic diagram showing cloning of three bacterial genes of xylose pathway into a binary vector. 1. Inserted XDT into pSAT4A. Transferred cassette to pRCS2-ocs-bar; 2. Inserted BFD into pSAT6A. Transferred cassette to pRCS2-ocs-bar/XDT; 3. Inserted XDG into pSAT3A. Transferred to XDT/BFD/pRCS2-ocs-bar. **Right:** PCR verification of XDG, XDT and BFD in pRCS2-ocs-bar (Clone); +Cont, PCR from plasmids; G- XDG, T- XDT, B- BFD,




Figure 6. Left: Schematic diagram showing cloning of synthetic genes of xylose pathway into a binary vector. 1. Inserted XDT into pSAT4A. Transferred cassette to pRCS2-ocs-bar; 2. Inserted BFD into pSAT6A. Transferred cassette to pRCS2-ocs-bar/XDT; 3. Inserted XDG into pSAT3A. Transferred to XDT/BFD/pRCS2-ocs-bar. **Right:** PCR verification of XDG, XDT and BFD in pRCS2-ocs-bar (Clone).




Figure 7. Left: Schematic diagram showing cloning of synthetic genes of arabinose pathway into a binary vector. Arabinose dehydrogenase (ADsyn), Arabinonate dehydratase (ADTsyn), Benzoylformate decarboxylase (BFDsyn) genes in a binary vector. **Right:** PCR verification of Arabinose dehydrogenase (A), Arabinonate dehydratase (T), Benzoylformate decarboxylase (B) in the construct.




Figure 8. Left: Transgenic plants containing three synthetic genes of xylose pathway **Right:** RT- PCR verification of expression of all three genes using gene-specific primers in three plants (7, 8 and 9)




Figure 9. Left: Transgenic plants containing three synthetic genes of arabinose pathway
Right: RT- PCR verification of expression of all three genes using gene-specific primers in two plants (11B, 14B).

Objective 3. To develop regeneration and robust stable transformation technologies for Miscanthus to introduce and express genes involved in synthesis of energetic materials.

Different plant parts (roots, leaves and inflorescence) from *Miscanthus* (*Miscanthus x giganteus*, a triploid species) were used for callus induction (Figure 10). We have tried 126 hormonal combinations for each tissue. Leaves did not produce callus in any hormonal combinations. Roots and inflorescence produced good calli with 80-90% frequency under a few hormonal combinations (Figure 11). We have developed methods for regeneration of plants from inflorescence callus with over 90% efficiency (see Figure 12)




Figure 10. Induction of callus from leaves (left), roots (middle) and inflorescence (right)




Figure 11. Calli induction from roots (left) and propagation of root calli (middle). Magnified view of two calli (right).




Figure 12. Regeneration of plants from callus derived from inflorescence. **From left to right:** 1) shoot induction, 2) close-up view of regenerating shoots, 3) root induction 4) regenerated plant.

We have constructed a fluorescent reporter fused nuclear protein (SR1) for transformation of *Miscanthus* (Fig. 13).

QuickTime™ and a
TIFF (Uncompressed) decompressor
are needed to see this picture.

YFP Light

Figure 13. Verification SR1-YFP construct. The fusion protein expression is verified in transgenic *Arabidopsis* plants (left, YFP). Bright field image is on the right (light).

E. Work Plan:

We will analyze bacterial and synthetic PhLD expression lines for the production of phloroglucinol. We will obtain homozygous lines for xylose pathway (for bacterial as well as synthetic genes) and arabinose pathway by screening T2 seeds. We will develop methods to detect and quantify D- and L-butanetriol from plant extracts. Homozygous lines expressing the xylose and arabinose pathway will be analyzed for the production of butanetriol. We will perform metabolomics analysis of transgenic lines of containing each construct. We will use the optimized regeneration conditions for transformation using fluorescent reporter systems. Transformation with different strains of *Agrobacterium* as well as direct gene transfer using gene gun method are being used. Once we establish robust transformation system we will then introduce all the constructs into *Miscanthus*.

F. Major Problems/Issues

No major problems were encountered. We were unable to clone the arabinose dehydrogenase and arabinonate dehydratase genes from the bacteria by PCR. We used these genes that are synthetically constructed, which are optimized for expression in Arabidopsis.

The cloning of the genes from the pSAT vectors into the pRCS2 vectors has proven to be a challenge. Due to the large size of the vectors, especially when they contained the cloned genes, cloning has been difficult. Different enzyme strategies have had to be developed and optimization of cloning has taken a longer time than expected. Furthermore, we also encountered problems with stability of large constructs in bacterial cells. Using different strains we solved this problem.

Callus induction from inflorescence took a while as it took 8 months for Miscanthus plants to flower. Because of this it has taken a long time to obtain calli from flowers.

G. Technology Transfer

No technology transfer as of now.

H. Foreign Collaborators and Supported Foreign National

Dr. Gul Shad Ali

Dr. Watanabe, Shizuoka University, Japan