

# **Linking Short and Long Term Sediment Delivery to Morphology and Acoustic Properties of Continental Margins**

James P.M. Syvitski  
INSTAAR Univ. of Colorado  
Boulder CO, 80309-0450  
phone: (303) 492-7909 fax: (303) 492-6388 email: [james.syvitski@colorado.edu](mailto:james.syvitski@colorado.edu)  
Robert Courtney  
Geological Survey of Canada-Atlantic  
BIO, Dartmouth NS, Canada, B2Y 4A2  
phone: (902) 426-5062 fax: (902) 426-6152 email: [courtney@agc.bio.ns.ca](mailto:courtney@agc.bio.ns.ca)

Award #: N00014-95-1-1281  
ASSERT: N00014-96-1-0994, DURIP: N00014-97-1-0380

## **LONG-TERM GOAL**

Develop numerical models useful for the simulation of sediment delivery and accumulation on continental margins over time scales of tens to thousands of years. Model predictions will help us understand the evolution of the sediment characteristics on continental margins through sea level fluctuations, climate change, and other relevant environmental factors. ONR interests include the development of a numerical predictor of the acoustic signature of remote margins based on a region's geological history.

## **OBJECTIVES**

- 1) Is the rate and direction of sealevel change coupled with hinterland climate? Does the evolving landscape control the dynamics of river plumes, and thus the physical properties of depositing sedimentary units within the slope environment?
- 2) Is there a separation of sand from mud during a river flood, with the mud fraction deposited on the outer shelf and upper continental slope?
- 3) Do closely-spaced river mouths contribute to plume intermingling such that a line source is created, particularly during low sea level stands?
- 4) Could a short lived hyperpycnal current develop during the flooding of the Eel River and might this explain the missing sediment mass observed in the STRATAFORM project.

## **APPROACH**

- Task B3) Develop a model of how sediment is delivered to, and accumulates on, a continental slope. Predict delivery to the shelf-slope break from fluvial point and line sources. Compare open-slope sediment delivery (distributed) with canyon-fed delivery (localized).
- Task B8) Model major influences (climate, sea-level, tectonics) on processes that control slope morphology, stratigraphy, and acoustic realizations. Model(s) to include the effects of: a) external forcing mechanisms; b) short-term and long-term delivery of sediment; c) 2-D simulations of individual mass movements (submarine slides, debris flows and turbidity currents); d) excess pore pressure on slope failure.

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>Linking Short and Long Term Sediment Delivery to Morphology and Acoustic Properties of Continental Margins</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of Colorado, INSTAAR, Boulder, CO, 80309</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## WORK COMPLETED

- evaluated USGS hydrology data of the Eel River (dynamics and trends),
- developed and verified a method to predict the rating coefficients of river loads from first principle,
- developed a world river data base of drainage basin and climate characteristics, hydrological parameters, long term sediment loads and yields,
- developed a method for predicting loads and yields in ungauged river basins,
- developed a 3-D model architecture (3D SEDFLUX) able to simulate multiple river sources from the coastline,
- developed transport models of both hyperpycnal (turbidity current) (INFLO) and hypopycnal (surface) (PLUME) events,
- developed a predictive understanding of the frequency of hyperpycnal flows generated during flood events of rivers and through sea level cycles,
- included the effects of coastal currents on plume dynamics on model simulations.
- completed the construction (DURIP funds) of a state-of-the-art Flocc Camera Assembly (with 250 engineered components).
- developed a pseudo-2D Eulerian model (INFLO) able to simulate an individual hyperpycnal event flowing across a continental margin at the dynamic level (1s time step) including erosion and deposition of a multigrain size load. The model was verified against observations on Saguenay Fjord and Var River events.
- developed an advanced pseudo-2D turbidity current model that has 'failed' sediment move from a debris flow to a dense turbidity current and eventually develop into to a turbidity current plume, Model was verified against Var data.
- developed a 2D multiprocess SEDFLUX model that combines the dynamics of ten individual processes that act in concert to shape continental margin. The model includes new dynamics for debris flow and turbidity currents, and storm-influenced diffusion.
- applied the 2D SEDFLUX model to the Eel River margin with runs of 2500 years assuming starting conditions of present bathymetry, modern Eel River discharge, and sea level fluctuations using observations from historical tide gauges.
- calibrated the synthetic seismic model (ECHO) with data on cores obtained from Emerald Basin, N.S.
- compiled a data base of acoustic parameters that extends modeling to a wider range of margin lithologies,
- applied seismic model (ECHO) to transport model predictions based on the dynamics of the Eel River discharge and compared results with existing seismic data sets.

## RESULTS

We now have models for simulating the discharge and the sediment load of ungauged rivers flowing into the ocean. Rating coefficients relate to the annual mean discharge, long term sediment yield and latitude (a parameter that captures the both frozen soil conditions, and snow cover duration). HYDROTREND results compare favorably with USGS observations on the Eel River, including our ability to predict the 400 yr return interval (e.g. 1964) flood of the river. With such a robust model we are able to simulate past or future climate effects on the production and delivery of sediment to the ocean. We have concluded from numerical experiments that fjord basins are able to capture, in terms of textural proxies, both the natural variability associated with precipitation and temperature as well as realistic scenarios of

abrupt climate change. Open ocean basins, like the Eel River, are less likely to record the proxy record of ambient climate variability, largely since the natural climate variability is strongly dominated by infrequent and large events that do not track climate events. PLUME can now be coupled with offshore buoy data (wind and currents), to simulate the basic features of coastal plumes under flood conditions. We predict that hyperpycnal flows will occur every 10 or so years off the Eel River, lasting for a few hours to a day. Longer return-interval floods will deliver hyperpycnal flows that could last 2 to 3 days. These events will deliver much sediment directly to the continental slope and could account for the missing sediment from surface plumes.

## **IMPACT/APPLICATION**

We are on the path to predict acoustic properties of the seafloor of continental margins, based on process based modeling and remote data input (i.e. satellites). Realistic numerical models offer the possibility of making predictions where field data is limited. The impact of floods and or storms on the acoustic character and features of continental margins can then be examined at will. SEDFLUX model is becoming an integral part of the reservoir characterization effort of MOBIL Technology Corporation. HYDROTREND and PLUME are models being considered by Raytheon-Hughes Corporation to fuse environmental data records (infrared and visible wavelengths) downloaded from satellites for the purpose of natural disaster mitigation efforts. A white paper is in place that would allow INSTAAR numerical models, such as HYDROTREND and PLUME to link with a variety of DSMP satellite data with other NOAA-generated microwave-meteorological models to provide environmental predictions in aid of the health of our coastal zone.

## **TRANSITIONS**

Our models have been shared with Old Dominion Observatory (Swift et al.) who have incorporated them into their own modeling efforts. Model runs of the Eel River Margin have been made available to USGS participants (Field, Lee). Mike Stekler (LDEO) helped us develop a predictive subsidence subroutine. INSTAAR models have been transferred to MOBIL Technology Center. We have helped Lincoln Pratson develop a 1-D Lagrangian turbidity current model (BANG). Our seismic simulator (ECHO) has been used in cooperative work with St. Anthony Falls Hydraulic Lab's flume efforts. Through our efforts, STRATAFORM has been accepted as a core project of the world IGBP-LOICZ (Land-Ocean Interaction Coastal Zone) program that includes participants from 80 countries. We have implemented Gary Parker's 1-D Lagrangian debris flow model (BING) into our 2D SEDFLUX seascape-evolution model. Provided Craig Fulthorpe (U Texas) and Greg Mountain (LDEO) an opportunity to have STRATAFORM multi-channel seismic data cleaned up by advanced industry algorithms (MOBIL Technology). Worked with Paul Hill (Dalhousie U.) to develop an advanced floc formation model using data obtained from Yukatat Bay Alaska.

## **RELATED PROJECTS**

- MOBIL Technology Center (Cullick, Sarg, Gouveis, Levin) supported INSTAAR efforts (Bahr, Pratson, Hutton, O'Grady) to develop a data base on continental margin morphology, sedimentology, oceanography and tectonics. Parameters are being linked into super-variables to determine the effective influence of past history, ocean energy, seismic energy and sediment delivery on continental margin morphology and deposit architecture. The two STRATAFORM margins are included in the data base.

- Work with G. Parker, C. Paola (U. Minneapolis), P. Heller (U. Wyoming), L. Pratson (Duke U.) in developing a continental margin experimental tank, through the support of a NSF grant and consortium support of oil companies (MOBIL, EXXON, TEXACO, CONOCO, JAL). It is anticipated that one of the experimental runs would be on a STRATAFORM margin.
- Raytheon-Hughes has linked with the INSTAAR modeling efforts to use STRATAFORM and other models for the development an algorithm related to fuse satellite data (infrared and visible wavelengths) and provide information on littoral sediment transport.

## REFERENCES

- Mulder, T., Syvitski, J.P.M. and Skene, K. 1997. Modelling of erosion and deposition by sediment gravity flows generated at river mouths. *Jour. Sedimentary Research*, 67(3): 571-584.
- Mulder, T., Savoye, B. and Syvitski, J.P.M. 1997. Numerical modelling of the sediment budget for a mid-sized gravity flow: the 1979 Nice turbidity current. *Sedimentology*, 44: 305-326.
- Syvitski, J., Pratson, L., Perlmutter, M., de Boer, P., Parker, G., Garcia, M., Wiberg, P., Steckler, M., Swift, D., and Lee, H.J., 1997. EARTHWORKS: A large scale and complex numerical model to understand the flux and deposition of sediment over various time scales. In: V. Pawlowsky-Glahn (Editor) *Proceedings of IAMG'97 The third annual conference of the International Association of Mathematical Geology*. CIMNE-Barcelona, 1997, v. 3 p. 29-33.
- Mulder T., Savoye B., Syvitski J.P.M. and Piper D.J.W. 1997. The Var Submarine Sedimentary System: understanding Holocene sediment delivery processes and their importance to the geological record. In M. S. Stoker, D. Evans, A. Cramp (editors): *Geological Processes on Continental Margins: Sedimentation, Mass Wasting and Stability*, Geol. Society, Spec. Publ., London, 145-166.
- Syvitski, J.P.M. and Hutton, E.W.H., 1997. FLOC: Image analysis of marine suspended particles. *Computers and Geoscience*, v. 23(9): 967-974.
- Skene, K., Mulder, T., and Syvitski, J.P.M. 1997. INFLO1: A model predicting the behaviour of turbidity currents generated at a river mouth. *Computers and Geoscience*, v. 23(9): 975-991.
- Wang Y., Ren, M.-e and J.P.M. Syvitski. 1998. Sediment Transport and Terrigenous Fluxes. In: K.H. Brink & A.R. Robinson (editors) *The Sea: Volume 10 - The Global Coastal Ocean: Processes and Methods*. John Wiley & Sons, New York, p. 253-292.
- Syvitski, J.P.M., Morehead, M. and Nicholson, M. 1998. HYDROTREND: Simulation of trends in river discharge and sediment transport from climate normal and GCM predictions. *Computers and Geoscience* v. 24(1): 51-68.
- Hill, P.; J P Syvitski, R D Powell, E A Cowan. 1998. Floc settling velocities under a buoyant discharge plume in Glacier Bay, Alaska. *Marine Geology* v.145 (1-2): p. 85-94.
- Syvitski, J.P.M., Nicholson, M., and Skene, K., Morehead, M.D. (in press). PLUME1.1: Deposition of sediment from a fluvial plume. *Computers and Geoscience*.
- Skene, K., Piper, D.J.W., Aksu, A.E., and Syvitski, J.P.M. (in press). Evaluation of the global oxygen isotope curve as a proxy for Quaternary sea level by modeling of delta progradation. *Journal of Sedimentary Research*.
- Syvitski, J.P. and Morehead, M.D. (in press). River predictions for oceanographers and paleoceanographers: Application to the Eel River Margin, Northern California. *Marine Geology*.
- Syvitski, J.P.M., Pratson, L. O'Grady, D. (in press) Stratigraphic Predictions of Continental Margins for the Navy. In: L. Whatney et al (eds.) *Numerical Experiments in Stratigraphy*. SEPM special publication.

- Perlmutter, M.A., DeBoer, P.L., and Syvitski, J.P. (in press) Geological Observations and Parameterizations. In: L. Whatney et al (eds.) Numerical Experiments in Stratigraphy. SEPM special publication.
- Morehead, M.D., and Syvitski, J.P. (in press) River plume sedimentation modeling for Sequence Stratigraphy: Application to the Eel River Margin, N. California. Marine Geology.
- Bahr, D., Hutton, E.W.H., Syvitski, J.P., and Pratson, L. (in review) Compaction: a theoretical analysis from first principle. Basin Research
- Syvitski, J.P.M., Bahr, D., Mulder, T., (in review). Sediment Load Dynamics: Predicting the Rating Coefficients. Water Resources Journal.

### **Conference Abstracts & Proceedings**

- Syvitski, J.P. (1997). Scaling Issues in Modeling Coastal Systems and Fluxes. In Abstracts of the Land-Ocean Interaction in the Coastal Zone (LOICZ) Open Science Meeting Oct 10-13, 1997 in Noordwijkerhout, The Netherlands.
- Syvitski, J.P. (1997). The STRATAFORM Project: Understanding Coastal Sediment Input, Distribution and Controls through Global Data Bases and Predictive Modeling. In Abstracts of the Land-Ocean Interaction in the Coastal Zone (LOICZ) Open Science Meeting Oct 10-13, 1997 in Noordwijkerhout, The Netherlands.
- Syvitski, J.P., Pratson, L., and Morehead, M. (1997). EARTHWORKS: A large spatial scale numerical model to study the flux of sediment to ocean basins and reworking of deposits over various time scales. AGU 1997 Fall meeting EOS supplement, v. 78(46), p. F258.
- Mullin, J. Ellis, C., Mohrig, D., Swenson, J., Paola, C., Parker, G., Syvitski, J.P., Pratson, L., (1997). Experimental study of stratigraphic response to changing base level. AGU 1997 Fall meeting EOS supplement, v. 78(46), p. F277.
- Syvitski, J.P., Hutton, E. W.H., Morehead, M.D., Pratson, L., O'Grady, D., Bahr, D., Sarg, R. (1998) Numerical Tools for Stratigraphers. SEPM-IAS Research Conference: "Strata and Sequences on Shelves and Slopes, Sept. 15-19, Sicily, Italy.
- Pratson, L., Gouveia, W., Courtney, R., Syvitski, J.P., Paola, C., and Parker, G. (1998) Constraining the stratigraphic information content of seismic data. SEPM-IAS Research Conference: "Strata and Sequences on Shelves and Slopes, Sept. 15-19, Sicily, Italy.
- Bahr D., Syvitski J. (1998) Predicting sediment stratigraphy on continental margins with limited data. In: Proceedings of IAMG'98 The fourth annual conference of the International Association of Mathematical Geology. Italy, 1998,
- Syvitski, J.P., (1998) Understanding Sedimentary Processes and Palaeoenvironments in Fjords: A Personal 25-Year Journey, In: International Workshop on Sedimentary Processes and Paleoenvironment in Fjords, University of Tromso, April 22-24, 1998.
- Syvitski, J.P., Meade, R.H., and Bobrovitskaya, N.N. (1998) Water and sediment discharge dynamics of Russian Arctic rivers. Program and Abstracts of the 28th Arctic Workshop, March 12-14, 1998. p. 145-148.
- Syvitski, J.P., (1998) The link between abrupt climate change and basin stratigraphy: A numerical approach. International Conference of Recognition of Abrupt Climate Change in Clastic Sedimentary Environment: Methods, Limitations and Potential. Program and Abstracts, June 8-10, 1998, Stockholm, Sweeden, p. 11-13.