

Upper Ocean Mixing, Coastal Mixing, LIWI, & CM AASERT

Michael C. Gregg, PI
University of Washington
Applied Physics Laboratory
1013 NE 40th St.
Seattle, WA 98105-6698

Phone: (206)543-1353; fax (206)543-6785; email: gregg@apl.washington.edu
Award #: N00014-94-1-0079, N00014-95-1-0406, N00014-97-1-0136, &
N00014-97-1-0647

Jack B. Miller, Co-PI
University of Washington
Applied Physics Laboratory
1013 NE 40th St.
Seattle, WA 98105-6698

Phone: (206) 543-9958; fax (206) 543-6785; email: miller@apl.washington.edu
Award #: N00014-94-1-0079, N00014-95-1-0406, N00014-97-1-0136, &
N00014-97-1-0647

LONG-TERM GOAL

My long-term goals are to identify and quantify the major mixing processes in the upper ocean and to relate them to the larger-scale processes producing the mixing as quantitatively as possible. When successful, this will take the form of parameterizations that can be used in numerical models.

OBJECTIVES

My scientific objectives are to measure mixing directly with microstructure sensors and to relate these measurements to the larger scales producing the mixing in such a way that the results can be compared with either the large-scale changes resulting from the mixing or with theoretical predictions of mixing rates. My technological objectives are to develop instruments and sensors to measure the major mixing parameters.

APPROACH

Because a first-order understanding of mixing in the open ocean is rapidly being developed, our approach has shifted to obtaining a similar understanding of mixing near coasts and in estuaries. Because dissipation rates are higher in these waters, we have also shifted our technological developments to improve the spatial resolution of microstructure sensors and to adapt open-ocean measurements of finescale velocity to shallow water.

WORK COMPLETED

Emin Ozsoy, our Turkish colleague, visited us in September 1998 to complete the first paper submitted about our 1994 observations in the Bosphorus. That paper was submitted (Gregg et al., 1998), and we

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Upper Ocean Mixing, Coastal Mixing, LIWI, & CM AASERT				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 NE 40th Street, Seattle, WA, 98195				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

are writing a second for submission late this year. We presented some of these results at a Symposium honoring Konostantin Fedorov in St. Petersburg in May 1998 and they are being published (Ozsoy et al, 1998).

Ren-Chieh Lien is the senior author on a draft manuscript analyzing observations of elevated shear and turbulence in an internal tidal beam observed on the continental Slope near Monterey, California in 1997 as part of the Littoral Internal Waves Initiative (LIWI).

All of the data from both cruises of the Coastal Mixing and Optics program have been processed and presented at the 1998 Ocean Sciences Meeting and the CMO workshop in Colorado in September 1998. Cd roms have been given to several of the CMO PIs and Jennifer MacKinnon is working on the analysis for her Ph.D. dissertation.

Jody Klymak has analyzed the three-dimensional structure of flow over the Knight Inlet sill for his Ph.D. dissertation and is now considering the dynamical implications of the results.

RESULTS

We demonstrated that at some times flow in the Bosphorus is strongly quasi-steady. During those times it makes a good natural laboratory for testing steady hydraulic theory. We were also able to estimate the surface slope along the strait from the observed mean density and velocity fields. As seen in Fig. 1, the surface slopes most steeply where the zero-flow surface dips, which is in the primary contraction of the strait.




Fig. 1. Observed pressure of noflow (lower) and surface (upper). The vertical line shows the range and mean sea level difference along the Bosphorus. The primary drop in surface height occurs between 10 and 15 km from the southern entrance, where the strait is narrowest.

The LIWI observations give the best evidence we know of to date for mixing in a tidal beam. Microstructure measurements were taken continuously along a fan ridge for five semidiurnal tidal periods. We observed a beam of enhanced turbulence mixing emanating from the shelf-break, extending for about four km (Fig. 2). The vertical scale of the beam is about 50 m. The average turbulence dissipation rate ϵ within the beam is about $10^{-6} \text{ W kg}^{-1}$. The beam of enhanced turbulence follows the slope of characteristics of semidiurnal internal tide. Strong turbulence mixing also exists within O (100 m) off the bottom on the continental slope.




Fig. 2. Contour of ϵ averaged over the along-ridge sections. Two dashed lines denote the average ship tracks along and across the fan ridge. Two solid lines show the characteristics of semidiurnal internal tides.

IMPACT/APPLICATION

Demonstration of low mixing rates in the upper thermocline of the open ocean by this project during previous years is leading many modelers to examine the role of mixing in their models and to develop models with more realistic mixing.

Our demonstration of low mixing rates most of the time during CMO is leading the optical and biological investigators to change their assumptions about how optical features evolve and to examine the role of advection.

TRANSITIONS

Development of a thinistor for measuring temperature microstructure, is funded by NSF with Prof. Fumio Ohuchi, University of Washington, Dept. of Materials Science and Engineering, based on preliminary work funded by ONR.

Observations of Mixing in the Banda Sea, a project funded by NSF, will use instruments developed with ONR funding to make the first mixing measurements in Indonesian waters in October 1998.

RELATED PROJECTS

1. Internal Waves and Mixing in Monterey Canyon with Eric Kunze and Leslie Rosenfeld, funded by NSF.
2. Hawaii Ocean Mixing Experiment (HOME) proposed to NSF by many investigators.

REFERENCES

Gregg, Michael C., Emin Özsoy and Mohammed A. Latif (1998), Quasi-steady exchange flow in the Bosphorus, *Geophys. Rev. Lett.*, submitted.

Ozsoy, Emin, M.C. Gregg, H. Fohrmann, and J. Backhaus (1998), The Bosphorus Strait and its shelf outflow in to the Black Sea: Experiments and Modelling, *Federov Symposium*, submitted.

PUBLICATIONS

Gregg, M.C. (1998), Uncertainties and limitations in measuring ε and χ_T , *J. Atmos. Ocean. Tech.*, in press.

Gregg, M. C. (1998), Estimation and geography of diapycnal mixing in the stratified ocean, in *Proceedings of the IUTAM Conference on Physical Limnology, Broome, Australia, 10-14 Sept., 1995*, edited by J. Imberger, in press.

Gregg, M.C. (1998), Turbulent mixing in the ocean, *Federov Symposium*, submitted.

Gregg, Michael C., Emin Özsoy and Mohammed A. Latif (1998), Quasi-steady exchange flow in the Bosphorus, *Geophys. Rev. Lett.*, submitted.

Ozsoy, Emin, M.C. Gregg, H. Fohrmann, and Jan Backhaus (1998), The Bosphorus Strait and its shelf outflow in to the Black Sea: Experiments and Modelling, *Federov Symposium*, submitted.

Seim, H.E., D.P. Winkel, G. Gawarkiewicz, and M.C. Gregg (1998), A benthic front in the Straits of Florida and its relationship to the structure of the Florida Current. *J. Phys. Oceanogr.*, submitted.

Winkel, D. P. (1998), Influences of mean shear in the Florida Current on turbulent production by internal waves, Ph.D. dissertation, *Univ. Wash. Press*, 137 pp.