

2011 Military Health System Conference

Findings of VA/DoD CPG on CAM Therapies for PTSD

The Quadruple Aim: Working Together, Achieving Success

Miguel Roberts, Ph. D., COL Charles Engel, M.D., Elizabeth Harper Cordova, M.A., Wayne Jonas, M.D.

January 26, 2011



Defense Centers of Excellence
for Psychological Health and Traumatic Brain Injury

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 26 JAN 2011	2. REPORT TYPE	3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Findings of VA/DoD CPG on CAM Therapies for PTSD		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Defense Centers of Excellence for Psychological Health & Traumatic Brain Injury, 2345 Crystal Drive, Crystal park 4, Suite 120, Arlington, VA, 22202		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			
13. SUPPLEMENTARY NOTES presented at the 2011 Military Health System Conference, January 24-27, National Harbor, Maryland			
14. ABSTRACT			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)
			18. NUMBER OF PAGES 30
			19a. NAME OF RESPONSIBLE PERSON

Overview & Agenda




- Objective: Review of CAM as Evidenced-Based Treatment for PTSD
 - Background
 - CPG recommended treatments of PTSD
 - Findings of PTSD VA/DoD CPG Working Group on CAM
 - Discussion / Q&A

Significant Findings



- PTSD Prevalence
 - Clinically diagnosed cases in OEF/OIF = 2.4%
 - Self-reported surveys of PTSD symptoms
 - Range from 1.4% (not exposed to combat) to 15% (populations exposed to sustained ground combat)
 - US general population = 7% to 8%, lifetime prevalence
 - peak prevalence ages 45-59 (9.2%)
 - Higher rates of PTSD are seen in :treatment seeking samples after discharge (e.g., veterans going to MH)
 - Injured during deployment (e.g., mTBI, wounded)
 - “High risk occupations” (e.g., EOD units), greater degree of combat
- Post-deployment problems wider-reaching than formal diagnoses
 - Suicide, adjustment problems, relationship & family problems, divorce, risky behaviors, etc.

PTSD Prevalence Studies



Complementary and Alternative Medicine



- 2008 CDC report: Increased CAM usage in the US general population
 - Almost 4 out of 10 adults had used CAM therapy in the past 12 months
 - Between 2002 and 2007 increased use was seen among adults for acupuncture, deep breathing exercises, massage therapy, meditation, naturopathy, and yoga

Complementary and Alternative Medicine




- White House commission CAM report (2002)
 - Need for “comprehensive, understandable **summary of current clinical evidence** in CAM for health care practitioners...”
 - Need for “**accurate and easily accessible information on CAM** practices and products”
 - Federally funded health care delivery programs, such as the VA, and **DoD should evaluate the applicability of CAM** wellness and prevention activities to their services.

VA/DoD CPG for the Management of Posttraumatic Stress



- CPG is an update of the 2004 version
- Systematically developed statements to assist practitioner and patient in choosing appropriate health care for specific clinical conditions
 - Actionable recommendations
 - Attempt to incorporate all issues relevant to a clinical question
 - Value judgments--weighting different outcomes, burdens, and costs
 - Guidance where evidence is lacking
 - In VA & DoD, driven by clinical algorithms
- www.healthquality.va.gov
- www.qmo.amedd.army.mil



CPG Assumptions




- Guidelines are developed by multidisciplinary groups
- They are based on a systematic review of the scientific evidence
- Recommendations are explicitly linked to the supporting evidence and graded according to the strength of that evidence

CPG Development Process



- One of the key questions
 - What interventions are effective in treatment of PTSD?
 - Resolution of symptoms and functional outcomes
 - Are any Complementary and Alternative Medicine (CAM) approaches more effective than no intervention?
 - Body-mind
 - Meditation (e.g., zen)
 - Herbal, food supplements
 - Energy (e.g., Reiki)
 - Tai Chi
 - Acupuncture

Level of the Evidence



EVIDENCE HIERARCHY

Strength of Recommendation



A Strongly Recommend to offer or provide...

- There is good evidence that the intervention improves important health outcomes – benefits substantially outweigh harm.

B Recommend to offer or provide...

- There is fair evidence that the intervention improves health outcomes – that benefits outweigh harm.

C Consider offering or providing...

- There is poor evidence that the intervention can improve health outcomes – balance of benefit and harm is too close to justify a general recommendation.

I Insufficient Evidence is to recommend for or against providing...

- Evidence that the intervention is effective is lacking or of poor quality, or conflicting – balance of benefit and harm cannot be determined.

Psychotherapy for PTSD



- **Revised CPG: “Strongly recommend... evidence-based trauma-focused psychotherapeutic interventions that include components of exposure and/or cognitive restructuring; or stress inoculation training [A]”**
- **Best evidence exists for:**
 - Prolonged Exposure (possibly also brief exposure)
 - Cognitive Processing Therapy
 - Stress Inoculation Training
 - EMDR

Psychopharmacology for PTSD



- **Strongly recommend selective serotonin reuptake inhibitors (SSRIs), for which fluoxetine, paroxetine or sertraline have the strongest support, or serotonin norepinephrine reuptake inhibitors (SNRIs), for which venlafaxine has the strongest support, for the treatment of PTSD. [A]**
- **Best evidence exists for:**
 - SSRIs (Fluoxetine, Paroxetine, or Sertraline)
and SNRIs (Venlafaxine)

Complementary and Alternative Medicine



- **Modalities reviewed in the PTSD CPG include:**
 - Body-Mind Approaches (e.g., Yoga, & Tai Chi)
 - RCTs show benefits in other areas (e.g. sleep, stress, anxiety, etc.), BUT no RCTs or comparison trials in PTSD
 - Meditation Training (e.g., zen)
 - Improves sleep, anxiety, and pain, BUT no RCTs in PTSD
 - Exercise (mostly aerobic exercise)
 - Rarely conducted in isolation from other interventions
 - Energy Medicine (e.g., Qi Gung, *Reiki*, *Johrei*)
 - Improvement in comorbid conditions, BUT not RCTs in PTSD
 - Acupuncture

Complementary and Alternative Medicine



- **VA/DoD PTSD CPG Recommendations on CAM:**
 - There is insufficient evidence to recommend as first line treatments for PTSD [I]
 - CAM approaches that facilitate a relaxation response (e.g. mindfulness, yoga, massage) may be considered for adjunctive treatment of hyperarousal symptoms, although there is no evidence that these are more effective than standard stress inoculation techniques [I]
 - May be considered as adjunctive approaches to address some co-morbid conditions (e.g. acupuncture for pain) [C]

Complementary and Alternative Medicine



VA/DoD PTSD CPG Findings on CAM:

- May facilitate engagement in care
- May be considered for some patients who refuse evidence-based treatments
- Providers should discuss the evidence for effectiveness and risk-benefits of different options, and ensure that the patient is appropriately informed

Acupuncture for PTSD and Related Conditions



- PTSD:
 - Hollifield et al. (2007).
 - Engel et al. (manuscript in prep).
- Anxiety, Depression, & Insomnia:
 - Zhang et al. (2010).
 - Leo & Ligot (2007).
 - Blitzer et al. (2004).
 - Spence et al. (2004).
 - Eich et al. (2000).
- Pain:
 - Weidenhammer et al. (2007).
 - Manheimer et al. (2005).
 - Birch et al. (2004).
 - Eshkevari (2003).

Acupuncture for PTSD: A Randomized Trial in a Military Population


Engel CC, Harper Cordova E, Benedek D, Jonas W, Ursano R.



- Two group, parallel arm, RCT in Active Duty service members at WRAMC
 - Group 1: Acupuncture treatment (ACU)
 - Group 2: Usual Care (UC)
- Two 90-min sessions per week for four weeks
- Acupuncturists blinded to study condition
- Follow-up for both groups at baseline, 4-wks, 8-wks, & 12-wks post-randomization
- Primary Outcome Measure: Posttraumatic Stress Disorder Checklist (PCL)



PTSD Symptoms (PCL)



Acupuncture: Conclusions



Compared to usual PTSD care, a four week course of twice weekly TCM acupuncture resulted in significantly greater improvements in...

- PTSD symptoms
pre-post ES 1.4-1.6 versus usual care ES 0.12-0.74
- Depression and pain symptoms
- Mental but not physical health functioning

- **VA/DoD PTSD CPG Recommendations:**
 - Acupuncture may be considered as treatment for patients with PTSD. [B]

Overall Conclusions and Take-Home Message



- Promoting evidence-based treatment ultimately enhances and optimizes treatment outcomes, including knowledge of state of the evidence for CAM modalities for PTSD
- Tools for providers and patients are needed—accurate and unbiased information on CAM is needed
- Use the current gaps in knowledge as a map for future research/improvements



Thank you!



Back-up slides

PTSD Diagnosis



- Exposure to a severely traumatic event involving actual or threatened death, serious injury, or threat to physical integrity of self or others
- Accompanied by fear, helplessness, or horror
- Must have:
 - Reexperiencing symptoms
 - Avoidance symptoms
 - Hyperarousal symptoms

PTSD Diagnosis



- Symptoms must persist for at least one month and cause significant distress or impairment

Guideline Development Process: Evidence Tables and Evidence Rating



Developed by *SUD Working Group* using USPSTF 2001 ratings process

High	I	RCT
High	II-1	Control Trial
Moderate	II-2	Case-Control Study
Moderate	II-3	Uncontrolled Experiment
Low	III	Opinion, Case Reports

Good	<i>High QE</i> → good health outcome
Fair	<i>High QE</i> → intermediate outcome <i>or</i> <i>Moderate QE</i> → good health outcome
Poor	<i>Low QE</i> <i>or</i> No linkage to health outcome

Substantial	> Small Impact
Moderate	Small impact
Small	Negligible impact
Zero/Negative	Negative / No Impact

Quality of Evidence

+

Overall Quality

+

Net Effect of Intervention

**Final
recommendation
grade**

Guideline Development Process: Strength of Recommendation



Developed by *SUD Working Group* using USPSTF 2001 ratings process

	<i>The net benefit of the intervention</i>			
<i>Quality of Evidence</i>	Substantial	Moderate	Small	Zero or Negative
<i>Good</i>	A	B	C	D
<i>Fair</i>	B	B	C	D
<i>Poor</i>	I	I	I	I

Key

- A** *Strong Recommendation* that clinicians provide intervention
- B** *Recommendation* that clinicians provide intervention
- C** *No Recommendation* for or against intervention
- D** *Recommendation Against* providing intervention
- I** *Insufficient Evidence* for recommendation

PTSD Treatment Psychotherapy Balance of Benefit and Harm



SR	Significant Benefit	Some Benefit	Unknown	NO Benefit
A	Trauma-focused psychotherapy that includes components of exposure and/or cognitive restructuring; or stress inoculation training			
B		<ul style="list-style-type: none"> •Imagery Rehearsal Therapy 		
C		<ul style="list-style-type: none"> •Patient Education •Psychodynamic Therapy •Hypnosis •Relaxation Techniques •Group Therapy 		
I		<ul style="list-style-type: none"> •Family Therapy 	<ul style="list-style-type: none"> •Web-based CBT •Dialectical Behavioral 	

PTSD Treatment: Pharmacotherapy



SR	Significant Benefit	Some Benefit	UNKNOWN	No Benefit
A	SSRIs SNRIs			
B		Atypical AP (as adjunct) TCAs MAOIs (phenelzine) Mirtazapine Nefazodone [Caution] Prazosin (sleep/nightmares)		
C			Prazosin (for CORE PTSD)	
D				Benzodiazepines [Harm] Guanfacine Valproate Tiagabine Topiramate
I			Antipsychotics [Harm] Risperidone	

Resilience



Building a Culture of Resilience



← EARLY INTERVENTION



← RECOVERY

MORNING

EVENING