

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 2012	2. REPORT TYPE	3. DATES COVERED 00-00-2012 to 00-00-2012			
4. TITLE AND SUBTITLE Draft Genome Sequence Of The Shrimp Pathogen Vibrio harveyi CAIM 1792		5a. CONTRACT NUMBER			
		5b. GRANT NUMBER			
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)		5d. PROJECT NUMBER			
		5e. TASK NUMBER			
		5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Center for Bio/Molecular Science and Engineering, Washington, DC, 81657		8. PERFORMING ORGANIZATION REPORT NUMBER			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)			
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)			
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Journal of Bacteriology, 2012, 194(8):2104, 1/20					
14. ABSTRACT Vibrio harveyi is a Gram-negative bacterium found in tropical and temperate marine environments as a free-living organism or in association with aquatic animals. We report the first sequenced genome of a Vibrio harveyi strain, CAIM 1792, the etiologic agent of the "bright red" syndrome of the Pacific white shrimp Litopenaeus vannamei.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Draft Genome Sequence of the Shrimp Pathogen *Vibrio harveyi* CAIM 1792

Iliana Espinoza-Valles, Sonia Soto-Rodríguez, Robert A. Edwards, Zheng Wang, Gary J. Vora and Bruno Gómez-Gil
J. Bacteriol. 2012, 194(8):2104. DOI: 10.1128/JB.00079-12.

Updated information and services can be found at:
<http://jb.asm.org/content/194/8/2104>

	<i>These include:</i>
REFERENCES	This article cites 7 articles, 2 of which can be accessed free at: http://jb.asm.org/content/194/8/2104#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://jb.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Draft Genome Sequence of the Shrimp Pathogen *Vibrio harveyi* CAIM 1792

Iliana Espinoza-Valles,^a Sonia Soto-Rodríguez,^a Robert A. Edwards,^{b,d} Zheng Wang,^c Gary J. Vora,^c and Bruno Gómez-Gil^a

CIAD A.C. Mazatlan Unit for Aquaculture and Environmental Management, Sinaloa, México^a; Department of Biology and Department of Computer Science, San Diego State University, San Diego, California, USA^b; Center for Bio/Molecular Science and Engineering, Naval Research Laboratory, Washington, DC, USA^c; and Mathematics and Computer Science Division, Argonne National Laboratory, Argonne, Illinois, USA^d

***Vibrio harveyi* is a Gram-negative bacterium found in tropical and temperate marine environments as a free-living organism or in association with aquatic animals. We report the first sequenced genome of a *Vibrio harveyi* strain, CAIM 1792, the etiologic agent of the “bright red” syndrome of the Pacific white shrimp *Litopenaeus vannamei*.**

Vibrio harveyi is a member of the *Vibrio* core group and an organism with an important role in the study of bacterial autoinduction (6) and bioluminescence (3). It is also an important pathogen of marine fish and invertebrates (1), especially for shrimp (5). Strain CAIM 1792 has caused recurrent outbreaks of vibriosis known as “bright red” syndrome in *Litopenaeus vannamei* shrimp farms in northwestern Mexico since 2005 (7). While it is accepted that genome sequence information can greatly enhance our understanding of the molecular pathogenicity mechanisms responsible for *V. harveyi*, the recent discovery of the misidentification of the two purported *V. harveyi* strains for which the genomes were sequenced demonstrated that there is a lack of a representative *V. harveyi* genome (4).

Strain CAIM 1792 was sequenced using a 454 GS-FLX pyrosequencing approach (Life Sciences/Roche, Branford, CT). The high-throughput sequencing resulted in ~18-fold coverage (104,105,476 bases sequenced), and the 529,955 reads were assembled using the Newbler software (version 2.5.3) into 87 contigs. A genome-scale assembly was constructed with Mauve Genome Alignment software (version 2.3.1; Genome Evolution Laboratory, University of Wisconsin—Madison [<http://asap.ahabs.wisc.edu>]). The ordering of the contigs was tested by using the Scaffold-folding builder program (G. G. Z. Silva, B. E. Dutilh, D. Matthews, K. Elkins, A. M. Segall, R. A. Edwards, and E. A. Dinsdale, submitted for publication) and with the completed *Vibrio campbellii* ATCC BAA-1116 genome as a reference. BlastX queries were performed to validate assemblies containing coding sequences (CDS). The draft genome described in this study contains 21 gaps that are in the process of closure via PCR amplicon sequencing.

The assembled genome of CAIM 1792 is 5.84 Mbp in size with 5,328 RAST server-annotated CDS (2), and it has two chromosomes and a plasmid. The 3.50-Mbp chromosome I (Chr I) is composed of seven supercontigs and four contigs with 11 gaps and contains eight rRNA operons and at least 47 tRNAs. In comparison, the 2.24-Mbp chromosome II (Chr II) is composed of four supercontigs with seven gaps and contains one rRNA operon and at least 12 tRNAs. Similar to most sequenced *Vibrio* spp. strains, *V. harveyi* CAIM 1792 has a superintegron located in Chr I that is 87.2 Kbp in size and contains ~130 gene cassettes, of which 97 encode hypothetical proteins. The genome sequence also revealed >100 genes that encode putative virulence factors (e.g., hemolysins, proteases, chitinases, collagenases, iron acquisition, type I, II, III, IV, and VI secretion systems, RTX toxins, and vibriolysin)

that, when combined with additional comparative genomic analyses with sequenced sister species such as *V. campbellii*, *V. parahaemolyticus*, *V. rotiferianus*, and *V. owensii*, will aid in revealing the underlying genetic assemblages responsible for pathogenicity, host range, environmental adaptation, and evolution within *V. harveyi* and the *Vibrio* core group.

Nucleotide sequence accession numbers. *Vibrio harveyi* CAIM 1792 has been deposited in the Collection of Aquatic Important Microorganisms (CAIM; www.ciad.mx/caim), and the genome sequence has been deposited in the GenBank database under accession number [AHHQ00000000](https://doi.org/10.1093/nar/nqs000). The version described in this study is the first version, [AHHQ01000000](https://doi.org/10.1093/nar/nqs000).

ACKNOWLEDGMENTS

This work was supported by CONACYT (I.E.-V., S.S.-R., and B.G.-G.), NSF grant DBI 0850356 from the Division of Biological Infrastructure (R.A.E.), and the Office of Naval Research via U.S. Naval Research Laboratory core funds (Z.W. and G.J.V.).

The opinions and assertions contained herein are those of the authors and are not to be construed as those of the U.S. Navy, military service at large, or U.S. Government.

REFERENCES

1. Austin B, Zhang XH. 2006. *Vibrio harveyi*: a significant pathogen of marine vertebrates and invertebrates. *Lett. Appl. Microbiol.* 43:119–124.
2. Aziz RK, et al. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75.
3. Belas R, et al. 1982. Bacterial bioluminescence: isolation and expression of the luciferase genes from *Vibrio harveyi*. *Science* 218:791–793.
4. Lin B, et al. 2010. Comparative genomic analyses identify the *Vibrio harveyi* genome sequenced strains BAA-1116 and HY01 as *Vibrio campbellii*. *Environ. Microbiol. Rep.* 2:81–89.
5. Moriarty DJW. 1998. Microbial ecology: protecting the prawns in polluted ponds. *Microbiol. Aust.* 19:22–27.
6. Neilson KH, Platt T, Hastings JW. 1970. Cellular control of the synthesis and activity of the bacterial luminescent system. *J. Bacteriol.* 104:313–322.
7. Soto-Rodríguez S, Gómez-Gil B, Lozano R. 2010. ‘Bright-red’ syndrome in Pacific white shrimp *Litopenaeus vannamei* is caused by *Vibrio harveyi*. *Dis. Aquat. Org.* 92:11–19.

Received 1 February 2012 Accepted 6 February 2012

Address correspondence to Bruno Gómez-Gil, bruno@ciad.mx.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00079-12